

## Epigenetic alterations in metastatic cutaneous carcinoma

Owen A. Darr, MD,<sup>1</sup> Justin A. Colacino, MPH,<sup>2</sup> Alice L. Tang, MD,<sup>3</sup> Jonathan B. McHugh, MD,<sup>4</sup> Emily L. Bellile, MS,<sup>5</sup> Carol R. Bradford, MD,<sup>1</sup> Mark P. Prince, MD,<sup>1</sup> Douglas B. Chepeha, MD,<sup>1</sup> Laura S. Rozek, PhD,<sup>1,2</sup> Jeffrey S. Moyer, MD<sup>1\*</sup>

<sup>1</sup>Department of Otolaryngology, University of Michigan Medical School, Ann Arbor, Michigan, <sup>2</sup>Department of Environmental Health Sciences, University of Michigan School of Public Health, Ann Arbor, Michigan, <sup>3</sup>Department of Otolaryngology, University of Cincinnati College of Medicine, Cincinnati, Ohio, <sup>4</sup>Department of Pathology, University of Michigan Medical School, Ann Arbor, Michigan, <sup>5</sup>Comprehensive Cancer Center, University of Michigan, Ann Arbor, Michigan.

Accepted 28 March 2014

Published online 27 June 2014 in Wiley Online Library (wileyonlinelibrary.com). DOI 10.1002/hed.23701

**ABSTRACT:** *Background.* Squamous cell carcinoma (SCC) and basal cell carcinoma (BCC) are the 2 most common cutaneous carcinomas. Molecular profiles predicting metastasis of these cancers have not been identified.

*Methods.* Epigenetic profiles of 37 primary cases of cutaneous SCC and BCC were quantified via the Illumina Goldengate Cancer Panel. Differential protein expression by metastatic potential was analyzed in 110 total cases by immunohistochemical (IHC) staining.

*Results.* Unsupervised hierarchical clustering analysis revealed that metastatic BCCs had a methylation profile resembling cutaneous SCCs. Metastatic cutaneous SCCs were found to be hypermethylated at *FRZB* (median methylation: 46.7% vs 4.7%;  $p = 4 \times 10^{-5}$ ). Metastatic BCCs

were found to be hypomethylated at *MYCL2* (median methylation: 3.8% vs 83.4%;  $p = 1.9 \times 10^{-6}$ ). Immunohistochemical staining revealed few differences between metastatic and nonmetastatic cancers.

*Conclusion.* Metastatic primary BCCs and cutaneous SCCs had distinct epigenetic profiles when compared to their nonmetastatic counterparts. Epigenetic profiling may prove useful in future diagnosis and prevention of advanced nonmelanoma skin cancers. © 2014 Wiley Periodicals, Inc. *Head Neck* 37: 994–1001, 2015

**KEY WORDS:** DNA methylation, metastasis, cutaneous carcinoma, epigenetic, squamous, basal

## INTRODUCTION

Cutaneous squamous cell carcinoma (SCC) and basal cell carcinoma (BCC) are the 2 most common cutaneous malignancies in the United States. Most nonmelanoma skin cancers remain localized and noninvasive; however, there is a risk of locoregional and distant spread of disease with both cutaneous SCC and BCC. Cutaneous SCC has a risk of metastasis, which has been reported between 2% and 9.9%.<sup>1–7</sup> Despite the high incidence of BCC, metastatic transformation is extremely rare with an estimated rate of metastasis between 0.0028% and 0.55%.<sup>8–11</sup> In recent decades, much has been learned about the meta-

static potential of cutaneous SCC. Risk factors, such as primary location, tumor size, depth of invasion, immunosuppression, and perineural invasion have all been associated with increased risk of metastasis.<sup>12–16</sup> Risk factors associated with the metastasis of BCC have been reported despite the rarity of its occurrence. Metastatic BCC tends to arise preferentially from tumors in the head and neck region, which account for 70% to 80% of reported cases in the literature, with ear, cheek, and forehead predominating.<sup>8,10,11</sup> The size of the primary tumor seems to be an important risk factor with large, neglected tumors comprising many of the reported cases in the literature. Snow et al,<sup>10</sup> in their series of metastatic BCC, found that 75% of metastatic tumors occurred in tumors >2 cm and/or with invasion of bone or other deep structures.

Epigenetic modifications, including DNA methylation, have been shown to play an integral role in carcinogenesis, cancer progression, and metastasis. Compared to normal tissues, tumors tend to be hypomethylated at intergenic distal regulatory regions and repetitive elements, which generally results in genomic instability, and hypermethylated at gene promoters, resulting in gene silencing.<sup>17</sup> Additionally, epigenetic modifications are known to regulate gene expression patterns and affect cell fate decisions.<sup>18</sup> Thus, quantification of epigenetic profiles across the genome can establish similarities or differences in tumor cell type of origin and can provide insight as to whether rare metastatic basal cancers actually represent a basal-like differentiation

\*Corresponding author: J. S. Moyer, Department of Otolaryngology, University of Michigan Medical School, Taubman Center, Floor 1, Reception A, 1500 E. Medical Center Drive, SPC 5312, Ann Arbor, MI 48109. E-mail: jmoyer@med.umich.edu

Owen A. Darr and Justin A. Colacino contributed equally to this work.

Contract grant sponsor: This study was funded as a part of the University of Michigan Head and Neck Specialized Program of Research Excellence, NIH/NCI CA097248, as well as through NIH/NCI R01CA158286. Support for Justin A. Colacino was provided by a Rackham Predoctoral Fellowship from the University of Michigan and Institutional Training Grants from the National Institute of Environmental Health Sciences (NIEHS) (T32 ES007062) and the National Human Genome Research Institute (NHGRI) (T32 HG00040).

Additional Supporting Information may be found in the online version of this article.

of SCCs. Epigenetic biomarkers of metastatic potential have previously been identified in malignant melanoma,<sup>19</sup> however, it is unknown if primary cutaneous SCCs and BCCs with metastatic potential have a distinct epigenetic profile. The high incidences of these cancers make the identification of novel biomarkers of metastasis in these tumors of high clinical utility.

Despite numerous reports citing risk factors associated with metastatic skin cancers, many patients with high-risk features, such as large size and invasion of bone, do not develop regional or distant metastases. A central question is how to more precisely identify the patients with cutaneous SCC or BCC that have a high potential to metastasize. To address this question, we undertook a study of the epigenetic profiles of a cohort of patients from the University of Michigan Cancer Center with documented metastasis of cutaneous SCC or BCC and compared their epigenetic signatures to an age-matched control group of nonmetastatic cutaneous SCC or BCC specimens, respectively, to determine unique patterns that might be associated with metastatic disease in each malignancy.

## MATERIALS AND METHODS

### Identification of cases

Institutional Review Board approval at the University of Michigan was prospectively obtained (HUM00028920) and cases of metastatic cutaneous SCC and BCC were identified in a systematic chart review of all surgical cases performed in the Department of Otolaryngology–Head and Neck Surgery at the University of Michigan between 1990 and 2010. Medical records for these patients were examined and pertinent data were collected regarding patient demographics, tumor pathology, and course of disease. Surgical specimens were collected to confirm histopathologic characteristics and presence of cutaneous SCC or BCC in primary and metastatic tumor by an experienced head and neck pathologist (J.B.M.). Medical records and specimens were scrutinized according to the criteria set forth by Lattes and Kessler<sup>20</sup> in 1951, and adapted by the Cotran<sup>21</sup> review in 1961, which states that an accurate diagnosis of metastatic BCC must satisfy the following criteria: (1) the primary lesion must be of cutaneous origin; (2) metastases must be distinguished from growth by direct extension of the original tumor; and (3) the primary and metastatic tissue must share similar histologic subtypes. These criteria were similarly applied for patients with metastatic cutaneous SCC.

### Formalin-fixed paraffin-embedded tissue, DNA isolation, and bisulfite conversion

Paraffin blocks containing pathologic tissue from both primary and metastatic cutaneous SCC and BCC tumors were obtained from the University of Michigan Pathology Core. In order to provide an experimental control group for patients with metastatic cutaneous SCC or BCC, paraffin blocks from age-matched patients that did not metastasize were randomly selected in a 1:2 ratio. Tumor blocks were recut for uniform histopathologic review and microdissection, with the first and last slides of a series of 12 reviewed by a qualified pathologist (J.M.) to confirm the original diagnosis and to circle areas for DNA

extraction. Percent cellularity was estimated for each tumor, and areas with >70% cancer cellularity were designated for use in the analyses.

Tumor regions that were identified for DNA extraction were cored from the formalin fixed paraffin embedded tissue blocks using an 18 gauge needle. DNA was isolated from the cored tissue samples using the QIAamp DNA formalin fixed paraffin embedded Tissue Kit (Qiagen, Valencia, CA) with a modification to the manufacturer's recommended lysis protocol (incubation overnight at 56°C in lysis buffer). DNA concentration and purity was confirmed via NanoDrop spectrophotometer (Thermo Scientific, Waltham, MA). We performed sodium bisulfite modification on 500 ng to 1 µg of DNA using the EZ DNA Methylation kit (Zymo Research, Orange, CA) in accord with the manufacturer's recommended protocol.

### Tissue microarray

Formalin-fixed, paraffin-embedded tissue blocks of 110 cases were obtained from the files of the Department of Pathology, University of Michigan Medical Center, Ann Arbor, Michigan. The University of Michigan Institutional Review Board provided a waiver of informed consent to obtain these samples. After pathological review, a tissue microarray (TMA) was constructed from the most representative area using the methodology of Nocito et al.<sup>22</sup> Each case was represented by two 1-mm diameter cores obtained from the most representative, non-necrotic area of the tumor.

### Immunohistochemistry

Immunohistochemical staining was performed on the DAKO Autostainer (DAKO, Carpinteria, CA) using diaminobenzidine as the chromogen. Serial sections of deparaffinized TMA sections were labeled with the antibodies listed in Supplementary Table 1, online only. Appropriate negative (no primary antibody) and positive controls were stained in parallel with each set of tumors studied. The immunoreactivity was scored on a 3-tier scale (negative, low- [1+] and high-positive [2+]). After titration of each antibody, and subsequent immunohistochemical (IHC) staining of the 2 TMA slides, the slides were taken to an experienced pathologist (J.B.M.) for review and scoring.

Scoring of each stained core on the TMA slides was carried out using a 2-variable system. The percentage of malignant cells with positive staining was represented by a number from 0 to 100. The intensity of staining was scored on a scale from 0 to 4, in which 0 = negative, 1+ = 0% to 25%, 2+ = 26% to 50%, 3+ = 51% to 75%, and 4+ = 76% to 100%. As each tumor core was represented on the TMA in duplicate, the numerical scores of the 2 cores were averaged separately by percentage and intensity. The resultant values were then analyzed statistically using chi-square contrast tests in a linear regression model to test differences across groups.

### Bead array methods

The Illumina Goldengate Methylation Cancer Panel was utilized to detect DNA methylation patterns in tumor samples. The Cancer Panel quantifies methylation at 1505

**TABLE 1.** Clinical features of patients with metastatic basal cell carcinoma or cutaneous squamous cell carcinoma.

Clinical characteristics	Patients with BCC (N = 5)	Patients with SCC (N = 46)
Male : female	4 M : 1 F	41 M : 5 F
Mean age at dx of metastasis (SD)	65 (8)	73 (11)
Median interval between primary and met	7 y	8.5 mo
Perineural invasion	5	16
Perivascular invasion	2	7
Bone involvement	5	3
Smoking status	3 current; 2 former	1 current; 27 former

Abbreviations: BCC, basal cell carcinoma; SCC, squamous cell carcinoma; dx, diagnosis; met, metastasis.

CpG sites located in known CpG islands across 807 genes related to cancer, including oncogenes, tumor suppressor genes, imprinted genes, and genes involved in cell cycle regulation, DNA repair, apoptosis, and metastasis. Bead arrays were run at the University of Michigan DNA Sequencing Core Facility, in accord with the manufacturer's protocol. Briefly, bisulfite converted tumor DNA was hybridized to the bead array as described previously,<sup>23</sup> and bead arrays were imaged using Illumina Cancer Panel Reader software. Raw bead array fluorescence data was initially analyzed using Illumina BeadStudio Methylation software, which converts fluorescence values of the methylated (Cy5) and unmethylated (Cy3) alleles into an average methylation value at a specific probe using the formula  $\beta = [\text{Max}(\text{Cy5},0)/[\text{Max}(\text{Cy5},0) + \text{Max}(\text{Cy3},0) + 100]]$  with  $\beta \in (0,1)$ .

Methylation at specific CpG probes on the Goldengate Cancer Panel has been previously shown to be biased by probe thermodynamic properties.<sup>24</sup> Known biases include probe length and guanine-cytosine content, which can affect the melting temperature of the probes as well as probe fluorescence intensities. Thus, we used the method proposed by Kuan et al<sup>24</sup> to normalize our average  $\beta$  values based on probe length and guanine-cytosine content.

### Statistical methods

Unsupervised hierarchical clustering of the methylation array data was performed using the Euclidean distance metric and the Ward clustering method in the *hclust* package in R version 2.10.1.<sup>25</sup> To minimize sex-specific effects, we excluded CpG sites on the sex chromosomes, and included the top 50% of CpG sites with the greatest variance in methylation across samples. For survival analyses, death was considered an "event"; survival time was censored at 3 years (1095 days). The log-rank test was used to test differences in survival distributions using the R Survival package.

### Methylation scores

We calculated a tumor suppressor gene methylation score using our previously described methodology.<sup>26</sup> Briefly, we identified genes associated with tumor suppression by con-

ducting a simple query of the National Center for Biotechnology Information gene database for "tumor suppressor" and extracted gene identifications. A total of 444 individual CpG sites across 237 genes on the Goldengate Cancer Panel were linked to tumor suppression. Next, for each study participant, the methylation score was defined as the number of tumor suppressor gene-associated CpG sites with normalized average  $\beta$  values above 0.5. Thus, the methylation score ranged from 0 to 444. We infer that a higher methylation score is associated with higher levels of DNA methylation of promoter regions of genes associated with tumor suppression and cell cycle regulation.

We also calculated a Polycomb Repressor Complex 2 (PRC2) gene occupancy methylation score for each study participant. PRC2 target genes were identified in human embryonic stem cells,<sup>27</sup> and are known to be important regulators of development<sup>28</sup> and dysregulated in cancer.<sup>29</sup> A total of 232 sites across 107 genes on the Cancer Panel were identified as PRC2 targets, as described by Lee et al<sup>27</sup>. A PRC2 occupancy methylation score was calculated by summing the number of PRC2-associated CpG sites with normalized average  $\beta$  values above 0.5. The PRC2 occupancy methylation scores ranged from 0 to 232. Twenty-six genes were found in both the PRC2 occupancy gene list and the tumor suppressor gene list.

### Site specific analyses


To determine whether specific sites of the genome may be differentially methylated in metastatic cancers, we calculated associations between DNA methylation and metastatic potential. The association between metastasis and individual CpG site DNA methylation at the 1505 CpG sites measured on the Goldengate Cancer Panel were examined using the Limma package in R 2.10.1,<sup>30</sup> adjusting for sex and age. Sample weights based on detection *p* values across samples were used in the *lmFit* function from the Limma package to downweight samples with higher detection *p* values. An empirical Bayes method (using the *eBayes* function in Limma) was used to shrink standard errors to a common value and to rank CpG sites in order of differential methylation. Multiple comparisons were accounted for using the *q*-value method previously described.<sup>31</sup>

## RESULTS

### Clinical findings of cohort

Our cutaneous SCC study cohort (Table 1) included 46 patients diagnosed with metastatic cutaneous SCC of the head and neck, including tumors from 41 men and 5 women. The mean age at diagnosis of metastatic disease was 73 years with a shorter time interval from primary diagnosis to metastasis of 8.5 months, as compared to BCC. The most common primary site resulting in regional metastasis was periauricular (14 of 46; 30%) followed by the frontotemporal region (8 of 46; 18%). Adverse pathological features included perineural invasion (16 of 46; 35%), perivascular invasion (7 of 46; 15%), and bone involvement (3 of 46; 6%). Sixty percent of patients (28 of 46) were either current or former smokers. The median overall survival for patients with metastatic cutaneous SCC was 4.4 years with no statistical difference at 3 years between metastatic and





nonmetastatic patients. There was, however, a trend at 2 years for worse survival in the metastatic cutaneous SCC cohort ( $p = .08$ ). The demographics of patients with metastatic cutaneous SCC are consistent with other reports in the literature.<sup>13–16</sup> The majority of patients were men (89%) and the most common primary sites involved with regional metastasis were from the ear (30%) or the frontotemporal region (18%). Perineural invasion was present in 35% of patients, perivascular invasion in 15%, and 18 of 46 patients (39%) of those with metastatic disease were poorly differentiated on histopathology, compared to 0% of the specimens in the nonmetastatic group.


Our BCC study cohort included 5 patients diagnosed with verified metastatic basal cell carcinoma (Table 1). The group consisted of 4 men and 1 woman. The mean age at the time of primary BCC diagnosis was 55.0, whereas the mean age at the diagnosis of metastatic disease was 65 years (range, 52–75 years). Sixty percent of the patients had multiply recurrent tumors before the diagnosis of a metastasis. The median interval between the development of the primary lesion and progression to clinically evident metastatic disease was 7 years. Eighty percent of primary sites originated in the facial region, including the temple, ear, chin, and forehead. Primary lesions ranged in size from 1.3 cm<sup>2</sup> to 70 cm<sup>2</sup>. Each of these cases developed metastatic disease in the neck: 1 patient with level 1 disease, 3 patients with level 2 disease, and the lower back primary had level IV and V disease along with extensive axillary disease (13 of 17 positive nodes). Histopathologic examination confirmed the presence of BCC in all metastatic tumor specimens, which were identical in subtype to the corresponding primary pathology. All specimens were representative of an aggressive growth pattern, and subtypes included nodular, infiltrative, metatypical, and infiltrating keratotic, with several cases including mixed features. Perineural invasion was present in all cases and

perivascular invasion was evident in 2 of 5 patients. Only 1 of the 5 patients received radiation before the development of metastasis because of the extent of the primary tumor. The other 4 patients received radiation after resection of the metastases, and only 1 of those underwent adjuvant chemotherapy. The mean overall survival time after the development of regional metastases for the cohort was 4.8 years. Two patients are still alive, but 1 has distant metastatic disease (lung). For all the deceased patients, the cause of death is unknown but each of the 3 was disease free at the date of the last documented follow-up.

The BCC case cohort of 5 patients is largely representative of the published literature on age, sex, and ethnic demographics<sup>8–11</sup>; we had a male predominance, with the primary diagnosis in the fifth decade and an extended interval before clinically apparent metastases were diagnosed. All patients had T4 disease based on current American Joint Committee on Cancer criteria (bone involvement) and were treated with radical surgical resection of both the primary disease and regional metastases. Four of these patients underwent adjuvant radiation, 1 of whom also received post-chemoradiotherapy (cisplatin). Despite the comorbidities of these procedures, as well as the reported median survival of 8 months,<sup>11</sup> the shortest observed survival after surgery was 18 months, and the median overall survival from the time of definitive surgical management was almost 5 years.

### Methylation patterns

Methylation patterns across all of the 1505 CpG sites assayed on the Goldengate Cancer Panel were compared between primary BCC and SCC tissues that did and did not metastasize. When the top 50% most variable sites were included in an unsupervised hierarchical clustering analysis, nonmetastatic BCCs tended to cluster separately




from metastatic BCCs and SCCs (Figure 1). For SCCs, there was no overall difference in mean methylation across all sites observed between the 2 groups (Wilcoxon,  $p = .18$ ), whereas for BCCs the primary tumors that metastasized were more likely to be hypomethylated (Wilcoxon,  $p = .07$ ; Figure 2). Additionally, comparing tumor suppressor methylation score as well as PRC2 occupancy score across metastatic and nonmetastatic primary BCCs revealed a strong trend that had a similar pattern to that seen with mean methylation, with both tumor suppressor genes and PRC2 target genes hypomethylated in metastatic tumors, however, this did not reach statistical significance (Figure 2). For SCCs, there was no significant difference in tumor suppressor methylation score comparing primary tumors that metastasized to those that did not (Wilcoxon,  $p = .83$ ). Nor was any difference observed between the 2 groups when comparing the PRC2 gene occupancy methylation score (Wilcoxon,  $p = .30$ ).

#### Site specific methylation analyses of cutaneous squamous cell carcinomas

To quantify site-specific epigenetic differences between primary cutaneous SCCs that metastasized and those that did not, we measured the associations between each of the 1505 CpG sites on the Goldengate Cancer Panel and metastatic potential of the tumor. The 10 most differentially methylated sites that were identified are listed in Table 2. The top hit, a CpG site in the promoter region of *FRZB*, a modulator of Wnt signaling known to be involved in regulation of bone development, was found to be hypermethylated in metastatic (median methylation: 46.7%) compared to nonmetastatic (median methylation: 4.7%) primary SCCs ( $p = 4 \times 10^{-5}$ ;  $q = 0.14$ ). Additionally, CpG sites associated with the developmental transcription factors *TFAP2C* and *ASCL2* were also found to be hypermethylated in metastatic primary SCCs, whereas CpG sites associated with the actin protein

**TABLE 2.** The 10 most differentially methylated CpG sites associated with metastatic potential of the primary cutaneous squamous cell carcinoma, adjusting for patient age and sex.

Gene	Chromosome	CpG coordinate	Log fold change	T value	$p$ value	Q value
<i>FRZB</i>	2	183439557	2.632	4.612	.0001	0.14
<i>ASCL2</i>	11	2249118	1.635	4.050	.0004	0.29
<i>TFAP2C</i>	20	54638025	1.115	3.840	.0007	0.29
<i>GSTM2</i>	1	110012367	2.501	3.717	.0010	0.29
<i>GSTM1</i>	1	110031602	-2.489	-3.699	.0010	0.29
<i>TFF1</i>	21	42659893	-2.570	-3.623	.0012	0.29
<i>RARRES1</i>	3	159933026	-1.883	-3.587	.0014	0.29
<i>ACTG2</i>	2	73973146	-1.565	-3.539	.0015	0.29
<i>ACTG2</i>	2	73973699	-1.943	-3.350	.0025	0.41
<i>NGFB</i>	1	115682393	1.085	3.154	.0040	0.46



*ACTG2* were found to be hypomethylated in metastatic primary SCCs. We also compared DNA methylation of 4 sites in the *FRZB* promoter sequence in a subset of the samples analyzed on the Goldengate Cancer Panel as well as 10 additional primary SCCs and observed a similar pattern of methylation based on metastatic potential as measured on the array (Wilcoxon,  $p = .08$ ; Figure 3).

**Site specific methylation analyses of basal cell carcinomas**


To quantify specific epigenetic differences seen in metastatic and nonmetastatic primary BCCs, we measured the associations between methylation at each of the 1505 CpG sites on the Goldengate array to assess the possible markers of metastatic potential in primary tumors. Two CpG sites, both associated with the putative tumor suppressor gene *MYCL2* (Figure 4), were found to be significantly hypomethylated in metastatic tumors compared to control primary BCC that did not metastasize ( $p = 1.9 \times 10^{-6}$ ;  $q = 0.002$  and  $p = 3.7 \times 10^{-5}$ ;  $q = 0.028$ ,

respectively). Although not reaching statistical significance when adjusting for multiple comparisons ( $q < 0.05$ ), a number of other genes were also found to be differentially methylated ( $p < .05$ ) in metastatic BCCs (Supplementary Table 2, online only). The majority of these sites were found to be hypomethylated, similar to the global measures of methylation presented above, including sites associated with *EGF*, *GNG7*, *GRB10*, and *FGF9*.

**Immunohistochemical staining**

During preliminary quantification of epigenetic differences between metastatic and nonmetastatic samples, numerous candidate genes were identified as potentially associated with metastatic potential. To investigate whether this differential methylation was reflected in protein expression, IHC staining was performed on a subset of the proteins for which IHC antibodies are commercially available (Supplementary Table 1, online only).

After the construction of a TMA that included pathological specimens from both cutaneous SCC and BCC



groups, tissue from metastatic and nonmetastatic tumors were used for IHC staining. Staining scores were then compared between the groups; however, a significant difference in protein expression that differentiated metastatic from nonmetastatic primary tumors was discovered in only 1 case. When comparing the intensity of staining across cutaneous SCC samples, a neurotrophin receptor kinase, TrkB (also known as NTRK2), was identified by this process as being more highly expressed in SCC primaries that went on to metastasize compared to those that did not. Notably, this data reflects the highly elevated staining intensity of only 3 specimens of 26 stained in the metastatic SCC group, all of which scored 2 on a 0 to 4 scale. This, when compared to the 6 cores of 70 with positive staining in the nonmetastatic group, all of which received an intensity score of 1, achieved statistical significance in our model. IHC staining of BCC samples did not yield any significant differences between metastatic and nonmetastatic tissues.

## DISCUSSION

In addition to reviewing our clinical experience with metastatic cutaneous SCC and BCC, we undertook a study of the epigenetic signatures in primary tumors that developed regional or distant metastatic disease to determine how to identify patients who might be at high risk of developing metastatic cutaneous SCC or BCC. We identified epigenetic predictors of metastatic potential in both cutaneous SCC and BCC, with metastatic cutaneous SCC primaries hypermethylated at *FRZB* and metastatic BCC primaries hypomethylated at *MYCL2*. Additionally, we identified that metastatic primary cutaneous SCCs were more likely to stain positive for TrkB, with a higher staining intensity, although this relationship was driven by a small number of cases.

Despite a small sample size, our comprehensive experimental analysis of epigenetic regulation in metastatic cutaneous SCC and BCC was adequately powered to detect significant differences in methylation signatures between groups. When comparing epigenetic profiles between metastatic versus nonmetastatic primary cutaneous SCCs, we did not see widespread differences in methylation patterns, but did identify differential methylation at a CpG site in the promoter region of *FRZB*. The protein product of *FRZB* is secreted and involved in regulation of bone morphogenesis during development<sup>32,33</sup> through antagonism of *Wnt* signaling.<sup>34,35</sup> Previous studies of the role of *FRZB* in cancer have identified that overexpression of *FRZB* is associated with a more differentiated tumor phenotype in gastric cancer,<sup>25</sup> as well as a less invasive phenotype in prostate cancer.<sup>36</sup> Additionally, promoter methylation of *FRZB* was associated with a higher grade in noninvasive bladder cancer cases.<sup>37</sup> Similarly, in this study, we observed increased promoter methylation in primary cutaneous SCC tumors that metastasized. Our results, combined with previous findings, suggest that *FRZB* methylation could serve as a biomarker of tumor aggressiveness or metastatic potential at multiple cancer sites.

We observed widespread methylation differences between metastatic and nonmetastatic primary BCCs. Metastatic BCCs were found to be hypomethylated across

the genome, as well as at tumor suppressor genes and PRC2 target genes (Figure 2). Additionally, we found that in an unsupervised hierarchical cluster analysis, nonmetastatic primary BCCs clustered separately from the rest of the tumors, suggesting a unique methylation profile (Figure 1). In gene specific analyses, *MYCL2* was the only gene to show a significantly different methylation score across metastatic and nonmetastatic primaries, and this was observed at both loci included on the array. This gene localizes to the X-chromosome, has a poorly characterized function, and has previously been described as highly and specifically expressed in testes.<sup>38</sup> *MYCL2* shares sequence homology with the tumor suppressor genes *MYCL1* and *MYC*. In these samples, *MYCL2* was unmethylated in metastatic BCCA and methylated in nonmetastatic BCCA, suggesting that methylation at these loci could be used as biomarkers of metastatic potential. Of interest in the gene list with borderline significance, *GNG7*, involved in transmembrane signaling, has been shown to be hypermethylated in head and neck SCC<sup>39</sup> and fibroblast growth factor 9 (*FGF9*) is upregulated in cutaneous SCCs.<sup>40</sup>

IHC staining for a range of proteins, including EGFR and RUNX3, did not reveal significant differences in expression when comparing primary tumors that metastasized and those that did not. Interestingly, despite the differences observed in methylation of *FRZB* in metastatic cutaneous SCCs, we did not observe a significant difference in protein expression between metastatic and nonmetastatic primaries. We did, however, observe a significant increase in staining intensity of TrkB in metastatic cutaneous SCCs. TrkB has been described previously as an important oncogene in head and neck cancer, medullary thyroid carcinoma, myelomas, prostate cancer, and lymphoid tumors.<sup>41–44</sup> The small number of metastatic cases that stained positively for TrkB suggests that there may be a subset of cutaneous SCCs in which metastasis is driven, in part, by TrkB overexpression.

There exists an ongoing debate as to whether BCC truly metastasizes, with some arguing that it is rather the cells of a primary lesion with squamous cell differentiation that initiate invasion and spread of disease, acting as a vehicle for BCC cells to disseminate. However, the pathology of our BCC specimens, both primary and metastatic, consistently demonstrates the features of BCC on histologic analysis. Interestingly, however, the primary BCCs that developed metastases had a methylation pattern that strongly clustered with cutaneous SCC (see Figure 1). This cluster profile was shared with both primary cutaneous SCC that metastasized and those that did not metastasize, which would suggest that metastatic BCCs, at least when viewed from a gene methylation perspective, more closely resembles cutaneous SCC. Perhaps analysis of the gene methylation profiles of high-risk BCC might offer clues to those primaries at greatest risk of regional and distant metastases and could offer the possibility of increased surveillance or the use of adjuvant treatments, such as the hedgehog pathway inhibitor (vismodegib) for high-risk lesions.

This study has elucidated definitive differences in epigenetic signatures among the metastatic forms of cutaneous SCC and BCC. *FRZB* and *MYCL2* may be putative



biomarkers of the metastatic phenotype and useful in future diagnosis and prevention of advanced disease. However, our data highlight the need for additional research on this subject; mechanistic and mutational data remains elusive because of the relative rare nature of this process. However, epigenetic modification of functionally important genes may hold special significance in the improved management of metastatic cutaneous disease.

## REFERENCES

- Chuang TY, Popescu NA, Su WP, Chute CG. Squamous cell carcinoma. A population-based incidence study in Rochester, Minn. *Arch Dermatol* 1990;126:185–188.
- Dinehart SM, Pollack SV. Metastases from squamous cell carcinoma of the skin and lip. An analysis of twenty-seven cases. *J Am Acad Dermatol* 1989; 21(2 Pt 1):241–248.
- Epstein E, Epstein NN, Bragg K, Linden G. Metastases from squamous cell carcinomas of the skin. *Arch Dermatol* 1968;97:245–251.
- Katz AD, Urbach F, Lilienfeld AM. The frequency and risk of metastases in squamous-cell carcinoma of the skin. *Cancer* 1957;10:1162–1166.
- Lund HZ. How often does squamous cell carcinoma of the skin metastasize? *Arch Dermatol* 1965;92:635–637.
- Rowe DE, Carroll RJ, Day CL Jr. Prognostic factors for local recurrence, metastasis, and survival rates in squamous cell carcinoma of the skin, ear, and lip. Implications for treatment modality selection. *J Am Acad Dermatol* 1992;26:976–990.
- Tavin E, Persky M. Metastatic cutaneous squamous cell carcinoma of the head and neck region. *Laryngoscope* 1996;106(2 Pt 1):156–158.
- Farmer ER, Helwig EB. Metastatic basal cell carcinoma: a clinicopathologic study of seventeen cases. *Cancer* 1980;46:748–757.
- Lo JS, Snow SN, Reizner GT, Mohs FE, Larson PO, Hruza GJ. Metastatic basal cell carcinoma: report of twelve cases with a review of the literature. *J Am Acad Dermatol* 1991;24(5 Pt 1):715–719.
- Snow SN, Sahl W, Lo JS, et al. Metastatic basal cell carcinoma. Report of five cases. *Cancer* 1994;73:328–335.
- von Domarus H, Stevens PJ. Metastatic basal cell carcinoma. Report of five cases and review of 170 cases in the literature. *J Am Acad Dermatol* 1984;10:1043–1060.
- Weinberg AS, Ogle CA, Shim EK. Metastatic cutaneous squamous cell carcinoma: an update. *Dermatol Surg* 2007;33:885–899.
- Turner SJ, Morgan GJ, Palme CE, Veness MJ. Metastatic cutaneous squamous cell carcinoma of the external ear: a high-risk cutaneous subsite. *J Laryngol Otol* 2010;124:26–31.
- Veness MJ, Palme CE, Morgan GJ. High-risk cutaneous squamous cell carcinoma of the head and neck: results from 266 treated patients with metastatic lymph node disease. *Cancer* 2006;106:2389–2396.
- Veness MJ, Porceddu S, Palme CE, Morgan GJ. Cutaneous head and neck squamous cell carcinoma metastatic to parotid and cervical lymph nodes. *Head Neck* 2007;29:621–631.
- Veness MJ. High-risk cutaneous squamous cell carcinoma of the head and neck. *J Biomed Biotechnol* 2007;2007:80572.
- Ehrlich M. DNA methylation in cancer: too much, but also too little. *Oncogene* 2002;21:5400–5413.
- Bracken AP, Dietrich N, Pasini D, Hansen KH, Helin K. Genome-wide mapping of Polycomb target genes unravels their roles in cell fate transitions. *Genes Dev* 2006;20:1123–1136.
- Tanemura A, Terando AM, Sim MS, et al. CpG island methylator phenotype predicts progression of malignant melanoma. *Clin Cancer Res* 2009; 15:1801–1807.
- Lattes R, Kessler RW. Metastasizing basal-cell epithelioma of the skin; report of two cases. *Cancer* 1951;4:866–878.
- Cotran RS. Metastasizing basal cell carcinomas. *Cancer* 1961;14:1036–1040.
- Nocito A, Kononen J, Kallioniemi OP, Sauter G. Tissue microarrays (TMAs) for high-throughput molecular pathology research. *Int J Cancer* 2001;94:1–5.
- Bibikova M, Lin Z, Zhou L, et al. High-throughput DNA methylation profiling using universal bead arrays. *Genome Res* 2006;16:383–393.
- Kuan PF, Wang S, Zhou X, Chu H. A statistical framework for Illumina DNA methylation arrays. *Bioinformatics* 2010;26:2849–2855.
- Qu Y, Li JF, Cai Q, et al. Over-expression of FRZB in gastric cancer cell suppresses proliferation and induces differentiation. *J Cancer Res Clin Oncol* 2008;134:353–364.
- Colacino JA, Arthur AE, Dolinoy DC, et al. Pretreatment dietary intake is associated with tumor suppressor DNA methylation in head and neck squamous cell carcinomas. *Epigenetics* 2012;7:883–891.
- Lee TI, Jenner RG, Boyer LA, et al. Control of developmental regulators by Polycomb in human embryonic stem cells. *Cell* 2006;125:301–313.
- Surface LE, Thornton SR, Boyer LA. Polycomb group proteins set the stage for early lineage commitment. *Cell Stem Cell* 2010;7:288–298.
- Bracken AP, Helin K. Polycomb group proteins: navigators of lineage pathways led astray in cancer. *Nat Rev Cancer* 2009;9:773–784.
- Smyth GK. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol* 2004;3:Article 3.
- Storey JD, Tibshirani R. Statistical significance for genome-wide studies. *Proc Natl Acad Sci U S A* 2003;100:9440–9445.
- Hoang B, Moos M Jr, Vukicevic S, Luyten FP. Primary structure and tissue distribution of FRZB, a novel protein related to Drosophila frizzled, suggest a role in skeletal morphogenesis. *J Biol Chem* 1996;271:26131–26137.
- Hoang BH, Thomas JT, Abdul-Karim FW, et al. Expression pattern of two Frizzled-related genes, Frzb-1 and Sfrp-1, during mouse embryogenesis suggests a role for modulating action of Wnt family members. *Dev Dyn* 1998;212:364–372.
- Enomoto-Iwamoto M, Kitagaki J, et al. The Wnt antagonist Frzb-1 regulates chondrocyte maturation and long bone development during limb skeletogenesis. *Dev Biol* 2002;251:142–156.
- Wang S, Krinks M, Lin K, Luyten FP, Moos M Jr. Frzb, a secreted protein expressed in the Spemann organizer, binds and inhibits Wnt-8. *Cell* 1997; 88:757–766.
- Zi X, Guo Y, Simoneau AR, et al. Expression of Frzb/secreted Frizzled-related protein 3, a secreted Wnt antagonist, in human androgen-independent prostate cancer PC-3 cells suppresses tumor growth and cellular invasiveness. *Cancer Res* 2005;65:9762–9770.
- Marsit CJ, Houseman EA, Christensen BC, et al. Identification of methylated genes associated with aggressive bladder cancer. *PLoS One* 2010;5: e12334.
- Robertson NG, Pomponio RJ, Mutter GL, Morton CC. Testis-specific expression of the human MYCL2 gene. *Nucleic Acids Res* 1991;19:3129–3137.
- Hartmann S, Szaumkessel M, Salaverria I, et al. Loss of protein expression and recurrent DNA hypermethylation of the GNG7 gene in squamous cell carcinoma of the head and neck. *J Appl Genet* 2012;53:167–174.
- Kathalia VP, Mussak EN, Chow SS, et al. Genome-wide transcriptional profiling in human squamous cell carcinoma of the skin identifies unique tumor-associated signatures. *J Dermatol* 2006;33:309–318.
- Kupferman ME, Jiffar T, El-Naggar A, et al. TrkB induces EMT and has a key role in invasion of head and neck squamous cell carcinoma. *Oncogene* 2010;29:2047–2059.
- Lee J, Jiffar T, Kupferman ME. A novel role for BDNF-TrkB in the regulation of chemotherapy resistance in head and neck squamous cell carcinoma. *PLoS One* 2012;7:e30246.
- Thiele CJ, Li Z, McKee AE. On Trk—the TrkB signal transduction pathway is an increasingly important target in cancer biology. *Clin Cancer Res* 2009;15:5962–5967.
- Yilmaz T, Jiffar T, de la Garza G, et al. Therapeutic targeting of Trk suppresses tumor proliferation and enhances cisplatin activity in HNSCC. *Cancer Biol Ther* 2010;10:644–653.