

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

Received Date : 26-Aug-2015

Revised Date : 01-Sep-2015

Accepted Date : 02-Sep-2015

Article type : Commentary

Commentary

Title: Novel Roles for CRF-Binding Protein and CRF Receptor 2 in Binge Drinking

Authors: Kyle D. Ketchesin^{1,2,a} and Audrey F. Seasholtz^{1,2,3,b}

Affiliations: ¹Neuroscience Graduate Program, ²Molecular and Behavioral Neuroscience Institute, and ³Department of Biological Chemistry, University of Michigan, Ann Arbor, MI 48109

Academic degrees: ^aM.S. and ^bPh.D.

Financial support: NIAAA INIA U01 AA013641 (AFS) and NSF Graduate Research Fellowship F031543 (KDK).

Corresponding author: Dr. Audrey F. Seasholtz, University of Michigan,
5035 Biomedical Science Research Building

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/acer.12897](https://doi.org/10.1111/acer.12897)

This article is protected by copyright. All rights reserved

23 109 Zina Pitcher Place, Ann Arbor, MI 48109

24 Tel: (734) 936-2072; Fax: (734) 936-2690

25 Email: aseashol@umich.edu

26 Stress is a key environmental factor in alcohol addiction, and the stress system has been linked
27 to alcohol use prior to dependence, after dependence, and in relapse to drinking after abstinence in
28 both humans and rodent model systems (reviewed in Lowery and Thiele, 2010; Phillips et al., 2015, Ray,
29 2011). The key central nervous system regulator of the mammalian stress response is Corticotropin-
30 Releasing Factor (CRF). This peptide mediates its primary effects through binding to G-protein coupled
31 receptors, CRF receptor 1 (CRF-R1) and CRF receptor 2 (CRF-R2), and its activity is modulated by the CRF-
32 binding protein (CRF-BP). Dysregulation of the CRF system has been observed in a number of rodent
33 models of binge drinking and alcohol dependence, with most studies focusing on the roles of CRF and
34 CRF receptors, particularly CRF-R1 (reviewed in Lowery and Thiele, 2010; Phillips et al., 2015). CRF-R1
35 antagonists decreased binge drinking and self-administration in alcohol-dependent animals (Lowery and
36 Thiele, 2010), suggesting potential therapeutic uses for CRF-R1 antagonists. However, fewer studies
37 have investigated the roles of CRF-R2 and CRF-BP in binge drinking and alcohol dependence.

38 In the recent study by Albrechet-Souza and colleagues (2015), a variety of drinking paradigms
39 and pharmacological tools were utilized to examine the roles of CRF-BP and CRF-R2 in the ventral
40 tegmental area (VTA) and central nucleus of the amygdala (CeA) in binge drinking and alcohol
41 dependence. To model binge drinking, the authors used the drinking in the dark (DID) paradigm, in
42 which mice were given limited access to 20% ethanol in the dark phase of their circadian cycle, resulting
43 in drinking to intoxication and pharmacologically relevant blood ethanol concentration. Two variations
44 of the DID paradigm were employed, both one-bottle and two-bottle choice (Rhodes et al., 2007).
45 Alcohol dependence was modeled using a 4-week two-bottle choice, intermittent access to alcohol (IAA)
46 paradigm. To investigate the role of CRF-BP in each of these paradigms, the CRF-BP ligand inhibitor
47 CRF₆₋₃₃ was utilized (Sutton et al., 1995). The authors found that CRF-BP in the VTA, but not CeA,
48 regulates one-bottle choice binge drinking, but not dependence-induced alcohol consumption. To
49 investigate the role of CRF-R2 in the VTA in binge drinking, the CRF-R2 selective antagonist astressin 2B
50 (A2B) was used. A2B injected alone into the VTA at a high dose led to a decrease in ethanol intake in
51 two-bottle choice DID, while low dose A2B had no effect. Together, these exciting results reveal novel
52 roles for CRF-BP and CRF-R2 in the VTA in binge drinking.

53 Roles for CRF-binding protein (CRF-BP):

54 The CRF-BP is a 37 kDa secreted glycoprotein that binds CRF and the CRF-like ligand, urocortin 1
55 (Ucn1), with very high affinity. It is co-localized with CRF or CRF receptors at numerous sites (i.e.
56 amygdala and BNST), suggesting potential sites of interaction in stress-reward pathways (reviewed in
57 Westphal and Seasholtz, 2006). In humans, it is estimated that 40-60% of CRF in the brain is bound by
58 CRF-BP, and *CRHBP* SNPs have been associated with alcohol use disorder and stress-induced alcohol
59 craving (Ray, 2011), suggesting a role for the CRF-BP in susceptibility to alcohol use and addiction.

60 The CRF-BP has been studied for over 20 years with many postulated roles. In an inhibitory role,
61 CRF-BP reduces CRF receptor activation, likely by sequestering CRF or Ucn1 and/or targeting them for
62 degradation. Consistent with this model, purified CRF-BP reduces CRF-mediated ACTH release from
63 anterior pituitary cultures or AtT-20 cells (Cortright et al., 1995; Potter et al., 1991; Sutton et al., 1995).
64 Recombinant CRF-BP also attenuates CRF-R1-mediated increases in cAMP (Boorse et al., 2006).
65 Similarly, CRF-BP-deficient mice show increased baseline anxiety-like behavior and slowed return to
66 homeostasis after lipopolysaccharide stress, consistent with elevated free levels of CRF in the absence of
67 an inhibitory CRF-BP (Karolyi et al., 1999). In contrast, the study by Albrechet-Souza and colleagues
68 (2015) suggests an enhancing or facilitatory role. In this role, CRF-BP may bind CRF and increase CRF
69 signaling, likely by delivering CRF to the receptor, extending the half-life of CRF, or modulating its
70 interaction with the receptor. Consistent with this hypothesis, VTA CRF-BP may normally facilitate
71 alcohol consumption, as administration of CRF₆₋₃₃ into VTA reduced binge drinking in mice (Albrechet-
72 Souza et al., 2015). Similarly, intra-VTA administration of CRF₆₋₃₃ attenuated CRF-induced relapse to
73 cocaine seeking in rats (Wang et al., 2007). Altogether, these results suggest that CRF-BP may have
74 different roles depending on the specific cell type or context in which it is expressed, consistent with
75 ultrastructural localization studies (Peto et al., 1999). Finally, the CRF-BP may exert ligand and/or
76 receptor independent actions. Administration (i.c.v.) of CRF₆₋₃₃ elicits c-fos activation not only in a
77 subset of CRF receptor-expressing cells, but also in CRF-BP-expressing cells that do not express ligand or
78 CRF receptors, suggesting additional actions of CRF₆₋₃₃ and/or CRF-BP (Chan et al., 2000). Hence, further
79 studies are needed to elucidate the multiple roles of CRF-BP *in vivo*, as the actions of CRF-BP are likely
80 brain region and cell type-dependent.

81 CRF Receptors in Binge Drinking and Dependence:

82 The CRF receptors are highly homologous G-protein coupled receptors with similar, yet distinct
83 ligand binding profiles. CRF binds with very high affinity to CRF-R1, but binds to CRF-R2 with about 30
84 fold lower affinity. Both CRF receptors couple predominantly to $G\alpha_s$, activating adenylyl cyclase and
85 increasing cAMP levels and protein kinase A activity. However, they have also been shown to act
86 through other G protein subtypes, including $G\alpha_{q/11}$, $G\alpha_o$, $G\alpha_i$, in certain cellular contexts (reviewed in
87 Hillhouse and Grammatopoulos, 2006).

88 There is a large body of literature implicating the role of CRF-R1 in excessive alcohol drinking
89 (reviewed in Lowery and Thiele, 2010; Phillips et al., 2015). Altered CRF-R1 expression has been
90 observed in rats with a history of alcohol dependence, and peripheral administration of CRF-R1
91 antagonists reduced both binge drinking and dependence-induced alcohol consumption (Lowery and
92 Thiele, 2010). Furthermore, CRF receptor regulation of alcohol consumption may be brain region-
93 specific, with CeA and VTA as major sites of action. Intra-CeA (but not basolateral amygdala)
94 administration of a CRF-R1 antagonist decreased binge drinking (Lowery-Gionta et al., 2012). Similarly,
95 intra-VTA CRF-R1 antagonist administration decreased binge drinking in a DID paradigm (Sparta et al.,
96 2013) and decreased excessive drinking during the two-bottle choice IAA paradigm (Hwa et al., 2013).
97 Although less studied, CRF-R2 has also been implicated in excessive alcohol drinking, but often in an
98 inverse fashion to CRF-R1. For example, i.c.v. administration of the CRF-R2-selective agonist Urocortin 3
99 dose-dependently decreased binge drinking, and intra-CeA administration of Urocortin 3 resulted in
100 decreased alcohol self-administration in alcohol dependent rats (reviewed in Lowery and Thiele, 2010;
101 Phillips et al., 2015). Hence, CRF-R1 antagonists and CRF-R2 agonists often elicit similar responses. In
102 contrast, in the recent study by Albrechet-Souza et al. (2015), intra-VTA CRF-R2 antagonist (astressin 2B)
103 administration decreased alcohol drinking in the two-bottle DID paradigm, suggesting that VTA CRF-R2 is
104 facilitating drinking. Additional studies that directly compare the effects of CRF-R1 and CRF-R2, via
105 genetic approaches and/or receptor-selective agonists and antagonists, in the VTA, CeA and other brain
106 regions will be required to fully elucidate the roles of these two receptors in alcohol binge drinking and
107 dependence.

108 Potential CRF-BP/CRF Receptor Interactions and Future Directions:

109 The CRF system is unique in that very few neuropeptides have a high affinity binding protein
110 that is totally distinct from its receptors. CRF and Ucn1 bind to CRF-BP with very high affinity ($K_d < 1nM$),
111 equal to or greater than the affinity of these ligands for CRF-R1 and/or CRF-R2. Interestingly, the CRF-BP
112 is widely expressed in brain, localizing to sites of CRF or Ucn1 expression or release, sites of CRF-R1 or

113 CRF-R2 expression, and sites where neither the receptors nor ligands are detected (Chan et al., 2000;
114 Potter et al., 1992). CRF-BP synthesis and release are also highly responsive to environmental cues, as
115 stress, CRF, glucocorticoids and IL-6 all regulate CRF-BP expression (Westphal and Seasholtz, 2006). Yet
116 we still know little about the molecular mechanisms of its actions. Purified CRF-BP attenuates CRF or
117 Ucn1 activity at endogenous CRF receptors in cultured cells (Westphal and Seasholtz, 2006). In contrast,
118 *in vivo* or slice studies suggest a facilitatory role for CRF-BP in VTA, with CRF₆₋₃₃ administration into the
119 VTA decreasing ethanol consumption, CRF-induced relapse to cocaine, and CRF-mediated potentiation
120 of NMDA EPSCs on dopamine neurons (Albrechet-Souza et al., 2015; Ungless et al., 2003; Wang et al.,
121 2007). Strikingly, many of the facilitatory effects of CRF-BP in VTA appear to be associated with CRF-R2
122 (Ungless et al. 2003; Wang et al. 2007). In fact, Ungless et al. (2003) has proposed that CRF-BP is
123 required for CRF activation of CRF-R2 and the downstream phospholipase C/protein kinase C, but not
124 PKA, signaling pathway in VTA dopamine neurons. The study by Albrechet-Souza et al. (2015) begins to
125 address this VTA CRF-BP/CRF-R2 interaction in the two-bottle choice DID paradigm, but does not test
126 CRF₆₋₃₃ alone in this protocol, making it difficult to ascertain whether CRF-BP and CRF-R2 are interacting.

127 Hence, while this interesting study by Albrechet-Souza et al. (2015) reveals new roles for the CRF
128 system, especially CRF-BP and CRF-R2, in alcohol binge drinking, many questions remain. We propose
129 that CRF-BP plays multiple roles in stress and alcohol addiction, likely in a brain-region and cell-type
130 specific manner, with CRF receptor subtype and signaling pathway as potential determinants. Clearly,
131 additional *in vitro* and *in vivo* studies are needed to define the mechanisms of CRF-BP/CRF/CRF receptor
132 interactions. Future studies should also include viral and genetic approaches allowing cell-type specific
133 overexpression or knockdown of CRF-BP, and/or CRF receptors, in various brain regions within the
134 stress-reward pathway, as the transition from pre-dependent to dependent states in alcohol addiction
135 involves different sites and circuits. Together, these multidisciplinary approaches will allow us to better
136 understand the multiple roles of the CRF-BP in inhibiting and facilitating CRF and/or Ucn1 interactions
137 with CRF-R1 and CRF-R2 and the functional outcomes of these interactions in stress and addiction.

138

139 Acknowledgements:

140 The authors thank Dr. Gwen Stinnett for critical reading of the manuscript. This work was supported by
141 NIAAA INIA U01 AA013641 (AFS) and NSF Graduate Research Fellowship F031543 (KDK).

142 References:

- 143 Albrechet-Souza, L., Hwa, L. S., Han, X., Zhang, E. Y., Debold, J. F. & Miczek, K. A. (2015) Corticotropin
144 Releasing Factor Binding Protein and CRF Receptors in the Ventral Tegmental Area: Modulation
145 of Ethanol Binge Drinking in C57BL/6J Mice. *Alcohol Clin Exp Res.* e-pub.
- 146 Boorse, G. C., Kholdani, C. A., Seasholtz, A. F. & Denver, R. J. (2006) Corticotropin-releasing factor is
147 cytoprotective in *Xenopus* tadpole tail: coordination of ligand, receptor, and binding protein in
148 tail muscle cell survival. *Endocrinology*, 147, 1498-507.
- 149 Chan, R. K., Vale, W. W. & Sawchenko, P. E. (2000) Paradoxical activational effects of a corticotropin-
150 releasing factor-binding protein "ligand inhibitor" in rat brain. *Neuroscience*, 101, 115-29.
- 151 Cortright, D. N., Nicoletti, A. & Seasholtz, A. F. (1995) Molecular and biochemical characterization of the
152 mouse brain corticotropin-releasing hormone-binding protein. *Mol Cell Endocrinol*, 111, 147-57.
- 153 Hillhouse, E. W. & Grammatopoulos, D. K. (2006) The molecular mechanisms underlying the regulation
154 of the biological activity of corticotropin-releasing hormone receptors: implications for
155 physiology and pathophysiology. *Endocr Rev*, 27, 260-86.
- 156 Hwa, L. S., Debold, J. F. & Miczek, K. A. (2013) Alcohol in excess: CRF(1) receptors in the rat and mouse
157 VTA and DRN. *Psychopharmacology (Berl)*, 225, 313-27.
- 158 Karolyi, I. J., Burrows, H. L., Ramesh, T. M., Nakajima, M., Lesh, J. S., Seong, E., Camper, S. A. & Seasholtz,
159 A. F. (1999) Altered anxiety and weight gain in corticotropin-releasing hormone-binding protein-
160 deficient mice. *Proc Natl Acad Sci U S A*, 96, 11595-600.
- 161 Lowery, E. G. & Thiele, T. E. (2010) Pre-clinical evidence that corticotropin-releasing factor (CRF)
162 receptor antagonists are promising targets for pharmacological treatment of alcoholism. *CNS*
163 *Neurol Disord Drug Targets*, 9, 77-86.
- 164 Lowery-Gionta, E. G., Navarro, M., Li, C., Pleil, K. E., Rinker, J. A., Cox, B. R., Sprow, G. M., Kash, T. L. &
165 Thiele, T. E. (2012) Corticotropin Releasing Factor Signaling in the Central Amygdala is Recruited
166 during Binge-Like Ethanol Consumption in C57BL/6J Mice. *J Neurosci*, 32, 3405-13.
- 167 Peto, C. A., Arias, C., Vale, W. W. & Sawchenko, P. E. (1999) Ultrastructural localization of the CRF-
168 binding protein in rat brain and pituitary. *J. Comparative Neurology*, 413, 241-154.
- 169 Phillips, T. J., Reed, C. & Pastor, R. (2015) Preclinical evidence implicating corticotropin-releasing factor
170 signaling in ethanol consumption and neuroadaptation. *Genes Brain Behav*, 14, 98-135.

- 171 Potter, E., Behan, D. P., Fischer, W. H., Linton, E. A., Lowry, P. J. & Vale, W. W. (1991) Cloning and
172 characterization of the cDNAs for human and rat corticotropin releasing factor-binding proteins.
173 *Nature*, 349, 423-6.
- 174 Potter, E., Behan, D. P., Linton, E. A., Lowry, P. J., Sawchenko, P. E. & Vale, W. W. (1992) The central
175 distribution of a corticotropin-releasing factor (CRF)-binding protein predicts multiple sites and
176 modes of interaction with CRF. *Proc Natl Acad Sci U S A*, 89, 4192-6.
- 177 Ray, L. A. (2011) Stress-induced and cue-induced craving for alcohol in heavy drinkers: Preliminary
178 evidence of genetic moderation by the OPRM1 and CRH-BP genes. *Alcohol Clin Exp Res*, 35, 166-
179 74.
- 180 Rhodes, J. S., Ford, M. M., Yu, C. H., Brown, L. L., Finn, D. A., Garland, T., Jr. & Crabbe, J. C. (2007) Mouse
181 inbred strain differences in ethanol drinking to intoxication. *Genes Brain Behav*, 6, 1-18.
- 182 Sparta, D. R., Hopf, F. W., Gibb, S. L., Cho, S. L., Stuber, G. D., Messing, R. O., Ron, D. & Bonci, A. (2013)
183 Binge ethanol-drinking potentiates corticotropin releasing factor R1 receptor activity in the
184 ventral tegmental area. *Alcohol Clin Exp Res*, 37, 1680-7.
- 185 Sutton, S. W., Behan, D. P., Lahrachi, S. L., Kaiser, R., Corrigan, A., Lowry, P., Potter, E., Perrin, M. H.,
186 Rivier, J. & Vale, W. W. (1995) Ligand requirements of the human corticotropin-releasing factor-
187 binding protein. *Endocrinology*, 136, 1097-102.
- 188 Ungless, M. A., Singh, V., Crowder, T. L., Yaka, R., Ron, D. & Bonci, A. (2003) CRF requires CRF-binding
189 protein to potentiate NMDA receptors via CRF receptor 2 in dopamine neurons. *Neuron*, 39,
190 401-407.
- 191 Wang, B., You, Z. B., Rice, K. C. & Wise, R. A. (2007) Stress-induced relapse to cocaine seeking: roles for
192 the CRF(2) receptor and CRF-binding protein in the ventral tegmental area of the rat.
193 *Psychopharmacology (Berl)*, 193, 283-94.
- 194 Westphal, N. J. & Seasholtz, A. F. (2006) CRH-BP: the regulation and function of a phylogenetically
195 conserved binding protein. *Front Biosci*, 11, 1878-91.

196

197