Running title: Sex differences in decision-making

Sex Differences in Animal Models of Decision-Making Caitlin A. Orsini¹; Barry Setlow^{1,2,3,4} Departments of ¹Psychiatry; ²Neuroscience; ³Psychology; ⁴Center for Addiction Research and Education, University of Florida, Gainesville, FL Associate Editor: Larry Cahill Please address all correspondence to: Dr. Caitlin Orsini, Department of Psychiatry, University of Florida College of Medicine, P.O. Box 100256, Gainesville, FL 32610-0256; Tel: 352-294-5047; email: orsini@ufl.edu Keywords: delay discounting, intertemporal choice, risk, probability, dopamine

Supported by NIH DA036534 (BS) and a Thomas H. Maren Fellowship (CAO)

Journal of Neuroscience Research

This article is protected by copyright. All rights reserved.

Abstract

The ability to weigh the costs and benefits of various options in order to make an adaptive decision is critical to an organism's survival and well-being. Many psychiatric diseases are characterized by maladaptive decision-making, indicating the need to better understand the mechanisms underlying this process and the ways in which it is altered in pathological conditions. Great strides have been made in uncovering these mechanisms, but the majority of what is known comes from studies conducted solely in male subjects. In recent years, decision-making research has begun to include females to determine whether sex differences exist and to identify the mechanisms that contribute to such differences. This review will begin by describing studies that have examined sex differences in animal (largely rodent) models of decision-making. Possible explanations, both theoretical and biological, for such differences in decision-making will then be considered. The review will conclude with a discussion of the implications of sex differences in decision-making for understanding psychiatric conditions.

Acc

2 Journal of Neuroscience Research This article is protected by copyright. All rights reserved.

Significance Statement:

Many psychiatric diseases affect one sex to a greater extent than the other. A common feature across these diseases is that decision-making abilities are impaired. Thus, sex differences in decision-making may contribute to the differential development or presentation of psychiatric diseases. This review discusses what is currently known about sex differences in animal models of decision-making and considers possible explanations for such differences. The review concludes by highlighting the need for inclusion of both male and female subjects to ensure that future scientific discoveries can be more readily translated to all human beings.

Accepted

Introduction

To make a decision, one must consider several variables before taking action. Information about the risks and rewards associated with each option must be integrated with internal cognitive and motivational drives as well as the environmental context in which the decision is made. This process happens on a daily basis and the majority of individuals are able to effectively calculate costs and benefits to engage in adaptive choice behavior. However, there are multiple psychiatric conditions that are characterized by maladaptive decision-making. For example, individuals suffering from substance abuse disorders (SUD) display heightened impulsive choice and risk-taking behavior. To date, the majority of studies that have assessed relationships between decision-making and psychiatric diseases such as SUD have only used male subjects; however, there is well-established evidence that the incidence and presentation of many of these pathological conditions differs between sexes (McCarthy et al. 2012). For instance, while males have higher rates of drug dependence, females develop dependence more rapidly and are at greater risk for relapse (Becker and Hu 2008; Lynch 2006). Thus, although previous work has been useful in beginning to understand how decision-making can be altered in psychiatric diseases, it is obviously not representative of the entire population and is therefore limited in its application.

The use of animal models of decision-making has allowed researchers to begin to address these gaps in knowledge. Using these models, scientists can answer fundamental questions about whether males and females differ in decision-making processes and what the neurobiological mechanisms are that mediate these differences. This review will present an overview of what is currently known about sex

differences in animal models of decision-making and discuss the implications of these findings for understanding sex differences in psychiatric disease.

Sex differences in animal models of decision-making

Intertemporal decision-making

One form of decision-making that is commonly assessed in the laboratory is intertemporal choice, which refers to choosing between options that differ in their time of arrival. These options usually differ in reward magnitude and as such, decisions often consist of choosing between a small reward available after a short delay and a larger reward available after a long delay. Consequently, this behavior provides a measure of impulsivity ("impulsive choice"), or the extent to which an individual is willing to wait to procure a greater reward, and reflects the degree to which the delay diminishes (or "discounts") the subjective value of the larger reward. Typical intertemporal decision-making performance in such a "delay discounting task" manifests as a decrease in the choice of larger, delayed rewards in favor of smaller, more immediate rewards as the delays increase in duration. Importantly, alterations in impulsive choice have been strongly linked with psychiatric diseases, such as SUD (Coffey et al. 2003; Johnson et al. 2015; Kirby and Petry 2004) and attention deficit hyperactivity disorder (Winstanley 2011).

While this form of decision-making has been well studied in males, it has not been as thoroughly characterized in females, and in the studies that have been conducted, the results are not always transparent. For example, in one of the earliest

animal studies assessing sex differences in impulsive choice, Perry et al. (2008) tested male and female rats in an "adjusting delays" intertemporal choice task (in which the delay to the large reward was adjusted based on the rat's previous choices) and found that choice behavior did not differ between sexes. More recent studies have replicated this lack of sex differences in impulsive choice in both rats and mice using delay discounting tasks in which the delays shift systematically within a test session (Doremus-Fitzwater et al. 2012; Eubig et al. 2014; Hamilton et al. 2015; Lukkes et al. 2016), and several studies in monkeys have found a similar absence of sex differences (Carroll et al. 2009; Hamilton et al. 2011; Latzman et al. 2015; Rosati and Hare 2013). Within some of the rodent studies, however, there were more nuanced behavioral differences between males and females, suggesting that there may be subtle sex differences in intertemporal decision-making. For instance, Eubig et al. (2014) reported that following acute administration of amphetamine, females were quicker to initiate trials and displayed more impulsive choices than males. In another study, males and females were characterized as "flat" or "steep" discounters based on their task performance; "steep" female discounters displayed a greater reduction in their preference for the large, delayed reward than their male counterparts at longer delays (Koot et al. 2009). Age of testing may also be critical in detecting sex differences; Lukkes et al. (2016) reported that adolescent females displayed less impulsive choice than adolescent males. Finally, Perry et al. (2007) showed that in rats selectively bred to be high saccharin (HiS) or low saccharin (LoS) preferring, female LoS displayed greater impulsive choice than male LoS whereas there were no sex differences in HiS rats (Perry et al. 2007). Together, these studies indicate that although females appear to be

more impulsive than males under some conditions, further work is needed to expand upon these findings. For example, individual differences in impulsive choice at baseline or relationships between impulsive choice and other behavioral variables (Koot et al. 2009; Perry et al. 2007) may be critical determinants of sex differences that could have implications for differential vulnerability of females and males in the development of psychiatric diseases.

Probabilistic decision-making

Many decisions involve making choices between options that differ in both their expected rewards and their potential for accompanying negative consequences. There are several different animal models of such probabilistic decision-making, all of which assess the extent to which the probability of an adverse consequence discounts the value of a rewarding outcome. These models have been instrumental in demonstrating sex differences in probabilistic decision-making, although these differences appear to depend on both the task and the type of adverse consequence involved. In an initial study examining sex differences in probabilistic decision-making, van den Bos et al. (2012) used a rodent version of the Iowa Gambling Task (r-IGT), in which rats made discrete choices between a long-term advantageous option and a long-term disadvantageous option (van den Bos et al. 2012). The former consisted of frequent small food rewards (sugar pellets) and infrequent punishments in the form of quininelaced sugar pellets. In contrast, the disadvantageous option consisted of occasional large food rewards intermixed with frequent punishments. Importantly, similar to the human lowa Gambling Task (IGT), this decision-making task specifically measures the

process by which subjects learn about the probability distributions of reward vs. punishment delivery across a ten day period (i.e., the transition from uncertainty to risk). Although both males and females chose the advantageous option over the disadvantageous option to the same extent by the end of the r-IGT, males developed this preference more rapidly than females. In addition, as the rats progressed through the task, males continued to choose the advantageous option irrespective of whether they were rewarded or punished on the previous trial. This suggests that males learned quickly that while punishment could occur, the advantageous option was the better choice in the long term. Females, however, tended to shift their choice to the disadvantageous option regardless of whether they were rewarded or punished for choosing the advantageous option. Importantly, choice behavior in females did not seem to be modulated by estrous cycle. Overall, these differences suggest that males and females use distinct information gathering strategies in the r-IGT to execute a decision: while males appear to use more global information to make decisions and settle on their preference, females use details obtained after assessment of both options to determine the most adaptive choice (as evidenced by their constant switching between the advantageous and disadvantageous options). These findings in rats are consistent with those in humans, which show that females take longer to develop a preferential strategy in the Iowa Gambling Task (IGT) than males (van den Bos et al. 2013b).

In a more recent study, Peak et al. (2015) used a different variant of the Iowa Gambling task (the rodent gambling task; rGT) to assess sex differences in decision-making. In contrast to the r-IGT, which only has two options from which to choose, the

rGT consists of four options that differ in both reward (and punishment) probability and reward magnitude. Over multiple training sessions, rats learn that of the four options, one is the most advantageous in the long term and one is the most disadvantageous in the long term. Contrary to the results of the van den Bos (2012) study described above, Peak et al. (2015) showed that females developed optimal choice behavior more rapidly than males. The differences in the outcomes of these two experiments are likely due to differences in the decision-making tasks employed, and consequently may have significant implications for how males and females process different types of probabilistic decision-making. As mentioned above, the r-IGT is conducted for ten days, irrespective of meeting certain behavioral criteria upon completion, whereas the rGT conducted by Peak et al. (2015) occurred in multiple phases to facilitate learning of the task contingencies. Specifically, rats were trained to learn the different rewardpunishment contingencies in a forced choice version of the rGT in which they experienced only one option at a time. Only after rats were trained in this version of the rGT (7 consecutive days) did they move on to the free choice rGT in which they could choose between the different options. This is an important distinction as performance in the r-IGT may more closely model uncertainty (involving an unknown probability distribution) whereas performance in the rGT may more closely model risk (involving a known probability distribution) given the greater opportunities for learning in the latter task. In addition, the punishment used in the r-IGT consists of guinine-treated sugar pellets whereas in the rGT, the punishment is that of lost reward opportunity (a timeout period during which no new trials can be initiated). Finally, while the r-IGT involves choosing between two options, the rGT consists of calculating the optimal choice

among four options that differ in both probability of reward delivery and reward magnitude. It is therefore conceivable that males and females learn about and process information about the rewards and probabilities inherent to the tasks differently depending on the structure of the decisions and the types of adverse consequence involved.

To complicate matters further, our laboratory recently evaluated sex differences in a third probabilistic decision-making task (the "Risky Decision-Making task", RDT) involving varying probabilities of explicit physical punishment (Orsini et al. 2016). In this task, rats make discrete choices between two levers, one which delivers a small safe food reward and the other which delivers a large food reward accompanied by varying probabilities (ranging from 0-100%) of mild footshock (Simon et al. 2009). Female rats showed a significantly greater preference for the small, safe reward than male rats (Orsini et al. 2016), a difference which could not be explained by disparities in body weight influencing shock perception nor by differences in reward motivation. Further, choice behavior in this task in females was not modulated by estrous cycle. On the surface, it seems as though the greater preference for the "safe" option in females conflicts with their performance in the r-IGT, in which females shifted between the advantageous and disadvantageous options frequently. Similarly, the greater preference for the large, probabilistically punished reward in males seems inconsistent with their performance in the r-IGT, in which males settled on the advantageous reward more rapidly than females. One difference that could account for this discrepancy is the type of punishment involved (quinine-laced food vs. shock). An alternate, and equally appealing, explanation for these conflicting effects of sex on decision-making in the

RDT and r-IGT is that, similar to the case of the rGT and r-IGT, the tasks assess distinct components of decision-making. Whereas the RDT is conducted until behavioral stability is obtained (~25-30 days), the r-IGT is conducted for a predetermined duration (10 days), irrespective of whether behavior is stable at the completion of training. Thus, performance in the former likely reflects informed choice and behavior driven by risk, while performance in the latter assesses learning about the reward-outcomes contingencies (taxing uncertainty to a greater extent). These distinct components of decision-making may therefore recruit different strategies to make decisions. In the RDT (and the rGT), rats must rely on their knowledge of task contingencies to make an adaptive choice. In the r-IGT, however, rats need to gather information about task contingencies as they proceed through the training. Indeed, in both the human IGT and r-IGT, females take longer than males to develop a preference for the most advantageous option (van den Bos et al. 2013b; van den Bos et al. 2012). While this may manifest as greater risk-seeking compared to males, it may actually be reflective of females taking longer to learn about the probability distribution of outcomes as they need to spend more time evaluating all of the options before deciding upon the most optimal. Consistent with this notion, female rats took longer to reach stable performance than males in the RDT (Orsini et al. 2016). They also omitted significantly more trials than males, which could be viewed as another strategy to evade punished outcomes, albeit different than actively avoiding the punished option by choosing the safe option. It is therefore critical that researchers recognize that males and females may use different strategies to make probability-based decisions as it may help explain some of the welldescribed sex differences in psychiatric diseases (e.g., SUD) in which altered decision-

making plays a prominent role.

Potential explanations for sex differences in decision-making

Evolutionary and behavioral mechanisms

I In a recent review, Cross, Coping and Campbell (2011) proposed a theoretical account for the well-established observation in human studies that females are more impulsive than males, an explanation that may be extended to understanding such differences in other forms of decision-making (e.g., probabilistic decision-making). They posited that differences between men and women in reward sensitivity, punishment sensitivity, and effortful control can explain sex differences in impulsivity (Cross et al. 2011). Deeply rooted in evolutionary theory, these authors suggest that each of these components contributes differently to ensure the reproductive success of men and women. For example, men may be more risk-taking because they are hypersensitive to reward and hyposensitive to punishment. Across the animal kingdom, males' reproductive success frequently depends on competition with other males to obtain mates and rise in social hierarchy. Further, in some species, males were traditionally responsible for securing food and resources in the face of potential danger. In contrast, females tend to be hyposensitive to rewards and hypersensitive to punishment. This may derive from the fact that the reproductive success of females often depends on avoiding harm and death not only for their sake, but also for their offspring. Because the young of many species depend more heavily on mothers than fathers, the energy expenditure for females is greater and thus limits the number of offspring. Hence, it may

be advantageous for females to avoid harm and injury to increase their offspring's chances of survival.

It is important to consider that motivation for reward, be it food or a mate, does differ between males and females, which can influence the choices they make and thus their reproductive success (Yoest et al. 2014). In contrast to males, females are motivated for different rewards depending on their sexual receptivity. Females are motivated to find a mate and reproduce only when conception is likely; this increase in sexual motivation, however, is accompanied by a decrease in motivation for food (Fessler 2003). Yoest et al. (2014) argue that these parallel changes in motivation for food and sex ensure reproductive success for females as less time spent finding food and eating means that more time can be devoted to finding an optimal mate and reproducing when chances of conception are high. Interestingly, these fluctuations in motivation for food and sex in females are modulated by estradiol, indicating that gonadal hormones can influence adaptive decision-making (see further discussion below under 'Biological Mechanisms').

Differences in effortful control between males and females can also have a large impact on their reproductive success (Cross et al. 2011). Behaviorally defined, effortful control refers to "the ability to inhibit a dominant response and perform a subdominant response" (Cross et al., 2011, p. 102). It is through effortful control that organisms can regulate impulsive choice and risk-taking to be able to make adaptive decisions that promote long-term survival. MacDonald (2008) argued that effortful control was necessary to inhibit innate and automatic responses that had evolved over time, such as behaviors related to mate selection. For example, the drive for intrasexual

competition is so strong in males that is difficult for them to inhibit this approach behavior (MacDonald 2008). While this might predict greater impulsive behavior in males, which is not necessarily consistent with preclinical and clinical literature, it does align with the fact that males tend to be more risk-seeking than females (Orsini et al. 2016) and are quicker to develop a preference for the more advantageous option in the r-IGT (van den Bos et al. 2012). Conversely, Bjorklund and Kipp (1996) proposed that females have to engage in more effortful control to ensure their reproductive success (Bjorklund and Kipp 1996). For instance, in order to find the best possible mate, females must inhibit the tendency to choose the first mate available so as to secure a more optimal long-term partner. Females must also exert effortful inhibitory control to prioritize the needs of their dependent offspring over their own needs. Finally, females need to inhibit behaviors that would place themselves or their offspring in danger. Again, this theory of increased inhibitory control in females does not readily explain sex differences in intertemporal choice, but could account for differences observed in probabilistic decision-making. For example, in both the r-IGT and RDT, females might need to exert more inhibitory control so as to evaluate all available choices rather than guickly developing a preference for one option, as is the case with males. In the RDT in particular, poor inhibitory control in males might explain their willingness to endure physical punishment to obtain the larger reward. Together with differences in reward and punishment sensitivity, variations in effortful control between males and females may thus be differentially adaptive for each sex; however, it is noteworthy that these differences may also predispose men and women to different psychiatric diseases.

Another potential interpretation of the observed sex differences in decision-

making, which is not mutually exclusive from those discussed above, is that females may be more flexible and exploratory in their behavior than males (Koot et al. 2009). As mentioned in previous sections, males and females appear to employ different strategies in making decisions. Males may initially use an exploratory strategy to determine the most advantageous option, but then switch to a strategy of exploitation to take advantage of this option (Koot et al. 2009; van den Bos et al. 2013a). In contrast, females may more readily shift between exploration and exploitation, allowing them to gather more information about each of the options. This theory can account for sex differences observed in the r-IGT and RDT: not only did females in both tasks take longer to develop a preferential choice across sessions, they were able to more rapidly shift their choices between the options within a session. In contrast, in the RDT, males began each session by choosing the large reward option and continued to do so throughout the sessions, even when the probability of punishment was high. In the r-IGT, males guickly settled on the advantageous option early in training and persisted with this choice behavior throughout the duration of the r-IGT. Similarly, Koot et al., (2009) reported that, in contrast to males, females that discounted delays steeply shifted their preference to the smaller, more immediate reward at longer delays. Notably, these differences in strategy utilization could also support the reproductive success of each sex, suggesting an evolutionary basis for the divergence in approach tactics.

Biological mechanisms

Given the wealth of evidence demonstrating that behavioral responses to drugs of abuse vary across the estrous cycle (Becker 1999; Becker and Hu 2008; Evans et al.

2002: Festa and Quinones-Jenab 2004: Jackson et al. 2006: Justice and de Wit 1999: Perry et al. 2013; Perry et al. 2015; Quinones-Jenab et al. 1999), it is conceivable that fluctuations in ovarian hormones in females may contribute to sex differences in decision-making. Evidence for this supposition, however, is mixed. Decision-making performance does not vary across the estrous cycle in females in either the RDT (Orsini et al. 2016) or the human IGT (van den Bos et al. 2013b). Interestingly, however, another study showed that while choice performance in an effort discounting task (in which rats decide between a small, low-effort reward and a large, high-effort reward) did not vary across the estrous cycle in intact females, it was affected by ovariectomy (OVX) (Uban et al. 2012). Compared to sham controls, OVX females exhibited an increase in choices of the large, high-effort reward. This increase appeared to be at least partially mediated by estradiol and estrogen receptors, as it was reversed by administration of either high dose estradiol or a combination of ERa and ERB agonists (although interestingly, ER α and ER β agonists administered alone had the opposite effect in OVX rats). These findings provide initial evidence that female gonadal hormones can affect decision-making, although it is unclear whether this extends to intertemporal or probabilistic decision-making. Notably, several recent studies showed that systemic administration of testosterone can modulate male rats' performance in the RDT, an effort discounting task, and a probability discounting task (in which subjects choose between a small guaranteed reward and a large reward associated with varying probabilities of omission (Cooper et al. 2014; Wallin et al., 2015), hinting that hormones can influence other forms of decision-making. Based on this accumulated evidence, it will be important in future studies to more rigorously determine how gonadal hormones

impact decision-making, perhaps by manipulating hormone levels rather than passively tracking estrous cycle.

Although it has not yet been thoroughly investigated, the sex differences in, and effects of hormonal manipulations on, decision-making described above may be attributable in part to interactions between gonadal hormones and dopamine signaling (Becker and Hu 2008). Indeed, performance in many if not all preclinical models of decision-making is sensitive to dopaminergic manipulations. For example, systemic administration of amphetamine decreases preference for the large, risky reward in the RDT (Mitchell et al. 2011; Orsini et al. 2015; Orsini et al. 2016; Simon et al. 2011) and decreases impulsive choice in intertemporal decision-making tasks (Setlow et al. 2009; van Gaalen et al. 2006; Wade et al. 2000; Winstanley et al. 2003). Manipulations of dopamine receptors, either systemically or within a specific brain region, also affect decision-making behavior (Barrus and Winstanley 2016; Di Ciano et al. 2015; Mitchell et al. 2014a; Simon et al. 2011; St Onge et al. 2011; St Onge and Floresco 2009; Stopper et al. 2013). Over a decade of research has shown that females are more sensitive to dopamine-induced changes in behavior and that estradiol seems to play a large role in this effect (Becker and Hu 2008; Becker et al. 2012). For example, intact females show greater behavioral sensitization to amphetamine and cocaine than males (Becker et al. 1982; Robinson 1984; Robinson et al. 1982; van Haaren and Meyer 1991). OVX females show little to no sensitization to these stimulants (Forgie and Stewart 1994; Robinson 1984; Robinson et al. 1982; Sircar and Kim 1999; van Haaren and Meyer 1991), but estradiol administration can restore normal behavioral sensitization to amphetamine (Forgie and Stewart 1994; Peris et al. 1991). Interestingly, amphetamine

administration causes a greater decrease in choice of the large, "risky" reward in the RDT in females relative to males (Orsini et al. 2016). Although the role of estradiol in this effect has yet to be tested, it is consistent with previous work showing that males and females differ in their responses to dopamine manipulations. This could be due to basal differences in extracellular levels of dopamine, dopamine receptor levels, and/or autoreceptor control, all of which are modulated by estradiol (Becker and Hu 2008; Becker et al. 2012). For example, males have more dopamine D1 receptors in the striatum relative to either intact or OVX females (Hruska et al. 1982). In contrast, there are no sex differences in levels of dopamine D2 receptors in the striatum of intact males and females (Levesque and Di Paolo 1990). However, there is greater D2 binding in OVX rats compared to castrated rats, and upon administration of estradiol, these receptors are rapidly downregulated in OVX females but not castrated males (Bazzett and Becker 1994). In light of these findings, it will be important to determine how gonadal hormones interact with dopamine signaling during decision-making as it could reveal mechanisms underlying the observed sex differences.

Despite the wealth of studies that have documented sex differences in decisionmaking at the behavioral level (see Intertemporal and Probabilistic Decision-making sections above), there is little information regarding the neural mechanisms that might underlie these differences. The only animal study conducted to date showed that the orbitofrontal cortex (OFC) is differentially activated in males and females (as assessed with c-fos expression) following testing in the r-IGT (van Hasselt et al. 2012). Specifically, c-fos expression in the lateral OFC was inversely correlated with the proportion of advantageous choices in the r-IGT in females (this relationship was absent

in males). The majority of studies that have investigated this question have used neuroimaging techniques in human subjects. Consistent with findings from van Hasselt, these studies also show that the OFC is differentially recruited for males and females in various types of decision-making tasks. For example, in the human IGT, the OFC is activated more robustly in males than females (Bolla et al. 2004). Another study used the Risky Gains Task to assess sex differences in neural activation during decisionmaking. In this task, participants choose among three options, one of which yields a certain reward (safe choice) and two that may or may not yield a larger reward (uncertain choice). The authors found that the OFC in females was more dynamically engaged than males during task performance (Lee et al. 2009): while there were no correlations between neural activity and behavior in males, there was a negative correlation between neural activity in the OFC and choice of the uncertain reward when preceded by a punished outcome (i.e., no reward delivery) and a positive correlation between OFC neural signal and choice of the uncertain outcome when preceded by a uncertain, but unpunished, outcome. Interestingly, in a recent study (Crowley et al. 2015), males had greater OFC activation than females prior to making safe choices in another risk-based decision-making task. These last two studies suggest that the OFC in females may be more selectively tuned to process punishment and uncertainty whereas the OFC in males may be more selectively recruited to process information regarding safe reinforcement. To our knowledge, there are no data on whether these sex differences in OFC recruitment extend to intertemporal choice behavior.

Other neuroimaging studies have reported that additional areas of the prefrontal cortex are recruited in a sex-dependent manner. Using the Balloon Analogue Risk Task,

Cazzell et al. (2012) reported that, compared to males, females had greater activation of the dorsolateral prefrontal cortex (dIPFC) in both hemispheres specifically during periods in which they experienced loss of monetary rewards. Interestingly, in another study, there were hemispheric differences in dIPFC activation between males and females in the IGT. Whereas there was increased activity in the right dIPFC in males relative to females, dIPFC activation was greater in the left dIPFC in females compared to males (Bolla et al. 2004). The insular cortex is also implicated in mediating risky choices in females, but not males (Lee et al. 2009). In the Risky Gains Task, Lee et al. (2009) showed that signal intensity in the insula was positively correlated with the number of choices of the uncertain outcome in female subjects when this choice type was followed by another choice of the uncertain outcome. Given the insula's role in encoding of aversive information and anticipated risk (Nagvi et al. 2014), it is conceivable that in females the insula is part of a network with the OFC and dIPFC that processes risk of uncertainty and punishment-related information associated with choices. All of the aforementioned studies are limited, however, in that they are correlational in nature; it will hence be useful to employ animal models to address the causal role of activity in these systems.

Brain regions that are known to be sexually dimorphic are involved in various forms of decision-making. For example, the amygdala is larger in males than females (Goldstein et al. 2001), is recruited in a sex-dependent manner during regulation of emotional memories (Cahill 2006; Cahill et al. 2001; Cahill et al. 2004; Kilpatrick et al. 2006), and is critically involved in both intertemporal and probabilistic decision-making tasks. Lesions of the basolateral amygdala in male rats cause an increase in impulsive

choice (Winstanley et al. 2004). Similarly, in the rGT and RDT, lesions of the BLA increase choice of a large reward associated with greater probabilities of punishment (Orsini et al. 2015; Zeeb and Winstanley 2011). In contrast, in a probability discounting task, temporary inactivation of the basolateral amygdala causes male rats to decrease their choice of the large, uncertain reward (Ghods-Sharifi et al. 2009). Together, these studies show that the amygdala is a key brain region in regulating adaptive decision-making. Given its sexually dimorphic structure and function, it stands to reason that the same manipulations of the amygdala in females may yield different results than those in males and thus suggests that the amygdala may contribute to sex-dependent differences in decision-making.

Clinical implications

Understanding the precise mechanisms underlying sex differences in decisionmaking may have significant clinical implications as many psychiatric diseases that are characterized by maladaptive decision-making are sex-biased. The prevalence of schizophrenia, which is associated with poor performance in the IGT (Kim et al. 2007; Kim et al. 2009; Shurman et al. 2005; Struglia et al. 2011) and increased impulsivity (Ahn et al. 2011), is greater in males than females (Abel et al. 2010). Greater risk-taking and impulsivity are characteristic symptoms of attention deficit hyperactivity disorder (Evenden 1999), which is diagnosed ten times more frequently in males than females (McCarthy et al. 2012). Anorexia nervosa is 13 times more prevalent in females than males (McCarthy et al. 2012) and is associated with pathological risk aversion (Kaye et al. 2013). Finally, there are considerable sex differences in SUD (Becker et al. 2012;

Journal of Neuroscience Research This article is protected by copyright. All rights reserved.

Carroll et al. 2004; Lynch 2006), which has been shown in both preclinical and human studies to be associated with increased impulsive choice and risk-taking behavior (Anker et al., 2009; Bechara et al. 2001; Gowin et al. 2013; Mitchell et al. 2014a; Mitchell et al. 2014b). Thus, differences in decision-making between males and females could be linked to each sex's predisposition to specific psychiatric conditions. For instance, the fact that female rats choose the small, safe reward more than males in the RDT (Orsini et al. 2016) could suggest that a similar behavioral phenotype in women renders them more vulnerable to development of eating disorders. Alternatively, it is possible that psychiatric diseases could impact decision-making in one sex more than the other. As an example, females, who at baseline appear to be more risk-averse (i.e., prefer options that are not associated with risk of punishment) than males, are quicker to escalate their drug use, progress from recreational drug use to dependence more rapidly, and are more vulnerable to relapse (Bobzean et al. 2014; Lynch 2006). Hence, it is possible that females are more sensitive to the effects of chronic drug use on decision-making than males. Consequently, females may display an increase in risky behavior associated with drug abuse, such as escalation of use and relapse. To date, however, the majority of preclinical studies that have examined relationships between drug use and risk-taking have exclusively used males. For example, Mitchell et al. (2014a) demonstrated that chronic cocaine self-administration causes an increase in risk-taking in the RDT in male rats. It is possible that females would show a different behavioral trajectory (e.g., more rapid transition to a risk-seeking phenotype) than males in this same experimental design. Overall, this underscores the importance of studying the mechanisms underlying decision-making in both males and females in both normal

and pathological conditions to determine whether tailored treatment is warranted for each sex.

Conclusions

This review outlines clear sex differences in decision-making, which may be due to different strategies that have evolved to ensure the reproductive success of each sex; however, this review also illustrates that there are still large gaps in our knowledge and understanding of these sex differences, largely due to the paucity of studies in female subjects. In a climate in which sex-dependent psychiatric diseases such as SUD and post-traumatic stress disorder are on the rise, it is exceedingly important that resources are devoted to research that addresses these major gaps in knowledge. The recent mandate by the National Institutes of Health that requires the inclusion of sex as a biological variable has brought this issue to the forefront of the neuroscience research community (Clayton and Collins 2014). Specifically, this new policy requires strong justification from the literature and/or preliminary data to only use one sex, clearly indicating that there should be few excuses for not including both sexes in a research program. Resistance to such efforts will only impede scientific discoveries that could benefit the health of both men and women. Thus, it is our hope that such mandates, in addition to educating the scientific community through lectures and publications, will encourage researchers to embrace the inclusion of both sexes in studies of decisionmaking to produce more representative and translatable scientific discoveries.

Conflict of Interest Statement

The authors have no conflicts of interest to disclose.

Role of authors

CAO drafted the manuscript. Both CAO and BS revised and finalized the manuscript.

epte Acce

> Journal of Neuroscience Research This article is protected by copyright. All rights reserved.

Table Legends

Table 1. Sex differences in commonly-used decision-making tasks in rodents.

. epted Acce

> Journal of Neuroscience Research This article is protected by copyright. All rights reserved.

References

- Abel KM, Drake R, Goldstein JM. 2010. Sex differences in schizophrenia. International review of psychiatry 22(5):417-428.
- Ahn WY, Rass O, Fridberg DJ, Bishara AJ, Forsyth JK, Breier A, Busemeyer JR, Hetrick WP, Bolbecker AR, O'Donnell BF. 2011. Temporal discounting of rewards in patients with bipolar disorder and schizophrenia. Journal of abnormal psychology 120(4):911-921.
- Ashenhurst JR, Seaman M, Jentsch JD. 2012. Responding in a test of decision-making under risk is under moderate genetic control in the rat. Alcoholism: Clinical and Experimental Research 36(6):941-949.
- Barrus MM, Winstanley CA. 2016. Dopamine D3 Receptors Modulate the Ability of Win-Paired Cues to Increase Risky Choice in a Rat Gambling Task. The Journal of neuroscience : the official journal of the Society for Neuroscience 36(3):785-794.
- Bazzett TJ, Becker JB. 1994. Sex differences in the rapid and acute effects of estrogen on striatal D2 dopamine receptor binding. Brain research 637(1-2):163-172.
- Bechara A, Dolan S, Denburg N, Hindes A, Anderson SW, Nathan PE. 2001. Decision-making deficits, linked to a dysfunctional ventromedial prefrontal cortex, revealed in alcohol and stimulant abusers. Neuropsychologia 39(4):376-389.
- Becker JB. 1999. Gender differences in dopaminergic function in striatum and nucleus accumbens. Pharmacology, biochemistry, and behavior 64(4):803-812.
- Becker JB, Hu M. 2008. Sex differences in drug abuse. Frontiers in neuroendocrinology 29(1):36-47.
- Becker JB, Perry AN, Westenbroek C. 2012. Sex differences in the neural mechanisms mediating addiction: a new synthesis and hypothesis. Biology of sex differences 3(1):14.
- Becker JB, Robinson TE, Lorenz KA. 1982. Sex differences and estrous cycle variations in amphetamine-elicited rotational behavior. European journal of pharmacology 80(1):65-72.
- Bjorklund DF, Kipp K. 1996. Parental investment theory and gender differences in the evolution of inhibition mechanisms. Psychological bulletin 120(2):163-188.
- Bobzean SA, DeNobrega AK, Perrotti LI. 2014. Sex differences in the neurobiology of drug addiction. Experimental neurology 259:64-74.
- Bolla KI, Eldreth DA, Matochik JA, Cadet JL. 2004. Sex-related differences in a gambling task and its neurological correlates. Cerebral cortex 14(11):1226-1232.
- Cahill L. 2006. Why sex matters for neuroscience. Nature reviews Neuroscience 7(6):477-484.
- Cahill L, Haier RJ, White NS, Fallon J, Kilpatrick L, Lawrence C, Potkin SG, Alkire MT. 2001. Sex-related difference in amygdala activity during emotionally influenced memory storage. Neurobiology of learning and memory 75(1):1-9.
- Cahill L, Uncapher M, Kilpatrick L, Alkire MT, Turner J. 2004. Sex-related hemispheric lateralization of amygdala function in emotionally influenced memory: an FMRI investigation. Learning & memory 11(3):261-266.
- Carroll ME, Lynch WJ, Roth ME, Morgan AD, Cosgrove KP. 2004. Sex and estrogen influence drug abuse. Trends in pharmacological sciences 25(5):273-279.
- Carroll ME, Mach JL, La Nasa RM, Newman JL. 2009. Impulsivity as a behavioral measure of withdrawal of orally delivered PCP and nondrug rewards in male and female monkeys. Psychopharmacology 207(1):85-98.
- Cazzell M, Li L, Lin ZJ, Patel SJ, Liu H. 2012. Comparison of neural correlates of risk decision making between genders: an exploratory fNIRS study of the Balloon Analogue Risk Task (BART). Neuroimage 62(3):1896-1911.
- Clayton JA, Collins FS. 2014. Policy: NIH to balance sex in cell and animal studies. Nature 509(7500):282-283.

- Coffey SF, Gudleski GD, Saladin ME, Brady KT. 2003. Impulsivity and rapid discounting of delayed hypothetical rewards in cocaine-dependent individuals. Experimental and clinical psychopharmacology 11(1):18-25.
- Cooper SE, Goings SP, Kim JY, Wood RI. 2014. Testosterone enhances risk tolerance without altering motor impulsivity in male rats. Psychoneuroendocrinology 40:201-212.
- Cross CP, Copping LT, Campbell A. 2011. Sex differences in impulsivity: a meta-analysis. Psychological bulletin 137(1):97-130.
- Crowley TJ, Dalwani MS, Mikulich-Gilbertson SK, Young SE, Sakai JT, Raymond KM, McWilliams SK, Roark MJ, Banich MT. 2015. Adolescents' Neural Processing of Risky Decisions: Effects of Sex and Behavioral Disinhibition. PloS one 10(7):e0132322.
- Di Ciano P, Pushparaj A, Kim A, Hatch J, Masood T, Ramzi A, Khaled MA, Boileau I, Winstanley CA, Le Foll B. 2015. The Impact of Selective Dopamine D2, D3 and D4 Ligands on the Rat Gambling Task. PloS one 10(9):e0136267.
- Doremus-Fitzwater TL, Barreto M, Spear LP. 2012. Age-related differences in impulsivity among adolescent and adult Sprague-Dawley rats. Behavioral neuroscience 126(5):735-741.
- Eubig PA, Noe TE, Floresco SB, Sable JJ, Schantz SL. 2014. Sex differences in response to amphetamine in adult Long-Evans rats performing a delay-discounting task. Pharmacology, biochemistry, and behavior 118:1-9.
- Evans SM, Haney M, Foltin RW. 2002. The effects of smoked cocaine during the follicular and luteal phases of the menstrual cycle in women. Psychopharmacology 159(4):397-406.
- Evenden JL. 1999. Varieties of impulsivity. Psychopharmacology 146(4):348-361.
- Fessler DM. 2003. No time to eat: an adaptationist account of periovulatory behavioral changes. The Quarterly review of biology 78(1):3-21.
- Festa ED, Quinones-Jenab V. 2004. Gonadal hormones provide the biological basis for sex differences in behavioral responses to cocaine. Hormones and behavior 46(5):509-519.
- Floresco SB, Tse MT, Ghods-Sharifi S. 2008. Dopaminergic and glutamatergic regulation of effort- and delay-based decision-making. Neuropsychopharmacology 33(8):1966-1979.
- Forgie ML, Stewart J. 1994. Six differences in the locomotor-activating effects of amphetamine: role of circulating testosterone in adulthood. Physiology & behavior 55(4):639-644.
- Ghods-Sharifi S, St Onge JR, Floresco SB. 2009. Fundamental contribution by the basolateral amygdala to different forms of decision making. The Journal of neuroscience : the official journal of the Society for Neuroscience 29(16):5251-5259.
- Goldstein JM, Seidman LJ, Horton NJ, Makris N, Kennedy DN, Caviness VS, Jr., Faraone SV, Tsuang MT. 2001. Normal sexual dimorphism of the adult human brain assessed by in vivo magnetic resonance imaging. Cerebral cortex 11(6):490-497.
- Gowin JL, Mackey S, Paulus MP. 2013. Altered risk-related processing in substance users: Imbalance of pain and gain. Drug and alcohol dependence 132(1-2):13-21.
- Hamilton KR, Mitchell MR, Wing VC, Balodis IM, Bickel WK, Fillmore M, Lane SD, Lejuez CW, Littlefield AK, Luijten M, Mathias CW, Mitchell SH, Napier TC, Reynolds B, Schutz CG, Setlow B, Sher KJ, Swann AC, Tedford SE, White MJ, Winstanley CA, Yi R, Potenza MN, Moeller FG. 2015. Choice impulsivity: Definitions, measurement issues, and clinical implications. Personality disorders 6(2):182-198.
- Hamilton LR, Czoty PW, Nader MA. 2011. Behavioral characterization of adult male and female rhesus monkeys exposed to cocaine throughout gestation. Psychopharmacology 213(4):799-808.
- Hruska RE, Ludmer LM, Pitman KT, De Ryck M, Silbergeld EK. 1982. Effects of Estrogen on Striatal Dopamine receptor function in male and female rats. Pharmacology, biochemistry, and behavior 16(2):285-291.
- Jackson LR, Robinson TE, Becker JB. 2006. Sex differences and hormonal influences on acquisition of cocaine self-administration in rats. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 31(1):129-138.

Johnson MW, Bruner NR, Johnson PS. 2015. Cocaine dependent individuals discount future rewards more than future losses for both cocaine and monetary outcomes. Addictive behaviors 40:132-136.

- Justice AJ, de Wit H. 1999. Acute effects of d-amphetamine during the follicular and luteal phases of the menstrual cycle in women. Psychopharmacology 145(1):67-75.
- Kaye WH, Wierenga CE, Bailer UF, Simmons AN, Bischoff-Grethe A. 2013. Nothing tastes as good as skinny feels: the neurobiology of anorexia nervosa. Trends in neurosciences 36(2):110-120.
- Kilpatrick LA, Zald DH, Pardo JV, Cahill LF. 2006. Sex-related differences in amygdala functional connectivity during resting conditions. NeuroImage 30(2):452-461.
- Kim H, Lee D, Shin YM, Chey J. 2007. Impaired strategic decision making in schizophrenia. Brain research 1180:90-100.
- Kim YT, Lee KU, Lee SJ. 2009. Deficit in decision-making in chronic, stable schizophrenia: from a reward and punishment perspective. Psychiatry Investig 6(1):26-33.
- Kirby KN, Petry NM. 2004. Heroin and cocaine abusers have higher discount rates for delayed rewards than alcoholics or non-drug-using controls. Addiction 99(4):461-471.
- Koot S, van den Bos R, Adriani W, Laviola G. 2009. Gender differences in delay-discounting under mild food restriction. Behavioural brain research 200(1):134-143.
- Latzman RD, Taglialatela JP, Hopkins WD. 2015. Delay of gratification is associated with white matter connectivity in the dorsal prefrontal cortex: a diffusion tensor imaging study in chimpanzees (Pan troglodytes). Proceedings Biological sciences / The Royal Society 282(1809):20150764.
- Lee TM, Chan CC, Leung AW, Fox PT, Gao JH. 2009. Sex-related differences in neural activity during risk taking: an fMRI study. Cerebral cortex 19(6):1303-1312.
- Levesque D, Di Paolo T. 1990. Effect of the rat estrous cycle at ovariectomy on striatal D-1 dopamine receptors. Brain research bulletin 24(2):281-284.
- Lukkes JL, Thompson BS, Freund N, Andersen SL. 2016. The developmental inter-relationships between activity, novelty preferences, and delay discounting in male and female rats. Developmental psychobiology 58(2):231-242.
- Lynch WJ. 2006. Sex differences in vulnerability to drug self-administration. Experimental and clinical psychopharmacology 14(1):34-41.
- MacDonald KB. 2008. Effortful control, explicit processing, and the regulation of human evolved predispositions. Psychological review 115(4):1012-1031.
- McCarthy MM, Arnold AP, Ball GF, Blaustein JD, De Vries GJ. 2012. Sex differences in the brain: the not so inconvenient truth. The Journal of neuroscience : the official journal of the Society for Neuroscience 32(7):2241-2247.
- Mitchell MR, Vokes CM, Blankenship AL, Simon NW, Setlow B. 2011. Effects of acute administration of nicotine, amphetamine, diazepam, morphine, and ethanol on risky decision-making in rats. Psychopharmacology 218(4):703-712.
- Mitchell MR, Weiss VG, Beas BS, Morgan D, Bizon JL, Setlow B. 2014a. Adolescent risk taking, cocaine self-administration, and striatal dopamine signaling. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 39(4):955-962.
- Mitchell MR, Weiss VG, Ouimet DJ, Fuchs RA, Morgan D, Setlow B. 2014b. Intake-dependent effects of cocaine self-administration on impulsive choice in a delay discounting task. Behavioral neuroscience 128(4):419-429.
- Naqvi NH, Gaznick N, Tranel D, Bechara A. 2014. The insula: a critical neural substrate for craving and drug seeking under conflict and risk. Annals of the New York Academy of Sciences 1316:53-70.

- Orsini CA, Trotta RT, Bizon JL, Setlow B. 2015. Dissociable Roles for the Basolateral Amygdala and Orbitofrontal Cortex in Decision-Making under Risk of Punishment. The Journal of neuroscience : the official journal of the Society for Neuroscience 35(4):1368-1379.
- Orsini CA, Willis ML, Gilbert RJ, Bizon JL, Setlow B. 2016. Sex differences in a rat model of risky decision making. Behavioral neuroscience 130(1):50-61.
- Peak JN, Turner KM, Burne TH. 2015. The effect of developmental vitamin D deficiency in male and female Sprague-Dawley rats on decision-making using a rodent gambling task. Physiology of Behavior 138:319-324.
- Peris J, Decambre N, Coleman-Hardee ML, Simpkins JW. 1991. Estradiol enhances behavioral sensitization to cocaine and amphetamine-stimulated striatal [3H]dopamine release. Brain research 566(1-2):255-264.
- Perry AN, Westenbroek C, Becker JB. 2013. The development of a preference for cocaine over food identifies individual rats with addiction-like behaviors. PloS one 8(11):e79465.
- Perry AN, Westenbroek C, Jagannathan L, Becker JB. 2015. The Roles of Dopamine and alpha1-Adrenergic Receptors in Cocaine Preferences in Female and Male Rats. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 40(12):2696-2704.
- Perry JL, Nelson SE, Anderson MM, Morgan AD, Carroll ME. 2007. Impulsivity (delay discounting) for food and cocaine in male and female rats selectively bred for high and low saccharin intake. Pharmacology, biochemistry, and behavior 86(4):822-837.
- Perry JL, Nelson SE, Carroll ME. 2008. Impulsive choice as a predictor of acquisition of IV cocaine self-administration ad reinstatement of cocaine-seeking behavior in male and female rats. Experimental Clinical Psychopharmacology 16(2):165-177.
- Quinones-Jenab V, Ho A, Schlussman SD, Franck J, Kreek MJ. 1999. Estrous cycle differences in cocaine-induced stereotypic and locomotor behaviors in Fischer rats. Behavioural brain research 101(1):15-20.
- Robinson TE. 1984. Behavioral sensitization: characterization of enduring changes in rotational behavior produced by intermittent injections of amphetamine in male and female rats. Psychopharmacology 84(4):466-475.
- Robinson TE, Becker JB, Presty SK. 1982. Long-term facilitation of amphetamine-induced rotational behavior and striatal dopamine release produced by a single exposure to amphetamine: sex differences. Brain research 253(1-2):231-241.
- Rosati AG, Hare B. 2013. Chimpanzees and bonobos exhibit emotional responses to decision outcomes. PloS one 8(5):e63058.
- Setlow B, Mendez IA, Mitchell MR, Simon NW. 2009. Effects of chronic administration of drugs of abuse on impulsive choice (delay discounting) in animal models. Behavioural pharmacology 20(5-6):380-389.
- Shurman B, Horan WP, Nuechterlein KH. 2005. Schizophrenia patients demonstrate a distinctive pattern of decision-making impairment on the Iowa Gambling Task. Schizophrenia research 72(2-3):215-224.
- Simon NW, Gilbert RJ, Mayse JD, Bizon JL, Setlow B. 2009. Balancing risk and reward: a rat model of risky decision making. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 34(10):2208-2217.
- Simon NW, Montgomery KS, Beas BS, Mitchell MR, LaSarge CL, Mendez IA, Banuelos C, Vokes CM, Taylor AB, Haberman RP, Bizon JL, Setlow B. 2011. Dopaminergic modulation of risky decision-making. The Journal of neuroscience : the official journal of the Society for Neuroscience 31(48):17460-17470.
- Sircar R, Kim D. 1999. Female gonadal hormones differentially modulate cocaine-induced behavioral sensitization in Fischer, Lewis, and Sprague-Dawley rats. The Journal of pharmacology and experimental therapeutics 289(1):54-65.

Journal of Neuroscience Research

- St Onge JR, Abhari H, Floresco SB. 2011. Dissociable contributions by prefrontal D1 and D2 receptors to risk-based decision making. The Journal of neuroscience : the official journal of the Society for Neuroscience 31(23):8625-8633.
- St Onge JR, Floresco SB. 2009. Dopaminergic modulation of risk-based decision making. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 34(3):681-697.
- Stopper CM, Khayambashi S, Floresco SB. 2013. Receptor-specific modulation of risk-based decision making by nucleus accumbens dopamine. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 38(5):715-728.
- Struglia F, Stratta P, Gianfelice D, Pacifico R, Riccardi I, Rossi A. 2011. Decision-making impairment in schizophrenia: relationships with positive symptomatology. Neuroscience letters 502(2):80-83.
- Uban KA, Rummel J, Floresco SB, Galea LA. 2012. Estradiol modulates effort-based decision making in female rats. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 37(2):390-401.
- van den Bos R, Davies W, Dellu-Hagedorn F, Goudriaan AE, Granon S, Homberg J, Rivalan M, Swendsen J, Adriani W. 2013a. Cross-species approaches to pathological gambling: a review targeting sex differences, adolescent vulnerability and ecological validity of research tools. Neuroscience and biobehavioral reviews 37(10 Pt 2):2454-2471.
- van den Bos R, Homberg J, de Visser L. 2013b. A critical review of sex differences in decisionmaking tasks: focus on the Iowa Gambling Task. Behavioural brain research 238:95-108.
- van den Bos R, Jolles J, van der Knaap L, Baars A, de Visser L. 2012. Male and female Wistar rats differ in decision-making performance in a rodent version of the Iowa Gambling Task. Behavioural brain research 234(2):375-379.
- van Gaalen MM, van Koten R, Schoffelmeer AN, Vanderschuren LJ. 2006. Critical involvement of dopaminergic neurotransmission in impulsive decision making. Biological psychiatry 60(1):66-73.
- van Haaren F, Meyer ME. 1991. Sex differences in locomotor activity after acute and chronic cocaine administration. Pharmacology, biochemistry, and behavior 39(4):923-927.
- van Hasselt FN, de Visser L, Tieskens JM, Cornelisse S, Baars AM, Lavrijsen M, Krugers HJ, van den Bos R, Joels M. 2012. Individual variations in maternal care early in life correlate with later life decision-making and c-fos expression in prefrontal subregions of rats. PloS one 7(5):e37820.
- Wade TR, de Wit H, Richards JB. 2000. Effects of dopaminergic drugs on delayed reward as a measure of impulsive behavior in rats. Psychopharmacology 150(1):90-101.
- Wallin KG, Alves JM, Wood RI. 2015. Anabolic-andronergic steroids and decision-making: Probability and effort discounting in male rats. Psychoneuroendocrinology 57:84-92.
- Winstanley CA. 2011. The utility of rat models of impulsivity in developing pharmacotherapies for impulse control disorders. British journal of pharmacology 164(4):1301-1321.
- Winstanley CA, Dalley JW, Theobald DE, Robbins TW. 2003. Global 5-HT depletion attenuates the ability of amphetamine to decrease impulsive choice on a delay-discounting task in rats. Psychopharmacology 170(3):320-331.
- Winstanley CA, Theobald DE, Cardinal RN, Robbins TW. 2004. Contrasting roles of basolateral amygdala and orbitofrontal cortex in impulsive choice. The Journal of neuroscience : the official journal of the Society for Neuroscience 24(20):4718-4722.
- Yoest KE, Cummings JA, Becker JB. 2014. Estradiol, dopamine and motivation. Central nervous system agents in medicinal chemistry 14(2):83-89.
- Zeeb FD, Winstanley CA. 2011. Lesions of the basolateral amygdala and orbitofrontal cortex differentially affect acquisition and performance of a rodent gambling task. The Journal of neuroscience : the official journal of the Society for Neuroscience 31(6):2197-2204.

Journal of Neuroscience Research

Table 1.

Task	Task measure	Sex tested	Sex differences	References
Rat lowa Gambling Task	Learning about probabilities of different reward outcomes	Males and females	Males develop preference for the advantageous option more quickly than females	van den Bos et al., 2012
Rat Gambling Task	Choice of optimal (more reward, less timeout punishment) over suboptimal options	Males and females	Females develop preference for the advantageous option more quickly than males	Peak et al., 2015
Risky Decision- Making Task	Choice of small reward vs. large reward associated with probabilistic footshock	Males and females	Females show greater preference for small, "safe" reward	Orsini et al., 2016
Delay discounting	Choice of small immediate vs. larger delayed rewards	Males and females	Females tend to be more impulsive than males	Eubig et al., 2014; Koot et al., 2009; Lukkes et al., 2016; Perry et al., 2007
Probability discounting	Choice of small guaranteed vs. large probabilistic rewards	Males only	Unknown	St. Onge and Floresco, 2009
Effort discounting	Choice of small low-effort vs. large, high-effort rewards	Males and females	No direct comparison between sexes	Uban et al., 2012; Floresco et al., 2008
Rat Balloon Analog Risk Task	Learning about the probability distribution of avoiding risk and obtaining rewards	Males and females	No sex differences	Ashenhurst et al., 2012

Accept