



Monitoring the Future

NATIONAL SURVEY RESULTS ON DRUG USE, 1975-2008

VOLUME II

**College Students &
Adults Ages 19-50**

2008

**National Institute on Drug Abuse
National Institutes of Health
U.S. Department of Health & Human Services**

MONITORING THE FUTURE
NATIONAL SURVEY RESULTS ON DRUG USE, 1975–2008

Volume II
College Students and Adults Ages 19–50

by

Lloyd D. Johnston, Ph.D.
Patrick M. O'Malley, Ph.D.
Jerald G. Bachman, Ph.D.
John E. Schulenberg, Ph.D.

The University of Michigan
Institute for Social Research

National Institute on Drug Abuse
6001 Executive Boulevard
Bethesda, Maryland 20892

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
National Institutes of Health

2009

This publication was written by the principal investigators and staff of the Monitoring the Future project, at the Institute for Social Research, the University of Michigan, under Research Grants No. R01 DA 01411 and R01 DA 016575 from the National Institute on Drug Abuse.

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the NIH.

Public Domain Notice

All material appearing in this volume is in the public domain and may be reproduced or copied without permission from the Institute or the authors. Citation of the source is appreciated.

Recommended Citation

Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (2009). *Monitoring the Future national survey results on drug use, 1975–2008: Volume II, College students and adults ages 19–50* (NIH Publication No. 09-7403). Bethesda, MD: National Institute on Drug Abuse.

ABBREVIATED CONTENTS*

Detailed Contents	v
List of Tables	ix
List of Figures.....	xiii
<i>Chapter 1</i> Introduction	1
<i>Chapter 2</i> Key Findings: An Overview and Integration across Five Populations	5
<i>Chapter 3</i> Study Design and Procedures	57
<i>Chapter 4</i> Prevalence of Drug Use in Early and Middle Adulthood	75
<i>Chapter 5</i> Trends in Drug Use in Early and Middle Adulthood	123
<i>Chapter 6</i> Attitudes and Beliefs about Drugs among Young Adults	183
<i>Chapter 7</i> The Social Context	203
<i>Chapter 8</i> Prevalence of Drug Use among College Students	247
<i>Chapter 9</i> Trends in Drug Use among College Students	258
Index	298

*See next page for Detailed Contents.

DETAILED CONTENTS

<i>Chapter 1</i>	Introduction	1
	Surveys of College Students and Noncollege Peers	1
	Surveys of Young Adults and Those Ages 35, 40, 45, and 50	2
	General Purposes of the Research.....	3
	Risk and Protective Behaviors Related to HIV/AIDS	4
<i>Chapter 2</i>	Key Findings: An Overview and Integration across Five Populations	5
	Trends in Illicit Drug Use—The Advent of Cohort Effects	5
	College–Noncollege Differences in Illicit Drug Use.....	21
	Male–Female Differences in Illicit Drug Use.....	22
	Trends in Alcohol Use	22
	College–Noncollege Differences in Alcohol Use.....	23
	Male–Female Differences in Alcohol Use.....	23
	Trends in Cigarette Smoking	24
	Age- and Cohort-Related Differences in Cigarette Smoking	26
	College–Noncollege Differences in Cigarette Smoking.....	27
	Male–Female Differences in Cigarette Smoking.....	27
	Racial/Ethnic Comparisons.....	28
	Drug Use in Eighth Grade.....	29
	Drug Use by Age 50	31
	Summary and Conclusions	32
<i>Chapter 3</i>	Study Design and Procedures	57
	Research Design and Procedures for the Twelfth-Grade Surveys.....	57
	The Population under Study.....	57
	The Omission of Dropouts.....	58
	Sampling Procedures and Sample Weights	58
	Questionnaire Administration.....	58
	Questionnaire Format.....	59
	Research Design and Procedures for the Eighth- and Tenth-Grade Surveys	59
	Mode of Administration.....	60
	Questionnaire Forms and Sample Proportions	61
	Research Design and Procedures for the Twelfth-Grade Follow-up Surveys	61
	Follow-Up Procedures	62
	Follow-Up Questionnaire Format.....	62
	Representativeness and Sample Accuracy.....	63
	School Participation	63
	Student Participation.....	65
	Sampling Accuracy of the Estimates	65
	Panel Retention.....	66

DETAILED CONTENTS (continued)

The Problem of Panel Attrition.....	66
Response Rates	66
The Impact of Panel Attrition on Research Results.....	67
Effects on Relational Analyses	68
Validity of the Measures of Self-Reported Drug Use.....	69
Consistency and the Measurement of Trends.....	70
<i>Chapter 4</i> Prevalence of Drug Use in Early and Middle Adulthood	75
A Note on Adjusted Lifetime Prevalence Estimates.....	76
Prevalence of Drug Use as a Function of Age.....	77
Prevalence Comparisons for Subgroups of Young Adults	83
Gender Differences	83
Regional Differences	85
Population Density Differences	88
<i>Chapter 5</i> Trends in Drug Use in Early and Middle Adulthood	123
Trends in Prevalence: Early and Middle Adulthood.....	124
Trends for Important Subgroups of Young Adults.....	133
Gender Differences in Trends.....	133
Regional Differences in Trends	138
Population Density Differences in Trends.....	142
<i>Chapter 6</i> Attitudes and Beliefs about Drugs among Young Adults	183
Perceived Harmfulness of Drugs	183
Personal Disapproval of Drug Use	188
Cohort Differences and Their Implications for Prevention and Theory.....	191
<i>Chapter 7</i> The Social Context	203
Peer Norms among Adults.....	203
Current Perceptions of Friends' Attitudes	204

DETAILED CONTENTS (continued)

Trends in Peer Norms	205
Adults' Exposure to Drug Use through Friends and Others	206
Exposure to Drug Use	207
Trends in Exposure to Drug Use	209
Perceived Availability of Drugs among Adults	211
Perceived Availability	211
Trends in Perceived Availability	212
<i>Chapter 8</i> Prevalence of Drug Use among College Students	247
Prevalence of Drug Use among College Students versus Those Not Enrolled in College	248
Gender Differences in Prevalence of Use among College Students	250
<i>Chapter 9</i> Trends in Drug Use among College Students	258
Trends in Prevalence 1980–2008: College Students versus Those Not Enrolled in College	259
Gender Differences in Trends among College Students	266
Index	298

LIST OF TABLES

Table 2-1.	Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th, and 12th Graders, College Students, and Young Adults (Ages 19–28).....	36
Table 2-2.	Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th, and 12th Graders, College Students, and Young Adults (Ages 19–28).....	43
Table 2-3.	Trends in 30-Day Prevalence of Use of Various Drugs for 8th, 10th, and 12th Graders, College Students, and Young Adults (Ages 19–28).....	49
Table 2-4.	Trends in 30-Day Prevalence of Daily Use of Various Drugs for 8th, 10th, and 12th Graders, College Students, and Young Adults (Ages 19–28)	54
Table 3-1.	Sample Sizes and Response Rates	71
Table 4-1.	Prevalence of Use of Various Types of Drugs by Gender among Respondents of Modal Ages 19–30, 2008	88
Table 4-2.	Lifetime Prevalence of Use of Various Types of Drugs by Subgroups among Respondents of Modal Ages 19–30, 2008	91
Table 4-3.	Annual Prevalence of Use of Various Types of Drugs by Subgroups among Respondents of Modal Ages 19–30, 2008	94
Table 4-4.	Thirty-Day Prevalence of Use of Various Types of Drugs by Subgroups among Respondents of Modal Ages 19–30, 2008	97
Table 4-5.	Thirty-Day Prevalence of Daily Use of Various Types of Drugs by Subgroups among Respondents of Modal Ages 19–30, 2008	100
Table 5-1.	Trends in Lifetime Prevalence of Various Types of Drugs among Respondents of Modal Ages 19–28	148
Table 5-2.	Trends in Annual Prevalence of Various Types of Drugs among Respondents of Modal Ages 19–28	151
Table 5-3.	Trends in 30-Day Prevalence of Various Types of Drugs among Respondents of Modal Ages 19–28	153
Table 5-4.	Trends in 30-Day Prevalence of Daily Use of Various Types of Drugs among Respondents of Modal Ages 19–28	155

LIST OF TABLES (continued)

Table 5-5.	Trends in Annual and 30-Day Prevalence of an Illicit Drug Use Index among Respondents of Modal Ages 19–28, Total and by Gender	156
Table 6-1.	Trends in Perceived Harmfulness as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30	193
Table 6-2.	Trends in Proportions Disapproving of Drug Use among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30	199
Table 7-1.	Trends in Proportions of Friends Disapproving of Drug Use among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30	216
Table 7-2.	Trends in Friends’ Use of Drugs as Estimated by Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50	218
Table 7-3.	Trends in Exposure to Drug Use among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30	234
Table 7-4.	Trends in Availability of Drugs as Perceived by Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50	240
Table 8-1.	Lifetime Prevalence of Use for Various Types of Drugs, 2008: Full-Time College Students vs. Others among Respondents 1 to 4 Years beyond High School	252
Table 8-2.	Annual Prevalence of Use for Various Types of Drugs, 2008: Full-Time College Students vs. Others among Respondents 1 to 4 Years beyond High School.....	254
Table 8-3.	Thirty-Day Prevalence of Use for Various Types of Drugs, 2008: Full-Time College Students vs. Others among Respondents 1 to 4 Years beyond High School.....	255
Table 8-4.	Thirty-Day Prevalence of Daily Use for Various Types of Drugs, 2008: Full-Time College Students vs. Others among Respondents 1 to 4 Years beyond High School	256
Table 9-1.	Trends in Lifetime Prevalence of Various Types of Drugs among College Students 1 to 4 Years beyond High School	269

LIST OF TABLES (continued)

Table 9-2.	Trends in Annual Prevalence of Various Types of Drugs among College Students 1 to 4 Years beyond High School	271
Table 9-3.	Trends in 30-Day Prevalence of Various Types of Drugs among College Students 1 to 4 Years beyond High School	272
Table 9-4.	Trends in 30-Day Prevalence of Daily Use of Various Types of Drugs among College Students 1 to 4 Years beyond High School	273
Table 9-5.	Trends in Lifetime, Annual, and 30-Day Prevalence of an Illicit Drug Use Index among College Students 1 to 4 Years beyond High School, by Gender.....	274

LIST OF FIGURES

Figure 2-1.	Trends in Annual Prevalence of an Illicit Drug Use Index across 5 Populations.....	55
Figure 3-1.	Schools included in 1 Year’s Data Collection: 8th, 10th and 12th Grades.....	72
Figure 3-2.	School Participation Rates	73
Figure 4-1.	Any Illicit Drug: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008.....	101
Figure 4-2.	Any Illicit Drug other than Marijuana: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008.....	102
Figure 4-3.	Marijuana: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008	103
Figure 4-4.	Amphetamines: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008.....	104
Figure 4-5.	Methamphetamine: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30 by Age Group, 2008.....	105
Figure 4-6.	Crystal Methamphetamine (Ice): Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30 by Age Group, 2008	106
Figure 4-7.	Cocaine: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008	107
Figure 4-8.	Crack Cocaine: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008.....	108
Figure 4-9.	Other Cocaine: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008.....	109
Figure 4-10.	Hallucinogens: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008.....	110
Figure 4-11.	LSD: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30 by Age Group, 2008	111

LIST OF FIGURES (continued)

Figure 4-12.	Hallucinogens other than LSD: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30 by Age Group, 2008.....	112
Figure 4-13.	Inhalants: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30 by Age Group, 2008	113
Figure 4-14.	Sedatives (Barbiturates): Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008	114
Figure 4-15.	Narcotics other than Heroin: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008.....	115
Figure 4-16.	Tranquilizers: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008.....	116
Figure 4-17.	Ecstasy (MDMA): Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30 by Age Group, 2008.....	117
Figure 4-18.	Steroids: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 30 by Age Group, 2008.....	118
Figure 4-19.	Heroin: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008.....	119
Figure 4-20a.	Alcohol: Lifetime, Annual, and 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008	120
Figure 4-20b.	Alcohol: 2-Week Prevalence of 5 or More Drinks in a Row and 30-Day Prevalence of Daily Use among Respondents of Modal Ages 18 through 50 by Age Group, 2008	121
Figure 4-21.	Cigarettes: Annual, 30-Day, Daily, and Half-Pack-a-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group, 2008.....	122

LIST OF FIGURES (continued)

Figure 5-1.	Any Illicit Drug: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group.....	157
Figure 5-2.	Any Illicit Drug other than Marijuana: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group.....	158
Figure 5-3a.	Marijuana: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group.....	159
Figure 5-3b.	Marijuana: Trends in 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group.....	160
Figure 5-3c.	Marijuana: Trends in 30-Day Prevalence of Daily Use among Respondents of Modal Ages 18 through 50 by Age Group.....	161
Figure 5-4.	Inhalants: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 32 by Age Group	162
Figure 5-5.	Hallucinogens: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group.....	163
Figure 5-6.	LSD: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 35 by Age Group	164
Figure 5-7.	Hallucinogens other than LSD: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 35 by Age Group.....	165
Figure 5-8.	Ecstasy (MDMA): Trends in Annual Prevalence among Respondents of Modal Ages 18 through 32 by Age Group	166
Figure 5-9.	Cocaine: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group	167
Figure 5-10.	Crack Cocaine: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group	168
Figure 5-11.	Heroin: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group	169
Figure 5-12.	Narcotics other than Heroin: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group.....	170

LIST OF FIGURES (continued)

Figure 5-13.	Amphetamines: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group	171
Figure 5-14.	Crystal Methamphetamine (Ice): Trends in Annual Prevalence among Respondents of Modal Ages 18 through 32 by Age Group.....	172
Figure 5-15.	Sedatives (Barbiturates): Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group	173
Figure 5-16.	Tranquilizers: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group	174
Figure 5-17.	Steroids: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 32 by Age Group	175
Figure 5-18a.	Alcohol: Trends in Annual Prevalence among Respondents of Modal Ages 18 through 50 by Age Group	176
Figure 5-18b.	Alcohol: Trends in 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group	177
Figure 5-18c.	Alcohol: Trends in 30-Day Prevalence of Daily Use among Respondents of Modal Ages 18 through 50 by Age Group.....	178
Figure 5-18d.	Alcohol: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row among Respondents of Modal Ages 18 through 50 by Age Group.....	179
Figure 5-19a.	Cigarettes: Trends in 30-Day Prevalence among Respondents of Modal Ages 18 through 50 by Age Group.....	180
Figure 5-19b.	Cigarettes: Trends in 30-Day Prevalence of Daily Use among Respondents of Modal Ages 18 through 50 by Age Group.....	181
Figure 5-19c.	Cigarettes: Trends in 30-Day Prevalence of Smoking a Half Pack or More per Day among Respondents of Modal Ages 18 through 50 by Age Group.....	182
Figure 9-1.	Any Illicit Drug: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	275

LIST OF FIGURES (continued)

Figure 9-2.	Any Illicit Drug other than Marijuana: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	276
Figure 9-3a.	Marijuana: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students.....	277
Figure 9-3b.	Marijuana: Trends in 30-Day Prevalence of Daily Use among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	278
Figure 9-4.	Inhalants: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students.....	279
Figure 9-5.	Hallucinogens: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	280
Figure 9-6.	LSD: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students.....	281
Figure 9-7.	Hallucinogens other than LSD: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students.....	282
Figure 9-8.	Ecstasy (MDMA): Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	283
Figure 9-9.	Cocaine: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students.....	284
Figure 9-10a.	Narcotics other than Heroin: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students.....	285

LIST OF FIGURES (continued)

Figure 9-10b.	Vicodin: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School	286
Figure 9-10c.	OxyContin: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School.....	287
Figure 9-11.	Amphetamines: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	288
Figure 9-12.	Sedatives (Barbiturates): Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	289
Figure 9-13.	Tranquilizers: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	290
Figure 9-14a.	Alcohol: Trends in Annual Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students.....	291
Figure 9-14b.	Alcohol: Trends in 30-day Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	292
Figure 9-14c.	Alcohol: Trends in 30-day Prevalence of Daily Use among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	293
Figure 9-14d.	Alcohol: Trends in 2-Week Prevalence of 5 or More Drinks in a Row among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	294
Figure 9-15a.	Cigarettes: Trends in 30-Day Prevalence among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	295
Figure 9-15b.	Cigarettes: Trends in 30-Day Prevalence of Daily Use among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students	296

LIST OF FIGURES (continued)

Figure 9-15c. Cigarettes: Trends in 30-Day Prevalence of Smoking a Half Pack or More per Day among College Students vs. Others 1 to 4 Years beyond High School and among Male vs. Female College Students297

Chapter 1

INTRODUCTION

This monograph presents findings from the Monitoring the Future (MTF) study on high school graduates who have completed follow-up surveys across young and middle adulthood, beginning after their senior year in high school. Now in its 34th year, MTF is a long-term program of research conducted at the University of Michigan's Institute for Social Research under a series of investigator-initiated research grants from the National Institute on Drug Abuse. The study is comprised of several ongoing series of annual surveys of nationally representative samples of 8th- and 10th-grade students (begun in 1991), 12th-grade students (begun in 1975), and high school graduates into adulthood (begun in 1976). As we report in this volume, several segments of the general adult population are covered in the follow-up surveys. One important segment is American college students; a second is comprised of their age peers who are not attending college, sometimes called the "forgotten half"; and a third is all young adult high school graduates of modal ages 19 to 30, which we refer to as the "young adult" sample. Finally, high school graduates at the specific modal ages of 35, 40, 45, and 50 are now included each year in the longer term follow-ups.

The follow-up surveys have been conducted by mail on representative subsamples of the previous participants from each high school senior class. This volume presents data from the 1977 through 2008 follow-up surveys of the graduating high school classes of 1976 through 2007, as these respondents have progressed beyond high school and into adulthood—now through age 50 for the oldest respondents.

To permit this volume to stand alone, we have repeated some material from *Volume I*. Specifically, chapter 2 in the present volume is the same as chapter 2 in *Volume I*; it provides an integrated overview of the key findings presented in both volumes. Chapter 3, "Study Design and Procedures," is also the same as chapter 3, *Volume I*. The reader already familiar with *Volume I* may wish to skip over these chapters here.

SURVEYS OF COLLEGE STUDENTS AND NONCOLLEGE PEERS

The MTF follow-up samples have provided excellent coverage of the national college student population for almost three decades. College students tend to be a difficult population to study. They are generally not well covered in household surveys, which typically exclude dormitories, fraternities, and sororities. Further, institution-based samples of college students must be quite large in order to attain accurate national representation because of the great heterogeneity in the types of student populations served in those institutions. Obtaining good samples and high response rates within many institutions also poses difficulties. The current study, which in essence draws the college sample prospectively, in senior year of high school, has considerable advantages for generating a broadly representative sample of college students who emerge from each graduating cohort, and does so at very low cost. Further, its "before," "during," and "after" college design permits examination of the many changes associated with the college experience.

Moreover, a particularly rare feature is that it generates comparable panel data on high school graduates who are not attending college, an important segment not only in its own right but also as a comparison group for college students.

As defined here, the college student population comprises all full-time students, one to four years post-high school, enrolled in a two- or four-year college in March during the year of the survey. More is said about this sample definition in chapters 3 and 8. Results on the *prevalence* of drug use among college students and their noncollege peers in 2008 are reported in chapter 8, and results on *trends* in substance use among college students and their noncollege peers over the past 29 national surveys are reported in chapter 9.

SURVEYS OF YOUNG ADULTS AND THOSE AGES 35, 40, 45, AND 50

The young adult sample considered in this volume is comprised of representative samples from each graduating class from 1996 to 2007, all surveyed in 2008, corresponding to modal ages 19 through 30.¹ (The college students are included as a part of this young adult sample.) The study design calls for annual follow-up surveys of each class cohort (though each individual participates in a follow-up survey only every two years) through modal age 30, after which surveys occur at five-year intervals beginning at modal age 35 (17 years after high school). In 2008 the graduating classes of 1976, 1981, 1986, and 1991 were sent the age-50, age-45, age-40, and age-35 questionnaires, respectively. Data were collected from 50-year-old respondents for the first time in 2008. Panel data into middle adulthood on nationally representative samples of the population are extremely rare and valuable. Panel data on successive class cohorts from the general population are particularly valuable, because they allow the differentiation of period-, age-, and cohort-related change.

In this volume, we have reweighted respondent data to adjust for the effects of panel attrition on measures such as drug use using a statistical technique called poststratification, which will be explained later. We are less able to adjust for the absence of high school dropouts who were not included in the original 12th-grade sample. Because nearly all college students have completed high school, the omission of dropouts should have almost no effect on the college student population estimates, but this omission does affect the estimates for entire age groups. Therefore, the reader is advised that the omission of about 15% of each cohort who drop out of high school will likely make the drug use estimates given here for the various young adult age bands somewhat low for the age group as a whole. The proportional effect may be greatest for some of the most dangerous drugs, such as heroin, crack, and methamphetamine, as well as for cigarettes—the use of which is highly correlated with educational aspirations and attainment.

¹While data exist in recent years on ages 19–30, trends for young adults have been presented for 19–28-year-olds in chapter 5 in order to allow for an additional two years of comparison.

GENERAL PURPOSES OF THE RESEARCH

The research purposes of MTF are extensive and can be outlined here only briefly.² One major purpose is to serve a social monitoring or social indicator function, to accurately characterize the levels and trends in certain behaviors, attitudes, beliefs, and environmental conditions in the population. Social indicators can have important agenda-setting functions for society, drawing attention to new threats to the public health and estimating the extent of those threats. They are especially useful for gauging progress toward national goals and indicating the impacts of major historical events, including social or policy changes. Another purpose of the study is to develop knowledge that increases our understanding of how and why changes in these behaviors, attitudes, beliefs, and so on are taking place. (In health-related disciplines, such work is usually labeled *epidemiology*.) These two broad purposes are addressed in the current series of volumes.

The many additional purposes for the research are addressed in other types of publications including books, journal articles, chapters, and occasional papers. These purposes include helping to discover what types of young people are at greatest risk for developing various patterns of drug abuse, gaining a better understanding of the lifestyles and value orientations associated with various patterns of drug use and monitoring how those orientations are shifting over time, determining the immediate and more general aspects of the social environment that are associated with drug use and abuse, and determining how drug use is affected by major transitions into and out of social environments (such as military service, civilian employment, college, unemployment) or social roles (marriage, pregnancy, parenthood). We also seek to determine the life course of the various drug-using behaviors during this period of development, distinguish such “age effects” from cohort and period effects that are influencing drug use, determine the effects of social legislation and changing regulations on various types of substance use, and determine the changing connotations of drug use and changing patterns of multiple drug use among youth. We believe that differentiating among period, age, and cohort effects on use of various types of substances has been a particularly important contribution of the project; the MTF cohort-sequential research design is especially well suited to allow the differentiation of changes with age common to all cohorts (age effects), differences among cohorts that tend to persist across time (cohort effects), and changes common to most or all ages in a given historical period (period effects). Knowing which type of change is occurring is important to determining what types of causes are accounting for the change, when in the life course they may have had their impact (in the case of cohort effects it may well have been in earlier historical periods than when the change is actually documented), and in being able to predict future change more accurately. A number of important cohort effects that first emerged in the 1990s in terms of both use and attitudes about use are featured in this volume, in addition to period effects and some well established age effects.

²For a more complete listing and discussion of MTF’s many objectives, see Johnston, L. D., O’Malley, P. M., Schulenberg, J. E., & Bachman, J. G. (2006). *The aims and objectives of the Monitoring the Future study and progress toward fulfilling them as of 2006* (Monitoring the Future Occasional Paper No. 65). Ann Arbor, MI: Institute for Social Research. Available online at www.monitoringthefuture.org/pubs.html#papers.

RISK AND PROTECTIVE BEHAVIORS RELATED TO HIV/AIDS

One important additional purpose, related to but distinct from the others, is no longer addressed in this monograph; instead, it is the subject of a separate short monograph presenting the 2008 results. In 2004, for the first time, data were included on the prevalence and interconnectedness of risk and risk-reduction behaviors related to the spread of the human immunodeficiency virus (HIV), which causes acquired immunodeficiency syndrome (AIDS). A set of questions about these behaviors was included in the 2004 through 2006 follow-up surveys in two of the six questionnaire forms that were administered to 21- to 30-year-olds. These questions were then included in three of the six questionnaire forms beginning in 2007. In addition to questions about injection drug use, and needle sharing in particular, there are also questions related to numbers of sexual partners, gender of partners, use of condoms, and blood donation. The results from that work are presented and discussed for the first time this year in a separate monograph that will be posted on the MTF Web site shortly after this volume is posted. See www.monitoringthefuture.org.

Readers interested in publications dealing with any of these areas are invited to visit the MTF Web site at www.monitoringthefuture.org. A complete listing of the many publications from the study is available there, as well as abstracts and/or complete manuscripts for many of those publications. Complete text of MTF press releases is also provided. For additional information, please e-mail us at MTFinfo@isr.umich.edu.

Chapter 2

KEY FINDINGS AN OVERVIEW AND INTEGRATION ACROSS FIVE POPULATIONS

Monitoring the Future, now in its 34th year, has become one of the nation's most relied-upon sources of information on changes taking place in licit and illicit psychoactive drug use among American adolescents, college students, young adults, and more recently, middle-aged adults. During the last three and a half decades, the study has tracked and reported on the use of an ever-growing array of such substances in these populations.

This annual series of monographs is one of the major vehicles by which the epidemiological findings from MTF are reported. Findings from the inception of the study in 1975 through 2008 are included—the results of 34 national in-school surveys and 32 national follow-up surveys.

MTF has conducted in-school surveys of nationally representative samples of (a) 12th-grade students each year since 1975 and (b) 8th- and 10th-grade students each year since 1991. In addition, beginning with the class of 1976, the project has conducted follow-up mail surveys on representative subsamples of the respondents from each previously participating 12th-grade class. These follow-up surveys now continue well into adulthood.

A number of important findings have been summarized in this chapter to provide the reader with an overview of the key results. Because so many populations, drugs, and prevalence intervals are discussed here, a single integrative set of tables (Tables 2-1 through 2-4) show the 1991–2008 trends for all drugs on five populations: 8th-grade students, 10th-grade students, 12th-grade students, full-time college students modal ages 19–22, and all young adults modal ages 19–28 who are high school graduates. (Note that the young adult group includes the college student population.) *Volume II* also contains data on older age bands based on the longer term follow-up surveys: specifically, ages 35, 40, 45, and 50.

TRENDS IN ILLICIT DRUG USE—THE ADVENT OF COHORT EFFECTS

Early in the 1990s, we noted an increase in use of several illicit drugs among secondary school students, and some important changes among the students in terms of certain key attitudes and beliefs related to drug use. In the volume reporting 1992 survey results, we noted the beginning of such reversals in both use and attitudes among 8th graders, the youngest respondents surveyed in this study, and also a reversal in attitudes among 12th graders. Specifically, the proportions seeing great risk in using drugs began to decline, as did the proportions saying they disapproved

of use. As we suggested then, those reversals indeed presaged “an end to the improvements in the drug situation that the nation may be taking for granted.” The use of illicit drugs rose sharply in all three grade levels after 1992 as negative attitudes and beliefs about drug use continued to erode. This pattern continued into the mid-1990s, and beyond that for some drugs.

In 1997, for the first time in six years, the overall rate of *illicit drug use* finally began to decline among 8th graders. Although marijuana use continued to rise that year among 10th and 12th graders, their use of several other drugs leveled off, and relevant attitudes and beliefs also began to reverse in many cases. In 1998, illicit drug use continued a gradual decline among 8th graders and started to decline at 10th and 12th grades. In 1999 and 2000, the decline continued for 8th graders, whereas use held fairly level among 10th and 12th graders. In 2002 and 2003, use by 8th and 10th graders decreased significantly, and use by 12th graders finally began to drop, albeit by less than a statistically significant amount. Nonsignificant declines continued for all three grades in 2004. The long-term decline in illicit drug use among 8th graders paused in 2005, but continued among 10th and 12th graders. Annual prevalence declined in all three grades in both 2006 and 2007, and the 8th-grade decline in 2007 was statistically significant. The decline continued among 10th graders in 2008; however, illicit drug use increased among 8th and 12th graders in 2008, possibly signaling an end to the immediately preceding period of decline. As we have noted previously, the gradual decline observed among 8th graders suggested an eventual further decline at the upper grades as the 8th graders aged. We are seeing those declines, though they are a bit erratic.

As will be illustrated below in the discussion of specific drugs, the increase in use of many drugs during the 1990s among secondary school students, combined with fairly level rates of use among college students and young adults, resulted in some unusual reversals in the usage rates by age (see Figure 2-1). In the early years of the epidemic, illicit drug use rates were clearly higher in the college-age group (and eventually the young adults) than they were among secondary school students. But by the late 1990s, the highest rates of active use (i.e., use within the prior year or prior 30 days) were found in the late secondary school years. In fact, in 1996 and 1997 both 10th and 12th graders actually had higher annual prevalence rates for illicit drug use (i.e., higher percentages reporting use within the prior year) than either college students or all young adults. This changed somewhat after 2001, as the earlier heavier using cohorts of adolescents began to comprise the college student and young adult populations, while at the same time use among the secondary school students was declining.

- In 2008, the rank order by age group for annual prevalence of using *any illicit drug* was 12th graders (37%), college students (35%), 19- to 28-year-olds (34%), 10th graders (27%), and 8th graders (14%). With respect to using *any illicit drug other than marijuana* in the past 12 months, the rank order was 19- to 28-year-olds (19%), 12th graders (18%), college students (15%), 10th graders (11%), and 8th graders (7%). As can be seen by this divergence of trends for the different age groups, something other than a simple secular trend in drug use was taking place; specifically, important cohort differences were emerging.

- From the early 1990s until 1997, *marijuana* use rose sharply among secondary school students, as did their use of a number of *other illicit drugs*, though more gradually. We have called this period a “relapse phase” in the longer term epidemic. An increase in marijuana use also occurred among American college students, largely reflecting “generational replacement” (i.e., a cohort effect), wherein earlier cohorts were replaced in the college population by more recent ones who were more drug-experienced before they left high school. This resurgence in illicit drug use spread *up* the age spectrum in a reversal of the way the epidemic spread several decades earlier. In the 1960s the epidemic began on the nation’s college campuses, and then diffused downward in age to high school students and eventually to middle school students. This time the increases began in middle schools and radiated up the age spectrum. The graduating class cohorts in the middle and late 1990s carried with them the pattern of heavier drug use that emerged while they were in secondary school in the early 1990s.

The increases during the 1990s in use of *any illicit drug* (including use of *marijuana* and use of *other illicit drugs* treated as a class) were substantially larger, in both proportional and absolute terms, in the three secondary school grades than in either the college or young adult populations. Among college students and young adults, the annual prevalence of use of *any illicit drug* held remarkably stable from 1991 through 1997, at the same time that use rose appreciably among adolescents (Figure 2-1). We projected that, as generational replacement continued to occur, we would likely see some increase in use of illicit drugs by the young adults. As would be expected given their younger age range (19–22), the increase happened sooner and more sharply among the college students than among the young adults in general (age range 19–28). Peak rates (since 1990) in annual prevalence of any illicit drug were reached in 1996 among 8th graders, in 1997 among 10th and 12th graders, in 2001 among college students, and in 2008 in the young adult segment. Similarly, the more recent declines in use among secondary students have thus far shown up only modestly among college students, and even less so among young adults.

Again, these diverging trends across age groups clearly show that changes during the 1990s reflected some important cohort effects rather than broad secular trends that would have appeared simultaneously in all of the age groups. During all of the previous years of the study, the use of most drugs moved in parallel across most age groups, indicating that secular change was prevailing.

- Similar to the use patterns for illicit drugs, the trend for *cigarette* smoking evidenced a generational replacement effect during the 1990s in that college students showed a sharp increase in smoking beginning in 1995, as the heavier smoking cohorts of adolescents from the early to mid-1990s entered college. This has been a more typical pattern of change for cigarettes, however, since differences in cigarette smoking rates among class cohorts tend to remain through the life course and also tend to account for much of the overall change in use observed at any given age.

In the early 1990s, cigarette smoking among 8th and 10th graders rose by about 50%—a particularly sharp and worrisome rise (based on 30-day prevalence rates shown in Table

2-3, and daily and half-pack rates shown in Table 2-4); MTF helped draw national attention to this worrisome situation. Smoking also rose among 12th graders, beginning a year later.

The increase in current smoking ended among 8th and 10th graders in 1996, among 12th graders in 1997, and among college students in 1999. The nation then entered a period of appreciable decline in smoking rates that first began among 8th graders in 1997 and then began radiating up the age spectrum as those cohorts aged. (The 8th-grade 30-day prevalence rate fell from 21% in 1996 to 7% in 2008.) But among the college and the young adult strata, the declines have been less clear. The 30-day smoking prevalence rate for college students in 2008 (18%) is down about four tenths from the recent peak of 31% in 1999, with the decline accelerating after 2005 as the cohort effect worked its way up the age bands. Smoking among the young adult subgroup, on the other hand, has only dropped by one fifth (to 25%) since its recent peak rate of 31% in 1998. The decline in smoking rates among secondary school students has been decelerated in all three grades in recent years.

- During the 1990s, the annual prevalence of *marijuana* use tripled among 8th graders (from 6% in 1991 to 18% in 1996), more than doubled among 10th graders (from 15% in 1992 to 35% in 1997), and nearly doubled among 12th graders (from 22% in 1992 to 39% in 1997). Among college students, however, the increase in marijuana use, presumably due to a generational replacement effect, was much more gradual. Annual prevalence of use rose by about one third, from 27% in 1991 to 36% in 1998. Marijuana use began to decline in 1997 among 8th graders and then did the same in 1998 among 10th and 12th graders. The rate of decline was rather modest, however, perhaps due in part to effects of the public debates over medical use of marijuana during that period. In 2001, use remained level in all three grades, but between 2001 and 2004 all three grades showed significant declines in their annual prevalence of marijuana use, with the proportional decline greatest among 8th graders. The 8th graders have shown the most steady long-term decline since their recent peak, which occurred in 1996, although the decline halted in 2008, for a 40% drop since 1996. Declines had been occurring in the upper grades after 1997, but mostly since about 2001, with their annual prevalence rates having fallen from recent peaks by 31% and 16% (since 1997) for 10th and 12th graders, respectively. The decline in annual marijuana use from recent peak levels among college students has so far been quite modest, declining from 36% in 2001 to 32% in 2008. Young adults showed very little change in that interval (see Table 2-2).

Daily marijuana use in all of these groups rose substantially after 1992, reaching peak levels in a somewhat staggered fashion as that just described (see Table 2-4). Daily use began a slow decline after 1999 among 8th graders, after 2001 among 10th graders, and after 2003 among 12th graders, consistent with a cohort effect pattern. College student and young adult rates have been fairly level in recent years. In general, prevalence of daily marijuana use has been slow to decline, even though annual and 30-day prevalence figures have been dropping. Still, the rates today are low in relation to the peaks reported in the late 1970s. For example, 12th graders' daily prevalence of use of 5.4% in 2008 is

half the 10.7% peak figure reached in 1978—at the height of the illicit drug epidemic—and a bit below the recent high of 6.0% recorded in 2003.

The amount of perceived risk associated with using marijuana fell during the earlier period of increased use in the late 1970s, and fell again during the more recent resurgence of use in the 1990s. Indeed, perceived risk among 12th graders began to decline a year *before* use began to rise in the upturn of the 1990s, making perceived risk a leading indicator of change in use. (The same may have happened in 8th grade, as well, but we do not have data starting early enough to check that possibility.) The decline in perceived risk halted after 1997 for 8th and 10th graders, and annual prevalence began to decline a year or two later. Again, perceived risk was a leading indicator of change in use, as it has proven to be for a number of drugs. As discussed in *Volume I*, chapter 8, these attitudes show evidence of cohort effects over the past decade and a half.

Personal disapproval of marijuana use slipped considerably among 8th graders between 1991 and 1996 and among 10th and 12th graders between 1992 and 1997. For example, the proportions of 8th, 10th, and 12th graders who said they disapproved of trying marijuana once or twice fell by 17, 21, and 19 percentage points, respectively, over those intervals of decline. There has since been some increase in disapproval among 8th graders and, beginning more recently, among 10th and 12th graders.

- Among 12th graders, the proportions using *any illicit drug other than marijuana* in the past year rose from a low of 15% in 1992 to 22% in 2001 (see Table 2-2); these levels are substantially below the 34% peak rate reached two decades earlier, in 1981. All of the younger groups showed significant increases between 1992 and 1997, with use beginning to increase in 1992 among 8th graders, in 1993 among 10th and 12th graders, and in 1995 among college students—again reflecting evidence of a cohort effect. Use peaked in 1996 among 8th and 10th graders and by 1997 among 12th graders; it appears to have peaked around 2004 among the college students and young adults. The 8th graders have shown some gradual decline in their use of the other illicit drugs, treated as a class, since 1996 with some leveling in 2008; the decline among 10th graders paused after 1998 and did not resume until after 2001, pausing again in 2007 and then decreasing significantly in 2008; 12th-grade use also showed some declines after 2001, and stands just three percentage points lower (at 18%) in 2008. College students have begun to show some decline in use over the past several years, whereas among young adults, there is no evidence yet of a decline in the proportions using any illicit drug other than marijuana.
- Between 1989 and 1992 we noted an increase among 12th graders, college students, and young adults in their use of *LSD*, a drug quite popular in the late 1960s and early 1970s. In 1992 the newly added populations (8th and 10th graders) were also showing an increase in LSD use; for several more years, modest increases persisted in all five populations. Use of LSD peaked in 1995 among college students and young adults and in 1996 among 8th, 10th, and 12th graders, after which LSD use gradually declined in all five populations until 2001, then it dropped sharply in 2002. Overall, the pattern for LSD use seems more consistent with secular change than a cohort effect. The different age

groups moved in parallel for the most part, likely in response to historical events in the environment, including a sharp reduction in LSD availability after 2001.

Prior to the significant increase in LSD use among 12th graders in 1993, there was a significant 4.3-percentage-point decline between 1991 and 1992 in the proportion seeing great risk associated with trying LSD. (Once again, perceived risk proved to be a leading indicator of change in use.) The decline in perceived risk continued through 1997 and halted in 1998. The proportion of 12th graders disapproving of LSD use began to decline in 1992, and continued to decline through 1996.

Because LSD was one of the earliest drugs to be popularly used in the American drug epidemic, young people in the 1990s may have been relatively unaware of the risks of use. They had less opportunity to learn vicariously about the consequences of use by observing others around them or to learn from intense media coverage of the issue, which occurred some years earlier. We were concerned that this type of generational forgetting of the dangers of a drug, which occurs as a result of generational replacement, could set the stage for a whole new epidemic of use. In fact, perceived harmfulness of LSD began to decline after 1991 among 12th graders. Perceived risk and disapproval among 8th and 10th graders, first measured in 1993, both showed declines until 1997 or 1998, after which they leveled and then declined some more. In 2004, twelfth graders' personal disapproval of trying LSD increased significantly, with no further change until 2008 when disapproval dropped back to the 2003 level. Because the decline in use in the last few years has generally *not* been accompanied by expected changes in these attitudes and beliefs, we suspected that some displacement by another drug might have been taking place, at least through 2001. The most logical candidate is *ecstasy (MDMA)*, which, like LSD, is used for its hallucinogenic effects; ecstasy was popular in the club and rave scenes, and was very much on the rise through 2001. After 2001, a sharp decline in the reported availability of LSD in all five populations (which corresponded to the closing of a major LSD lab by the DEA) very likely played a major role in the sharp decline in use among all of them. However, we want to caution that young people's attitudes are changing such as to make them receptive to LSD use some time in the future, should a plentiful supply re-emerge.

- Questions about the use of *ecstasy (MDMA)* have been included in the follow-up surveys of college students and young adults since 1989; however, because of our concern about stimulating interest in an attractive-sounding and little-known drug, these questions were not added to the secondary school surveys until 1996. From 1989 to 1994, the annual prevalence rates tended to be quite low in the older age groups for whom we had data, but in 1995 these rates increased substantially—from 0.5% in 1994 to 2.4% in 1995 among college students, and from 0.7% to 1.6% over the same time span among young adults generally.

When usage data were first gathered on secondary school students in 1996, the 10th and 12th graders actually showed higher rates of annual use (both 4.6%) than the college students (2.8%). Ecstasy use then fell steadily at all three grades between 1996 and 1998, though it did not fall in the older age groups. But between 1998 and 2001, use rose

sharply in all five populations. In fact, annual prevalence more than doubled in that three-year period among 12th graders, college students, and young adults, and nearly doubled in the lower grades. In 2000 even the 8th graders showed a significant increase in use. Ecstasy use for all five age groups declined slightly in 2002, but significantly only for 10th graders; declined again in 2003, with significant drops for all groups except the college students; and showed some decline again in 2004, with the largest decreases among college students and young adults. This pattern suggests that both cohort effects and a secular trend were at work. Once again, this decline in use among 12th graders was predicted by an increase in perceived risk in 2001—an increase that continued through 2005. The annual prevalence fell by half in 2004 alone among college students, and all five groups are at rates that range from just under one half to three quarters lower than their recent peaks in 2001. Since 2004, use has been fairly stable among secondary school students, but continued to decline some among college students and young adults generally.

Ecstasy use among all five populations has been moving fairly synchronously since 1999, which suggests that a secular trend (some change in events in the social environment) has affected everyone. An important change during this period was the increasing availability of information on the adverse effects of ecstasy use via stories in the popular media, dissemination of the scientific evidence by the National Institute on Drug Abuse, and an anti-ecstasy media campaign by the Partnership for a Drug-Free America and the Office of National Drug Control Policy, initiated in 2002.

The quite dramatic increase in reported availability of ecstasy through 2001, reported by 12th graders, was substantiated by law enforcement data on ecstasy seizures. Of the 12th graders surveyed in 1991, only 22% thought they could get ecstasy fairly easily, but a decade later (in 2001) 62% thought that they could. After 2001, however, the perceived availability of ecstasy began decreasing in all three grades, possibly due in part to the steep decline in the number of users, who serve as supply points for others. The decreases continued into 2007 among 8th graders and then halted in 2008. In the upper grades, decline in perceived availability halted in 2007, followed in 2008 by some further downturn among 10th graders and some increase among 12th graders. See Figure 8-6 in *Volume I*, chapter 8 for a graphic presentation of the trends in ecstasy use, availability, and perceived risk for 12th graders.

- Between 1982 and 1992, annual prevalence rates for *amphetamine* use (other than use that was ordered by a physician) among 12th graders fell by nearly two thirds, from 20.3% to 7.1%. Rates among college students fell even more over the same interval, from 21.1% to 3.6%. During the relapse phase in the drug epidemic in the 1990s, annual amphetamine use increased by about half among 8th and 10th graders between 1991 and 1996, and also increased among 12th graders and college students between 1992 and 1996. After 1996 the age groups diverged, with amphetamine use declining gradually among 8th and 10th graders and continuing to rise among 12th graders, college students, and young adults until about 2002. The decline continued through 2004 for 8th graders and through 2005 for 10th graders, while the rise among 12th graders and college students finally halted by 2003. The 12th graders finally exhibited a significant decline in

annual prevalence of amphetamine use in 2005, and college students showed a leveling after 2006. Young adults have not shown a clear pattern of decline; their rates of amphetamine use have been stable since 2000. In 2008 all except the 8th graders showed further declines in use, although only the 10th-grade decline reached statistical significance. This pattern of cross-age-group change suggests a cohort effect at work for amphetamine use.

The increase in nonmedical use of amphetamines (and a concurrent decrease in disapproval) began among 12th graders in 1993; it followed a sharp drop in perceived risk a year earlier (which, as we have noted for a number of drugs, often serves as a leading indicator). Following a period of decline, both perceived risk and disapproval among 12th graders generally increased from 1995 through 2007 for perceived risk and through 2008 for disapproval. Nonmedical use of amphetamines among 12th graders remained fairly steady from 1997 through 2004, and then decreased from 2004 to 2008.

- Use of the amphetamine *Ritalin* outside of medical supervision showed a distinct increase around 1997—with annual prevalence among 12th graders going from 0.1% in 1992 to 2.8% in 1997—and then stayed level for a few years (see appendix E in *Volume I*, Table E-2³). Because of its increasing importance, a differently structured question was introduced for Ritalin use in 2001. This new question, which we prefer to the original, does not use a prior branching question and produced somewhat higher prevalence rates. Results from the new question suggest an ongoing, gradual decline in Ritalin use in all five populations, which continued into 2008 in all populations except 10th graders, whose use leveled that year.
- *Methamphetamine* questions were introduced in 1999 because of rising concern about use of this drug; but a decline in use has been observed among all five populations in the years since then, although young adults did not show declines until 2005. In 2007 this decline continued in all five populations, and was significant in grades 8 and 12. In 2008, there was some decrease for 12th graders and young adults; use in the other populations leveled. These declines occurred during a period in which there were many stories in the media suggesting that methamphetamine use was a *growing* problem—an example of the importance of having objective epidemiological data available against which to test conventional wisdom.
- Measures on the use of *crystal methamphetamine (ice)* (a crystallized form of methamphetamine that can be smoked, much like crack) have been included in MTF since 1990. The use of crystal methamphetamine increased between the early and late 1990s among the three populations asked about their use: 12th graders, college students, and young adults. The estimates are less stable than usual due to the relatively small sample sizes asked about this drug, but it appears that crystal methamphetamine use held

³As discussed in appendix E of *Volume I*, the absolute prevalence rates for Ritalin were probably higher than these statistics indicate, but the trend story is likely quite accurate. See Table 2-2 for more accurate estimates of the absolute annual prevalence rates in recent years; these estimates are based on a new question that does not require the respondent to first indicate some amphetamine use before asking about his or her Ritalin use.

fairly steady from 1999 through 2005 among 12th graders, after which it began to decline, including a significant drop in 2008. Use rose somewhat among college students and other young adults until 2005, before trailing off some since then.

- **Inhalants** are defined as fumes or gases that are inhaled to get high, and they include common household substances such as glues, aerosols, butane, and solvents of various types. Among 12th graders there was a long-term gradual increase in the use of inhalants (unadjusted for nitrite inhalants) from 1976 to 1987, followed by a leveling for a few years and then a further increase in the 1990s. In the early 1990s, there was a troublesome increase in inhalant use among secondary school students generally, followed by a reversal after 1995. After reaching a low point in 2002 or 2003 in grades 8, 10, and 12, use of inhalants increased some in all grades, but then declined in all grades more recently, at least through 2007. Perceived risk among 8th and 10th graders was declining fairly steadily after 2001, quite possibly as a result of generational forgetting of the dangers of these drugs; this decline halted among the 8th graders, but then resumed in 2008. A new anti-inhalant campaign might well be effective in offsetting this decline in perceived risk in recent years, much as a similar campaign did in the mid-1990s.
- One class of inhalants, **amyl and butyl nitrites**, became somewhat popular in the late 1970s, but their use has been almost eliminated. The annual prevalence rate among 12th-grade students was 6.5% in 1979 but only 0.6% in 2008. When nitrites were included in the definition of inhalants, they masked the increase that was occurring in the use of other inhalants, because their use was declining (Figure 5-4c in *Volume I*).
- **Crack cocaine** use spread rapidly from the early to mid-1980s. Still, among 12th graders, the use of crack remained relatively low during this period (3.9% annual prevalence in 1987). Clearly, crack had quickly attained a reputation as a dangerous drug, and by the time of our first measurement of perceived risk in 1987, it was seen as the most dangerous of all drugs. Annual prevalence dropped sharply in the next few years, reaching 1.5% by 1991, where it remained through 1993. Perceived risk began a long and substantial decline after 1990; use began to rise gradually after 1993, from 1.5% to 2.7% by 1999, before finally declining slightly in 2000 and then leveling until 2008, when there was a significant decline in use.

Among 8th and 10th graders, crack use rose gradually in the 1990s: from 0.7% in 1991 to 2.1% by 1998 among 8th graders, and from 0.9% in 1992 to 2.5% in 1998 among 10th graders. And, as just discussed, use among 12th graders peaked in 1999 at 2.7% and among young adults at 1.4%. Since those peak years, crack use has declined appreciably—by about half among 8th and 10th graders, and two fifths among 12th graders—yet it has held fairly steady among college students and young adults, at least until 2007, when use among college students finally appeared to decline. In general, the prevalence rates for this drug are relatively low—between 0.5% and 1.6% in all five groups. Twelfth graders have the highest prevalence rate, and within that group annual crack prevalence among the college-bound is considerably lower than among those not bound for college (1.1% for college-bound versus 3.3% for non-college-bound in 2008).

All five populations showed small declines in crack use in 2008, with that at 12th grade being significant.

We believe that the particularly intense and early media coverage of the hazards of crack cocaine likely had the effect of capping an epidemic early by deterring many would-be users and motivating many experimenters to desist use. As has been mentioned, when we first measured crack use in 1987, it had the highest level of perceived risk of any illicit drug. Also, it did not turn out to be “instantly addicting” upon first-time use, as had been widely reported. In recent years, roughly 3% of 12th graders reported trying crack; however, only about 1% reported any use in the prior month, and of these recent users, only about half (i.e., about 0.5% of the total sample) reported using crack more than one or two times in the prior month. It thus appears that, among the small numbers of 12th graders who have ever tried crack, the great majority did not establish a pattern of continued use, let alone develop an addiction.

In 1993 the levels of perceived risk and disapproval associated with crack dropped in all three grade levels, foretelling the rise in use that occurred in all three grades between 1994 and 1998. Because more than a decade had passed since the 1986 media frenzy over crack and its dangers, it is quite possible that generational forgetting of the risks of this drug contributed to the declines in risk and disapproval. Indeed, perceived risk of crack use eroded steadily at all grade levels from 1991 (or 1992 for 12th graders) through 2000. There has not been much systematic change in risk or disapproval of crack since then.

- Use of *cocaine*⁴ in general began to decline a year earlier than crack, probably because crack was still in the process of diffusing to new parts of the country, being still quite new. Between 1986 and 1987 the annual prevalence rate for cocaine dropped dramatically, by about one fifth in all three populations being studied at that time—12th graders, college students, and young adults. The decline occurred when young people finally began to view experimental and occasional use—the type of use in which they are most likely to engage—as more dangerous. This change was probably influenced by the extensive media campaigns that began in the preceding year, but also almost surely by the highly publicized cocaine-related deaths in 1986 of sports stars Len Bias and Don Rogers. By 1992 the annual prevalence of cocaine use had fallen by about two thirds among the three populations for which long-term data are available (12th graders, college students, and young adults).

During the resurgence of illicit drug use in the 1990s, however, cocaine use in all five age populations increased some, both beginning and ending in a staggered pattern by age, consistent with a cohort effect. Use rose among 8th graders from 1991 to 1998, among 10th and 12th graders from 1992 to 1999, among college students from 1994 to 2004, and among young adults from 1996 through 2004. As with crack, all five populations showed some decline in cocaine use in 2008, though none reached statistical significance.

⁴Unless otherwise specified, all references to “cocaine” refer to the use of cocaine in any form, including crack.

The story regarding attitudes and beliefs about cocaine use is informative. Having risen substantially after 1986, the perceived risk of using cocaine actually showed some (nonsignificant) decline in 1992 among 12th graders. In 1993, perceived risk for cocaine powder fell sharply in all grades and disapproval began to decline in all grades, though not as sharply as perceived risk. The decline in perceived risk had virtually ended by 1995 among 8th graders, by 1998 among 10th graders, and by 2001 among 12th graders, suggesting a cohort effect at work in this important belief, which tends to drive use. Disapproval declined between 1991 and 1996 among 8th graders, before leveling (until 2007, when a significant increase was observed followed by some decline in 2008); it also declined from 1992 through 1998 among 10th and 12th graders, with the exception of increases for 12th graders in 1995 and 2008. These changes foretold a subsequent leveling of use at each grade level. Use has since drifted down gradually in the lower grades and more recently among college students and young adults generally.

The perceived availability of cocaine among 12th graders rose steadily from 1983 to 1989, suggesting that availability played *no* role in the substantial downturn in use that occurred after 1986. After 1989, however, perceived availability fell some among 12th graders—which may be explained in part by the greatly reduced proportions of 12th graders who said they have any friends who use, because friendship circles are an important part of the supply system. From 1992 through 1998 or 1999, perceived availability of cocaine powder changed little in the three grades, but after 1998 it declined fairly steadily among 8th graders through 2004 and among 10th and 12th graders through 2003, after which it leveled in 8th grade and eventually in 10th grade followed by a drop in 2008 among both 10th and 12th graders.

As with all the illicit drugs, lifetime cocaine prevalence climbs with age; in 2008 it reached 40% among 50-year-olds. Unlike all of the other illicit drugs, active use of cocaine—i.e., annual or monthly prevalence—holds fairly steady after high school (and, until recent years, its use actually increased after high school) rather than declining (see Figure 4-7 in *Volume II*). Nearly all of the other illicit drugs show a decline in active use with age.

- **PCP** use fell sharply among 12th graders between 1979 and 1982, from an annual prevalence of 7.0% to 2.2%. It reached a low point of 1.2% in 1988, rose some in the 1990s to 2.6% in 1996 during the relapse period in the drug epidemic, and then declined to 1.1% by 2002, with little change thereafter. For young adults, the annual prevalence rate has fluctuated between 0.1% and 0.6%.
- Looking at the long-term trends, we see that the annual prevalence of **heroin** use among 12th graders fell by half between 1975 (1.0%) and 1979 (0.5%), then stabilized for 15 years, through 1994. Heroin use was also stable in the early 1990s among the other four populations covered here. Then, in 1994 for 8th graders and in 1995 for all other groups, use suddenly increased, with rates doubling or tripling in one or two years for 12th graders, college students, and young adults, and then remaining at the new higher levels among all five populations for the rest of the decade. Between 1999 and 2000, however, use significantly decreased among 8th graders (from 1.4% to 1.1%) and significantly

increased among 12th graders (from 1.1% to 1.5%), with the latter change due entirely to an increase in noninjection use. Use of heroin declined significantly among 10th and 12th graders in 2001, as did use of heroin without a needle. In 2002 little change took place among the secondary school students, but young adults showed a significant decline in their reported heroin use. A significant decline in use of heroin overall, as well as use of heroin without a needle, occurred among 10th graders in 2003. In sum, all age groups except for the young adults have annual prevalence rates of heroin use in 2008 that are below recent peaks (by roughly one third to one half in the case of 8th, 10th, and 12th graders, but by less among college students); there is no evidence of any ongoing trends at present. In 2008 the young adults showed a nonsignificant increase back to their peak level of 0.5%, last seen in 2001. Next year's survey may indicate whether the 2008 finding is just a sampling anomaly or an indication of a real trend.

Two factors very likely contributed to the upturn in heroin use in the 1990s. One is a long-term decline in the perceived risk of harm, probably due to generational forgetting, because it had been a long time since the country had experienced a heroin epidemic along with publicity about its accompanying casualties. The second factor, not unrelated to the first, is that in the 1990s the greatly increased purity of heroin allowed it to be used by means other than injection. This may have lowered an important psychological barrier for some potential users, making heroin use less aversive and seemingly less addictive and less risky in general, because avoiding injection reduces the likelihood of transmission of HIV, hepatitis, or other serious blood-borne diseases. By introducing some new questions on heroin use in 1995, we were able to show that significant proportions of past-year users in all five populations were indeed taking heroin by means other than injection (see Table 2-2 and chapter 4 in both *Volume I* and *Volume II* for details).

The risk perceived to be associated with heroin fell for more than a decade after the study began, with 60% of the 1975 twelfth graders seeing a great risk of trying heroin once or twice, and only 46% of the 1986 twelfth graders saying the same. Between 1986 and 1991, perceived risk rose some, from 46% to 55%, undoubtedly reflecting the newly recognized threat of HIV infection associated with heroin injection. After 1991, however, perceived risk fell again (to 51% by 1995), this time perhaps reflecting the fact that the newer heroin available on the street could be administered by methods other than injection. Between 1996 and 1998, perceived risk among 12th graders rose—possibly as the result of an antiheroin campaign launched by the Partnership for a Drug-Free America in June 1996, as well as the visibility of heroin-related deaths of some celebrities in the entertainment and fashion design worlds. The perceived risk of trying heroin decreased among 12th graders in 1999, however, foretelling a significant increase in their use of the drug in 2000. In 2001, as the perceived risk of trying heroin increased slightly, 12th-graders' use declined significantly. In recent years there has been little change in the perceived risk of heroin use, at least until it fell by three percentage points in 2008.

Questions about the degree of risk perceived to be associated with heroin use were introduced into the questionnaires for 8th and 10th graders in 1995. The questions asked

specifically and only about use “without using a needle” because we thought this was the form of heroin use of greatest concern at that point. (Similar questions were asked of 12th graders, as well, in one of the six questionnaire forms.) In general, perceived risk for heroin use without a needle rose some in 12th grade in 1996 and 1997, before leveling, and has held quite steady among 8th and 10th graders since it was first measured.

- The use of *narcotics other than heroin* is reported only for 12th graders and older populations because we believe that younger students are not accurately discriminating among the drugs that should be included or excluded from this general class. Use declined gradually over most of the first half of the study in these groups. Twelfth graders had an annual prevalence rate in 1977 of 6.4%, which fell to 3.3% by 1992. But after about 1992 or 1993, all of the older age groups showed continuing increases for a decade or more, through 2003 or 2004, before stabilizing. An updating of the list of examples given in the question stem in 2002 led to an increase in reported prevalence. After a considerable increase in use from 1992 through 2001 during the relapse phase in the general epidemic and going beyond it, the use of narcotics other than heroin has remained relatively constant, but at the highest levels ever recorded by the study.

The specific drugs in this class are listed in Table E-4 in appendix E of *Volume I*. Among these, *Vicodin*, *codeine*, *OxyContin*, and *Percocet* are commonly mentioned by 12th graders in recent years. They also account for much of the increase in use of the general class, though reported use of other substances in the class have increased as well.

- In 2002, specific questions were added for two drugs in this class—*Vicodin* and *OxyContin*—and the observed prevalence rates suggest that these two drugs very likely help to account for the upturn in use of the general class of narcotics other than heroin. In 2003, Vicodin had attained surprisingly high prevalence rates in the five populations under study here—an annual prevalence of 2.8% in 8th grade, 7.2% in 10th grade, 10.5% in 12th grade, 7.5% among college students, and 8.6% among young adults. In 2008 the rates were similar, at 2.9%, 6.7%, 9.7%, 6.7%, and 9.1% (respectively). Lower annual prevalence rates were found for OxyContin than Vicodin in 2003 across all age groups—1.7%, 3.6%, 4.5%, 2.2%, and 2.6%, respectively—but given that it is a highly addictive narcotic drug, the rates are not inconsequential. In 2008 the respective annual prevalence rates were generally a little higher: 2.1%, 3.6%, 4.7%, 3.6%, and 3.9%. Because OxyContin has received considerable adverse publicity in recent years, it is possible that perceived risk (which we do not measure) will increase. But because its use appears to have originated in several fairly delimited geographic areas, it seems likely that OxyContin was diffusing to new communities for some time, which may have delayed its turnaround overall, as we judge to have happened earlier for crack and ecstasy.
- Annual prevalence of *tranquilizer* use among 12th graders saw a long and substantial decline from 11% in 1977 to 2.8% in 1992. After 1992, use increased significantly among 12th graders (as has been true with most drugs), reaching 7.7% in 2002 (but the question was revised slightly in 2001 to include Xanax as an example of a tranquilizer, so a small portion of the increase may be an artifact). Since then annual prevalence has

leveled or even dropped a bit (6.2% in 2008). Reported tranquilizer use also increased modestly among 8th graders, from 1.8% in 1991 to 3.3% in 1996, before declining to 2.6% in 1998 and leveling since then. As with a number of other drugs, the downturn in use began considerably earlier among 8th graders compared to their older counterparts. Among 10th graders, annual prevalence remained stable between 1991 and 1994 at around 3.3%, and increased significantly to 5.6% in 2000 and 7.3% in 2001 (possibly including some artifact, as noted above). Use declined a bit after 2001, reaching 4.6% in 2008. After a period of stability, college student use showed an increase between 1994 and 2003, more than tripling in that period. For the young adult sample, after a long period of decline, annual prevalence more than doubled between 1997 and 2002, with little change thereafter. Most of the reported tranquilizer use in recent years has involved *Valium* and *Xanax* (see Table E-3 in appendix E of *Volume I*).

- The long-term gradual decline in *sedative (barbiturate)* use among 12th graders, which has been observed since the start of the study in 1975, halted in 1992. (Data are not included here for 8th and 10th graders, again because we believe that these students have more problems with proper classification of the relevant drugs.) Use among 12th graders then rose during the relapse phase in the drug epidemic, from 2.8% in 1992 to 6.7% by 2002—still well below the peak rate of 10.7% in 1975; use has shown only a small improvement since 2002 (5.8% in 2008). The 2008 annual prevalence of this class of drugs is lower among young adults (4.7%) and college students (3.7%) than among 12th graders. Use among college students began to rise a few years later than it did among 12th graders, likely reflecting a cohort effect. Among young adults, sedative (barbiturate) use has increased since the early 1990s, rising from 1.6% in 1992 to 4.4% in 2004. It stands at 4.7% in 2008.
- *Methaqualone*, another sedative drug, has shown a trend pattern quite different from barbiturates. Methaqualone use rose among 12th graders from 1975 to 1981, when annual prevalence reached 7.6%. Its use then fell very sharply, declining to 0.2% by 1993 before rising significantly during the general drug resurgence in the 1990s, to 1.1% by 1996. Prevalence rates have shown little consistent change since then, with use standing at 0.5% in 2008. Use also fell in the 1980s among young adults and college students, who had annual prevalence rates by 1989—the last year they were asked about this drug—of only 0.3% and 0.2%, respectively. In the late 1980s, shrinking availability may well have played a role in the decline, as legal manufacture and distribution of methaqualone ceased. Because of very low usage rates, only 12th graders are now asked about use of this drug.
- Clearly use of most of the several classes of *psychotherapeutic drugs*—sedatives (barbiturates), tranquilizers, and narcotics other than heroin—has become a larger part of the nation’s drug abuse problem. While the rise appears to have halted, most rates remain near recent peak levels. During much of the 1990s and into the 2000s, we were seeing a virtually uninterrupted increase among 12th graders, college students, and young adults in the use of all of these drugs, which had fallen from favor from the mid-1970s through the early 1990s. Use then began rising in the early 1990s and continued to rise after the increase in use of most of the illegal drugs had ended in the late 1990s.

- For many years, five classes of illicitly used drugs—*marijuana*, *amphetamines*, *cocaine*, *LSD*, and *inhalants*—had an impact on appreciable proportions of young Americans in their late teens and 20s. In 2008, twelfth graders showed annual prevalence rates for these drugs of 32.4%, 6.8%, 4.4%, 2.7%, and 3.8% (respectively), reflecting declines in most of them, especially for LSD. Among college students in 2008, the comparable annual prevalence rates are 32.3%, 5.7%, 4.4%, 2.6%, and 1.1%; for all young adults the rates are 28.6%, 5.3%, 6.0%, 1.4%, and 1.4%. Because *LSD* use has fallen so precipitously since 2001 in all five populations, it no longer ranks as one of the major drugs of abuse, whereas *narcotics other than heroin* have become quite important due to the long-term rise in use that began in the 1990s. These narcotics now have annual prevalence rates of 7–9% among 12th graders, college students, and young adults. *Tranquilizers* have also become more important due to a similar rise in use, with prevalence rates in 2008 of about 5–7% across the same three populations, as have *sedatives (barbiturates)*, with rates of 5.8%, 3.7% and 4.7%, respectively. The increase in use of these prescription-type drugs, combined with the decline in use of many illegal drugs, means that the use of prescription-type drugs clearly has become a more important part of the nation’s drug problem.
- *Ecstasy (MDMA)* joined this set of long-established, more prevalent drugs for a period of time. However, annual prevalence rates for ecstasy dropped considerably between 2000 and 2008, from 3.1% to 1.7% for 8th graders, from 5.4% to 2.9% for 10th graders, from 8.2% to 4.3% for 12th graders, from 9.1% to 3.7% among college students, and from 7.2% to 3.3% among young adults.
- In 8th grade, *inhalants* rank second only to marijuana among the illicitly used drugs in terms of annual prevalence, and they actually rank first in lifetime use. Because the use of inhalants reflects a form of illicit psychoactive drug use, and because of its importance among the younger adolescents, an additional index of “any illicit drug use including inhalants” was introduced in Tables 2-1 through 2-3. The inclusion of inhalants makes relatively little difference in the illicit drug index prevalence rates for the older age groups, but considerable difference for the younger ones. For example, in 2008 the proportion of 8th graders reporting any illicit drug use in their lifetime, exclusive of inhalants, was 20%, whereas including inhalants raised the figure to 28%.
- Several drugs have been added to MTF’s coverage in recent years, and they are all discussed in chapter 4 of *Volume I*. These include *ketamine*, *GHB*, and *Rohypnol*, which are so-called “club drugs” (in addition to LSD and ecstasy). In general, these drugs have low prevalence rates that have declined over the past several years among 8th, 10th, and 12th graders: the 2008 annual prevalence rates for *ketamine* are 1.2%, 1.0%, and 1.5%, respectively; for *GHB*, 1.1%, 0.5%, and 1.2%; and for *Rohypnol*, 0.5% and 0.4% for 8th and 10th graders (the Rohypnol question for 12th graders was changed in 2002 and in 2008 stands at 1.3%). There was little change this year in the use of these three drugs, with the exception that GHB showed a further significant decline. The two narcotic drugs added to our coverage in 2002—*OxyContin* and *Vicodin*—show considerably higher prevalence rates, as noted earlier.

- Two new substances used primarily by males to develop physique and physical strength were added to the question set in 2001. One is **androstenedione** (a precursor to anabolic steroids), which could be purchased over the counter until early 2005. Among males, where use is heavily concentrated, the 2008 annual prevalence rates are 1.3%, 1.3%, and 1.7% in grades 8, 10, and 12. (Among females, the rates are 0.5%, 0.6%, and 0.7%.) As discussed in chapter 10 of *Volume I*, the proportion of young males who report past-year use of **androstenedione** and/or **steroids** was appreciable. In 2001, when the “andro” question was introduced, the annual prevalence rate was 8.0% for 12th-grade boys. The rate has fallen considerably in all three grades since then; in 2008 it was 3.2% among 12th-grade boys, reflecting a drop of three fifths.
- Another physique-enhancing substance that is not a drug, but rather a type of protein supplement, is **creatine**. Because we thought its use was often combined with the use of steroids and androstenedione, we included a question on it in 2001 and found prevalence of use to be very high. Among males, who again are the primary users, the 2008 annual prevalence for creatine is 3.2%, 11.5%, and 15.7% in grades 8, 10, and 12. In other words, one in every six 12th-grade boys had used creatine in the prior year. (For girls, the rates are far lower at 0.7%, 0.9%, and 1.5%, respectively.)
- Beginning in 1982, MTF included a set of questions about the use of *nonprescription* stimulants, including stay-awake pills, diet pills, and the so-called “look-alikes” (see chapter 10 of *Volume I* for more detailed findings). The annual prevalence among 12th graders of over-the-counter **stay-awake pills**, which usually contain caffeine as their active ingredient, nearly doubled between 1982 and 1990, increasing from 12% to 23%. After 1990 this statistic fell considerably, reaching 6.3% by 2008, the lowest level ever reported. Use has also declined among the college-age young adult population (ages 19 to 22), from a peak of 26% in 1989 to a low of 7.3% in 2007 (up to 8.7% in 2008). (Data for young adults are not shown.) The **look-alikes** have also shown some falloff in recent years. Among 12th graders, annual prevalence decreased slightly from 6.8% in 1995 to 5.0% in 1999, increased to 7.1% in 2001, and then decreased to 2.8% by 2007, the lowest level ever reported. It stands at 3.1% in 2008. Among young adults ages 19 to 22, use of look-alikes also declined from 6.0% in 1995 to 2.0% in 2008. Among 12th graders, annual prevalence rates for over-the-counter **diet pills** declined from 15% to 10% between 1986 and 1995, increased to 15% by 2002, then declined to 7.2% in 2008. (Among 12th-grade girls in 2008, 15% had tried diet pills by the end of senior year, 10% used them in the past year, and 5% used them in just the past 30 days.) Among young adults ages 19 to 22, annual prevalence rates declined from 17% to 7% between 1986 and 1995, rose back to 17% by 2002, and then declined again to 7% by 2008.
- One additional type of over-the-counter drug was added to the 8th-, 10th-, and 12th-grade questionnaires in 2006—**dextromethorphan**, a cough suppressant found in many cough and cold medications. Respondents were asked, “How often have you taken cough or cold medicines to get high?” The proportions indicating such use in the prior 12 months were 4%, 5%, and 7% in grades 8, 10, and 12 in 2006—not inconsequential proportions. In 2008, the rates were about the same (4%, 5%, and 6% for grades 8, 10, and 12, respectively).

College–Noncollege Differences in Illicit Drug Use

- For analytic purposes, “college students” are defined here as those respondents one to four years past high school who are actively enrolled full-time in a two- or four-year college in March of the year of the survey. For nearly all categories of illicit drugs, college students show *lower* rates of use than their age-mates not in college. For a few categories of drugs—including *any illicit drug*, *marijuana*, and *hallucinogens*—college students also show annual usage rates that are about *average* for all high school graduates their age. (College students are about average on the index of any illicit drug use because they have average rates of marijuana use, which largely drives the index.)
- Although college-bound 12th graders have generally had below-average rates of use on *all* of the illicit drugs while they were in high school, these students’ eventual use of some illicit drugs attained equivalence with, or even exceeded, the rates of those who do not attend college. As MTF results published in two recent books have shown, this college effect of “catching up” is largely explainable in terms of differential rates of leaving the parental home after high school graduation and of getting married. College students are more likely than their age peers to have left the parental home and its constraining influences, and less likely to have entered marriage with its constraining influences.⁵
- In general, the substantial decline in illicit substance use among American college students after 1980 has paralleled that of their age peers not in college. Further, from 1980 until 1992, all young adult high school graduates through age 28, as well as college students taken separately, showed trends that were highly parallel (for the most part) to trends among 12th graders (see chapter 9 of *Volume II*). After 1992, a number of drugs showed an increase in use among 12th graders (as well as 8th and 10th graders), but not among college students and young adults for some period of time.

This divergence, combined with the fact that the upturn began first among 8th graders (in 1992), suggests that cohort effects were emerging for illicit drug use, as discussed earlier. Indeed, as those heavier using cohorts of 12th graders entered the college years, we saw a lagged increase in the use of several drugs in college. For example, annual prevalence reached a low point among 12th graders in 1992 for a number of drugs (e.g., *cocaine*, *amphetamines*, *sedatives [barbiturates]*, *tranquilizers*, *narcotics other than heroin*, and *any illicit drug other than marijuana*) before rising thereafter; among college students, those same drugs reached a low two years later in 1994, and then began to rise gradually. Then, in 1998, as *marijuana* use was declining in secondary school, we saw a sharp increase in use among college students. Consistent with our earlier predictions, the evidence for cohort effects resulting from generational replacement is impressive.

⁵Bachman, J. G., Wadsworth, K. N., O’Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates. See also Bachman, J. G., O’Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002). *The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs*. Mahwah, NJ: Lawrence Erlbaum Associates.

Male–Female Differences in Illicit Drug Use

- Regarding gender differences in the three older populations (12th graders, college students, and young adults), males are more likely to use most *illicit drugs*, and the differences tend to be largest at the higher frequency levels. For example, 2008 *daily marijuana* use rates among 12th graders are 7.6% for males versus 3.0% for females; among all young adults (ages 19 to 28) the rates are 7.3% for males versus 3.6% for females; and among college students the rates are 7.3% for males versus 1.7% for females.
- The 8th- and 10th-grade samples evidence fewer and smaller gender differences in the use of drugs—perhaps because girls tend to date and then emulate older boys, who are in age groups considerably more likely to use drugs. While the rate of prior-year *marijuana* use is slightly higher for males, the rate for the use of *any illicit drug other than marijuana* tends to be slightly higher for females. There is little gender difference in 8th and 10th grades in the use of *cocaine*, *crack*, *other cocaine*, *heroin*, or *Ritalin*. The use of *inhalants*, *amphetamines*, *methamphetamine*, and *tranquilizers* is slightly higher among females.

TRENDS IN ALCOHOL USE

- Several findings about *alcohol* use in these age groups are noteworthy. First, despite the fact that it is illegal for virtually all secondary school students and most college students to purchase alcoholic beverages, their experience with alcohol is widespread. Alcohol has been tried by 39% of current 8th graders, 58% of 10th graders, 72% of 12th graders, and 85% of college students; active use is also widespread. Most important, perhaps, is the prevalence of *occasions of heavy drinking*—five or more drinks in a row at least once in the prior two-week period—which was reported by 8% of 8th graders, 16% of 10th graders, 25% of 12th graders, and 40% of college students surveyed in 2008. Heavy drinking peaks in the early 20s, and recedes with age after that, reflected by the 32% rate found among 29- to 30-year-olds.

Alcohol use did not increase as use of other illicit drugs decreased among 12th graders from the late 1970s to the early 1990s, although it was common to hear such a displacement hypothesis asserted. MTF demonstrates that the opposite seems to be true. After 1980, when illicit drug use was declining, the *monthly prevalence of alcohol use* among 12th graders also declined gradually, but substantially, from 72% in 1980 to 51% in 1992. *Daily alcohol use* declined by half over the same interval, from a peak of 6.9% in 1979 to 3.4% in 1992; the prevalence of drinking *five or more drinks in a row* during the prior two-week interval fell from 41% in 1983 to 28% in 1993—nearly a one-third decline. When illicit drug use rose again in the 1990s, alcohol use (particularly binge drinking) rose some as well—albeit not as sharply as marijuana use. In the late 1990s, as illicit drug use leveled in secondary schools and began a gradual decline, similar trends were observed for alcohol. Therefore, the long-term evidence indicates that alcohol use moves much more in concert with illicit drug use than counter to it.

College–Noncollege Differences in Alcohol Use

- Trends in *alcohol* use among college students are quite different than those for 12th graders or noncollege respondents of the same age (see Figure 9-14 in *Volume II*). From 1980 to 1993, college students showed considerably less drop-off in monthly prevalence of *alcohol* use (82% to 70%) than did 12th graders (72% to 51%), and also less decline in *occasions of heavy drinking* (from 44% to 40%) than either 12th graders (41% to 28%) or their noncollege age-mates (41% to 34%). Because both the noncollege 19- to 22-year-olds and high school students were showing greater declines, the college students stood out as having maintained a high rate of heavy (or binge) drinking. Since 1993, this behavior has changed little among college students—their rate of binge drinking in 2008, 40%, is almost the same as their 1993 rate—while the rate among noncollege age-mates decreased to 30% in 2008 and the 12th graders’ rate, after increasing to 32% in 1998, dropped to 25% by 2006. So, college students continue to stand out as having a relatively high rate of binge drinking.

Although college-bound 12th graders are consistently less likely than their non-college-bound counterparts to report occasions of heavy drinking, the higher rates of such drinking among college students compared to noncollege peers indicate that these 12th graders catch up to and pass their peers in binge drinking after high school graduation. As stated above, we have shown that this differential change after high school is largely attributable to college students’ greater likelihood of leaving the parental home and smaller likelihood of getting married in the four years after graduating from high school. A recent MTF publication also shows that membership in a fraternity or sorority tends to increase heavy episodic drinking and marijuana use.⁶

- Since 1980, college students have generally had *daily drinking* rates that were slightly lower than their age peers, suggesting that they were more likely to confine their drinking to weekends, when they tend to drink a lot. The rate of daily drinking among the noncollege group fell from 8.3% in 1980 to 3.2% in 1994, rose to 5.8% by 2000, and dropped to 3.7% in 2008. Daily drinking by the college group also dropped in approximately the same time period, from 6.5% in 1980 to 3.0% in 1995, then increased to 5.0% in 2002; since then it has remained at 4–5%.

Male–Female Differences in Alcohol Use

- College males report considerably higher rates of *daily drinking* than college females (5.1% versus 3.3% in 2008). This gender difference also exists in the noncollege group (5.0% versus 2.7% in 2008).
- Given that the physiological impacts of five drinks are considerably greater for the typical young female versus the typical young male, it is not surprising that we find substantial gender differences in the prevalence of having *five or more drinks in a row*.

⁶McCabe, S. E., Schulenberg, J. E., Johnston, L. D., O’Malley, P. M., Bachman, J. G., & Kloska, D. D. (2005). Selection and socialization effects of fraternities and sororities on U.S. college student substance use: A multi-cohort national longitudinal study. *Addiction, 100*, 512–524.

Among 12th graders, the rates in 2008 are 21% for females versus 28% for males. This difference has generally been diminishing since the study began; in 1975 there was a 23-percentage-point difference, versus a 7-point difference by 2008.

- Among college students and young adults generally, there are also substantial gender differences in alcohol use, with college males drinking the most. In 2008, for example, half (49%) of all college males reported having *five or more drinks in a row* over the previous two weeks versus one third (34%) of college females. Since MTF began, this gender difference has narrowed gradually, with the rate declining somewhat for males and increasing somewhat for females.

TRENDS IN CIGARETTE SMOKING

A number of very important findings about *cigarette smoking* among American adolescents and young adults have emerged during the life of the study, and we believe that one of the study's more important contributions to the long-term health of the nation has been to document and call public attention to these trends. Despite the demonstrated health risks associated with smoking, young people have continued to establish regular cigarette habits during late adolescence in sizeable proportions, and, during the first half of the 1990s, in growing proportions. In fact, since MTF began in 1975, cigarettes have consistently remained the class of abusable substances most frequently used on a daily basis by high school students.

- During most of the 1980s, when smoking rates were falling steadily among adults, we reported that smoking among adolescents was not declining. Then the situation went from bad to worse. Among 8th and 10th graders, the *current (past 30-day) smoking* rate increased by about half between 1991 (when their use was first measured) and 1996; among 12th graders, the current smoking rate rose by nearly one third between 1992 and 1997. MTF played an important role in bringing these disturbing increases in adolescent smoking to public attention during those years, which was the historical period in which major social action was initiated in the White House, the FDA, the Congress, and eventually the state attorneys general, culminating in the tobacco settlement between the industry and the states.

Fortunately—and largely as a result of that settlement, we believe—there have been some important declines in current smoking since 1996 among 8th and 10th graders, and since 1997 among 12th graders. In fact, the declines have more than offset the increases observed earlier in the 1990s. In 2008, 7% of 8th graders (down from 14% in 1991 and 21% in 1996) reported smoking one or more cigarettes in the prior 30 days—a decline of two thirds from the recent peak rate. Some 12% of 10th graders were current smokers in 2008 (down from 21% in 1991 and 30% in 1996), representing a drop of six tenths from the recent peak rate. And in 2008, 20% of 12th graders were current smokers (versus 28% in 1991 and 37% in 1997), representing a drop of between four and five tenths from the recent peak. In recent years these declines have decelerated though, importantly, they appear not to have ended. Despite these very important improvements in the past decade, one fifth of today's young Americans are current smokers by the time they

complete high school. Other research consistently shows that smoking rates are substantially higher among those who drop out before graduating.⁷

Among college students, the peak rate in current smoking was not reached until 1999 (31%), but after that it declined only moderately (to 24% in 2005) until 2006, when a significant decline brought it down to 19%; with an 18% rate in 2008, this reflects a decline of four tenths from the recent peak so far. Young adults 19 to 28 years old have shown only modest change in rates of current smoking between 2001 (30%) and 2008 (25%)—a decline of about one fifth so far. However, we would expect that, as the cohort effects work their way up the age spectrum, smoking will decrease more in this age group as well.

- The dangers that survey participants perceive to be associated with *pack-a-day smoking* differ greatly by grade level, and seem to be unrealistically low at all grade levels. Currently, about three quarters of 12th graders (74%) report that pack-a-day smokers run a great risk of harming themselves physically or in other ways, but only 60% of the 8th graders say the same. All three grades showed a decrease in perceived risk between 1993 and 1995, as use was rising rapidly, but a slightly larger and offsetting increase in perceived risk occurred between 1995 and 2000, presaging the subsequent downturn in smoking. Between 2000 and 2003, perceived risk stayed relatively level in all grades. In 2004, it increased in all grades, but since then it has risen only modestly in the upper grades, and decreased significantly in 2008 in grade 12.
- Disapproval of cigarette smoking was in decline for a longer period: from 1991 through 1996 among 8th and 10th graders, and from 1992 to 1996 among 12th graders. Since then there has been a fairly steady increase in disapproval of cigarette smoking in all three grades—at least until 2007, when the increase halted among 8th and 12th graders. Undoubtedly the heavy media coverage of the tobacco issue (the settlement with the state attorneys general, the congressional debate, the eventual state settlements, etc.) had an important influence on these attitudes and beliefs. However, that coverage diminished considerably in 1998, raising the question of whether those changes in youth attitudes would continue. It may well be, of course, that the removal of certain kinds of cigarette advertising and promotion, combined with national- and state-level antismoking campaigns and more recent significant increases in cigarette prices, have served to sustain and prolong these changes. In terms of media effects, MTF has shown important changes in reported recall of antismoking ads resulting from both state and national campaigns.⁸

⁷For a recent analysis showing much higher smoking rates among 8th graders who later dropped out before completing high school, see Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education-drug use connection: How successes and failures in school relate to adolescent smoking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

⁸Johnston, L. D., Terry-McElrath, Y. M., O'Malley, P. M., & Wakefield, M. (2005). Trends in recall and appraisal of anti-smoking advertising among American youth: National survey results, 1997–2001. *Prevention Science*, 6, 1–19.

Age- and Cohort-Related Differences in Cigarette Smoking

- Initiation of smoking occurs most often in grades 6 through 9 (i.e., at modal ages 11–12 to 14–15), although according to the 2008 eighth graders, 8% had already initiated smoking before grade 6. The initiation rate trails off considerably by 12th grade, although a number of the light smokers in 12th grade make the transition to heavy smoking in the first two years after high school. Analyses presented in this volume and elsewhere have shown that cigarette smoking evidences a clear cohort effect. That is, if a class (or birth) cohort establishes an unusually high rate of smoking at an early age relative to other cohorts, the rate is likely to remain high throughout the life cycle relative to that of other birth cohorts at equivalent ages.
- As we reported in the “Other Findings from the Study” chapter in the 1986 volume in this series, some 53% of 12th graders who were half-pack-a-day (or more) smokers in senior year in 1985 said that they had tried to quit smoking but could not. Of those who had been daily smokers in 12th grade, nearly three quarters were still daily smokers seven to nine years later (based on the 1985 follow-up survey), despite that in high school only 5% thought they would “definitely” be smoking five years hence. A subsequent analysis, based on the 1995 follow-up survey, showed similar results. Nearly two thirds (63%) of those who had been daily smokers in 12th grade were still daily smokers seven to nine years later, although in high school only 3% of them had thought they would “definitely” be smoking five years hence. Clearly, the smoking habit is established at an early age, is difficult to break for those young people who have initiated use, and young people greatly overestimate their own ability to quit. Additional data from 8th- and 10th-grade students show us that younger adolescents are even more likely than older ones to seriously underestimate the dangers of smoking.
- MTF surveys of 8th and 10th graders also show that cigarettes are readily available to teens in 2008, even though perceived availability has been dropping for some years for these age groups: 57% of 8th graders and 77% of 10th graders say that cigarettes would be “fairly easy” or “very easy” for them to get, if they wanted them. Between 1992 (when these questions were first asked) and 1997, there was little change in reported availability. After that, however, perceived availability of cigarettes decreased significantly for 8th and 10th graders, quite likely reflecting the impact of new regulations and related enforcement efforts aimed at reducing the sale of cigarettes to children (including the Synar amendment, which required states to pass and enforce laws prohibiting the sale and distribution of tobacco products to persons under 18).⁹ (Twelfth graders are not asked this question.) In 2008 the perceived availability of cigarettes increased significantly for 8th graders, while 10th graders showed a significant decrease.

⁹For a more detailed examination of recent changes in youth access to cigarettes, see Johnston, L. D., O’Malley, P. M., & Terry-McElrath, Y. M. (2004). Methods, locations, and ease of cigarette access for American youth, 1997–2002. *American Journal of Preventive Medicine*, 27, 267–276.

College–Noncollege Differences in Cigarette Smoking

- A striking difference in smoking rates has long existed between college-bound and non-college-bound 12th graders. For example, in 2008, smoking a half pack or more per day is about three times as prevalent among the non-college-bound 12th graders as among the college bound (11.9% versus 3.8%). Among respondents of college age (one to four years past high school), those not in college also show dramatically higher rates of half-pack-a-day smoking than those who are in college—15.2% versus 4.3%, respectively. Clearly, these important differences precede college attendance.
- In the first half of the 1990s, smoking rose among college students and their same-age peers, although the increases were not as steep for either group as they were among 12th graders. But in 1998 and 1999, while smoking was declining among secondary school students at all grades, smoking increased significantly for college students, no doubt reflecting the cohort effect from earlier, heavier smoking classes of 12th graders moving into the older age groups. Between 1991 and 1999, the 30-day prevalence of cigarette smoking by college students rose from 23% to 31%, or by about one third, and daily smoking rose from 14% to 19%, also by about one third. The year 2000 showed, for the first time in several years, a decline in college student smoking; that continued with a significant decline to 23% in 2003, and another significant decline to 19% in 2006. The rate in 2008 was 18%. (Because of the smaller numbers of cases in the college student samples, the trend lines are not always as smooth as they are for most of the other groups discussed here.) A much more modest decline has also been observed among their noncollege peers, but only since 2001. A number of in-depth analyses of MTF panel data have revealed that the differences in smoking rates between those who do and do not attend college are evident by the end of 12th grade and have their roots in earlier educational successes and failures.¹⁰

Male–Female Differences in Cigarette Smoking

- In the 1970s, 12th-grade females caught up to and passed 12th-grade males in rates of *current smoking*. Both genders then showed a decline in use followed by a long, fairly level period, with use by females consistently higher, but with the gender difference diminishing. In the early 1990s, another crossover occurred when rates rose more among males than females, and males have been consistently slightly higher in rates of current smoking since 1991 among 12th graders. In the lower grades, the genders have had similar smoking rates since their use was first measured in 1991.
- Among college students, females had a slightly higher probability of being daily smokers from 1980 through 1994—although this long-standing gender difference was not seen among their age peers who were not in college. However, a crossover occurred between

¹⁰Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates. Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002). *The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs*. Mahwah, NJ: Lawrence Erlbaum Associates. Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education–drug use connection: How successes and failures in school relate to adolescent smoking, drinking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

1994 and 2001, with college males exceeding college females in daily smoking—an echo of the crossover among 12th graders in 1991. Since about 2001 there has been little consistent gender difference in smoking among college students.

RACIAL/ETHNIC COMPARISONS

The three largest ethnic groups in the population—Whites, African Americans, and Hispanics—are examined here for 8th, 10th, and 12th graders. (Sample size limitations simply do not allow finer subgroup breakdowns unless data from many years are combined. Separate publications from the study have done just that.) A number of interesting findings emerge from the comparison of these three groups; the reader is referred to chapters 4 and 5 of *Volume I* for a full discussion and to appendix D of *Volume I* for a tabular documentation across all drugs.¹¹ The trends for these three subgroups are also presented graphically in an occasional paper available online.¹²

- African-American 12th graders have consistently shown lower usage rates than White 12th graders for most drugs, both licit and illicit. At the lower grade levels, where few have yet dropped out of school, African-American students also have lower usage rates for many drugs, though not all. The differences are quite large for some drugs, including *inhalants*, *LSD*, *hallucinogens other than LSD*, *ecstasy*, *powder cocaine*, *amphetamines*, *methamphetamine*, *tranquilizers*, *OxyContin*, and *Vicodin*.
- African-American students currently have a much lower 30-day prevalence rate of *cigarette smoking* than White students (10% versus 25% among 12th graders in 2008) because their smoking rate declined from 1983 to 1992, while the rate for White students stabilized for some years. After 1992, smoking rates rose among both White and African-American 12th graders, but by 1998 there was a leveling, and since then a reversal, in both groups in all grades. The White students showed a continuing decline since 2003 in all three grades, while smoking rates among African-American students have stayed about level. In recent years the 8th and 10th grades have shown some decline in smoking rates among African-American students.

¹¹We periodically publish comparisons that contain a number of the smaller racial/ethnic groups in the population, based on data combined for a number of contiguous years in order to attain adequate sample sizes. The first was Bachman, J. G., Wallace, J. M., Jr., O'Malley, P. M., Johnston, L. D., Kurth, C. L., & Neighbors, H. W. (1991). Racial/ethnic differences in smoking, drinking, and illicit drug use among American high school seniors, 1976–1989. *American Journal of Public Health*, *81*, 372–377. More recent articles are: Wallace, J. M., Jr., Bachman J. G., O'Malley, P. M., Johnston, L. D., Schulenberg, J. E., & Cooper, S. M. (2002). Tobacco, alcohol and illicit drug use: Racial and ethnic differences among U.S. high school seniors, 1976–2000. *Public Health Reports*, *117* (Supplement 1), S67–S75; Wallace, J. M., Jr., Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Cooper, S. M., & Johnston, L. D. (2003). Gender and ethnic differences in smoking, drinking, and illicit drug use among American 8th, 10th, and 12th grade students, 1976–2000. *Addictions*, *98*, 225–234; and Delva, J., Wallace, J. M., Jr., O'Malley, P. M., Bachman, J. G., Johnston, L. D., & Schulenberg, J. E. (2005). The epidemiology of alcohol, marijuana, and cocaine use among Mexican American, Puerto Rican, Cuban American, and other Latin American 8th-grade students in the United States: 1991–2002. *American Journal of Public Health*, *95*, 696–702.

¹²Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (2009). *Demographic subgroup trends for various licit and illicit drugs, 1975–2008* (Monitoring the Future Occasional Paper No. 71) [Online]. Ann Arbor, MI: Institute for Social Research. Available: www.monitoringthefuture.org.

- In 12th grade, *occasions of heavy drinking* are much less likely to be reported by African-American students (11%) than White (30%) or Hispanic students (22%).
- In 12th grade, of the three racial/ethnic groups, Whites tend to have the highest rates of use on a number of drugs, including *marijuana*, *hallucinogens*, *LSD* specifically, *hallucinogens other than LSD*, *ecstasy (MDMA)*, *narcotics other than heroin*, *OxyContin* specifically, *Vicodin* specifically, *amphetamines*, *Ritalin* specifically, *sedatives (barbiturates)*, *tranquilizers*, *alcohol*, *getting drunk*, *cigarettes*, and *smokeless tobacco*.
- Hispanics have tended to have the highest usage rate in 12th grade for a number of the most dangerous drugs, such as *heroin* in general and *heroin with a needle* (though Whites are at the same level this year), *crack*, and *crystal methamphetamine (ice)*. Further, in 8th grade, Hispanics have the highest rates for most drugs (though not for amphetamines, Vicodin, OxyContin, or steroids). For example, in 8th grade, the 2008 annual prevalence of *marijuana* use for Hispanics is 13%, versus 10% for Whites and 11% for African Americans; the two-week prevalence of *binge drinking* is 12% for Hispanics, 8% for Whites, and 6% for African Americans. Hispanics have the highest rates of use for many drugs in 8th grade, but not for as many in 12th, which suggests that their considerably higher dropout rate (compared to Whites and African Americans) may change their relative ranking by 12th grade.
- With regard to trends, 12th graders in all three racial/ethnic groups exhibited a decline in *cocaine* use from 1986 through 1992, although the decline was less steep among African-American 12th graders because their earlier increase in use was not as large as the increase among White and Hispanic students.
- For virtually *all of the illicit drugs*, the three groups have tended to trend in parallel. Because White 12th graders had the highest level of use on a number of drugs—including *amphetamines*, *sedatives (barbiturates)*, and *tranquilizers*—they also had the largest declines; African Americans have had the lowest rates and, therefore, the smallest declines.

DRUG USE IN EIGHTH GRADE

It is useful to focus specifically on the youngest age group in the study—the 8th graders, most of whom are 13 or 14 years old—because the worrisome levels of both licit and illicit drug use that they report help illustrate the nation’s urgent need to continue to address the substance abuse problems among its youth.

- Among 8th graders in 2008, 39% report having tried *alcohol* (more than just a few sips), and nearly one in five (18%) say they have already been *drunk* at least once.

- One fifth of 8th graders in 2008 (21%) have tried *cigarettes*, and one in fifteen (6.8%) say they have smoked in the prior month. Shocking to most adults is the fact that only 60% of 8th graders recognize that there is great risk associated with smoking one or more packs of cigarettes per day. While an increasing proportion will recognize the risk by 12th grade, for many this is too late, because they will have developed a smoking habit by then.
- *Smokeless tobacco* has been tried by 14% of male 8th graders in 2008, was used in the past month by 5.5%, and is used daily by 1.2%. (Rates are much lower among females.)
- One 8th grader in six (16%) reported using *inhalants*, and 1 in 24 (4.1%) reported inhalant use in just the past month. This is the only class of drugs for which use is substantially higher in 8th grade than in 10th or 12th grade.
- *Marijuana* has been tried by one in every seven 8th graders (15%) and has been used in the prior month by about 1 in every 17 (5.8%).
- A surprisingly large number of 8th graders (6.8%) say they have tried prescription-type *amphetamines* without medical instruction; 2.2% say they have used them in the prior 30 days.
- For most of the *other illicit drugs*, relatively few 8th graders in 2008 say they have tried them. (This is consistent with the retrospective reports from 12th graders concerning the grades in which they first used the various drugs.) But the proportions having at least some experience with them is not inconsequential. Even a rate as low as 3% represents about one child in every 30-student classroom. The 2008 eighth-grade proportions reporting any lifetime experience with the other illicit drugs are: *tranquilizers* (3.9%), *hallucinogens other than LSD* (2.5%), *cocaine other than crack* and *ecstasy* (both at 2.4%), *methamphetamine* (2.3%), *crack* (2.0%), *LSD* (1.9%), *steroids* (1.4% overall, and 1.8% among males), *heroin* (1.4%), and *Rohypnol* (0.7%).
- In total, 28% of all 8th graders in 2008 have tried some *illicit drug other than marijuana* (including inhalants), while 11%, or about one in nine, have tried *some illicit drug other than marijuana or inhalants*. Put another way, in an average 30-student classroom of 8th graders, about eight have used some illicit drug other than marijuana including inhalants, and three or four have used some drug other than marijuana or inhalants.
- The very large number of 8th graders who have already begun using the so-called “gateway drugs” (*tobacco, alcohol, inhalants, and marijuana*) suggests that a substantial number are also at risk of proceeding further to such drugs as LSD, cocaine, amphetamines, and heroin.

DRUG USE BY AGE 50

Because we have now followed graduating 12th graders into their 50s, we can characterize the drug-using history of today's 50-year-olds (at least those who are high school graduates). This is important, not only because it shows how use by these respondents has developed over the three decades since they left high school, but also because most of them are now themselves the parents of adolescents and young adults. Their own past experiences with drug use may complicate communications with their children regarding drugs; worse, the continuing active use of substances by some of them may set an unfortunate example. The level of lifetime use they have attained is striking (see chapter 4 of *Volume II* for greater detail and discussion).

- Among 50-year-old high school graduates in 2008, we estimate that about three out of four (74%) have tried *marijuana*, and that almost two out of three (63%) have tried an *illicit drug other than marijuana*. (These estimates are adjusted to correct for panel attrition, as described in chapter 4 of *Volume II*.)

Their current behavior is far less extreme than those statistics might suggest, but it is not by any means negligible. One in eight (12%) indicates using marijuana in the last 12 months, while 1 in 10 (10%) affirms use of any other illicit drug in that time period (details provided below). Their past-month prevalence rates are lower—7.2% and 5.9%, respectively, for marijuana and any other illicit drug. About 1 in 50 (2.0%) is a *current daily marijuana* user, though substantially more indicate that they have used marijuana daily at some time in the past.

- Quite high proportions of 50-year-old respondents in 2008 have had some experience during their lifetime with nonmedical use of several of the specific illicit drugs other than marijuana. These include *cocaine* in any form (40%), *amphetamines* (37%), *tranquilizers* (31%), *sedatives (barbiturates)* (26%), *hallucinogens* of any type (22%), and *narcotics other than heroin* (22%). In sum, today's adults in their 50s tend to be a very drug-experienced segment of the population, as might be expected due to the fact that they graduated from high school near the peak of the drug epidemic. To repeat, 74% have tried marijuana and 63% have tried some illicit drug other than marijuana.
- Illicit drugs other than marijuana that have been used in just the prior 12 months by this age group (outside of medical regimen) include *narcotics other than heroin* (3.3%), *tranquilizers* (4.2%), *cocaine* (2.0%), and *noncrack forms of cocaine* (1.7%). Little active use is reported by these respondents for *amphetamines*, *crack*, or *heroin*. (Of course, we would not expect *heavy* heroin or crack users to have remained in the panel studies.)
- *Alcohol* consumption is relatively high among 50-year-olds, with almost two thirds (64%) indicating that they consumed at least one alcoholic drink in the prior 30 days, 11% reporting current *daily drinking* (defined as drinking on 20 or more occasions in the prior 30 days), and 20% indicating *recent heavy drinking* (defined as five or more drinks on at least one occasion in the prior two weeks). The rate of recent heavy drinking is

much lower than was exhibited by members of this cohort when they were of high school and college ages.

- Nearly one in five (19%) 50-year-old high school graduates currently smokes *cigarettes*. Almost all of those are current *daily smokers* (17%).

SUMMARY AND CONCLUSIONS

We can summarize the findings on trends as follows: For more than a decade—from the late 1970s to the early 1990s—the use of a number of *illicit drugs* declined appreciably among 12th-grade students, and declined even more among American college students and young adults. These substantial improvements—which seem largely explainable in terms of changes in attitudes about drug use, beliefs about the risks of drug use, and peer norms against drug use—have some extremely important policy implications. One clear implication is that these various substance-using behaviors among American young people are malleable—they *can* be changed. It has been done before. The second is that demand-side (rather than supply-side) factors appear to have been pivotal in bringing about most of those changes. The levels of marijuana availability, as reported by 12th graders, have held fairly steady throughout the life of the study. (Moreover, both abstainers and quitters rank availability and price very low on their list of reasons for not using.) And, in fact, the perceived availability of cocaine was actually rising during the beginning of the sharp decline in cocaine and crack use in the mid- to late 1980s, which occurred when the perceived risk associated with that drug rose sharply. (See the last section of chapter 9, *Volume I*, for more examples and further discussion of this point.)

However, improvements are surely not inevitable; and when they occur, they should not be taken for granted. Relapse is always possible and, indeed, just such a relapse in the longer term epidemic occurred during the early to mid-1990s, as the country let down its guard on many fronts. (See chapter 8 of *Volume I* for a more detailed discussion.)

In 1992, eighth graders exhibited a significant increase in annual use of *marijuana*, *cocaine*, *LSD*, and *hallucinogens other than LSD*, as well as an increase in *inhalant* use. (In fact, all five populations showed some increase in *LSD* use, continuing a longer term trend for college students and young adults.) Further, the attitudes and beliefs of 12th graders regarding drug use began to soften.

In 1993, use of several drugs began to rise among 10th and 12th graders as well, fulfilling our earlier predictions based on their eroding beliefs about the dangers of drugs and their decreasing disapproval of drug use. Increases occurred in a number of the so-called “gateway drugs”—*marijuana*, *cigarettes*, and *inhalants*—that we argued boded ill for the later use of other drugs in the usual sequence of drug use involvement. Indeed, the proportion of students reporting the use of *any illicit drug other than marijuana* rose steadily after 1991 among 8th and 10th graders and after 1992 among 12th graders. (This proportion increased by more than half among 8th graders, with annual prevalence rising from 8.4% in 1991 to 13.1% in 1996.) The softening attitudes

about *crack* and *other forms of cocaine* also provided a basis for concern—and indeed the use of both increased fairly steadily through 1998.

Over the years, MTF has demonstrated that changes in perceived risk and disapproval have been important causes of change in the use of a number of drugs. These beliefs and attitudes are almost certainly influenced by the amount and nature of public attention paid to the drug issue in the historical period during which young people are growing up. A substantial decline in attention to this issue in the early 1990s very likely explains why the increases in perceived risk and disapproval among students ceased and began to backslide. News coverage of the drug issue plummeted between 1989 and 1993 (although it made a considerable comeback as surveys—including MTF—began to document that the nation’s drug problem was worsening again), and the media’s *pro bono* placement of ads from the Partnership for a Drug-Free America also fell considerably. (During that period, MTF 12th graders showed a steady decline in their recalled exposure to such ads, and in the judged impact of such ads on their own drug-taking behavior.)

Also, the deterioration in the drug abuse situation first began among our youngest cohorts—perhaps because they had not had the same opportunities for vicarious learning from the adverse drug experiences of people around them and people portrayed in the media—those we have called the “unfortunate role models.” Clearly, there was a danger that, as the drug epidemic subsided in the 1980s and early 1990s, newer cohorts would have far less opportunity to learn through informal means about the dangers of drugs—that what we have called a generational forgetting of those risks would occur through a process of generational replacement of older, more drug-savvy cohorts with newer, more naive ones. This suggests that as drug use subsides, as it did by the early 1990s, the nation must redouble its efforts to ensure that such naive cohorts learn these lessons about the dangers of drugs through more formal means—from schools, parents, and focused messages in the media, for example—and that this more formalized prevention effort be *institutionalized* so that it will endure for the long term. Clearly, for the foreseeable future, American young people will be aware of the psychoactive potential of a host of drugs and will continue to have access to them—a situation quite different from the one that preceded the late 1960s. That means that each new generation of young people must learn the reasons that they should *not* use drugs. Otherwise, their natural curiosity and desires for new experiences will lead a great many to use.

Another lesson that derives from the MTF epidemiological data is that social influences that tend to reduce the *initiation* of substance use also have the potential to deter *continuation* by those who have already begun to use, particularly if they are not yet habitual users. Chapter 5 of *Volume I* shows how increased quitting rates have contributed importantly to downturns in the use of a number of drugs at different historical periods. The lesson is that primary prevention should not be the only goal of intervention programs; early-stage users may be persuaded to quit when their beliefs and attitudes regarding drugs are changed.

The following facts help to put into perspective the magnitude and variety of substance use problems that presently remain among American young people:

- More than a quarter (28%) of today's 8th graders have tried an *illicit drug* (if inhalants are included as an illicit drug), and nearly half (49%) of 12th graders have done so.
- By their late 20s, nearly three of five (59%) of today's young adults have tried an *illicit drug*, and more than a third (35%) have tried some *illicit drug other than marijuana* (usually in addition to marijuana). (These figures do not include inhalants.)
- Today, over one in seven young adults (15% in 2008) have tried *cocaine*, and 7% have tried it by their senior year of high school (i.e., by age 17 or 18). More than 1 in every 36 twelfth graders (2.8%) has tried *crack*. In the young adult sample, 1 in 23 (4.4%) has tried crack by age 29–30.
- More than 1 in every 19 twelfth graders (5.4%) in 2008 smokes *marijuana daily*. Among young adults ages 19 to 28, the percentage is about the same (5.1%) and very close to the recent peak level. Among those same 12th graders in 2008, one in every seven (15%) has been a daily marijuana smoker at some time for at least a month, and among young adults the comparable figure is one in six (17%).
- One in four 12th graders (25%) consumed *five or more drinks in a row* at least once in the two weeks prior to the survey, and we know that such behavior tends to increase among young adults one to four years past high school—that is, in the peak college years. Indeed, half (49%) of all male college students report such binge drinking.
- Despite considerable improvements in smoking rates among American adolescents over more than a decade, a fifth (20%) of 12th graders in 2008 were current *cigarette* smokers, and one in nine (11%) were already current *daily* smokers. In addition, we know from studying previous cohorts that many young adults increase their rates of smoking within a year or so after they leave high school.

Despite the substantial improvement in this country's drug situation in the 1980s and early 1990s, and then some further improvement beginning in the late 1990s, American secondary school students and young adults show a level of involvement with *illicit drugs* that is among the highest in the world's industrialized nations.¹³ Even by longer term historical standards in this country, these rates remain extremely high, though in general they are not as high as in the peak years of the epidemic in the late 1970s. *Heavy drinking* also remains widespread and troublesome, and certainly the continuing initiation to *cigarette smoking* of a large, albeit declining, proportion of young people remains a matter of the greatest public health concern. Unfortunately, the declines in youth smoking have decelerated sharply in all grades in recent years, indicating that the improvements in youth smoking overall may be nearing an end.

¹³A published report from an international collaborative study, modeled largely after MTF, provides comparative data from national school surveys of 15- to 16-year-olds that was completed in 2007 in 35 European countries. It also includes 2007 MTF data from 10th graders in the United States. See Hibell, B., Andersson, B., Bjarnasson, T., Ahlström, S., Balakireva, O., Kokkevi, A., & Morgan, M. (Eds.). (2009). *The 2007 ESPAD report (The European School Survey Project on Alcohol and Other Drugs): Substance Use among Students in 35 European countries*. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs, The European Monitoring Centre for Drugs and Drug Addiction, the Council of Europe, and the Co-operation Group to Combat Drug Abuse and Illicit Trafficking in Drugs.

- Of particular note, despite the gradual (and in some cases sharp) declines in the use of many of the illegal drugs in the middle to late 1990s and early 2000s, the prescription-type, abusable drugs have shown very limited declines (with the notable exception of amphetamines). The use of *tranquilizers*, *sedatives (barbiturates)*, and *narcotics other than heroin* are all at or near their recent peak levels. None of them is exhibiting any further increase, however, and some gradual declines have begun.
- Finally, we note the seemingly unending capacity of pharmacological experts and amateurs to discover new substances with abuse potential that can be used to alter mood and consciousness, and of young people to discover the abuse potential of existing products, such as Roflutussin, and to rediscover older drugs, such as *LSD* and *heroin*. While as a society we have made significant progress on a number of fronts in the fight against drug abuse, we must remain vigilant against the opening of new fronts, as well as the reemergence of trouble on older ones. In particular, we must guard against generational forgetting in our newest cohorts of adolescents due to a lack of public attention to the issue during the time that they are growing up.

One of the dynamics that keeps the drug epidemic rolling is the emergence of new drugs whose hazards are little known. In 1999 we saw this happen with the drug *ecstasy (MDMA)*. Other drugs like *Rohypnol*, *ketamine*, *GHB*, and *OxyContin* have appeared in the past decade and have been added to the list of drugs under study. The spread of such new drugs appears to be facilitated and hastened today by young people's widespread use of chat rooms and other sites on the Internet. We predict a continuous flow of such new substances onto the scene, and believe that the task of rapidly documenting their emergence, establishing their adverse consequences, and quickly demystifying them will remain an important means by which policymakers, researchers, and educators deal with the continuing threats posed by such drugs. We also anticipate that there will be rediscoveries of older substances, as has been occurring in recent years with respect to the various psychotherapeutic prescription drugs, including tranquilizers, sedatives (barbiturates), and narcotic drugs.

The drug problem is not an enemy that can be vanquished. It is more a recurring and relapsing problem that must be contained to the greatest extent possible on an ongoing basis. Therefore, it is a problem that requires an ongoing, dynamic response—one that takes into account the continuing generational replacement of our children, the generational forgetting of the dangers of drugs that can occur with that replacement, and the perpetual stream of new abusable substances that will threaten to lure our young people into involvement with drugs.

TABLE 2-1
Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Any Illicit Drug^a																			
8th Grade	18.7	20.6	22.5	25.7	28.5	31.2	29.4	29.0	28.3	26.8	26.8	24.5	22.8	21.5	21.4	20.9	19.0	19.6	+0.6
10th Grade	30.6	29.8	32.8	37.4	40.9	45.4	47.3	44.9	46.2	45.6	45.6	44.6	41.4	39.8	38.2	36.1	35.6	34.1	-1.6
12th Grade	44.1	40.7	42.9	45.6	48.4	50.8	54.3	54.1	54.7	54.0	53.9	53.0	51.1	51.1	50.4	48.2	46.8	47.4	+0.6
College Students	50.4	48.8	45.9	45.5	45.5	47.4	49.0	52.9	53.2	53.7	53.6	51.8	53.9	52.2	52.3	50.6	50.5	49.5	-0.9
Young Adults	62.2	60.2	59.6	57.5	57.4	56.4	56.7	57.0	57.4	58.2	58.1	59.0	60.2	60.5	60.4	59.7	59.8	59.3	-0.4
Any Illicit Drug other than Marijuana^{a,b}																			
8th Grade	14.3	15.6	16.8	17.5	18.8	19.2	17.7	16.9	16.3	15.8†	17.0	13.7	13.6	12.2	12.1	12.2	11.1	11.2	+0.1
10th Grade	19.1	19.2	20.9	21.7	24.3	25.5	25.0	23.6	24.0	23.1†	23.6	22.1	19.7	18.8	18.0	17.5	18.2	15.9	-2.2 s
12th Grade	26.9	25.1	26.7	27.6	28.1	28.5	30.0	29.4	29.4	29.0†	30.7	29.5	27.7	28.7	27.4	26.9	25.5	24.9	-0.6
College Students	25.8	26.1	24.3	22.0	24.5	22.7	24.4	24.8	25.5	25.8†	26.3	26.9	27.6	28.0	26.5	26.3	25.3	22.6	-2.7
Young Adults	37.8	37.0	34.6	33.4	32.8	31.0	30.5	29.9	30.2	31.3†	31.6	32.8	33.9	35.2	34.0	34.8	34.2	34.7	+0.5
Any Illicit Drug including Inhalants^{a,c,d}																			
8th Grade	28.5	29.6	32.3	35.1	38.1	39.4	38.1	37.8	37.2	35.1	34.5	31.6	30.3	30.2	30.0	29.2	27.7	28.3	+0.6
10th Grade	36.1	36.2	38.7	42.7	45.9	49.8	50.9	49.3	49.9	49.3	48.8	47.7	44.9	43.1	42.1	40.1	39.8	38.7	-1.2
12th Grade	47.6	44.4	46.6	49.1	51.5	53.5	56.3	56.1	56.3	57.0	56.0	54.6	52.8	53.0	53.5	51.2	49.1	49.3	+0.3
College Students	52.0	50.3	49.1	47.0	47.0	49.1	50.7	55.4	54.4	54.6	53.1	52.3	54.1	52.9	53.9	53.3	52.5	51.0	-1.5
Young Adults	63.4	61.2	61.2	58.5	59.0	58.2	58.4	58.5	58.5	59.5	59.0	59.6	60.6	62.5	61.4	61.2	61.2	60.2	-1.1
Marijuana/Hashish																			
8th Grade	10.2	11.2	12.6	16.7	19.9	23.1	22.6	22.2	22.0	20.3	20.4	19.2	17.5	16.3	16.5	15.7	14.2	14.6	+0.4
10th Grade	23.4	21.4	24.4	30.4	34.1	39.8	42.3	39.6	40.9	40.3	40.1	38.7	36.4	35.1	34.1	31.8	31.0	29.9	-1.1
12th Grade	36.7	32.6	35.3	38.2	41.7	44.9	49.6	49.1	49.7	48.8	49.0	47.8	46.1	45.7	44.8	42.3	41.8	42.6	+0.8
College Students	46.3	44.1	42.0	42.2	41.7	45.1	46.1	49.9	50.8	51.2	51.0	49.5	50.7	49.1	49.1	46.9	47.5	46.8	-0.7
Young Adults	58.6	56.4	55.9	53.7	53.6	53.4	53.8	54.4	54.6	55.1	55.7	56.8	57.2	57.4	57.0	56.7	56.7	55.9	-0.8
Inhalants^{c,d}																			
8th Grade	17.6	17.4	19.4	19.9	21.6	21.2	21.0	20.5	19.7	17.9	17.1	15.2	15.8	17.3	17.1	16.1	15.6	15.7	+0.2
10th Grade	15.7	16.6	17.5	18.0	19.0	19.3	18.3	18.3	17.0	16.6	15.2	13.5	12.7	12.4	13.1	13.3	13.6	12.8	-0.9
12th Grade	17.6	16.6	17.4	17.7	17.4	16.6	16.1	15.2	15.4	14.2	13.0	11.7	11.2	10.9	11.4	11.1	10.5	9.9	-0.6
College Students	14.4	14.2	14.8	12.0	13.8	11.4	12.4	12.8	12.4	12.9	9.6	7.7	9.7	8.5	7.1	7.4	6.3	4.9	-1.4
Young Adults	13.4	13.5	14.1	13.2	14.5	14.1	14.1	14.2	14.2	14.3	12.8	12.4	12.2	11.6	10.3	10.9	9.1	9.5	+0.3
Nitrites^e																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.6	1.5	1.4	1.7	1.5	1.8	2.0	2.7	1.7	0.8	1.9	1.5	1.6	1.3	1.1	1.2	1.2	0.6	-0.7
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	1.4	1.2	1.3	1.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Hallucinogens^{b,f}																			
8th Grade	3.2	3.8	3.9	4.3	5.2	5.9	5.4	4.9	4.8	4.6†	5.2	4.1	4.0	3.5	3.8	3.4	3.1	3.3	+0.2
10th Grade	6.1	6.4	6.8	8.1	9.3	10.5	10.5	9.8	9.7	8.9†	8.9	7.8	6.9	6.4	5.8	6.1	6.4	5.5	-0.9
12th Grade	9.6	9.2	10.9	11.4	12.7	14.0	15.1	14.1	13.7	13.0†	14.7	12.0	10.6	9.7	8.8	8.3	8.4	8.7	+0.4
College Students	11.3	12.0	11.8	10.0	13.0	12.6	13.8	15.2	14.8	14.4†	14.8	13.6	14.5	12.0	11.0	10.6	9.1	8.5	-0.6
Young Adults	15.7	15.7	15.4	15.4	16.1	16.4	16.8	17.4	18.0	18.4†	18.3	19.6	19.7	19.3	17.6	17.2	16.0	14.8	-1.3

(Table continued on next page.)

TABLE 2-1 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
LSD																			
8th Grade	2.7	3.2	3.5	3.7	4.4	5.1	4.7	4.1	4.1	3.9	3.4	2.5	2.1	1.8	1.9	1.6	1.6	1.9	+0.3
10th Grade	5.6	5.8	6.2	7.2	8.4	9.4	9.5	8.5	8.5	7.6	6.3	5.0	3.5	2.8	2.5	2.7	3.0	2.6	-0.4
12th Grade	8.8	8.6	10.3	10.5	11.7	12.6	13.6	12.6	12.2	11.1	10.9	8.4	5.9	4.6	3.5	3.3	3.4	4.0	+0.5
College Students	9.6	10.6	10.6	9.2	11.5	10.8	11.7	13.1	12.7	11.8	12.2	8.6	8.7	5.6	3.7	3.5	3.3	4.3	+1.0
Young Adults	13.5	13.8	13.6	13.8	14.5	15.0	15.0	15.7	16.2	16.4	16.0	15.1	14.6	13.4	11.2	10.1	9.6	8.1	-1.5 ss
Hallucinogens other than LSD^b																			
8th Grade	1.4	1.7	1.7	2.2	2.5	3.0	2.6	2.5	2.4	2.3‡	3.9	3.3	3.2	3.0	3.3	2.8	2.6	2.5	-0.1
10th Grade	2.2	2.5	2.8	3.8	3.9	4.7	4.8	5.0	4.7	4.8‡	6.6	6.3	5.9	5.8	5.2	5.5	5.7	4.8	-0.9 s
12th Grade	3.7	3.3	3.9	4.9	5.4	6.8	7.5	7.1	6.7	6.9‡	10.4	9.2	9.0	8.7	8.1	7.8	7.7	7.8	+0.1
College Students	6.0	5.7	5.4	4.4	6.5	6.5	7.5	8.7	8.8	8.2‡	10.7	11.0	12.8	10.1	10.6	10.1	8.5	8.2	-0.4
Young Adults	8.4	8.0	7.6	7.4	7.8	7.9	8.5	9.4	9.3	9.9‡	12.0	15.0	16.4	15.6	15.4	14.9	14.1	13.0	-1.1
PCP^g																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	2.9	2.4	2.9	2.8	2.7	4.0	3.9	3.9	3.4	3.4	3.5	3.1	2.5	1.6	2.4	2.2	2.1	1.8	-0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	3.1	2.0	1.9	2.0	2.2	1.9	2.4	2.7	2.3	2.3	3.1	2.5	3.0	2.7	2.0	2.4	2.1	2.2	+0.1
Ecstasy (MDMA)^h																			
8th Grade	—	—	—	—	—	3.4	3.2	2.7	2.7	4.3	5.2	4.3	3.2	2.8	2.8	2.5	2.3	2.4	+0.1
10th Grade	—	—	—	—	—	5.6	5.7	5.1	6.0	7.3	8.0	6.6	5.4	4.3	4.0	4.5	5.2	4.3	-1.0
12th Grade	—	—	—	—	—	6.1	6.9	5.8	8.0	11.0	11.7	10.5	8.3	7.5	5.4	6.5	6.5	6.2	-0.3
College Students	2.0	2.9	2.3	2.1	3.1	4.3	4.7	6.8	8.4	13.1	14.7	12.7	12.9	10.2	8.3	6.9	5.4	6.2	+0.7
Young Adults	3.2	3.9	3.8	3.8	4.5	5.2	5.1	7.2	7.1	11.6	13.0	14.6	15.3	16.0	14.9	14.4	13.1	13.1	-0.1
Cocaine																			
8th Grade	2.3	2.9	2.9	3.6	4.2	4.5	4.4	4.6	4.7	4.5	4.3	3.6	3.6	3.4	3.7	3.4	3.1	3.0	-0.1
10th Grade	4.1	3.3	3.6	4.3	5.0	6.5	7.1	7.2	7.7	6.9	5.7	6.1	5.1	5.4	5.2	4.8	5.3	4.5	-0.8
12th Grade	7.8	6.1	6.1	5.9	6.0	7.1	8.7	9.3	9.8	8.6	8.2	7.8	7.7	8.1	8.0	8.5	7.8	7.2	-0.6
College Students	9.4	7.9	6.3	5.0	5.5	5.0	5.6	8.1	8.4	9.1	8.6	8.2	9.2	9.5	8.8	7.7	8.5	7.2	-1.3
Young Adults	21.0	19.5	16.9	15.2	13.7	12.9	12.1	12.3	12.8	12.7	13.1	13.5	14.7	15.2	14.3	15.2	14.7	14.8	0.0
Crackⁱ																			
8th Grade	1.3	1.6	1.7	2.4	2.7	2.9	2.7	3.2	3.1	3.1	3.0	2.5	2.5	2.4	2.4	2.3	2.1	2.0	-0.1
10th Grade	1.7	1.5	1.8	2.1	2.8	3.3	3.6	3.9	4.0	3.7	3.1	3.6	2.7	2.6	2.5	2.2	2.3	2.0	-0.3
12th Grade	3.1	2.6	2.6	3.0	3.0	3.3	3.9	4.4	4.6	3.9	3.7	3.8	3.6	3.9	3.5	3.5	3.2	2.8	-0.5
College Students	1.5	1.7	1.3	1.0	1.8	1.2	1.4	2.2	2.4	2.5	2.0	1.9	3.1	2.0	1.7	2.3	1.3	1.4	+0.1
Young Adults	4.8	5.1	4.3	4.4	3.8	3.9	3.6	3.8	4.3	4.6	4.7	4.3	4.7	4.2	4.1	4.4	3.9	4.3	+0.5
Other Cocaine^j																			
8th Grade	2.0	2.4	2.4	3.0	3.4	3.8	3.5	3.7	3.8	3.5	3.3	2.8	2.7	2.6	2.9	2.7	2.6	2.4	-0.2
10th Grade	3.8	3.0	3.3	3.8	4.4	5.5	6.1	6.4	6.8	6.0	5.0	5.2	4.5	4.8	4.6	4.3	4.8	4.0	-0.8
12th Grade	7.0	5.3	5.4	5.2	5.1	6.4	8.2	8.4	8.8	7.7	7.4	7.0	6.7	7.3	7.1	7.9	6.8	6.5	-0.4
College Students	9.0	7.6	6.3	4.6	5.2	4.6	5.0	7.4	7.8	8.1	8.3	8.6	8.5	9.3	8.1	6.2	8.0	7.1	-0.8
Young Adults	19.8	18.4	15.1	13.9	12.4	11.9	11.3	11.5	11.8	11.7	12.1	12.8	13.5	14.4	13.3	14.4	14.0	13.9	-0.1

(Table continued on next page.)

TABLE 2-1 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Heroin^k																			
8th Grade	1.2	1.4	1.4	2.0	2.3	2.4	2.1	2.3	2.3	1.9	1.7	1.6	1.6	1.6	1.5	1.4	1.3	1.4	0.0
10th Grade	1.2	1.2	1.3	1.5	1.7	2.1	2.1	2.3	2.3	2.2	1.7	1.8	1.5	1.5	1.5	1.4	1.5	1.2	-0.4 s
12th Grade	0.9	1.2	1.1	1.2	1.6	1.8	2.1	2.0	2.0	2.4	1.8	1.7	1.5	1.5	1.5	1.4	1.5	1.3	-0.3
College Students	0.5	0.5	0.6	0.1	0.6	0.7	0.9	1.7	0.9	1.7	1.2	1.0	1.0	0.9	0.5	0.7	0.5	0.7	+0.2
Young Adults	0.9	0.9	0.9	0.8	1.1	1.3	1.3	1.6	1.7	1.8	2.0	1.8	1.9	1.9	1.7	1.9	1.6	1.9	+0.3
With a Needle^l																			
8th Grade	—	—	—	—	1.5	1.6	1.3	1.4	1.6	1.1	1.2	1.0	1.0	1.1	1.0	1.0	0.9	0.9	0.0
10th Grade	—	—	—	—	1.0	1.1	1.1	1.2	1.3	1.0	0.8	1.0	0.9	0.8	0.8	0.9	0.9	0.7	-0.2
12th Grade	—	—	—	—	0.7	0.8	0.9	0.8	0.9	0.8	0.7	0.8	0.7	0.7	0.9	0.8	0.7	0.7	0.0
College Students	—	—	—	—	0.4	0.1	0.2	0.5	0.8	0.7	0.2	0.3	0.1	0.1	0.3	0.3	0.1	0.0	-0.1
Young Adults	—	—	—	—	0.4	0.4	0.3	0.4	0.6	0.4	0.6	0.4	0.5	0.4	0.6	0.6	0.5	0.5	0.0
Without a Needle^l																			
8th Grade	—	—	—	—	1.5	1.6	1.4	1.5	1.4	1.3	1.1	1.0	1.1	1.0	0.9	0.9	0.7	0.9	+0.2
10th Grade	—	—	—	—	1.1	1.7	1.7	1.7	1.6	1.7	1.3	1.3	1.0	1.1	1.1	1.0	1.1	0.8	-0.2
12th Grade	—	—	—	—	1.4	1.7	2.1	1.6	1.8	2.4	1.5	1.6	1.8	1.4	1.3	1.1	1.4	1.1	-0.3
College Students	—	—	—	—	0.5	1.0	1.2	2.1	1.0	2.5	1.3	1.2	1.1	1.0	0.3	0.8	0.4	0.7	+0.2
Young Adults	—	—	—	—	0.9	1.3	1.5	1.7	1.9	2.1	2.1	1.8	2.2	2.1	1.8	2.4	1.9	2.1	+0.2
Narcotics other than Heroin^{m,n}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	6.6	6.1	6.4	6.6	7.2	8.2	9.7	9.8	10.2	10.6	9.9‡	13.5	13.2	13.5	12.8	13.4	13.1	13.2	+0.1
College Students	7.3	7.3	6.2	5.1	7.2	5.7	8.2	8.7	8.7	8.9	11.0‡	12.2	14.2	13.8	14.4	14.6	14.1	12.4	-1.7
Young Adults	9.3	8.9	8.1	8.2	9.0	8.3	9.2	9.1	9.5	10.0	11.5‡	13.9	16.8	17.6	17.8	18.7	18.8	19.5	+0.6
Amphetamines^m																			
8th Grade	10.5	10.8	11.8	12.3	13.1	13.5	12.3	11.3	10.7	9.9	10.2	8.7	8.4	7.5	7.4	7.3	6.5	6.8	+0.3
10th Grade	13.2	13.1	14.9	15.1	17.4	17.7	17.0	16.0	15.7	15.7	16.0	14.9	13.1	11.9	11.1	11.2	11.1	9.0	-2.1 ss
12th Grade	15.4	13.9	15.1	15.7	15.3	15.3	16.5	16.4	16.3	15.6	16.2	16.8	14.4	15.0	13.1	12.4	11.4	10.5	-0.9
College Students	13.0	10.5	10.1	9.2	10.7	9.5	10.6	10.6	11.9	12.3	12.4	11.9	12.3	12.7	12.3	10.7	11.2	9.1	-2.1
Young Adults	22.4	20.2	18.7	17.1	16.6	15.3	14.6	14.3	14.1	15.0	15.0	14.8	15.2	15.9	14.6	15.6	15.3	14.6	-0.7
Methamphetamine^{o,p}																			
8th Grade	—	—	—	—	—	—	—	—	4.5	4.2	4.4	3.5	3.9	2.5	3.1	2.7	1.8	2.3	+0.5
10th Grade	—	—	—	—	—	—	—	—	7.3	6.9	6.4	6.1	5.2	5.3	4.1	3.2	2.8	2.4	-0.4
12th Grade	—	—	—	—	—	—	—	—	8.2	7.9	6.9	6.7	6.2	6.2	4.5	4.4	3.0	2.8	-0.3
College Students	—	—	—	—	—	—	—	—	7.1	5.1	5.3	5.0	5.8	5.2	4.1	2.9	1.9	1.9	0.0
Young Adults	—	—	—	—	—	—	—	—	8.8	9.3	9.0	9.1	8.9	9.0	8.3	7.3	6.7	6.3	-0.4
Crystal Meth. (Ice)^p																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.3	2.9	3.1	3.4	3.9	4.4	4.4	5.3	4.8	4.0	4.1	4.7	3.9	4.0	4.0	3.4	3.4	2.8	-0.6
College Students	1.3	0.6	1.6	1.3	1.0	0.8	1.6	2.2	2.8	1.3	2.3	2.0	2.9	2.2	2.4	1.7	1.3	1.1	-0.2
Young Adults	2.9	2.2	2.7	2.5	2.1	3.1	2.5	3.4	3.3	3.9	4.0	4.1	4.7	4.7	4.4	4.7	3.7	3.6	-0.1

(Table continued on next page.)

TABLE 2-1 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Sedatives																			
(Barbiturates)^m																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	6.2	5.5	6.3	7.0	7.4	7.6	8.1	8.7	8.9	9.2	8.7	9.5	8.8	9.9	10.5	10.2	9.3	8.5	-0.9
College Students	3.5	3.8	3.5	3.2	4.0	4.6	5.2	5.7	6.7	6.9	6.0	5.9	5.7	7.2	8.5	6.3	5.9	6.4	+0.5
Young Adults	8.2	7.4	6.5	6.4	6.7	6.6	6.5	6.9	7.4	8.1	7.8	8.0	8.7	9.7	10.0	9.5	9.8	10.6	+0.8
Methaqualone^{m,q}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.3	1.6	0.8	1.4	1.2	2.0	1.7	1.6	1.8	0.8	1.1	1.5	1.0	1.3	1.3	1.2	1.0	0.8	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers^{b,m}																			
8th Grade	3.8	4.1	4.4	4.6	4.5	5.3	4.8	4.6	4.4	4.4†	5.0	4.3	4.4	4.0	4.1	4.3	3.9	3.9	0.0
10th Grade	5.8	5.9	5.7	5.4	6.0	7.1	7.3	7.8	7.9	8.0‡	9.2	8.8	7.8	7.3	7.1	7.2	7.4	6.8	-0.6
12th Grade	7.2	6.0	6.4	6.6	7.1	7.2	7.8	8.5	9.3	8.9‡	10.3	11.4	10.2	10.6	9.9	10.3	9.5	8.9	-0.6
College Students	6.8	6.9	6.3	4.4	5.4	5.3	6.9	7.7	8.2	8.8‡	9.7	10.7	11.0	10.6	11.9	10.0	9.1	8.6	-0.4
Young Adults	11.8	11.3	10.5	9.9	9.7	9.3	8.6	9.6	9.6	10.5‡	11.9	13.4	13.8	14.9	14.5	15.0	14.5	15.8	+1.3
Rohypnol^f																			
8th Grade	—	—	—	—	—	1.5	1.1	1.4	1.3	1.0	1.1	0.8	1.0	1.0	1.1	1.0	1.0	0.7	-0.3
10th Grade	—	—	—	—	—	1.5	1.7	2.0	1.8	1.3	1.5	1.3	1.0	1.2	1.0	0.8	1.3	0.9	-0.4
12th Grade	—	—	—	—	—	1.2	1.8	3.0	2.0	1.5	1.7	—	—	—	—	—	—	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Alcohol^s																			
Any Use																			
8th Grade	70.1	69.3‡	55.7	55.8	54.5	55.3	53.8	52.5	52.1	51.7	50.5	47.0	45.6	43.9	41.0	40.5	38.9	38.9	+0.1
10th Grade	83.8	82.3‡	71.6	71.1	70.5	71.8	72.0	69.8	70.6	71.4	70.1	66.9	66.0	64.2	63.2	61.5	61.7	58.3	-3.4 ss
12th Grade	88.0	87.5‡	80.0	80.4	80.7	79.2	81.7	81.4	80.0	80.3	79.7	78.4	76.6	76.8	75.1	72.7	72.2	71.9	-0.2
College Students	93.6	91.8	89.3	88.2	88.5	88.4	87.3	88.5	88.0	86.6	86.1	86.0	86.2	84.6	86.6	84.7	83.1	85.3	+2.1
Young Adults	94.1	93.4	92.1	91.2	91.6	91.2	90.7	90.6	90.2	90.7	89.9	90.2	89.3	89.4	89.1	88.9	87.9	88.4	+0.5
Been Drunk^t																			
8th Grade	26.7	26.8	26.4	25.9	25.3	26.8	25.2	24.8	24.8	25.1	23.4	21.3	20.3	19.9	19.5	19.5	17.9	18.0	+0.1
10th Grade	50.0	47.7	47.9	47.2	46.9	48.5	49.4	46.7	48.9	49.3	48.2	44.0	42.4	42.3	42.1	41.4	41.2	37.2	-4.0 sss
12th Grade	65.4	63.4	62.5	62.9	63.2	61.8	64.2	62.4	62.3	62.3	63.9	61.6	58.1	60.3	57.5	56.4	55.1	54.7	-0.4
College Students	79.6	76.8	76.4	74.4	76.6	76.2	77.0	76.8	75.1	74.7	76.1	75.1	74.9	73.4	72.9	73.1	71.6	72.5	+0.9
Young Adults	82.9	81.1	81.4	80.7	82.1	80.7	81.4	79.8	81.6	80.4	81.1	81.2	80.9	80.1	79.9	80.9	80.1	80.1	0.0
Flavored Alcoholic Beverages^{g,o}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	37.9	35.5	35.5	34.0	32.8	-1.3
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	58.6	58.8	58.1	55.7	53.5	-2.3
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	71.0	73.6	69.9	68.4	65.5	-2.9
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	79.0	84.5	80.9	80.6	78.6	-2.0
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	83.2	84.6	84.4	84.0	82.6	-1.4

(Table continued on next page.)

TABLE 2-1 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Cigarettes																			
Any Use																			
8th Grade	44.0	45.2	45.3	46.1	46.4	49.2	47.3	45.7	44.1	40.5	36.6	31.4	28.4	27.9	25.9	24.6	22.1	20.5	-1.7
10th Grade	55.1	53.5	56.3	56.9	57.6	61.2	60.2	57.7	57.6	55.1	52.8	47.4	43.0	40.7	38.9	36.1	34.6	31.7	-2.9 ss
12th Grade	63.1	61.8	61.9	62.0	64.2	63.5	65.4	65.3	64.6	62.5	61.0	57.2	53.7	52.8	50.0	47.1	46.2	44.7	-1.5
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Smokeless Tobacco^u																			
8th Grade	22.2	20.7	18.7	19.9	20.0	20.4	16.8	15.0	14.4	12.8	11.7	11.2	11.3	11.0	10.1	10.2	9.1	9.8	+0.8
10th Grade	28.2	26.6	28.1	29.2	27.6	27.4	26.3	22.7	20.4	19.1	19.5	16.9	14.6	13.8	14.5	15.0	15.1	12.2	-2.9 ss
12th Grade	—	32.4	31.0	30.7	30.9	29.8	25.3	26.2	23.4	23.1	19.7	18.3	17.0	16.7	17.5	15.2	15.1	15.6	+0.5
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids^{v,w}																			
8th Grade	1.9	1.7	1.6	2.0	2.0	1.8	1.8	2.3	2.7	3.0	2.8	2.5	2.5	1.9	1.7	1.6	1.5	1.4	-0.2
10th Grade	1.8	1.7	1.7	1.8	2.0	1.8	2.0	2.0	2.7	3.5	3.5	3.5	3.0	2.4	2.0	1.8	1.8	1.4	-0.4
12th Grade	2.1	2.1	2.0	2.4	2.3	1.9	2.4	2.7	2.9	2.5	3.7	4.0	3.5	3.4	2.6	2.7	2.2	2.2	0.0
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	1.7	1.9	1.5	1.3	1.5	1.5	1.4	1.4	1.9	1.4	1.4	1.6	1.8	1.9	1.8	1.8	1.7	1.8	+0.1

Source. The Monitoring the Future study, the University of Michigan.

Footnotes for Tables 2-1 through 2-4

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$.

“—” indicates data not available.

“**” indicates less than 0.05% but greater than 0%.

“‡” indicates some change in the question. See relevant footnote for that drug. See relevant figure to assess the impact of the wording changes.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

Approximate Weighted Ns	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
8th Graders	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	16,500	16,100	15,700
10th Graders	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	16,200	16,100	15,100
12th Graders	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200	14,500	14,000
College																		
Students	1,410	1,490	1,490	1,410	1,450	1,450	1,480	1,440	1,440	1,350	1,340	1,260	1,270	1,400	1,360	1,280	1,250	1,270
Young Adults	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900

^aFor 12th graders, college students, and young adults only: Use of “any illicit drug” includes any use of marijuana, LSD, other hallucinogens, crack, other cocaine, or heroin; or any use of narcotics other than heroin, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor’s orders. For 8th and 10th graders only: The use of narcotics other than heroin and sedatives (barbiturates) has been excluded because these younger respondents appear to overreport use (perhaps because they include the use of nonprescription drugs in their answers).

^bIn 2001 the question text was changed on half of the questionnaire forms for each age group. “Other psychedelics” was changed to “other hallucinogens” and “shrooms” was added to the list of examples. For the tranquilizer list of examples, Miltown was replaced with Xanax. For 8th, 10th, and 12th graders only: The 2001 data presented here are based on the changed forms only; N is one half of N indicated. In 2002 the remaining forms were changed to the new wording. The data are based on all forms beginning in 2002. Data for any illicit drug other than marijuana and data for hallucinogens are also affected by these changes and have been handled in a parallel manner.

^cFor 12th graders, college students, and young adults only: Data based on five of six forms in 1991–1998; N is five sixths of N indicated. Data based on three of six forms beginning in 1999; N is three sixths of N indicated.

^dInhalants are unadjusted for underreporting of amyl and butyl nitrites.

^eFor 12th graders only: Data based on one of six forms; N is one sixth of N indicated. For young adults only: Data based on one of six forms; N is one sixth of N indicated. Questions about nitrite use were dropped from the young adult questionnaires in 1995.

^fHallucinogens are unadjusted for underreporting of PCP.

^gFor 12th graders, college students, and young adults only: Data based on one of six forms; N is one sixth of N indicated.

^hFor 8th and 10th graders only: Data based on one of two forms in 1996; N is one half of N indicated. Data based on one third of N indicated in 1997–2001 due to changes in the questionnaire forms. Data based on two of four forms beginning in 2002; N is one half of N indicated. For 12th graders only: Data based on one of six forms in 1996–2001; N is one sixth of N indicated. Data based on two of six forms beginning in 2002; N is two sixths of N indicated. For college students and young adults only: Data based on two of six forms in 1991–2001; N is two sixths of N indicated. Data based on three of six forms beginning in 2002; N is three sixths of N indicated.

ⁱFor college students and young adults only: Data based on five of six forms beginning in 2002; N is five sixths of N indicated.

^jFor 12th graders only: Data based on four of six forms; N is four sixths of N indicated. For college students and young adults only: Data based on four of six forms; N is four sixths of N indicated.

^kIn 1995, the heroin question was changed in one of two forms for 8th and 10th graders, in three of six forms for 12th graders, and in two of six forms for college students and young adults. Separate questions were asked for use with and without injection. In 1996, the heroin question was changed in all remaining 8th- and 10th-grade forms. Data presented here represent the combined data from all forms.

^lFor 8th and 10th graders only: Data based on one of two forms in 1995; N is one half of N indicated. Data based on all forms beginning in 1996. For 12th graders only: Data based on three of six forms; N is three sixths of N indicated. For college students and young adults only: Data based on two of six forms; N is two sixths of N indicated.

^mOnly drug use not under a doctor’s orders is included here.

ⁿIn 2002 the question text was changed in half of the questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced with Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only; N is one half of N indicated. In 2003, the remaining forms were changed to the new wording. The data are based on all forms beginning in 2003.

^oFor 8th and 10th graders only: Data based on one of four forms; N is one third of N indicated.

^pFor 12th graders, college students, and young adults only: Data based on two of six forms; N is two sixths of N indicated.

^qFor 12th graders only: Data based on one of six forms; N is one sixth of N indicated.

^rFor 8th and 10th graders only: Data based on one of two forms in 1996; N is one half of N indicated. Data based on three of four forms in 1997–1998; N is two thirds of N indicated. Data based on two of four forms in 1999–2001; N is one third of N indicated. Data based on one of four forms beginning in 2002; N is one sixth of N indicated.

For 12th graders only: Data based on one of six forms in 1996–2001; N is one sixth of N indicated. Data based on two of six forms beginning in 2002; N is two sixths of N indicated. Data for 2001 and 2002 are not comparable due to changes in the questionnaire forms. For college students and young adults only: Data based on two of six forms; N is two sixths of N indicated.

^sFor 8th, 10th, and 12th graders only: In 1993, the question text was changed slightly in half of the forms to indicate that a “drink” meant “more than just a few sips.” The 1993 data are based on the changed forms only; N is one half of N indicated for these groups. In 1994 the remaining forms were changed to the new wording. The data are based on all forms beginning in 1994. In 2004, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. The remaining forms were changed in 2005. For college students and young adults: The revision of the question text resulted in rather little change in the reported prevalence of use. The data for all forms are used to provide the most reliable estimate of change.

^tFor 12th graders only: Data based on two of six forms; N is two sixths of N indicated. For college students and young adults only: Data based on three of six forms; N is three sixths of N indicated.

^uFor 8th and 10th graders only: Data based on one of two forms for 1991–1996 and on two of four forms beginning in 1997; N is one half of N indicated. For 12th graders only: Data based on one of six forms; N is one sixth of N indicated. For college students and young adults only: Questions about smokeless tobacco use were dropped from the analyses in 1989.

^vFor 8th, 10th, and 12th graders only: In 2006, the question text was changed slightly in some of the forms. An examination of the data did not show any effect from the wording change. For 12th graders only: Data based on two of six forms in 1991–2005; *N* is two sixths of *N* indicated. Data based on three of six forms in 2006; *N* is three sixths of *N* indicated.

^wFor college students and young adults only: Data based on two of six forms; *N* is two sixths of *N* indicated. In 2008, the question text was changed slightly.

^xFor 12th graders only: Data based on two of six forms in 2002–2005; *N* is two sixths of *N* indicated. Data based on three of six forms beginning in 2006; *N* is three sixths of *N* indicated.

^yFor 12th graders only: Data based on two of six forms in 2000; *N* is two sixths of *N* indicated. Data based on three of six forms in 2001; *N* is three sixths of *N* indicated. Data based on one of six forms beginning in 2002; *N* is one sixth of *N* indicated. For college students and young adults only: Data based on two of six forms; *N* is two sixths of *N* indicated.

^zFor 12th graders only: Data based on two of six forms in 2000; *N* is two sixths of *N* indicated. Data based on three of six forms beginning in 2001; *N* is three sixths of *N* indicated. For college students and young adults only: Data based on two of six forms; *N* is two sixths of *N* indicated.

^{aa}For 12th graders only: The 2003 flavored alcoholic beverage data were created by adjusting the 2004 data to reflect the observed 2003 to 2004 change in a slightly different version of the flavored alcoholic beverage question. In 2004 the original question was revised to include wine coolers among the examples—a change that had very little effect on the observed prevalence-of-use rate.

^{bb}Daily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes and smokeless tobacco, for which actual daily use is measured, and for 5+ drinks, for which the prevalence of having five or more drinks in a row in the last two weeks is measured.

^{cc}For 12th graders only: Due to a coding error, previously released versions of this table contained values that were slightly off for the measure of five or more drinks in a row for 2005 and 2006. These have been corrected here. For 8th and 10th graders only: The 1991–2007 estimates for five or more drinks in a row differ slightly from some previous reports due to an error in the data editing process prior to 2008. The revised estimates average about 2 percentage points lower than the estimates previously reported. Those previous overestimates have been corrected in this table.

TABLE 2-2
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Any Illicit Drug^a																			
8th Grade	11.3	12.9	15.1	18.5	21.4	23.6	22.1	21.0	20.5	19.5	19.5	17.7	16.1	15.2	15.5	14.8	13.2	14.1	+0.8
10th Grade	21.4	20.4	24.7	30.0	33.3	37.5	38.5	35.0	35.9	36.4	37.2	34.8	32.0	31.1	29.8	28.7	28.1	26.9	-1.2
12th Grade	29.4	27.1	31.0	35.8	39.0	40.2	42.4	41.4	42.1	40.9	41.4	41.0	39.3	38.8	38.4	36.5	35.9	36.6	+0.6
College Students	29.2	30.6	30.6	31.4	33.5	34.2	34.1	37.8	36.9	36.1	37.9	37.0	36.5	36.2	36.6	33.9	35.0	35.2	+0.2
Young Adults	27.0	28.3	28.4	28.4	29.8	29.2	29.2	29.9	30.3	30.8	32.1	32.4	33.0	33.7	32.8	32.1	32.5	33.8	+1.2
Any Illicit Drug other than Marijuana^{a,b}																			
8th Grade	8.4	9.3	10.4	11.3	12.6	13.1	11.8	11.0	10.5	10.2†	10.8	8.8	8.8	7.9	8.1	7.7	7.0	7.4	+0.4
10th Grade	12.2	12.3	13.9	15.2	17.5	18.4	18.2	16.6	16.7	16.7†	17.9	15.7	13.8	13.5	12.9	12.7	13.1	11.3	-1.7 s
12th Grade	16.2	14.9	17.1	18.0	19.4	19.8	20.7	20.2	20.7	20.4†	21.6	20.9	19.8	20.5	19.7	19.2	18.5	18.3	-0.3
College Students	13.2	13.1	12.5	12.2	15.9	12.8	15.8	14.0	15.4	15.6†	16.4	16.6	17.9	18.6	18.5	18.1	17.3	15.3	-2.0
Young Adults	14.3	14.1	13.0	13.0	13.8	13.2	13.6	13.2	13.7	14.9†	15.4	16.3	18.1	18.8	18.5	18.4	18.1	18.9	+0.8
Any Illicit Drug including Inhalants^{a,c,d}																			
8th Grade	16.7	18.2	21.1	24.2	27.1	28.7	27.2	26.2	25.3	24.0	23.9	21.4	20.4	20.2	20.4	19.7	18.0	19.0	+1.0
10th Grade	23.9	23.5	27.4	32.5	35.6	39.6	40.3	37.1	37.7	38.0	38.7	36.1	33.5	32.9	31.7	30.7	30.2	28.8	-1.3
12th Grade	31.2	28.8	32.5	37.6	40.2	41.9	43.3	42.4	42.8	42.5	42.6	42.1	40.5	39.1	40.3	38.0	37.0	37.3	+0.3
College Students	29.8	31.1	31.7	31.9	33.7	35.1	35.5	39.1	37.4	37.0	38.2	37.7	36.0	35.9	37.9	35.5	36.8	35.7	-1.1
Young Adults	27.8	29.2	28.9	29.2	30.4	30.2	30.1	30.6	30.6	31.2	33.2	32.4	32.7	34.9	32.8	32.6	33.2	33.5	+0.3
Marijuana/Hashish																			
8th Grade	6.2	7.2	9.2	13.0	15.8	18.3	17.7	16.9	16.5	15.6	15.4	14.6	12.8	11.8	12.2	11.7	10.3	10.9	+0.6
10th Grade	16.5	15.2	19.2	25.2	28.7	33.6	34.8	31.1	32.1	32.2	32.7	30.3	28.2	27.5	26.6	25.2	24.6	23.9	-0.6
12th Grade	23.9	21.9	26.0	30.7	34.7	35.8	38.5	37.5	37.8	36.5	37.0	36.2	34.9	34.3	33.6	31.5	31.7	32.4	+0.7
College Students	26.5	27.7	27.9	29.3	31.2	33.1	31.6	35.9	35.2	34.0	35.6	34.7	33.7	33.3	33.3	30.2	31.8	32.3	+0.5
Young Adults	23.8	25.2	25.1	25.5	26.5	27.0	26.8	27.4	27.6	27.9	29.2	29.3	29.0	29.2	28.2	27.7	28.5	28.6	+0.1
Inhalants^{c,d}																			
8th Grade	9.0	9.5	11.0	11.7	12.8	12.2	11.8	11.1	10.3	9.4	9.1	7.7	8.7	9.6	9.5	9.1	8.3	8.9	+0.6
10th Grade	7.1	7.5	8.4	9.1	9.6	9.5	8.7	8.0	7.2	7.3	6.6	5.8	5.4	5.9	6.0	6.5	6.6	5.9	-0.7
12th Grade	6.6	6.2	7.0	7.7	8.0	7.6	6.7	6.2	5.6	5.9	4.5	4.5	3.9	4.2	5.0	4.5	3.7	3.8	+0.1
College Students	3.5	3.1	3.8	3.0	3.9	3.6	4.1	3.0	3.2	2.9	2.8	2.0	1.8	2.7	1.8	1.5	1.5	1.1	-0.4
Young Adults	2.0	1.9	2.1	2.1	2.4	2.2	2.3	2.1	2.3	2.1	1.7	1.6	1.4	1.7	1.3	1.3	0.8	1.4	+0.6
Nitrites^e																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.9	0.5	0.9	1.1	1.1	1.6	1.2	1.4	0.9	0.6	0.6	1.1	0.9	0.8	0.6	0.5	0.8	0.6	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.2	0.1	0.4	0.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Hallucinogens^{b,f}																			
8th Grade	1.9	2.5	2.6	2.7	3.6	4.1	3.7	3.4	2.9	2.8†	3.4	2.6	2.6	2.2	2.4	2.1	1.9	2.1	+0.2
10th Grade	4.0	4.3	4.7	5.8	7.2	7.8	7.6	6.9	6.9	6.1†	6.2	4.7	4.1	4.1	4.0	4.1	4.4	3.9	-0.5
12th Grade	5.8	5.9	7.4	7.6	9.3	10.1	9.8	9.0	9.4	8.1†	9.1	6.6	5.9	6.2	5.5	4.9	5.4	5.9	+0.5
College Students	6.3	6.8	6.0	6.2	8.2	6.9	7.7	7.2	7.8	6.7†	7.5	6.3	7.4	5.9	5.0	5.6	4.9	5.1	+0.2
Young Adults	4.5	5.0	4.5	4.8	5.6	5.6	5.9	5.2	5.4	5.4†	5.4	4.7	5.2	4.7	4.5	4.1	3.8	3.8	0.0

(Table continued on next page.)

TABLE 2-2 (cont.)
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
LSD																			
8th Grade	1.7	2.1	2.3	2.4	3.2	3.5	3.2	2.8	2.4	2.4	2.2	1.5	1.3	1.1	1.2	0.9	1.1	1.3	+0.2
10th Grade	3.7	4.0	4.2	5.2	6.5	6.9	6.7	5.9	6.0	5.1	4.1	2.6	1.7	1.6	1.5	1.7	1.9	1.8	-0.1
12th Grade	5.2	5.6	6.8	6.9	8.4	8.8	8.4	7.6	8.1	6.6	6.6	3.5	1.9	2.2	1.8	1.7	2.1	2.7	+0.6
College Students	5.1	5.7	5.1	5.2	6.9	5.2	5.0	4.4	5.4	4.3	4.0	2.1	1.4	1.2	0.7	1.4	1.3	2.6	+1.3 s
Young Adults	3.8	4.3	3.8	4.0	4.6	4.5	4.4	3.5	4.0	3.7	3.4	1.8	1.2	0.9	0.8	1.2	1.1	1.4	+0.3
Hallucinogens other than LSD^b																			
8th Grade	0.7	1.1	1.0	1.3	1.7	2.0	1.8	1.6	1.5	1.4†	2.4	2.1	2.1	1.9	2.0	1.8	1.6	1.6	0.0
10th Grade	1.3	1.4	1.9	2.4	2.8	3.3	3.3	3.4	3.2	3.1†	4.3	4.0	3.6	3.7	3.5	3.7	3.8	3.3	-0.5
12th Grade	2.0	1.7	2.2	3.1	3.8	4.4	4.6	4.6	4.3	4.4†	5.9	5.4	5.4	5.6	5.0	4.6	4.8	5.0	+0.2
College Students	3.1	2.6	2.7	2.8	4.0	4.1	4.9	4.4	4.5	4.4†	5.5	5.8	7.1	5.6	5.0	5.4	4.7	4.4	-0.4
Young Adults	1.7	1.9	1.9	2.0	2.5	2.8	3.1	3.0	3.0	3.4†	3.5	4.0	4.9	4.5	4.2	3.8	3.6	3.4	-0.2
PCP^g																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.4	1.4	1.4	1.6	1.8	2.6	2.3	2.1	1.8	2.3	1.8	1.1	1.3	0.7	1.3	0.7	0.9	1.1	+0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.3	0.3	0.2	0.3	0.3	0.2	0.5	0.6	0.6	0.3	0.6	0.3	0.3	0.1	0.6	0.2	0.3	0.4	+0.1
Ecstasy (MDMA)^h																			
8th Grade	—	—	—	—	—	2.3	2.3	1.8	1.7	3.1	3.5	2.9	2.1	1.7	1.7	1.4	1.5	1.7	+0.2
10th Grade	—	—	—	—	—	4.6	3.9	3.3	4.4	5.4	6.2	4.9	3.0	2.4	2.6	2.8	3.5	2.9	-0.6
12th Grade	—	—	—	—	—	4.6	4.0	3.6	5.6	8.2	9.2	7.4	4.5	4.0	3.0	4.1	4.5	4.3	-0.2
College Students	0.9	2.0	0.8	0.5	2.4	2.8	2.4	3.9	5.5	9.1	9.2	6.8	4.4	2.2	2.9	2.6	2.2	3.7	+1.6
Young Adults	0.8	1.0	0.8	0.7	1.6	1.7	2.1	2.9	3.6	7.2	7.5	6.2	4.5	3.5	3.0	3.0	2.5	3.3	+0.8
Cocaine																			
8th Grade	1.1	1.5	1.7	2.1	2.6	3.0	2.8	3.1	2.7	2.6	2.5	2.3	2.2	2.0	2.2	2.0	2.0	1.8	-0.2
10th Grade	2.2	1.9	2.1	2.8	3.5	4.2	4.7	4.7	4.9	4.4	3.6	4.0	3.3	3.7	3.5	3.2	3.4	3.0	-0.5
12th Grade	3.5	3.1	3.3	3.6	4.0	4.9	5.5	5.7	6.2	5.0	4.8	5.0	4.8	5.3	5.1	5.7	5.2	4.4	-0.8
College Students	3.6	3.0	2.7	2.0	3.6	2.9	3.4	4.6	4.6	4.8	4.7	4.8	5.4	6.6	5.7	5.1	5.4	4.4	-1.0
Young Adults	6.2	5.7	4.7	4.3	4.4	4.1	4.7	4.9	5.4	5.4	5.8	5.8	6.6	7.1	6.9	6.6	6.2	6.0	-0.3
Crackⁱ																			
8th Grade	0.7	0.9	1.0	1.3	1.6	1.8	1.7	2.1	1.8	1.8	1.7	1.6	1.6	1.3	1.4	1.3	1.3	1.1	-0.2
10th Grade	0.9	0.9	1.1	1.4	1.8	2.1	2.2	2.5	2.4	2.2	1.8	2.3	1.6	1.7	1.7	1.3	1.3	1.3	-0.1
12th Grade	1.5	1.5	1.5	1.9	2.1	2.1	2.4	2.5	2.7	2.2	2.1	2.3	2.2	2.3	1.9	2.1	1.9	1.6	-0.4 s
College Students	0.5	0.4	0.6	0.5	1.1	0.6	0.4	1.0	0.9	0.9	0.9	0.4	1.3	1.3	0.8	1.0	0.6	0.5	-0.1
Young Adults	1.2	1.4	1.3	1.1	1.1	1.1	1.0	1.1	1.4	1.2	1.3	1.0	1.0	1.3	1.2	1.1	1.0	0.9	-0.1
Other Cocaine^l																			
8th Grade	1.0	1.2	1.3	1.7	2.1	2.5	2.2	2.4	2.3	1.9	1.9	1.8	1.6	1.6	1.7	1.6	1.5	1.4	-0.1
10th Grade	2.1	1.7	1.8	2.4	3.0	3.5	4.1	4.0	4.4	3.8	3.0	3.4	2.8	3.3	3.0	2.9	3.1	2.6	-0.5
12th Grade	3.2	2.6	2.9	3.0	3.4	4.2	5.0	4.9	5.8	4.5	4.4	4.4	4.2	4.7	4.5	5.2	4.5	4.0	-0.5
College Students	3.2	2.4	2.5	1.8	3.3	2.3	3.0	4.2	4.2	4.1	4.1	5.0	5.1	6.3	5.0	3.8	5.3	4.2	-1.1
Young Adults	5.4	5.1	3.9	3.6	3.9	3.8	4.3	4.5	4.8	4.8	5.3	5.6	6.1	6.4	6.3	5.9	5.6	5.5	-0.1

(Table continued on next page.)

TABLE 2-2 (cont.)
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Heroin^k																			
8th Grade	0.7	0.7	0.7	1.2	1.4	1.6	1.3	1.3	1.4	1.1	1.0	0.9	0.9	1.0	0.8	0.8	0.8	0.9	0.0
10th Grade	0.5	0.6	0.7	0.9	1.1	1.2	1.4	1.4	1.4	1.4	0.9	1.1	0.7	0.9	0.9	0.9	0.8	0.8	0.0
12th Grade	0.4	0.6	0.5	0.6	1.1	1.0	1.2	1.0	1.1	1.5	0.9	1.0	0.8	0.9	0.8	0.8	0.9	0.7	-0.2
College Students	0.1	0.1	0.1	0.1	0.3	0.4	0.3	0.6	0.2	0.5	0.4	0.1	0.2	0.4	0.3	0.3	0.2	0.3	+0.2
Young Adults	0.1	0.2	0.2	0.1	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.2	0.4	0.3	0.4	0.4	0.3	0.5	+0.2
With a Needle^l																			
8th Grade	—	—	—	—	0.9	1.0	0.8	0.8	0.9	0.6	0.7	0.6	0.6	0.7	0.6	0.5	0.6	0.5	0.0
10th Grade	—	—	—	—	0.6	0.7	0.7	0.8	0.6	0.5	0.4	0.6	0.5	0.5	0.5	0.5	0.5	0.5	0.0
12th Grade	—	—	—	—	0.5	0.5	0.5	0.4	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.5	0.4	0.4	0.0
College Students	—	—	—	—	0.1	*	0.1	0.2	0.1	0.1	0.1	*	0.1	0.1	0.3	0.3	*	0.0	0.0
Young Adults	—	—	—	—	0.1	0.1	0.1	0.1	0.1	*	0.3	*	*	0.1	0.2	0.3	0.1	0.1	0.0
Without a Needle^l																			
8th Grade	—	—	—	—	0.8	1.0	0.8	0.8	0.9	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.6	+0.1
10th Grade	—	—	—	—	0.8	0.9	1.1	1.0	1.1	1.1	0.7	0.8	0.5	0.7	0.7	0.6	0.6	0.6	0.0
12th Grade	—	—	—	—	1.0	1.0	1.2	0.8	1.0	1.6	0.8	0.8	0.8	0.7	0.8	0.6	1.0	0.5	-0.5 ss
College Students	—	—	—	—	0.0	0.8	0.4	0.9	0.3	0.8	0.6	0.2	0.1	0.6	0.2	0.3	0.2	0.3	+0.1
Young Adults	—	—	—	—	0.3	0.4	0.4	0.7	0.6	0.5	0.9	0.2	0.4	0.3	0.4	0.5	0.3	0.4	+0.1
Narcotics other than Heroin^{m,n}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.5	3.3	3.6	3.8	4.7	5.4	6.2	6.3	6.7	7.0	6.7‡	9.4	9.3	9.5	9.0	9.0	9.2	9.1	0.0
College Students	2.7	2.7	2.5	2.4	3.8	3.1	4.2	4.2	4.3	4.5	5.7‡	7.4	8.7	8.2	8.4	8.8	7.7	6.5	-1.2
Young Adults	2.5	2.5	2.2	2.5	3.0	2.9	3.3	3.4	3.8	4.1	5.0‡	7.1	8.5	9.0	8.7	9.1	8.7	9.1	+0.4
OxyContin^{m,o,w,x}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	1.3	1.7	1.7	1.8	2.6	1.8	2.1	+0.3
10th Grade	—	—	—	—	—	—	—	—	—	—	—	3.0	3.6	3.5	3.2	3.8	3.9	3.6	-0.3
12th Grade	—	—	—	—	—	—	—	—	—	—	—	4.0	4.5	5.0	5.5	4.3	5.2	4.7	-0.5
College Students	—	—	—	—	—	—	—	—	—	—	—	1.5	2.2	2.5	2.1	3.0	2.8	3.6	+0.8
Young Adults	—	—	—	—	—	—	—	—	—	—	—	1.9	2.6	3.1	3.1	3.1	2.9	3.9	+0.9
Vicodin^{m,o,w,x}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	2.5	2.8	2.5	2.6	3.0	2.7	2.9	+0.2
10th Grade	—	—	—	—	—	—	—	—	—	—	—	6.9	7.2	6.2	5.9	7.0	7.2	6.7	-0.5
12th Grade	—	—	—	—	—	—	—	—	—	—	—	9.6	10.5	9.3	9.5	9.7	9.6	9.7	+0.1
College Students	—	—	—	—	—	—	—	—	—	—	—	6.9	7.5	7.4	9.6	7.6	6.7	6.7	0.0
Young Adults	—	—	—	—	—	—	—	—	—	—	—	8.2	8.6	8.9	9.3	9.1	8.9	9.1	+0.2
Amphetamines^m																			
8th Grade	6.2	6.5	7.2	7.9	8.7	9.1	8.1	7.2	6.9	6.5	6.7	5.5	5.5	4.9	4.9	4.7	4.2	4.5	+0.3
10th Grade	8.2	8.2	9.6	10.2	11.9	12.4	12.1	10.7	10.4	11.1	11.7	10.7	9.0	8.5	7.8	7.9	8.0	6.4	-1.6 ss
12th Grade	8.2	7.1	8.4	9.4	9.3	9.5	10.2	10.1	10.2	10.5	10.9	11.1	9.9	10.0	8.6	8.1	7.5	6.8	-0.7
College Students	3.9	3.6	4.2	4.2	5.4	4.2	5.7	5.1	5.8	6.6	7.2	7.0	7.1	7.0	6.7	6.0	6.9	5.7	-1.2
Young Adults	4.3	4.1	4.0	4.5	4.6	4.2	4.6	4.5	4.7	5.4	5.8	5.9	5.8	6.2	5.1	5.6	5.6	5.3	-0.3

(Table continued on next page.)

TABLE 2-2 (cont.)
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Ritalin^{m,o,p}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	2.9	2.8	2.6	2.5	2.4	2.6	2.1	1.6	-0.4
10th Grade	—	—	—	—	—	—	—	—	—	—	4.8	4.8	4.1	3.4	3.4	3.6	2.8	2.9	+0.2
12th Grade	—	—	—	—	—	—	—	—	—	—	5.1	4.0	4.0	5.1	4.4	4.4	3.8	3.4	-0.4
College Students	—	—	—	—	—	—	—	—	—	—	—	5.7	4.7	4.7	4.2	3.9	3.7	3.2	-0.5
Young Adults	—	—	—	—	—	—	—	—	—	—	—	2.9	2.9	2.7	2.5	2.6	2.4	2.4	0.0
Methamphetamine^{o,p}																			
8th Grade	—	—	—	—	—	—	—	—	3.2	2.5	2.8	2.2	2.5	1.5	1.8	1.8	1.1	1.2	+0.1
10th Grade	—	—	—	—	—	—	—	—	4.6	4.0	3.7	3.9	3.3	3.0	2.9	1.8	1.6	1.5	-0.1
12th Grade	—	—	—	—	—	—	—	—	4.7	4.3	3.9	3.6	3.2	3.4	2.5	2.5	1.7	1.2	-0.5
College Students	—	—	—	—	—	—	—	—	3.3	1.6	2.4	1.2	2.6	2.9	1.7	1.2	0.4	0.5	+0.1
Young Adults	—	—	—	—	—	—	—	—	2.8	2.5	2.8	2.5	2.7	2.8	2.4	1.9	1.5	1.0	-0.5
Crystal Meth. (Ice)^p																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.4	1.3	1.7	1.8	2.4	2.8	2.3	3.0	1.9	2.2	2.5	3.0	2.0	2.1	2.3	1.9	1.6	1.1	-0.6 s
College Students	0.1	0.2	0.7	0.8	1.1	0.3	0.8	1.0	0.5	0.5	0.6	0.8	0.9	1.1	1.4	0.6	0.7	0.1	-0.6
Young Adults	0.3	0.4	0.8	0.9	1.2	0.9	0.9	1.1	0.9	1.2	1.1	1.4	1.3	1.5	1.6	1.1	1.1	0.8	-0.3
Sedatives (Barbiturates)^m																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.4	2.8	3.4	4.1	4.7	4.9	5.1	5.5	5.8	6.2	5.7	6.7	6.0	6.5	7.2	6.6	6.2	5.8	-0.4
College Students	1.2	1.4	1.5	1.2	2.0	2.3	3.0	2.5	3.2	3.7	3.8	3.7	4.1	4.2	3.9	3.4	3.6	3.7	+0.1
Young Adults	1.8	1.6	1.9	1.8	2.1	2.2	2.4	2.5	2.8	3.4	3.7	3.9	3.9	4.4	4.2	3.9	4.2	4.7	+0.5
Methaqualone^{m,q}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.5	0.6	0.2	0.8	0.7	1.1	1.0	1.1	1.1	0.3	0.8	0.9	0.6	0.8	0.9	0.8	0.5	0.5	0.0
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers^{b,m}																			
8th Grade	1.8	2.0	2.1	2.4	2.7	3.3	2.9	2.6	2.5	2.6†	2.8	2.6	2.7	2.5	2.8	2.6	2.4	2.4	0.0
10th Grade	3.2	3.5	3.3	3.3	4.0	4.6	4.9	5.1	5.4	5.6†	7.3	6.3	5.3	5.1	4.8	5.2	5.3	4.6	-0.7
12th Grade	3.6	2.8	3.5	3.7	4.4	4.6	4.7	5.5	5.8	5.7†	6.9	7.7	6.7	7.3	6.8	6.6	6.2	6.2	0.0
College Students	2.4	2.9	2.4	1.8	2.9	2.8	3.8	3.9	3.8	4.2†	5.1	6.7	6.9	6.7	6.4	5.8	5.5	5.0	-0.5
Young Adults	3.5	3.4	3.1	2.9	3.4	3.2	3.1	3.8	3.7	4.6†	5.5	7.0	6.8	7.4	6.7	6.5	7.1	6.8	-0.4
OTC Cough/Cold Medicines^{o,p}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4.2	4.0	3.6	-0.5
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.3	5.4	5.3	-0.1
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.9	5.8	5.5	-0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page.)

TABLE 2-2 (cont.)
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Rohypnol^f																			
8th Grade	—	—	—	—	—	1.0	0.8	0.8	0.5	0.5	0.7	0.3	0.5	0.6	0.7	0.5	0.7	0.5	-0.2
10th Grade	—	—	—	—	—	1.1	1.3	1.2	1.0	0.8	1.0	0.7	0.6	0.7	0.5	0.5	0.7	0.4	-0.3
12th Grade	—	—	—	—	—	1.1	1.2	1.4	1.0	0.8	0.9‡	1.6	1.3	1.6	1.2	1.1	1.0	1.3	+0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	0.7	0.4	0.3	0.1	0.2	0.1	0.3	+0.2
Young Adults	—	—	—	—	—	—	—	—	—	—	—	0.3	0.5	0.1	0.1	0.2	0.3	0.2	0.0
GHB^{o,y}																			
8th Grade	—	—	—	—	—	—	—	—	—	1.2	1.1	0.8	0.9	0.7	0.5	0.8	0.7	1.1	+0.4 s
10th Grade	—	—	—	—	—	—	—	—	—	1.1	1.0	1.4	1.4	0.8	0.8	0.7	0.6	0.5	0.0
12th Grade	—	—	—	—	—	—	—	—	—	1.9	1.6	1.5	1.4	2.0	1.1	1.1	0.9	1.2	+0.3
College Students	—	—	—	—	—	—	—	—	—	—	—	0.6	0.3	0.7	0.4	*	0.1	0.2	+0.1
Young Adults	—	—	—	—	—	—	—	—	—	—	—	0.8	0.6	0.5	0.3	0.2	0.4	0.3	-0.1
Ketamine^{o,z}																			
8th Grade	—	—	—	—	—	—	—	—	—	1.6	1.3	1.3	1.1	0.9	0.6	0.9	1.0	1.2	+0.2
10th Grade	—	—	—	—	—	—	—	—	—	2.1	2.1	2.2	1.9	1.3	1.0	1.0	0.8	1.0	+0.2
12th Grade	—	—	—	—	—	—	—	—	—	2.5	2.5	2.6	2.1	1.9	1.6	1.4	1.3	1.5	+0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	1.3	1.0	1.5	0.5	0.9	0.2	0.4	+0.1
Young Adults	—	—	—	—	—	—	—	—	—	—	—	1.2	0.9	0.6	0.5	0.5	0.3	0.4	+0.1
Alcohol^s																			
Any Use																			
8th Grade	54.0	53.7‡	45.4	46.8	45.3	46.5	45.5	43.7	43.5	43.1	41.9	38.7	37.2	36.7	33.9	33.6	31.8	32.1	+0.3
10th Grade	72.3	70.2‡	63.4	63.9	63.5	65.0	65.2	62.7	63.7	65.3	63.5	60.0	59.3	58.2	56.7	55.8	56.3	52.5	-3.8 sss
12th Grade	77.7	76.8‡	72.7	73.0	73.7	72.5	74.8	74.3	73.8	73.2	73.3	71.5	70.1	70.6	68.6	66.5	66.4	65.5	-0.9
College Students	88.3	86.9	85.1	82.7	83.2	82.9	82.4	84.6	83.6	83.2	83.0	82.9	81.7	81.2	83.0	82.1	80.9	82.1	+1.3
Young Adults	86.9	86.2	85.3	83.7	84.7	84.0	84.3	84.0	84.1	84.0	84.3	84.9	83.3	84.4	83.8	84.4	84.0	83.6	-0.4
Been Drunk^t																			
8th Grade	17.5	18.3	18.2	18.2	18.4	19.8	18.4	17.9	18.5	18.5	16.6	15.0	14.5	14.5	14.1	13.9	12.6	12.7	+0.1
10th Grade	40.1	37.0	37.8	38.0	38.5	40.1	40.7	38.3	40.9	41.6	39.9	35.4	34.7	35.1	34.2	34.5	34.4	30.0	-4.3 sss
12th Grade	52.7	50.3	49.6	51.7	52.5	51.9	53.2	52.0	53.2	51.8	53.2	50.4	48.0	51.8	47.7	47.9	46.1	45.6	-0.5
College Students	69.1	67.3	65.6	63.1	62.1	64.2	66.8	67.0	65.4	64.7	68.8	66.0	64.7	67.1	64.2	66.2	64.8	66.8	+2.0
Young Adults	62.0	60.9	61.1	58.8	61.6	59.9	63.2	59.6	63.2	60.6	63.1	61.8	62.9	63.8	63.5	65.7	65.8	66.0	+0.3
Flavored Alcoholic Beverages^{g,o,aa}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	30.4	27.9	26.8	26.0	25.0	-1.0
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	49.7	48.5	48.8	45.9	43.4	-2.4
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	55.2	55.8	58.4	54.7	53.6	51.8	-1.9
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	63.2	67.0	63.5	62.6	65.0	+2.4
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	62.7	58.4	58.5	58.9	58.3	-0.6
Cigarettes																			
Any Use																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
College Students	35.6	37.3	38.8	37.6	39.3	41.4	43.6	44.3	44.5	41.3	39.0	38.3	35.2	36.7	36.0	30.9	30.7	30.0	-0.6
Young Adults	37.7	37.9	37.8	38.3	38.8	40.3	41.8	41.6	41.1	40.9	41.1	39.1	38.6	39.0	39.1	36.9	36.2	35.0	-1.2

(Table continued on next page.)

TABLE 2-2 (cont.)
Trends in Annual Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Bidis^{o,p}																			
8th Grade	—	—	—	—	—	—	—	—	—	3.9	2.7	2.7	2.0	1.7	1.6	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	6.4	4.9	3.1	2.8	2.1	1.6	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	9.2	7.0	5.9	4.0	3.6	3.3	2.3	1.7	1.9	+0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Kreteks^{o,p}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	2.6	2.6	2.0	1.9	1.4	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	6.0	4.9	3.8	3.7	2.8	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	10.1	8.4	6.7	6.5	7.1	6.2	6.8	6.8	0.0
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids^{v,w}																			
8th Grade	1.0	1.1	0.9	1.2	1.0	0.9	1.0	1.2	1.7	1.7	1.6	1.5	1.4	1.1	1.1	0.9	0.8	0.9	+0.1
10th Grade	1.1	1.1	1.0	1.1	1.2	1.2	1.2	1.2	1.7	2.2	2.1	2.2	1.7	1.5	1.3	1.2	1.1	0.9	-0.1
12th Grade	1.4	1.1	1.2	1.3	1.5	1.4	1.4	1.7	1.8	1.7	2.4	2.5	2.1	2.5	1.5	1.8	1.4	1.5	+0.1
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.5	0.4	0.3	0.4	0.5	0.3	0.5	0.4	0.6	0.4	0.4	0.4	0.5	0.5	0.5	0.3	0.7	0.4	-0.3

Source. The Monitoring the Future study, the University of Michigan.

Note. See relevant footnotes at the end of Table 2-1.

TABLE 2-3
Trends in 30-Day Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Any Illicit Drug^a																			
8th Grade	5.7	6.8	8.4	10.9	12.4	14.6	12.9	12.1	12.2	11.9	11.7	10.4	9.7	8.4	8.5	8.1	7.4	7.6	+0.2
10th Grade	11.6	11.0	14.0	18.5	20.2	23.2	23.0	21.5	22.1	22.5	22.7	20.8	19.5	18.3	17.3	16.8	16.9	15.8	-1.1
12th Grade	16.4	14.4	18.3	21.9	23.8	24.6	26.2	25.6	25.9	24.9	25.7	25.4	24.1	23.4	23.1	21.5	21.9	22.3	+0.5
College Students	15.2	16.1	15.1	16.0	19.1	17.6	19.2	19.7	21.6	21.5	21.9	21.5	21.4	21.2	19.5	19.2	19.3	18.9	-0.3
Young Adults	15.1	14.8	14.9	15.3	15.8	15.8	16.4	16.1	17.1	18.1	18.8	18.9	19.9	19.1	18.6	18.5	18.9	19.3	+0.4
Any Illicit Drug other than Marijuana^{a,b}																			
8th Grade	3.8	4.7	5.3	5.6	6.5	6.9	6.0	5.5	5.5	5.6‡	5.5	4.7	4.7	4.1	4.1	3.8	3.6	3.8	+0.1
10th Grade	5.5	5.7	6.5	7.1	8.9	8.9	8.8	8.6	8.6	8.5‡	8.7	8.1	6.9	6.9	6.4	6.3	6.9	5.3	-1.6 sss
12th Grade	7.1	6.3	7.9	8.8	10.0	9.5	10.7	10.7	10.4	10.4‡	11.0	11.3	10.4	10.8	10.3	9.8	9.5	9.3	-0.2
College Students	4.3	4.6	5.4	4.6	6.3	4.5	6.8	6.1	6.4	6.9‡	7.5	7.8	8.2	9.1	8.2	8.2	8.1	7.3	-0.8
Young Adults	5.4	5.5	4.9	5.3	5.7	4.7	5.5	5.5	6.0	6.4‡	7.0	7.7	8.3	8.5	8.2	8.1	8.6	8.9	+0.2
Any Illicit Drug including Inhalants^{a,c,d}																			
8th Grade	8.8	10.0	12.0	14.3	16.1	17.5	16.0	14.9	15.1	14.4	14.0	12.6	12.1	11.2	11.2	10.9	10.1	10.4	+0.3
10th Grade	13.1	12.6	15.5	20.0	21.6	24.5	24.1	22.5	23.1	23.6	23.6	21.7	20.5	19.3	18.4	17.7	18.1	16.8	-1.3
12th Grade	17.8	15.5	19.3	23.0	24.8	25.5	26.9	26.6	26.4	26.4	26.5	25.9	24.6	23.3	24.2	22.1	22.8	22.8	0.0
College Students	15.1	16.5	15.7	16.4	19.6	18.0	19.6	21.0	21.8	22.6	21.9	21.9	21.6	21.7	19.0	19.7	18.1	18.9	+0.8
Young Adults	15.4	15.3	15.1	16.1	16.1	16.4	16.9	16.7	17.4	18.8	19.2	19.5	20.1	19.6	18.0	18.4	19.1	19.3	+0.2
Marijuana/Hashish																			
8th Grade	3.2	3.7	5.1	7.8	9.1	11.3	10.2	9.7	9.7	9.1	9.2	8.3	7.5	6.4	6.6	6.5	5.7	5.8	+0.1
10th Grade	8.7	8.1	10.9	15.8	17.2	20.4	20.5	18.7	19.4	19.7	19.8	17.8	17.0	15.9	15.2	14.2	14.2	13.8	-0.3
12th Grade	13.8	11.9	15.5	19.0	21.2	21.9	23.7	22.8	23.1	21.6	22.4	21.5	21.2	19.9	19.8	18.3	18.8	19.4	+0.7
College Students	14.1	14.6	14.2	15.1	18.6	17.5	17.7	18.6	20.7	20.0	20.2	19.7	19.3	18.9	17.1	16.7	16.8	17.0	+0.2
Young Adults	13.5	13.3	13.4	14.1	14.0	15.1	15.0	14.9	15.6	16.1	16.7	16.9	17.3	16.5	15.8	15.7	16.0	16.0	0.0
Inhalants^{c,d}																			
8th Grade	4.4	4.7	5.4	5.6	6.1	5.8	5.6	4.8	5.0	4.5	4.0	3.8	4.1	4.5	4.2	4.1	3.9	4.1	+0.2
10th Grade	2.7	2.7	3.3	3.6	3.5	3.3	3.0	2.9	2.6	2.6	2.4	2.4	2.2	2.4	2.2	2.3	2.5	2.1	-0.4
12th Grade	2.4	2.3	2.5	2.7	3.2	2.5	2.5	2.3	2.0	2.2	1.7	1.5	1.5	1.5	2.0	1.5	1.2	1.4	+0.2
College Students	0.9	1.1	1.3	0.6	1.6	0.8	0.8	0.6	1.5	0.9	0.4	0.7	0.4	0.4	0.3	0.4	0.1	0.4	+0.2
Young Adults	0.5	0.6	0.7	0.5	0.7	0.5	0.5	0.7	0.8	0.5	0.4	0.5	0.3	0.3	0.2	0.3	0.2	0.4	+0.3
Nitrites^e																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.4	0.3	0.6	0.4	0.4	0.7	0.7	1.0	0.4	0.3	0.5	0.6	0.7	0.7	0.5	0.3	0.5	0.3	-0.2
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	*	0.1	0.2	0.1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Hallucinogens^{b,f}																			
8th Grade	0.8	1.1	1.2	1.3	1.7	1.9	1.8	1.4	1.3	1.2‡	1.6	1.2	1.2	1.0	1.1	0.9	1.0	0.9	0.0
10th Grade	1.6	1.8	1.9	2.4	3.3	2.8	3.3	3.2	2.9	2.3‡	2.1	1.6	1.5	1.6	1.5	1.5	1.7	1.3	-0.4
12th Grade	2.2	2.1	2.7	3.1	4.4	3.5	3.9	3.8	3.5	2.6‡	3.3	2.3	1.8	1.9	1.9	1.5	1.7	2.2	+0.5 s
College Students	1.2	2.3	2.5	2.1	3.3	1.9	2.1	2.1	2.0	1.4‡	1.8	1.2	1.8	1.3	1.2	0.9	1.3	1.7	+0.4
Young Adults	1.1	1.5	1.2	1.4	1.7	1.2	1.5	1.4	1.3	1.2‡	1.2	0.9	1.2	0.9	0.8	0.7	0.9	0.9	0.0

(Table continued on next page.)

TABLE 2-3 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
 (Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
LSD																			
8th Grade	0.6	0.9	1.0	1.1	1.4	1.5	1.5	1.1	1.1	1.0	1.0	0.7	0.6	0.5	0.5	0.4	0.5	0.5	0.0
10th Grade	1.5	1.6	1.6	2.0	3.0	2.4	2.8	2.7	2.3	1.6	1.5	0.7	0.6	0.6	0.6	0.7	0.7	0.7	0.0
12th Grade	1.9	2.0	2.4	2.6	4.0	2.5	3.1	3.2	2.7	1.6	2.3	0.7	0.6	0.7	0.7	0.6	0.6	1.1	+0.5 ss
College Students	0.8	1.8	1.6	1.8	2.5	0.9	1.1	1.5	1.2	0.9	1.0	0.2	0.2	0.2	0.1	0.3	0.3	0.8	+0.5
Young Adults	0.8	1.1	0.8	1.1	1.3	0.7	0.9	1.0	0.8	0.8	0.7	0.3	0.2	0.1	0.1	0.2	0.2	0.4	+0.2 s
Hallucinogens other than LSD^b																			
8th Grade	0.3	0.4	0.5	0.7	0.8	0.9	0.7	0.7	0.6	0.6‡	1.1	1.0	1.0	0.8	0.9	0.7	0.7	0.7	0.0
10th Grade	0.4	0.5	0.7	1.0	1.0	1.0	1.2	1.4	1.2	1.2‡	1.4	1.4	1.2	1.4	1.3	1.3	1.4	1.0	-0.5 ss
12th Grade	0.7	0.5	0.8	1.2	1.3	1.6	1.7	1.6	1.6	1.7‡	1.9	2.0	1.5	1.7	1.6	1.3	1.4	1.6	+0.3
College Students	0.6	0.7	1.1	0.8	1.6	1.2	1.2	0.7	1.2	0.8‡	0.8	1.1	1.7	1.2	1.1	0.7	1.1	1.3	+0.2
Young Adults	0.3	0.5	0.6	0.6	0.6	0.6	0.7	0.5	0.6	0.7‡	0.6	0.8	1.2	0.9	0.8	0.6	0.8	0.7	-0.1
PCP^g																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.5	0.6	1.0	0.7	0.6	1.3	0.7	1.0	0.8	0.9	0.5	0.4	0.6	0.4	0.7	0.4	0.5	0.6	+0.1
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.1	0.2	0.2	0.1	*	0.1	0.1	0.2	0.2	*	*	0.1	0.1	0.1	*	*	*	0.1	-0.1
Ecstasy (MDMA)^h																			
8th Grade	—	—	—	—	—	1.0	1.0	0.9	0.8	1.4	1.8	1.4	0.7	0.8	0.6	0.7	0.6	0.8	+0.2
10th Grade	—	—	—	—	—	1.8	1.3	1.3	1.8	2.6	2.6	1.8	1.1	0.8	1.0	1.2	1.2	1.1	-0.1
12th Grade	—	—	—	—	—	2.0	1.6	1.5	2.5	3.6	2.8	2.4	1.3	1.2	1.0	1.3	1.6	1.8	+0.2
College Students	0.2	0.4	0.3	0.2	0.7	0.7	0.8	0.8	2.1	2.5	1.5	0.7	1.0	0.7	0.8	0.6	0.4	0.6	+0.2
Young Adults	0.1	0.3	0.3	0.2	0.4	0.3	0.6	0.8	1.3	1.9	1.8	1.3	0.8	0.6	0.6	0.7	0.5	0.6	+0.1
Cocaine																			
8th Grade	0.5	0.7	0.7	1.0	1.2	1.3	1.1	1.4	1.3	1.2	1.2	1.1	0.9	0.9	1.0	1.0	0.9	0.8	-0.1
10th Grade	0.7	0.7	0.9	1.2	1.7	1.7	2.0	2.1	1.8	1.8	1.3	1.6	1.3	1.7	1.5	1.5	1.3	1.2	0.0
12th Grade	1.4	1.3	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.1	2.1	2.3	2.1	2.3	2.3	2.5	2.0	1.9	-0.1
College Students	1.0	1.0	0.7	0.6	0.7	0.8	1.6	1.6	1.2	1.4	1.9	1.6	1.9	2.4	1.8	1.8	1.7	1.2	-0.6
Young Adults	2.0	1.8	1.4	1.3	1.5	1.2	1.6	1.7	1.9	1.7	2.2	2.2	2.4	2.2	2.2	2.3	2.1	1.9	-0.2
Crackⁱ																			
8th Grade	0.3	0.5	0.4	0.7	0.7	0.8	0.7	0.9	0.8	0.8	0.8	0.8	0.7	0.6	0.6	0.6	0.6	0.5	-0.2
10th Grade	0.3	0.4	0.5	0.6	0.9	0.8	0.9	1.1	0.8	0.9	0.7	1.0	0.7	0.8	0.7	0.7	0.5	0.5	0.0
12th Grade	0.7	0.6	0.7	0.8	1.0	1.0	0.9	1.0	1.1	1.0	1.1	1.2	0.9	1.0	1.0	0.9	0.9	0.8	-0.1
College Students	0.3	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.1	0.3	0.4	0.4	0.1	*	0.1	0.1	0.0
Young Adults	0.4	0.4	0.4	0.3	0.2	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.4	+0.1
Other Cocaine^j																			
8th Grade	0.5	0.5	0.6	0.9	1.0	1.0	0.8	1.0	1.1	0.9	0.9	0.8	0.7	0.7	0.7	0.7	0.6	0.6	0.0
10th Grade	0.6	0.6	0.7	1.0	1.4	1.3	1.6	1.8	1.6	1.6	1.2	1.3	1.1	1.5	1.3	1.3	1.1	1.0	-0.1
12th Grade	1.2	1.0	1.2	1.3	1.3	1.6	2.0	2.0	2.5	1.7	1.8	1.9	1.8	2.2	2.0	2.4	1.7	1.7	0.0
College Students	1.0	0.9	0.6	0.3	0.8	0.6	1.3	1.5	1.0	0.9	1.5	1.4	1.9	2.2	1.8	1.3	1.6	1.1	-0.5
Young Adults	1.8	1.7	1.1	1.0	1.3	1.1	1.5	1.5	1.6	1.5	1.8	2.0	2.1	2.1	1.9	1.9	2.0	1.7	-0.3

(Table continued on next page.)

TABLE 2-3 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Heroin^k																			
8th Grade	0.3	0.4	0.4	0.6	0.6	0.7	0.6	0.6	0.6	0.5	0.6	0.5	0.4	0.5	0.5	0.3	0.4	0.4	0.0
10th Grade	0.2	0.2	0.3	0.4	0.6	0.5	0.6	0.7	0.7	0.5	0.3	0.5	0.3	0.5	0.5	0.5	0.4	0.4	-0.1
12th Grade	0.2	0.3	0.2	0.3	0.6	0.5	0.5	0.5	0.5	0.7	0.4	0.5	0.4	0.5	0.5	0.4	0.4	0.4	0.0
College Students	0.1	*	*	*	0.1	*	0.2	0.1	0.1	0.2	0.1	*	*	0.1	0.1	0.2	0.1	*	-0.1
Young Adults	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.3	*	0.1	0.1	0.1	0.2	0.1	0.1	0.0
With a Needle^l																			
8th Grade	—	—	—	—	0.4	0.5	0.4	0.5	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.0
10th Grade	—	—	—	—	0.3	0.3	0.3	0.4	0.3	0.3	0.2	0.3	0.2	0.3	0.3	0.3	0.3	0.2	-0.1
12th Grade	—	—	—	—	0.3	0.4	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.2	0.0
College Students	—	—	—	—	*	*	0.1	*	0.1	0.1	*	*	0.1	0.1	0.1	0.1	*	0.0	0.0
Young Adults	—	—	—	—	*	*	0.1	*	0.1	*	0.2	*	*	0.1	0.1	0.1	*	*	0.0
Without a Needle^l																			
8th Grade	—	—	—	—	0.3	0.4	0.4	0.3	0.4	0.3	0.4	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.0
10th Grade	—	—	—	—	0.3	0.3	0.4	0.5	0.5	0.4	0.2	0.4	0.2	0.3	0.3	0.3	0.2	0.3	0.0
12th Grade	—	—	—	—	0.6	0.4	0.6	0.4	0.4	0.7	0.3	0.5	0.4	0.3	0.5	0.3	0.4	0.2	-0.2
College Students	—	—	—	—	*	0.1	0.2	0.2	0.3	0.4	0.3	*	*	0.3	*	0.2	0.1	0.1	0.0
Young Adults	—	—	—	—	0.1	*	0.1	0.2	0.2	0.2	0.4	*	0.1	0.1	0.1	0.3	0.2	*	-0.1
Narcotics other than Heroin^{m,n}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.1	1.2	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.9	3.0‡	4.0	4.1	4.3	3.9	3.8	3.8	3.8	0.0
College Students	0.6	1.0	0.7	0.4	1.2	0.7	1.3	1.1	1.0	1.7	1.7‡	3.2	2.3	3.0	3.1	3.1	2.2	2.3	+0.1
Young Adults	0.6	0.7	0.7	0.6	0.9	0.7	0.9	0.9	1.2	1.4	1.7‡	2.9	2.9	3.0	3.5	3.2	3.4	3.6	+0.2
Amphetamines^m																			
8th Grade	2.6	3.3	3.6	3.6	4.2	4.6	3.8	3.3	3.4	3.4	3.2	2.8	2.7	2.3	2.3	2.1	2.0	2.2	+0.2
10th Grade	3.3	3.6	4.3	4.5	5.3	5.5	5.1	5.1	5.0	5.4	5.6	5.2	4.3	4.0	3.7	3.5	4.0	2.8	-1.2 sss
12th Grade	3.2	2.8	3.7	4.0	4.0	4.1	4.8	4.6	4.5	5.0	5.6	5.5	5.0	4.6	3.9	3.7	3.7	2.9	-0.8 ss
College Students	1.0	1.1	1.5	1.5	2.2	0.9	2.1	1.7	2.3	2.9	3.3	3.0	3.1	3.2	2.9	2.5	3.1	2.8	-0.3
Young Adults	1.5	1.5	1.5	1.7	1.7	1.5	1.7	1.7	1.9	2.3	2.4	2.5	2.5	2.4	2.1	2.2	2.3	2.2	-0.2
Methamphetamine^{o,p}																			
8th Grade	—	—	—	—	—	—	—	—	1.1	0.8	1.3	1.1	1.2	0.6	0.7	0.6	0.6	0.7	+0.1
10th Grade	—	—	—	—	—	—	—	—	1.8	2.0	1.5	1.8	1.4	1.3	1.1	0.7	0.4	0.7	+0.4 s
12th Grade	—	—	—	—	—	—	—	—	1.7	1.9	1.5	1.7	1.7	1.4	0.9	0.9	0.6	0.6	-0.1
College Students	—	—	—	—	—	—	—	—	1.2	0.2	0.5	0.2	0.6	0.2	0.1	0.2	0.1	0.0	-0.1
Young Adults	—	—	—	—	—	—	—	—	0.8	0.7	1.0	1.0	0.7	0.6	0.7	0.5	0.6	0.3	-0.3
Crystal Meth. (Ice)^p																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.6	0.5	0.6	0.7	1.1	1.1	0.8	1.2	0.8	1.0	1.1	1.2	0.8	0.8	0.9	0.7	0.6	0.6	0.0
College Students	*	*	0.3	0.5	0.3	0.1	0.2	0.3	*	*	0.1	*	0.3	0.1	0.2	*	0.1	0.0	-0.1
Young Adults	*	0.1	0.3	0.5	0.3	0.3	0.3	0.3	0.4	0.4	0.4	0.5	0.4	0.4	0.6	0.3	0.3	0.3	0.0

(Table continued on next page.)

TABLE 2-3 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Sedatives																			
(Barbiturates)^m																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.4	1.1	1.3	1.7	2.2	2.1	2.1	2.6	2.6	3.0	2.8	3.2	2.9	2.9	3.3	3.0	2.7	2.8	+0.1
College Students	0.3	0.7	0.4	0.4	0.5	0.8	1.2	1.1	1.1	1.1	1.5	1.7	1.7	1.5	1.3	1.3	1.4	1.4	+0.1
Young Adults	0.5	0.5	0.6	0.6	0.8	0.8	0.9	0.9	1.1	1.3	1.7	1.5	1.5	1.8	1.7	1.5	1.6	1.9	+0.3
Methaqualone^{m,q}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.2	0.4	0.1	0.4	0.4	0.6	0.3	0.6	0.4	0.2	0.5	0.3	0.4	0.5	0.5	0.4	0.4	0.2	-0.1
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers^{b,m}																			
8th Grade	0.8	0.8	0.9	1.1	1.2	1.5	1.2	1.2	1.1	1.4†	1.2	1.2	1.4	1.2	1.3	1.3	1.1	1.2	+0.1
10th Grade	1.2	1.5	1.1	1.5	1.7	1.7	2.2	2.2	2.2	2.5†	2.9	2.9	2.4	2.3	2.3	2.4	2.6	1.9	-0.7 sss
12th Grade	1.4	1.0	1.2	1.4	1.8	2.0	1.8	2.4	2.5	2.6†	2.9	3.3	2.8	3.1	2.9	2.7	2.6	2.6	0.0
College Students	0.6	0.6	0.4	0.4	0.5	0.7	1.2	1.3	1.1	2.0†	1.5	3.0	2.8	2.7	2.2	2.1	1.8	1.6	-0.3
Young Adults	0.9	1.0	1.0	0.8	1.1	0.7	1.1	1.2	1.3	1.8†	2.1	2.8	2.4	2.7	2.6	2.3	2.8	2.7	-0.2
Rohypnol^f																			
8th Grade	—	—	—	—	—	0.5	0.3	0.4	0.3	0.3	0.4	0.2	0.1	0.2	0.2	0.4	0.3	0.1	-0.3
10th Grade	—	—	—	—	—	0.5	0.5	0.4	0.5	0.4	0.2	0.4	0.2	0.3	0.2	0.2	0.2	0.2	0.0
12th Grade	—	—	—	—	—	0.5	0.3	0.3	0.3	0.4	0.3	—	—	—	—	—	—	—	—
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Alcohol^s																			
Any Use																			
8th Grade	25.1	26.1†	24.3	25.5	24.6	26.2	24.5	23.0	24.0	22.4	21.5	19.6	19.7	18.6	17.1	17.2	15.9	15.9	+0.1
10th Grade	42.8	39.9†	38.2	39.2	38.8	40.4	40.1	38.8	40.0	41.0	39.0	35.4	35.4	35.2	33.2	33.8	33.4	28.8	-4.7 sss
12th Grade	54.0	51.3†	48.6	50.1	51.3	50.8	52.7	52.0	51.0	50.0	49.8	48.6	47.5	48.0	47.0	45.3	44.4	43.1	-1.3
College Students	74.7	71.4	70.1	67.8	67.5	67.0	65.8	68.1	69.6	67.4	67.0	68.9	66.2	67.7	67.9	65.4	66.6	69.0	+2.4
Young Adults	70.6	69.0	68.3	67.7	68.1	66.7	67.5	66.9	68.2	66.8	67.0	68.3	67.0	68.4	68.6	68.7	69.5	68.9	-0.5
Been Drunk^t																			
8th Grade	7.6	7.5	7.8	8.7	8.3	9.6	8.2	8.4	9.4	8.3	7.7	6.7	6.7	6.2	6.0	6.2	5.5	5.4	-0.1
10th Grade	20.5	18.1	19.8	20.3	20.8	21.3	22.4	21.1	22.5	23.5	21.9	18.3	18.2	18.5	17.6	18.8	18.1	14.4	-3.7 sss
12th Grade	31.6	29.9	28.9	30.8	33.2	31.3	34.2	32.9	32.9	32.3	32.7	30.3	30.9	32.5	30.2	30.0	28.7	27.6	-1.1
College Students	45.0	45.0	43.8	42.8	37.9	40.3	46.4	44.3	44.6	43.9	44.7	44.4	40.4	47.4	43.1	47.6	46.8	45.3	-1.5
Young Adults	35.4	35.6	34.2	34.3	33.0	33.2	35.6	34.2	37.7	35.7	36.8	37.1	37.8	39.0	39.0	42.1	41.4	40.7	-0.7
Flavored Alcoholic Beverages^{g,o}																			
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	14.6	12.9	13.1	12.2	10.2	-2.0 ss
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	25.1	23.1	24.7	21.8	20.2	-1.6
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	31.1	30.5	29.3	29.1	27.4	-1.8
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	34.1	30.9	26.2	27.5	35.8	+8.3
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	29.5	27.6	24.9	25.9	26.7	+0.9

(Table continued on next page.)

TABLE 2-3 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Cigarettes																			
Any Use																			
8th Grade	14.3	15.5	16.7	18.6	19.1	21.0	19.4	19.1	17.5	14.6	12.2	10.7	10.2	9.2	9.3	8.7	7.1	6.8	-0.2
10th Grade	20.8	21.5	24.7	25.4	27.9	30.4	29.8	27.6	25.7	23.9	21.3	17.7	16.7	16.0	14.9	14.5	14.0	12.3	-1.7 s
12th Grade	28.3	27.8	29.9	31.2	33.5	34.0	36.5	35.1	34.6	31.4	29.5	26.7	24.4	25.0	23.2	21.6	21.6	20.4	-1.2
College Students	23.2	23.5	24.5	23.5	26.8	27.9	28.3	30.0	30.6	28.2	25.7	26.7	22.5	24.3	23.8	19.2	19.9	17.9	-2.0
Young Adults	28.2	28.3	28.0	28.0	29.2	30.1	29.9	30.9	30.3	30.1	30.2	29.2	28.4	29.2	28.6	27.0	26.2	24.6	-1.6
Smokeless Tobacco^u																			
8th Grade	6.9	7.0	6.6	7.7	7.1	7.1	5.5	4.8	4.5	4.2	4.0	3.3	4.1	4.1	3.3	3.7	3.2	3.5	+0.3
10th Grade	10.0	9.6	10.4	10.5	9.7	8.6	8.9	7.5	6.5	6.1	6.9	6.1	5.3	4.9	5.6	5.7	6.1	5.0	-1.1
12th Grade	—	11.4	10.7	11.1	12.2	9.8	9.7	8.8	8.4	7.6	7.8	6.5	6.7	6.7	7.6	6.1	6.6	6.5	-0.1
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids^{v,w}																			
8th Grade	0.4	0.5	0.5	0.5	0.6	0.4	0.5	0.5	0.7	0.8	0.7	0.8	0.7	0.5	0.5	0.5	0.4	0.5	+0.1
10th Grade	0.6	0.6	0.5	0.6	0.6	0.5	0.7	0.6	0.9	1.0	0.9	1.0	0.8	0.8	0.6	0.6	0.5	0.5	0.0
12th Grade	0.8	0.6	0.7	0.9	0.7	0.7	1.0	1.1	0.9	0.8	1.3	1.4	1.3	1.6	0.9	1.1	1.0	1.0	0.0
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	0.2	0.1	0.0	0.1	0.2	0.2	0.2	0.2	0.3	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.4	0.2	-0.2

Source. The Monitoring the Future study, the University of Michigan.

Note. See relevant footnotes at the end of Table 2-1.


TABLE 2-4
Trends in 30-Day Prevalence of Daily Use of Various Drugs for 8th, 10th,
and 12th Graders, College Students, and Young Adults (Ages 19–28)
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
Marijuana/Hashish																			
Daily ^{bb}																			
8th Grade	0.2	0.2	0.4	0.7	0.8	1.5	1.1	1.1	1.4	1.3	1.3	1.2	1.0	0.8	1.0	1.0	0.8	0.9	+0.1
10th Grade	0.8	0.8	1.0	2.2	2.8	3.5	3.7	3.6	3.8	3.8	4.5	3.9	3.6	3.2	3.1	2.8	2.8	2.7	-0.1
12th Grade	2.0	1.9	2.4	3.6	4.6	4.9	5.8	5.6	6.0	6.0	5.8	6.0	6.0	5.6	5.0	5.0	5.1	5.4	+0.4
College Students	1.8	1.6	1.9	1.8	3.7	2.8	3.7	4.0	4.0	4.6	4.5	4.1	4.7	4.5	4.0	4.3	3.5	3.9	+0.4
Young Adults	2.3	2.3	2.4	2.8	3.3	3.3	3.8	3.7	4.4	4.2	5.0	4.5	5.3	5.0	4.9	5.0	5.0	5.1	+0.1
Alcohol^{s,bb}																			
Any Daily Use																			
8th Grade	0.5	0.6†	1.0	1.0	0.7	1.0	0.8	0.9	1.0	0.8	0.9	0.7	0.8	0.6	0.5	0.5	0.6	0.7	+0.2
10th Grade	1.3	1.2†	1.8	1.7	1.7	1.6	1.7	1.9	1.9	1.8	1.9	1.8	1.5	1.3	1.3	1.4	1.4	1.0	-0.4 ss
12th Grade	3.6	3.4†	3.4	2.9	3.5	3.7	3.9	3.9	3.4	2.9	3.6	3.5	3.2	2.8	3.1	3.0	3.1	2.8	-0.3
College Students	4.1	3.7	3.9	3.7	3.0	3.2	4.5	3.9	4.5	3.6	4.7	5.0	4.3	3.7	4.6	4.8	4.3	4.0	-0.3
Young Adults	4.9	4.5	4.5	3.9	3.9	4.0	4.6	4.0	4.8	4.1	4.4	4.7	5.1	4.5	5.2	5.4	5.6	5.3	-0.3
Been Drunk																			
Daily ^{bb}																			
8th Grade	0.1	0.1	0.2	0.3	0.2	0.2	0.2	0.3	0.4	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.0
10th Grade	0.2	0.3	0.4	0.4	0.6	0.4	0.6	0.6	0.7	0.5	0.6	0.5	0.5	0.4	0.4	0.5	0.5	0.3	-0.2 ss
12th Grade	0.9	0.8	0.9	1.2	1.3	1.6	2.0	1.5	1.9	1.7	1.4	1.2	1.6	1.8	1.5	1.6	1.3	1.4	+0.1
College Students	0.5	0.2	0.3	0.8	0.5	0.1	1.3	0.8	1.0	0.7	0.5	0.8	1.1	0.8	0.5	0.6	0.7	0.5	-0.2
Young Adults	0.5	0.4	0.4	0.5	0.3	0.4	0.9	0.5	0.9	0.5	0.4	0.6	0.8	0.7	0.5	0.6	0.6	0.5	-0.1
5+ Drinks in a Row																			
in Last 2 Weeks ^{cc}																			
8th Grade	10.9	11.3	11.3	12.1	12.3	13.3	12.3	11.5	13.1	11.7	11.0	10.3	9.8	9.4	8.4	8.7	8.3	8.1	-0.2
10th Grade	21.0	19.1	21.0	21.9	22.0	22.8	23.1	22.4	23.5	24.1	22.8	20.3	20.0	19.9	19.0	19.9	19.6	16.0	-3.7 sss
12th Grade	29.8	27.9	27.5	28.2	29.8	30.2	31.3	31.5	30.8	30.0	29.7	28.6	27.9	29.2	27.1	25.4	25.9	24.6	-1.3
College Students	42.8	41.4	40.2	40.2	38.6	38.3	40.7	38.9	40.0	39.3	40.9	40.1	38.5	41.7	40.1	40.2	41.1	40.0	-1.1
Young Adults	34.7	34.2	34.4	33.7	32.6	33.6	34.4	34.1	35.8	34.7	35.9	35.9	35.8	37.1	37.0	37.6	37.8	37.9	+0.1
Cigarettes																			
Any Daily Use																			
8th Grade	7.2	7.0	8.3	8.8	9.3	10.4	9.0	8.8	8.1	7.4	5.5	5.1	4.5	4.4	4.0	4.0	3.0	3.1	0.0
10th Grade	12.6	12.3	14.2	14.6	16.3	18.3	18.0	15.8	15.9	14.0	12.2	10.1	8.9	8.3	7.5	7.6	7.2	5.9	-1.3 s
12th Grade	18.5	17.2	19.0	19.4	21.6	22.2	24.6	22.4	23.1	20.6	19.0	16.9	15.8	15.6	13.6	12.2	12.3	11.4	-0.9
College Students	13.8	14.1	15.2	13.2	15.8	15.9	15.2	18.0	19.3	17.8	15.0	15.9	13.8	13.8	12.4	9.2	9.3	9.2	-0.1
Young Adults	21.7	20.9	20.8	20.7	21.2	21.8	20.6	21.9	21.5	21.8	21.2	21.2	20.3	20.8	19.6	18.6	17.3	16.7	-0.6
1/2 Pack+/Day																			
8th Grade	3.1	2.9	3.5	3.6	3.4	4.3	3.5	3.6	3.3	2.8	2.3	2.1	1.8	1.7	1.7	1.5	1.1	1.2	0.0
10th Grade	6.5	6.0	7.0	7.6	8.3	9.4	8.6	7.9	7.6	6.2	5.5	4.4	4.1	3.3	3.1	3.3	2.7	2.0	-0.7 s
12th Grade	10.7	10.0	10.9	11.2	12.4	13.0	14.3	12.6	13.2	11.3	10.3	9.1	8.4	8.0	6.9	5.9	5.7	5.4	-0.2
College Students	8.0	8.9	8.9	8.0	10.2	8.4	9.1	11.3	11.0	10.1	7.8	7.9	7.6	6.8	6.7	4.9	4.3	4.3	0.0
Young Adults	16.0	15.7	15.5	15.3	15.7	15.3	14.6	15.6	15.1	15.1	14.6	14.2	13.9	13.5	12.5	11.9	11.1	10.2	-0.9
Smokeless Tobacco																			
Daily ^d																			
8th Grade	1.6	1.8	1.5	1.9	1.2	1.5	1.0	1.0	0.9	0.9	1.2	0.8	0.8	1.0	0.7	0.7	0.8	0.8	-0.1
10th Grade	3.3	3.0	3.3	3.0	2.7	2.2	2.2	2.2	1.5	1.9	2.2	1.7	1.8	1.6	1.9	1.7	1.6	1.4	-0.2
12th Grade	—	4.3	3.3	3.9	3.6	3.3	4.4	3.2	2.9	3.2	2.8	2.0	2.2	2.8	2.5	2.2	2.8	2.7	-0.1
College Students	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Young Adults	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

Source. The Monitoring the Future study, the University of Michigan.

Note. See relevant footnotes at the end of Table 2-1.

FIGURE 2-1
Trends in Annual Prevalence of an Illicit Drug Use Index
across 5 Populations



Source. The Monitoring the Future study, the University of Michigan.

Notes. "Illicit drug use index" includes any use of marijuana, LSD, other hallucinogens, crack, other cocaine, or heroin; or any use which is not under a doctor's orders of narcotics other than heroin, stimulants, sedatives (barbiturates), methaqualone (excluded since 1990), or tranquilizers. Beginning in 1982, the question about stimulant use (i.e., amphetamines) was revised to get respondents to exclude the inappropriate reporting of nonprescription stimulants. The prevalence rate dropped slightly as a result of this methodological change.

Chapter 3

STUDY DESIGN AND PROCEDURES

Monitoring the Future (MTF) incorporates several types of surveys into one study, yielding analytic power beyond the sum of those component parts. The components include cross-sectional studies, repeated cross-sectional studies, and panel studies of particular cohorts. The annual cross-sectional studies provide point estimates of various behaviors and conditions at any given year for a number of subpopulations (e.g., 8th graders, 10th graders, 12th graders, college students, etc.). Repeating these cross-sectional studies over time allows an assessment of change across years in those same segments of the population. The panel-study feature permits the examination of change over time in the same individuals comprising a class cohort, in this case, as they enter adult roles and environments, assume adult responsibilities, and continue further into adulthood.

With a *series* of panel studies of sequential graduating class cohorts, in what is known as a cohort-sequential design, we are able to distinguish among, and explain, three fundamentally different types of change: period, age, and cohort. It is this last feature, the cohort-sequential design aspect, which creates the synergistic effect in terms of analytic power.

This chapter describes this complex research design, including the sampling and field procedures used in both the annual in-school cross-sectional surveys of 8th-, 10th-, and 12th-grade students and the follow-up surveys into adulthood. Related methodological issues such as response rates, population coverage, and the validity of the measures are also discussed.^{14, 15}

RESEARCH DESIGN AND PROCEDURES FOR THE TWELFTH-GRADE SURVEYS

Twelfth graders have been surveyed in the spring of each year since 1975. Each year's data collection takes place in approximately 120 to 146 public and private high schools selected to provide an accurate representative cross-section of 12th graders throughout the coterminous United States (see Figure 3-1).

The Population under Study

We chose the senior year of high school because, for several reasons, it is an optimal point at which to monitor the drug use and related attitudes of youth. First, completion of high school

¹⁴For a more detailed description of the study design, see Bachman, J. G., Johnston, L. D., O'Malley, P. M., & Schulenberg, J. E. (2006). *The Monitoring the Future project after thirty-two years: Design and procedures* (Monitoring the Future Occasional Paper No. 64). Ann Arbor, MI: Institute for Social Research, available online at www.monitoringthefuture.org/pubs/occpapers/occ64.pdf.

¹⁵For a more detailed description of the full range of research objectives of Monitoring the Future, see Johnston, L. D., O'Malley, P. M., Schulenberg, J. E., & Bachman, J. G. (2006). *The aims and objectives of the Monitoring the Future study and progress toward fulfilling them as of 2006* (Monitoring the Future Occasional Paper No. 65). Ann Arbor, MI: Institute for Social Research, available online at www.monitoringthefuture.org/pubs/occpapers/occ65.pdf.

represents the end of an important developmental stage in this society, demarcating both the end of universal education and, for many, the end of living full-time in the parental home. Therefore, it is a logical point at which to take stock of the cumulated influences of these two major environments on American youth. Further, completion of high school represents the jumping-off point from which young people diverge into widely differing social environments and experiences. Thus senior year represents a good time to take a “before” measure, allowing calculation of changes that may be attributable to the many environmental and role transitions occurring in young adulthood, including college attendance. Finally, there were some important practical advantages to building the original system of data collections around samples of 12th graders. The need for systematically repeated, large-scale samples from which to make reliable estimates of change requires that considerable emphasis be put on cost efficiency as well as feasibility. The last year of high school constitutes the final point at which a reasonably good national sample of an age-specific cohort can be drawn and studied economically.

The Omission of Dropouts

One limitation in the study design is the exclusion of those young men and women who drop out of high school before graduation—between 13% and 20% of each age cohort nationally, according to U.S. Census statistics. Clearly, the omission of high school dropouts introduces biases in the estimation of certain characteristics of the entire age group; however, for most purposes, the small proportion of students who drop out sets outer limits on the bias. Further, since the bias from missing dropouts should remain just about constant from year to year, their omission should introduce little or no bias in *change* estimates. Indeed, we believe the changes observed over time for those who finish high school are likely to parallel the changes for dropouts in most instances. Appendix A to *Volume I* addresses in detail the likely effects of the exclusion of dropouts (as well as absentees from school) on estimates of drug use prevalence and trends among the entire age cohort.

Sampling Procedures and Sample Weights

A multistage random sampling procedure is used to secure the nationwide sample of 12th graders each year. Stage 1 is the selection of particular geographic areas, Stage 2 is the selection (with probability proportionate to size) of one or more high schools in each area, and Stage 3 is the selection of 12th graders within each high school. Up to about 350 twelfth graders in each school may be included. In schools with fewer 12th graders, the usual procedure is to include all of them in the data collection, though a smaller sample is sometimes taken to accommodate the needs of the school. When a subset of 12th graders is to be selected, it is done either by randomly sampling entire classrooms or by some other unbiased, random method. Weights are assigned to compensate for differential probabilities of selection at each stage of sampling. Final weights are normalized to average 1.0 (so that the weighted number of cases equals the unweighted number of cases overall). This three-stage sampling procedure has yielded the numbers of participating schools and students shown in Table 3-1.

Questionnaire Administration

About three weeks prior to the questionnaire administration date, parents of the target respondents are sent a letter by first-class mail, usually from the principal, announcing and describing the study and providing them an opportunity to decline participation of their son or daughter if they wish. A flyer describing the study in more detail is enclosed with the letter.

Copies of the same flyers are also given to the students by the teachers in the target classrooms in advance of the date of administration. The flyers make clear that participation is entirely voluntary. Local Institute for Social Research representatives and their assistants conduct the actual questionnaire administrations following standardized procedures detailed in an instruction manual. The questionnaires are administered in classrooms during a normal class period whenever possible; however, circumstances in some schools require the use of larger group administrations. Teachers are asked to remain present in the classroom to help maintain order, but to remain at their desks so that they cannot see students' answers.

Questionnaire Format

Because many questions are needed to cover all of the topic areas in the study, much of the questionnaire content for 12th graders is divided into six different questionnaire forms distributed to participants in an ordered sequence that ensures six virtually identical random subsamples. (Five questionnaire forms were used between 1975 and 1988.) About one third of each questionnaire form consists of key, or "core," variables common to all forms. All demographic variables, and nearly all of the drug *use* variables included in this report, are contained in this core set of measures. Many of the questions on attitudes, beliefs, and perceptions of relevant features of the social environment are in a single form only, and the data are thus based on one fifth as many cases in 1975–1988 (approximately 3,300 per year) and on one sixth as many cases beginning in 1989 (approximately 2,600 per year). All tables in this report list the sample sizes upon which the statistics are based, stated in terms of the weighted number of cases (which, as explained above, is roughly equivalent to the actual number of cases).

RESEARCH DESIGN AND PROCEDURES FOR THE EIGHTH- AND TENTH-GRADE SURVEYS

In 1991, the study was expanded to include nationally representative samples of 8th- and 10th-grade students surveyed on an annual basis. In general, the procedures used for the annual in-school surveys of 8th- and 10th-grade students closely parallel those used for 12th graders, including the procedures for selecting schools and students, questionnaire administration, and questionnaire formats. A major exception is that only two different questionnaire forms were used from 1991 to 1996, expanding to four forms beginning in 1997, rather than the six used for 12th graders. The 8th and 10th graders receive the same questionnaire forms; most of the content is drawn from the 12th-grade surveys, including the core section. Thus, key demographic variables and measures of drug use and related attitudes and beliefs are generally identical for all three grades. Many fewer questions about other values and attitudes are included in the 8th- and 10th-grade forms, in part because we think that many of them are likely to be more fully formed by 12th grade and, therefore, are best monitored there.

About 17,000 eighth-grade students in approximately 150 schools (mostly middle schools) and about 15,000 tenth-grade students in approximately 130 schools are surveyed each year (see Table 3-1).¹⁶

Mode of Administration

From 1991 to 1993, follow-ups for 8th and 10th graders were administered similarly to those for 12th graders (see Footnote 3). When follow-up surveys of new 8th- and 10th-grade cohorts were discontinued, the collection of personal identification information was no longer necessary. For confidentiality reasons, this personal information had been gathered on a tear-off sheet at the back of each questionnaire. We believed that there were potential advantages in moving toward a fully anonymous procedure for these grade levels, including the following: (a) school cooperation might be easier to obtain; (b) any suppression effect on self-reported substance use that the confidential mode of administration might have could be both eliminated and quantified; and (c) if there *were* any mode of administration effect, it would be removed from the national data, which are widely compared with results of state and local surveys (nearly all of which use anonymous questionnaires), thus making those comparisons more valid. Therefore, in 1998, the half-sample of schools beginning their two-year participation in MTF received fully anonymous questionnaires, while the half-sample participating in the study for their second and final year continued to get the confidential questionnaires. Since 1999, all questionnaires administered to 8th and 10th graders have been fully anonymous.

A careful examination of the 1998 results, based on the two equivalent half-samples at grades 8 and 10, revealed that there was no effect of this methodological change among 10th graders, and, at most, only a very modest effect in the self-reported substance use rates among 8th graders (with prevalence rates slightly higher in the anonymous condition). The net effect of this methodological change is a possible increase in the observed 8th-grade prevalence estimates for marijuana, alcohol, and cigarettes in 1998 from what they would have been without a change in questionnaire administration; thus, the declines in use in 1998 may be slightly understated for 8th graders. For example, the annual prevalence of marijuana use among 8th graders is shown to have fallen by 0.8 percentage points between 1997 and 1998; however, the half-sample of 8th-grade schools receiving exactly the same type of questionnaire that was used in 1997 showed a slightly greater decline of 1.5 percentage points.¹⁷

¹⁶The research design originally called for follow-up surveys of subsamples of the 8th- and 10th-grade participants, carried out at two-year intervals, similar to the 12th-grade follow-up samples. From 1991 to 1994, this plan influenced the design of the cross-sectional studies of 8th and 10th graders in an important way. In order to recapture many of the 8th-grade participants two years later in the normal 10th-grade cross-sectional study for that year, we selected the 8th-grade schools by drawing a sample of high schools and then selecting a sample of their feeder schools that contained 8th graders. This extra stage in the sampling process meant that many of the 8th-grade participants in, say, the 1991 cross-sectional survey were also participants as 10th graders in the 1993 cross-sectional survey. Thus, a fair amount of panel data was generated at no additional cost. However, having followed this design from 1991 through 1993, we concluded that the savings in follow-up costs did not justify the complexities in sampling, administration, and interpretation. Therefore, since 1994, we have used a simplified design in which 8th-grade schools are drawn independently of the 10th-grade school sample. Further follow-ups (at two-year intervals) have been conducted only on respondent panels drawn from the first three cohorts of 8th and 10th graders—that is, those surveyed in school in 1991, 1992, and 1993. A book reporting results from analyses of these panels was published recently: Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education–drug use connection: How successes and failures in school relate to adolescent smoking, drinking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

¹⁷We have examined in detail the effects of administration mode using multivariate controls to assess the effects of the change on 8th-grade self-report data. Our findings generally show even less effect than is to be found without such controls. See O'Malley, P. M., Johnston, L. D., Bachman, J. G., & Schulenberg, J. E. (2000). A comparison of confidential versus anonymous survey procedures: Effects on reporting of drug use and related attitudes and beliefs in a national study of students. *Journal of Drug Issues*, 30, 35–54.

All tables and figures in *Volume I* use data from both half-samples of 8th graders surveyed in a given year, combined. This is also true for 10th graders (for whom we found no methodological effect) and 12th graders (for whom it is assumed there is no such effect, since none was found among the 10th graders). (See this chapter's later section entitled "Representativeness and Sample Accuracy" under the subheading "School Participation," for a further discussion of half-samples among all three grades.)

Questionnaire Forms and Sample Proportions

Another benefit of not interlocking the 8th- and 10th-grade samples was that we could consider having more forms of the questionnaire. Beginning in 1997, the number of forms was expanded to four, but the four forms are not distributed in equal numbers. Forms 1, 2, 3, and 4 are assigned to one third, one third, one sixth, and one sixth of the students, respectively. Thus, if a question appears on only one form, it is administered to either one third or one sixth of the sample. A question in two forms may be assigned to one third of the sample (one sixth plus one sixth), one half of the sample (one third plus one sixth), or two thirds of the sample (one third plus one third). No questions appear on exactly three forms. Footnotes to the tables indicate what proportion of all respondents in each grade complete the question, if that proportion is other than the entire sample.

The two additional forms were introduced to allow for more questions. The new Forms 1 and 2 substantially follow the content of the previous Forms 1 and 2, but each is now assigned to a third of the sample instead of half. Form 3 builds on Form 1, with some questions omitted to make room for more content; and Form 4 builds on the content of Form 2 in a similar manner. Much of the new content was placed in both of the new forms (Forms 3 and 4), each of which is administered to one sixth of the sample, in order to assign one third of the total sample to those new measures.

RESEARCH DESIGN AND PROCEDURES FOR THE TWELFTH-GRADE FOLLOW-UP SURVEYS

Beginning with the graduating class of 1976, some members of each 12th-grade class have been selected to be surveyed by mail after high school graduation. From the roughly 13,000 to 17,000 twelfth graders originally participating in a given senior class, a representative sample of 2,400 individuals is randomly chosen for follow-up. In order to ensure sufficient numbers of drug users in the follow-up surveys, 12th graders reporting 20 or more occasions of marijuana use in the previous 30 days (i.e., "daily users"), or *any* use of the other illicit drugs in the previous 30 days, are selected with higher probability (by a factor of 3.0) than the remaining 12th graders. Differential weighting is then used in all follow-up analyses to compensate for these differential sampling probabilities. Because those in the drug-using stratum receive a weight of only 0.33 in the calculation of all statistics to correct for their overrepresentation at the selection stage, there are actually more follow-up respondents than are reported in the weighted *Ns* given in the tables.

The 2,400 participants selected from each 12th-grade class are randomly split into two groups of 1,200 each—one group to be surveyed on even-numbered calendar years, and the other group to be surveyed on odd-numbered years. This two-year cycle is intended to reduce the burden on

individual respondents, thus yielding a better retention rate across the years. By alternating the two half-samples, we have data from a given graduating class every year, even though any given respondent participates only every other year.

Until 2002, each respondent was surveyed biennially up to seven times; at the seventh follow-up, which would occur either 13 or 14 years after graduation, the respondents had reached modal age 31 or 32. Beginning in 2002, the seventh biennial follow-up was discontinued, and each respondent was surveyed every other year until modal age 29 or 30. Additional follow-ups occur at modal ages 35, 40, 45, and 50. Data like these, gathered on representative national samples over such a large portion of the life span, are extremely rare and can provide needed insight into the etiology and life-course history of substance use and related behaviors, including those related to the risk of HIV transmission.

Follow-Up Procedures

Using information provided by 12th-grade respondents on a tear-off card (containing the respondent's name, address, and phone number, and the name and address of someone who would always know how to reach them), mail contact is maintained with the subset of people selected for inclusion in the follow-up panels. Newsletters are sent to them each year, and name and address corrections are requested from both the U.S. Postal Service and the individual. Questionnaires are sent to each individual biennially in the spring. A check for \$20, made payable to the respondent, is attached to the front of each questionnaire.¹⁸ Reminder letters and postcards are sent at fixed intervals thereafter; finally, those who have not responded receive a prompting phone call from the Survey Research Center's phone interviewing facility in Ann Arbor, Michigan. If requested, a second copy of the questionnaire is sent; but no questionnaire content is administered by phone. If a respondent asks not to be contacted further, that wish is honored.

Follow-Up Questionnaire Format

The questionnaires used in the follow-up surveys of 19- to 30-year-olds parallel those used in 12th grade. Many of the questions are the same (including the core section on drug use, background, and demographics), and respondents are consistently mailed the same version (or form) of the questionnaire that they first received in 12th grade, so that *changes over time* in their behaviors, attitudes, experiences, and so forth can be measured. Questions specific to high school status and experiences are dropped in the follow-up, of course, and questions relevant to post-high school status and experiences are added. Thus, there are questions about college, military service, civilian employment, marriage, parenthood, and so on. Most of these are added to the core section. For the five-year surveys beginning at age 35, much of the questionnaire content is maintained, but streamlined (only one form is used) with more focus on the major family and work issues relevant to respondents ages 35 to 50; we have also added measures of substance use disorders.

¹⁸For the class of 1991 and all prior classes, the follow-up checks were for \$5. The rate was raised to \$10, beginning with the class of 1992, to compensate for the effects of inflation. An experiment was first conducted that suggested that the increased payment was justified based on the increased panel retention it achieved. Payment increased to \$20 in 2004 for the same reason.

For the early follow-up cohorts, the numbers of cases on single-form questions were one fifth the size of the total follow-up sample because five different questionnaire forms were used. Beginning with the class of 1989, a sixth form was introduced in 12th grade. That new questionnaire form was first sent to follow-up respondents in 1990. Single-form data since then have *N*s one sixth the total follow-up sample size. In the follow-up studies, single-form samples from a single cohort are too small to make reliable estimates; therefore, in most cases where they are reported, the data from several adjacent cohorts are combined.

REPRESENTATIVENESS AND SAMPLE ACCURACY

School Participation

Schools are invited to participate in the study for a two-year period. For each school that declines to participate, a similar school (in terms of size, geographic area, urbanicity, etc.) is recruited as a replacement. In 2008, either an original school or a replacement school was obtained in 96% of the sample units. With very few exceptions, each school participating in the first year has agreed to participate in the second year as well. Figure 3-2 provides the year-specific school participation rates and the percentage of units filled since 1977. As shown in the figure, replacements for declining schools are obtained in the vast majority of cases.

Two questions are sometimes raised with respect to school participation rates: (a) Are participation rates sufficient to ensure the representativeness of the sample? (b) Does variation in participation rates over time contribute to changes in estimates of drug use?

With respect to the first issue, the selection of replacement schools (which occurs in practically all instances of an original school refusal) almost entirely removes problems of bias in region, urbanicity, and the like that might result from certain schools refusing to participate. Other potential biases could be more subtle, however. If, for example, it turned out that most schools with “drug problems” refused to participate, the sample would be seriously biased. And if any other single factor were dominant in most refusals, that reason for refusal also might suggest a source of serious bias. However, the reasons given for a school refusing to participate tend to be varied and are often a function of happenstance specific to that particular year; only a very small proportion specifically object to the drug-related or “sensitive” nature of the survey content.

If it were the case that schools differed substantially in drug use, then which particular schools participated could have a greater effect on estimates of drug use. However, the great majority of variance in drug use lies within schools, not between schools.¹⁹ For example, between 1991 and 2002, the between-schools variance for 12-month marijuana use was 4.0–5.3% of the total variance for each of the three grades; for inhalant use, 1.6–2.7%; for cocaine use, 1.2–2.2%; for alcohol use, 3.5–6.1%; and for cigarette use, 2.1–5.2%. To the extent that schools tend to be fairly similar in drug use, which particular schools participate (within a selection framework that seeks national representation) has a small effect on estimates of drug use. The fact that the overwhelming majority of variance in drug use lies within schools implies that, at least with

¹⁹O'Malley, P. M., Johnston, L. D., Bachman, J. G., Schulenberg, J. E., & Kumar, R. (2006). How substance use differs among American secondary schools. *Prevention Science*, 7, 409–420.

respect to drug use, schools are for the most part fairly similar.²⁰ Further, some, if not most, of the between-schools variance is due to differences related to region, urbanicity, etc.—factors that remain well controlled in the present sampling design.

With respect to the second issue, it is extremely unlikely that results have been significantly affected by changes in school participation rates. If changes in participation rates seriously affected prevalence estimates, there would be noticeable bumps up or down in concert with the changing rates. But in fact this series of surveys produces results that are very smooth and change in an orderly fashion from one year to the next. Moreover, different substances trend in distinctly different ways, thus further refuting any likelihood that changes in participation rates are affecting prevalence estimates generally. We have observed, for example, marijuana use decreasing while cocaine use was stable (in the early 1980s), alcohol use declining while cigarette use was stable (in the mid- to late 1980s), and marijuana use increasing while inhalant use was decreasing (from 1994 to 1997). All of these patterns are explainable in terms of psychological, social, and cultural factors and cannot be explained by the common factor of changes in school participation rates.

Of course, there could be some sort of constant bias across the years; but even in the unlikely event that there is, it seems highly improbable that it would be of much consequence for policy purposes, given that it would not affect trends and likely would have a very modest effect on prevalence rates. Thus we have a high degree of confidence that school refusal rates have not seriously biased the survey results.

Nevertheless, securing the cooperation of schools has become more difficult in recent years. This is a problem common to the field, not specific to MTF. Therefore, beginning with the 2003 survey, we have provided payment to schools as a means of increasing their incentive to participate. (Several other ongoing school-based survey studies also use payments to schools.)

At each grade level, half of each year's sample comprises schools that started their participation the previous year, and half comprises schools that began participating in the current year. (Both samples are national replicates, meaning that each is drawn to be nationally representative by itself.) This staggered half-sample design is used to check on possible errors in the year-to-year trend estimates due to school turnover. For example, separate sets of one-year trend estimates are computed based on students in the half-sample of schools that participated in both 2006 *and* 2007, then based on the students in the half-sample that participated in both 2007 *and* 2008, and so on. Thus, each one-year *matched half-sample* trend estimate derived in this way is based on a constant set of schools (about 65 in 12th grade, for example, over a given one-year interval). When the trend data derived from the matched half-sample (examined separately for each class of drugs) are compared with trends based on the total sample of schools, the results are usually highly similar, indicating that the trend estimates are affected little by school turnover or shifting

²⁰Among participating schools, there is very little difference in substance use rates between the schools that were original selections, taken as a set, and the schools that were replacements. Averaged over the years 1991 through 2000, for grades 8, 10, and 12 combined, the difference between original schools and replacement schools averaged 0.03% in the observed prevalence rates averaged across a number of drug use measures: two indexes of annual illicit drug use, the annual prevalence of each of the major illicit drug classes, and several measures of alcohol and cigarette use. For the individual drugs and drug indexes, the differences between the original and replacement schools, averaged across grades and years, fell within $\pm 0.9\%$.

participation rates. As would be expected, the *absolute* prevalence-of-use estimates for a given year are not as accurate using just the half-sample because the sample size is only half as large.

Student Participation

In 2008, completed questionnaires were obtained from 90% of all sampled students in 8th grade, 88% in 10th grade, and 79% in 12th grade (see Table 3-1 for response rates in earlier years). In most cases, students are missed due to absence from class at the time of data collection; for reasons of cost efficiency, we typically do not schedule special follow-up data collections for absent students. Because students with fairly high rates of absenteeism also report above-average rates of drug use, some degree of bias is introduced into the prevalence estimates by missing the absentees. Much of that bias could be corrected through the use of special weighting based on the reported absentee rates provided by the students who *did* respond; however, we decided not to use such a weighting procedure because the bias in overall drug use estimates was determined to be quite small *and* the necessary weighting procedures would have introduced greater sampling variance in the estimates.²¹ Appendix A in *Volume I* illustrates the changes in trend and prevalence estimates that would result if corrections for absentees had been included. Of course, some students simply refuse, when asked, to complete a questionnaire. However, the proportion of explicit refusals amounts to less than 1.5% of the target sample for each grade.

Sampling Accuracy of the Estimates

Confidence intervals (95%) are provided in Tables 4-1a through 4-1d for lifetime, annual, 30-day, and daily prevalence of use for 8th-, 10th-, and 12th-grade students. As can be seen in Table 4-1a, confidence intervals for lifetime prevalence for 12th graders average less than $\pm 1.5\%$ across a variety of drug classes. That is, if we took a large number of samples of this size from the universe of all schools containing 12th graders in the coterminous United States, 95 times out of 100 the sample would yield a result that would be less than 1.5 percentage points divergent from the result we would get from a comparable massive survey of *all* 12th graders in *all* schools. This is a high level of sampling accuracy, permitting detection of fairly small changes from one year to the next. Confidence intervals for the other prevalence periods (last 12 months, last 30 days, and current daily use) are generally smaller than those for lifetime use. In general, confidence intervals for 8th and 10th graders are very similar to those observed for 12th graders. Some drugs (smokeless tobacco, PCP, nitrites, and others, as indicated in the footnotes for Tables 2-1 to 2-4) are measured on only one or two questionnaire forms; these drugs will have somewhat larger confidence intervals due to their smaller sample sizes. Appendix C of *Volume I* provides information on how to calculate confidence intervals around other point estimates, as well as information needed to compare trends across time or to test the significance of differences between subgroups in any given year.

²¹See appendix A in the following publication for a discussion of this point: Johnston, L. D., O'Malley, P. M., & Bachman, J. G. (1984). *Drugs and American high school students: 1975–1983* (DHHS (ADM) 85-1374). Washington, DC: U.S. Government Printing Office.

PANEL RETENTION

We discuss here the nature of the panel attrition problem generally, the response rates for MTF panel surveys in recent years, and evidence relevant to assessing the impact of attrition on the study's research results.

The Problem of Panel Attrition

Virtually all longitudinal studies of drug use—including MTF—experience attrition, which is often differential with respect to substance use.²² In addition, survey response rates in general have been declining over the past few decades,²³ highlighting an important challenge in the conduct of population-based research.

A vital feature of the MTF panel studies is their very low cost per respondent. There are many advantages to collecting panel data through low-cost mail surveys, as we have done since the outset of the study. Indeed, given the number of questionnaires sent each year (roughly 18,000) across the entire coterminous United States, using low-cost mail surveys is our best (and really the only) cost-effective option. One disadvantage of this mode of data collection is that attrition rates tend to be higher than those that might be obtained with much more expensive methods, such as intensive personal tracking and interviewing. Certainly there exist a few large epidemiological/etiological surveys that have better retention rates, but their procedures are extremely expensive and not realistic for an ongoing effort like this one. Nevertheless, our retention rates compare favorably with those of most longitudinal studies (including interview studies) reported in the field.

Response Rates

The series of survey data on American college students now encompasses 29 years. We know about our respondents' actual college attendance only from those who are invited and do complete follow-up questionnaires. We can use 12th-grade questionnaire answers (i.e., college intentions and program of study) to predict college attendance with a high degree of accuracy. The study's retention of college-bound 12th graders remains quite good. Among those participants in high school who were targeted for follow-up, *and* who reported planning to attend college and being enrolled in a college-prep curriculum, the follow-up retention rates for the three most recent classes surveyed at each follow-up point were: 61% in the first follow-up, one to two years past high school (based on the classes of 2005–2007); 59% in the second follow-up, three to four years past high school (based on the classes of 2003–2005); and 56% in the third follow-up, five to six years past high school (based on the classes of 2001–2003). These rates compare quite favorably with another national survey of substance use among college students, the Harvard College Alcohol Study, which had cross-sectional response rates of 59% in 1997 and 1999, and 52% in 2001.²⁴ To date in *Volume II*, we have reported only on college students who are one to four years past high school graduation. As the average age of attendance rises,

²²McGuigan, K. A., Ellickson, P. L., Hays, R. D., & Bell, R. M. (1997). Adjusting for attrition in school-based samples: Bias, precision, and cost trade-off of three methods. *Evaluation Review*, 21, 554–567.

²³Groves, R. M., Dillman, D. A., Eltinge, J. L., & Little, R. J. A. (Eds.) (2002). *Survey nonresponse*. New York: Wiley.

²⁴Wechsler, H., Lee, J. E., Kuo, M., Seibring, M., Nelson, T. F., & Lee, H. (2002). Trends in college binge drinking during a period of increased prevention efforts: Findings from 4 Harvard School of Public Health College Alcohol Study surveys: 1993–2001. *Journal of American College Health*, 50, 203–217.

having the extended age coverage will be of growing importance. Retention rates in the biennial follow-ups of *all* panel members modal ages 19–30 (corresponding to the first six follow-ups) decline with the length of the follow-up interval, of course. For the five surveys from 2004 to 2008, the response rate in the first follow-up (corresponding to one to two years past high school) averaged 56%; and for the second through sixth follow-ups (corresponding to 3–12 years past high school) response rates averaged 52%. Among long-term respondents—the 35-, 40-, 45-, and 50-year-olds—the retention rates are quite good, apparently because some of the decline with age in retention rates reflects cohort differences. Among the 35-year-old respondents surveyed from 2004 to 2008, corresponding to 17 years past high school, the average response rate was 48%. Among the 40-year-old respondents surveyed from 2004 to 2008, corresponding to a 22-year follow-up interval, the average retention rate was 53%. Among 45-year-olds surveyed in 2004 to 2008, the average retention rate was 59%. In sum, the response rates attained under the current design range from respectable to quite good, especially when the low-cost nature of the procedures, the long time intervals, and the substantial length of the questionnaires are taken into account. More importantly, the evidence leaves us confident that the data resulting from these follow-up panels are reasonably accurate, which brings us to our adjustments for panel attrition and the comparison of our results with those from other sources.

The Impact of Panel Attrition on Research Results

An important purpose of the MTF follow-ups is to allow estimation of drug prevalence rates among American high school graduates at various age levels. Thus, we have always been concerned about making the appropriate adjustments to account for panel attrition. In essence, our standard adjustment process is a poststratification procedure in which we reweight the data obtained from the follow-up samples so that their reweighted 12th-grade distribution on a given drug reproduces the original distribution of use observed for that drug based on all participating seniors. This procedure is carried out separately for cigarettes, alcohol, and marijuana, as well as other illicit drugs (combined). As expected, this produces prevalence estimates that are somewhat higher than those uncorrected for attrition, indicating that there is indeed some positive association between drug use and panel attrition. However, the adjustments are relatively modest, as is documented next.

One reason the adjustments are modest is that attrition rates do not differ greatly by levels of 12th-grade substance use; they differ some, but less than one might expect. For example, among all respondents who had never used marijuana, an average of 79% of the classes of 1976–1998 participated in the first follow-up. The proportion responding is somewhat lower among those who had used marijuana once or twice in the last 12 months: 75%. This proportion decreases gradually with increasing levels of marijuana use; but even among those who used marijuana on 20 or more occasions in the last 30 days in 12th grade, 67% participated in the first follow-up. The corresponding participation rates for the same drug use strata at the fourth follow-up (i.e., at modal ages 25–26) were 66%, 63%, and 56%, respectively. Thus, even among those who were quite heavy users of marijuana in high school, response rates at the fourth follow-up were only 10 percentage points lower than among those who had never used marijuana by 12th grade. That is not to say that we assume all types of drug users remain in the panels at comparably high rates. We believe that people who become dependent on or addicted to heroin or cocaine are unlikely to be retained in reasonable proportions. That is why we are careful not to quantify or characterize these special segments of the population. But we note that they constitute very low

proportions of the drug-using portion of the population, and even lower proportions of the entire adult population.

The National Survey on Drug Use and Health (NSDUH) seems to provide the best available data against which to validate the estimates generated for adult age groups in MTF because it is also based on national samples, but uses cross-sectional surveys that do not carry the burden of panel attrition. Their results, of course, may be affected by their own nonresponse rates; but that will be true of any comparison survey. The overall response rate for the NSDUH in 2006 was 74%.

In some earlier analyses, we compared the prevalence rates on a set of drugs—cigarettes, alcohol, marijuana, and cocaine—for which there was reasonable similarity in question wording across the two studies. The comparisons that follow are for the age group 19–28 in the MTF panel data, and for 19–28 (or 19–29 for 1999 only) in the NSDUH cross-sectional data. We used the most recent readily available comparable data—2004, but similar results are found in a number of prior years. Other things equal, NSDUH should have higher rates than MTF because it includes school dropouts. Nevertheless, the MTF estimates for the 30-day prevalence of marijuana are actually higher (14.4% without poststratification and 16.5% with it) than the NSDUH estimate (14.0%). The same is true for the 12-month cocaine prevalence estimate (6.3% without poststratification and 7.1% with it, vs. 6.3% in the NSDUH estimate). The higher MTF estimates for both marijuana and cocaine suggest that attrition does not produce substantially lower estimates of drug use than would be obtained if response rates were higher.

Comparisons for alcohol and cigarettes show larger differences, with alcohol use consistently higher in MTF and cigarette use consistently higher in NSDUH. We believe it likely that both are due to definitional differences in the exact question wording. In 2004, MTF estimates of 30-day alcohol prevalence were 67.8% and 68.4% (with poststratification) versus 62.6% in NSDUH. For cigarettes, the 30-day MTF prevalence estimates were 27.1% and 29.2%, respectively, versus 39.4% in NSDUH. (Because cigarette smoking rates are particularly high among dropouts, some of this difference should be explainable by differences in the populations covered by the two studies.) It is worth noting that the nature and magnitude of the differences between MTF and NSDUH estimates tend to be quite consistent for each of the four drugs at least as far back as 1992.

Even with attrition, substantial proportions of recent drug users remain in the MTF follow-up samples. In recent years, about 15–17% of the 19–28-year-old respondents reported marijuana use in just the prior 30 days, and about 5–7% reported past 12-month use of cocaine. These proportions and the underlying numbers of actual cases are quite adequate for analytic purposes.

An important point worth emphasizing here is that in the MTF panel, attrition is not necessarily as great a problem as in a cross-sectional study. This is because we already know a great deal about each of the follow-up nonrespondents, including their substance use, based on a lengthy questionnaire in senior year (and, for many, in subsequent years as well). Thus, adjustments can be made utilizing data that are highly informative about the missing individuals.

Effects on Relational Analyses

While differential attrition (uncorrected) may contribute to some bias in point estimates and other univariate statistics, such attrition tends to have less influence on bivariate and multivariate

statistics. This was found to be true in a secondary analysis of data from seven panel studies that followed adolescents over time,²⁵ and we have found this to be true in MTF panel analyses²⁶ and in analyses with other panel data sets.²⁷ Thus, differential attrition may be of less concern in multivariate panel analyses focused on understanding the course, causes, and consequences of substance use. Still, as we summarized above, correcting for attrition can be important, and we continue to do it.

VALIDITY OF THE MEASURES OF SELF-REPORTED DRUG USE

Are sensitive behaviors such as drug use honestly reported? Like most studies dealing with sensitive behaviors, we have no direct, totally objective validation of the present measures; however, the considerable amount of existing inferential evidence strongly suggests that the self-report questions used in MTF produce largely valid data. A more complete discussion of the contributing evidence leading to this conclusion may be found in other publications.²⁸ Here we only briefly summarize the evidence.

First, using a three-wave panel design, we established that the various measures of self-reported drug use have a high degree of reliability—a necessary condition for validity.²⁹ In essence, respondents were highly consistent in their self-reported behaviors over a three- to four-year time interval. Second, we found a high degree of consistency among logically related measures of use within the same questionnaire administration. Third, the proportion of 12th graders reporting some illicit drug use by senior year has reached two thirds of all respondents in peak years and over 80% in some follow-up years, constituting *prima facie* evidence that the degree of underreporting must be very limited. Fourth, 12th graders' reports of use by their unnamed friends—about whom they would presumably have considerably less reason to conceal information about use—have been highly consistent with self-reported use in the aggregate, in terms of both prevalence *and* trends in prevalence, as will be discussed in chapter 9, *Volume I*.

²⁵Cordray, S., & Polk, K. (1983). The implication of respondent loss in panel studies of deviant behavior. *Journal of Research in Crime and Delinquency*, 20, 214–242.

²⁶Bryant, A. L., Schulenberg, J. E., Bachman, J. G., O'Malley, P. M., & Johnston, L. D. (2000). Understanding the links among school misbehavior, academic achievement, and cigarette use: A national panel study of adolescents. *Prevention Science*, 1(2), 71–87; Schulenberg, J. E., Bachman, J. G., O'Malley, P. M., & Johnston, L. D. (1994). High school educational success and subsequent substance use: A panel analysis following adolescents into young adulthood. *Journal of Health and Social Behavior*, 35, 45–62.

²⁷Bachman, J. G., O'Malley, P. M., & Johnston, J. (1978). *Youth in Transition: Vol. 6. Adolescence to adulthood: A study of change and stability in the lives of young men*. Ann Arbor, MI: Institute for Social Research; Schulenberg, J. E., Bryant, A. L., & O'Malley, P. M. (2004). Taking hold of some kind of life: How developmental tasks relate to trajectories of well-being during the transition to adulthood. *Development and Psychopathology*, 16, 1119–1140.

²⁸Johnston, L. D., & O'Malley, P. M. (1985). Issues of validity and population coverage in student surveys of drug use. In B. A. Rouse, N. J. Kozel, & L. G. Richards (Eds.), *Self-report methods of estimating drug use: Meeting current challenges to validity* (NIDA Research Monograph No. 57 (ADM) 85-1402). Washington, DC: U.S. Government Printing Office; Johnston, L. D., O'Malley, P. M., & Bachman, J. G. (1984). *Drugs and American high school students: 1975–1983* (DHHS (ADM) 85-1374). Washington, DC: U.S. Government Printing Office; Wallace, J. M., Jr., & Bachman, J. G. (1993). Validity of self-reports in student-based studies on minority populations: Issues and concerns. In M. de LaRosa (Ed.), *Drug abuse among minority youth: Advances in research and methodology* (NIDA Research Monograph No. 130). Rockville, MD: National Institute on Drug Abuse.

²⁹O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (1983). Reliability and consistency in self-reports of drug use. *International Journal of the Addictions*, 18, 805–824.

Fifth, we have found self-reported drug use to relate in consistent and expected ways to a number of other attitudes, behaviors, beliefs, and social situations—in other words, there is strong evidence of construct validity. Sixth, the missing data rates for the self-reported use questions are only very slightly higher than for the preceding nonsensitive questions, in spite of explicit instructions to respondents immediately preceding the drug section to leave blank those drug use questions they felt they could not answer honestly. Seventh, an examination of consistency in reporting of lifetime use conducted on the long-term panels of graduating seniors found quite low levels of recanting of earlier reported use of the illegal drugs.³⁰ There was a higher level of recanting for the psychotherapeutic drugs, suggesting that adolescents may actually overestimate their use of some drugs because of misinformation about definitions that is corrected as they get older. Finally, the great majority of respondents, when asked, say they would answer such questions honestly if they were users.³¹

This is not to argue that self-reported measures of drug use are valid in all cases. In the present study we have gone to great lengths to create a situation and set of procedures in which students recognize that their confidentiality will be protected. We have also tried to present a convincing case as to why such research is needed. The evidence suggests that a high level of validity has been obtained. Nevertheless, insofar as any remaining reporting bias exists, we believe it to be in the direction of underreporting. Thus, with the possible exception of the psychotherapeutic drugs, we believe our estimates to be lower than their true values, even for the obtained samples, but not substantially so.

As an additional step to assure the validity of the data, we check for logical inconsistencies in the triplets of answers about the use of each drug (i.e., about lifetime, annual, and past 30-day use), and if a respondent exceeds a minimum number of inconsistencies across the drug use questions, his or her record is deleted from the data set. Similarly, we check for improbably high rates of use of multiple drugs and delete such cases, on the assumption that the respondents are not taking the task seriously. Fortunately, relatively few cases have to be eliminated for these reasons.

Consistency and the Measurement of Trends

MTF is designed to be sensitive to changes from one time period to another. A great strength of this study, in our opinion, is that the measures and procedures have been standardized and applied consistently across many years. To the extent that any biases remain because of limits in school and/or student participation, and to the extent that there are distortions (lack of validity) in the responses of some students, it seems very likely that such problems will exist in much the same proportions from one year to the next. In other words, biases in the survey estimates will tend to be consistent from one year to another, meaning that our measurement of *trends* should be affected very little. The smooth and consistent nature of most trend curves reported for the various drugs provides rather compelling empirical support for this assertion.

³⁰Johnston, L. D., & O'Malley, P. M. (1997). The recanting of earlier reported drug use by young adults. In L. Harrison (Ed.), *The validity of self-reported drug use: Improving the accuracy of survey estimates* (NIDA Research Monograph No. 167, pp. 59–80). Rockville, MD: National Institute on Drug Abuse.


³¹For a discussion of reliability and validity of student self-report measures of drug use like those used in MTF across varied cultural settings, see Johnston, L. D., Driessen, F. M. H. M., & Kokkevi, A. (1994). *Surveying student drug misuse: A six-country pilot study*. Strasbourg, France: Council of Europe.

TABLE 3-1
Sample Sizes and Response Rates

Grade:	Number of Public Schools			Number of Private Schools			Total Number of Schools				Total Number of Students				Student Response Rate (%)		
	8th	10th	12th	8th	10th	12th	8th	10th	12th	Total	8th	10th	12th	Total	8th	10th	12th
1975	—	—	111	—	—	14	—	—	125	—	—	—	15,791	—	—	—	78
1976	—	—	108	—	—	15	—	—	123	—	—	—	16,678	—	—	—	77
1977	—	—	108	—	—	16	—	—	124	—	—	—	18,436	—	—	—	79
1978	—	—	111	—	—	20	—	—	131	—	—	—	18,924	—	—	—	83
1979	—	—	111	—	—	20	—	—	131	—	—	—	16,662	—	—	—	82
1980	—	—	107	—	—	20	—	—	127	—	—	—	16,524	—	—	—	82
1981	—	—	109	—	—	19	—	—	128	—	—	—	18,267	—	—	—	81
1982	—	—	116	—	—	21	—	—	137	—	—	—	18,348	—	—	—	83
1983	—	—	112	—	—	22	—	—	134	—	—	—	16,947	—	—	—	84
1984	—	—	117	—	—	17	—	—	134	—	—	—	16,499	—	—	—	83
1985	—	—	115	—	—	17	—	—	132	—	—	—	16,502	—	—	—	84
1986	—	—	113	—	—	16	—	—	129	—	—	—	15,713	—	—	—	83
1987	—	—	117	—	—	18	—	—	135	—	—	—	16,843	—	—	—	84
1988	—	—	113	—	—	19	—	—	132	—	—	—	16,795	—	—	—	83
1989	—	—	111	—	—	22	—	—	133	—	—	—	17,142	—	—	—	86
1990	—	—	114	—	—	23	—	—	137	—	—	—	15,676	—	—	—	86
1991	131	107	117	31	14	19	162	121	136	419	17,844	14,996	15,483	48,323	90	87	83
1992	133	106	120	26	19	18	159	125	138	422	19,015	14,997	16,251	50,263	90	88	84
1993	126	111	121	30	17	18	156	128	139	423	18,820	15,516	16,763	51,099	90	86	84
1994	116	116	119	34	14	20	150	130	139	419	17,708	16,080	15,929	49,717	89	88	84
1995	118	117	120	34	22	24	152	139	144	435	17,929	17,285	15,876	51,090	89	87	84
1996	122	113	118	30	20	21	152	133	139	424	18,368	15,873	14,824	49,065	91	87	83
1997	125	113	125	27	18	21	152	131	146	429	19,066	15,778	15,963	50,807	89	86	83
1998	122	110	124	27	19	20	149	129	144	422	18,667	15,419	15,780	49,866	88	87	82
1999	120	117	124	30	23	19	150	140	143	433	17,287	13,885	14,056	45,228	87	85	83
2000	125	121	116	31	24	18	156	145	134	435	17,311	14,576	13,286	45,173	89	86	83
2001	125	117	117	28	20	17	153	137	134	424	16,756	14,286	13,304	44,346	90	88	82
2002	115	113	102	26	20	18	141	133	120	394	15,489	14,683	13,544	43,716	91	85	83
2003	117	109	103	24	20	19	141	129	122	392	17,023	16,244	15,200	48,467	89	88	83
2004	120	111	109	27	20	19	147	131	128	406	17,413	16,839	15,222	49,474	89	88	82
2005	119	107	108	27	20	21	146	127	129	402	17,258	16,711	15,378	49,347	90	88	82
2006	122	105	116	29	18	20	151	123	136	410	17,026	16,620	14,814	48,460	91	88	83
2007	119	103	111	32	17	21	151	120	132	403	16,495	16,398	15,132	48,025	91	88	81
2008	116	103	103	28	19	17	144	122	120	386	16,253	15,518	14,577	46,348	90	88	79


Source. The Monitoring the Future study, the University of Michigan.

FIGURE 3-1
Schools included in 1 Year's Data Collection
8th, 10th, and 12th Grades



Source. The Monitoring the Future study, the University of Michigan.
Note. One dot equals one school.

FIGURE 3-2
School Participation Rates*



Percent of slots filled by...	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05	'06	'07	'08
Original	59	63	62	63	71	71	66	72	72	67	66	72	71	68	70	59	55	60	52	53	51	51	57	62	56	49	53	62	63	59	58	53
Replacements	39	36	35	32	25	26	32	26	29	33	26	26	30	29	29	39	43	39	44	44	43	47	42	35	42	48	37	34	40	39	43	
Total	98	99	97	95	96	97	99	98	96	99	99	98	99	99	99	98	98	99	96	96	98	99	99	97	98	97	98	99	97	99	97	96

Source. The Monitoring the Future study, the University of Michigan.
 *The data for the years prior to 1991 are for 12th grade only; beginning in 1991, the data are for 8th, 10th, and 12th grades combined.

Chapter 4

PREVALENCE OF DRUG USE IN EARLY AND MIDDLE ADULTHOOD

Longitudinal panel studies such as those included in MTF are most typically used to examine maturational changes with age, by tracking the same individuals across years. But another potential use is to look across multiple cohorts in a given year in order to get a cross-age snapshot of the prevalence of using various drugs in that year. Such snapshots are examined in this chapter. We will see that use tends to be higher in the early post-high school age groups, corresponding to the new freedoms associated with leaving high school and often moving away from the parental home.³² But sometimes there are also strong cohort effects that are reflected in differences among age groups at a given point in time; in this chapter we will see evidence of both types of effects.

Estimates of drug use in the adult population are most often generated through household survey interviews of cross-sections of the general population. In the present study, our estimates come from self-completed mail questionnaires from respondents in the follow-up surveys. These are representative samples of previous classes of high school students who started their participation in MTF in their senior year. As described in more detail in chapter 3, MTF has conducted ongoing panel studies on representative samples from each graduating high school senior class beginning with the class of 1976. From each class, two matched subpanels of roughly 1,200 students each are randomly selected to comprise long-term follow-up panels—one of these two panels is surveyed every even-numbered year after graduation, and the other is surveyed every odd-numbered year, up through age 30. So, while each cohort participates every year, each respondent participates only every other year. This alternating panel design was chosen to reduce the repetitiveness of participating in the panel study, because the questionnaire remains the same between ages 19 and 30. Thus, in a given year, the study encompasses one of the two panels from each of the last 12 senior classes previously participating in MTF.³³ In 2008, representative samples of the classes of 1996 through 2007—ages 19–30—were surveyed, using the standard young adult survey instruments. For brevity, we refer to this 19- through 30-year-old age group as “young adults” in this chapter.

To build on these important national panels of young adults, we have extended the surveys into middle adulthood. The middle adulthood surveys are conducted at modal age 35 (that is, 17 years after high school graduation) and at five-year intervals thereafter. In these later follow-ups, both of the half panels from the relevant graduating class are surveyed using a single questionnaire form (one for age 35, a different one for ages 40 and 45, and another for age 50) instead of six

³²Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates; see also Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education-drug use connection: How successes and failures in school relate to adolescent smoking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

³³Through 2001, the follow-ups also included modal ages 31 and 32. This seventh follow-up was dropped in 2002 because we believed that the costs were no longer justified by the marginal benefits of having these follow-up data, given that an age-35 survey is being conducted. Throughout the time between surveys, we send a newsletter to respondents in order to help maintain contact.

forms. The results of the 2008 follow-up surveys should accurately characterize approximately 87% of all young adults ages 19–30 as well as about 88% of adults ages 35, 40, 45 and 50). The remaining 12% or so—the high school dropout segment—were missing from the senior year surveys and, as a result, were missing from all of the follow-up surveys as well. Thus, the results presented here are not necessarily generalizable to that small segment of the population.

Figures 4-1 through 4-21 contain 2008 *prevalence* data by age, corresponding to respondents ages 19–30, as well as 35-, 40-, 45-, and 50-year-olds. For comparison purposes, data are also included for the 2008 high school senior class, listed as 18-year-olds. Figures provided in chapter 5 contain the *trend* data for each age group derived from the repeated cross-sectional surveys, including 12th graders and high school graduates through age 50. In the figures in chapters 4 and 5, age groups spanning the young adult years have been paired into two-year intervals in order to increase the number of cases, and thus the precision, for each point estimate. The data for ages 35, 40, 45, and 50 are, of necessity, based on a single age in each case. As indicated above, *both* half samples from a given class cohort are included in each year's samples of 35-, 40-, 45-, and 50-year-olds, so in 2008 the two half samples come from the graduating classes of 1991, 1986, 1981, and 1976, respectively. Their respective weighted numbers of cases are 901, 899, 875, and 897. (Actual unweighted numbers are somewhat higher.)

It is worth noting that the pattern of age-related differences showing up in any one year can be checked in an adjacent year (i.e., the previous year's volume or the succeeding year's) for replicability, because two nonoverlapping half samples of follow-up respondents in the 19-to-30 age band are surveyed on alternating years. In the case of the 35-, 40-, 45-, and 50-year-olds, two entirely different graduating classes make up the samples for any two adjacent years.

A NOTE ON ADJUSTED LIFETIME PREVALENCE ESTIMATES

In Figures 4-1 through 4-21, two different estimates of *lifetime* prevalence are provided. One estimate is based on the respondent's *most recent* statement of whether he or she ever used the drug in question (the light gray bar). The other estimate takes into account the respondent's answers regarding lifetime use gathered in *all* of the previous data collections in which he or she participated (the white bar). *To be categorized as one who has used the drug based on all past answers regarding that drug, the respondent must have reported either lifetime use in the most recent data collection and/or some use in his or her lifetime on at least two earlier data collections.* Because respondents of ages 18 through 20 cannot have their responses adjusted on the basis of two earlier data collections, adjusted prevalence rates are reported only for ages 21 and up. Most other epidemiological studies can present only an unadjusted estimate because they have data from a single cross-sectional survey. An adjusted estimate of the type used here is possible only when panel data have been gathered so that a respondent can be classified as having used a drug at some time in his or her life, based on earlier answers, even though he or she no longer indicates lifetime use in the most recent survey.

The divergence of these two estimates as a function of age shows that there is more inconsistency as time passes. Obviously, there is more opportunity for inconsistency as the number of data collections increases. Our judgment is that “the truth” lies somewhere between the two estimates: the lower estimate may be depressed by tendencies to forget, forgive, or

conceal earlier use, and the upper estimate may include earlier response errors or incorrect definitions of drugs that respondents appropriately corrected in later surveys as they became more knowledgeable. It should be noted that a fair proportion of those giving inconsistent answers across time had earlier reported having used the given drug only once or twice in their lifetime.

As we have reported elsewhere, the cross-time stability of self-reported usage measures, taking into account both prevalence and frequency of self-reported use, is still very high.³⁴ Note that the divergence between the two lifetime prevalence estimates is greatest for the psychotherapeutic drugs and for the derivative index of “use of an illicit drug other than marijuana,” which is heavily affected by the psychotherapeutic estimates. We believe this is due to respondents having greater difficulty accurately categorizing psychotherapeutic drugs (usually taken in pill form) with a high degree of certainty—especially if such a drug was used only once or twice. We expect higher inconsistency across time when the event—and in many of these cases, a single event—is reported with a relatively low degree of certainty at quite different points in time. Those who have gone beyond simple experimentation with one of these drugs would undoubtedly be able to categorize them with a higher degree of certainty. Also, those who have experimented more recently, in the past month or year, should have a higher probability of recall, as well as fresher information for accurately categorizing the drug.

We provide both estimates to make clear that a full use of respondent information provides a possible range for lifetime prevalence estimates, not a single point. However, by far the most important use of the prevalence data is to track *trends* in *current* (as opposed to lifetime) use. Thus, we are much less concerned about the nature of the variability in the lifetime estimates than we might otherwise be. The lifetime prevalence estimates are of importance primarily in showing the degree to which a drug class has penetrated the general population overall as well as particular cohorts; we believe that the evidence from the lifetime estimates suggests that cross-sectional surveys of adults are subject to underreporting, and that to a degree such underreporting increases with age, because adolescence and early adulthood are the periods in the life course during which most drug use occurs.³⁵

The reader is reminded that the reweighting procedures used to correct the panel data for the effects of panel attrition are described in chapter 3.

PREVALENCE OF DRUG USE AS A FUNCTION OF AGE

Figures 4-1 through 4-21 provide 2008 prevalence rates for each class of drugs, covering respondents ages 18 to 50. For virtually all drugs, available age comparisons show much higher

³⁴O’Malley, P. M., Bachman, J. G., & Johnston, L. D. (1983). Reliability and consistency in self-reports of drug use. *International Journal of the Addictions*, 18, 805–824.

³⁵For a more detailed analysis and discussion, see Johnston, L. D., & O’Malley, P. M. (1997). The recanting of earlier-reported drug use by young adults. In L. Harrison & A. Hughes (Eds.), *The validity of self-reported drug use: Improving the accuracy of survey estimates* (NIDA Research Monograph No. 97-4147). Washington, DC: National Institute on Drug Abuse.

lifetime prevalence for the older age groups, as would be expected. In fact, the figures reach surprisingly high levels among adults in their early 30s through their 50s.

- *The adjusted lifetime prevalence figures are most striking for today's 50-year-olds (the class of 1976), who were passing through adolescence near the peak of the drug epidemic. Some 86% reported trying an illicit drug (lifetime prevalence, adjusted), leaving only 14% or about one in every seven who reported never having done so (see Figure 4-1). Some 79% of 50-year-olds said they had tried marijuana, and almost three quarters (73%) said they had tried some other illicit drug, including 46% who have tried cocaine specifically. The adjusted lifetime prevalences for 45-year-olds (the class of 1981) are similar to 50-year-olds. Clearly, the parents of today's teenagers and young adults are themselves a very drug-experienced generation.*
- In 2008, the adjusted lifetime prevalence figures among 29- to 30-year-olds reach 71% for *any illicit drug*, 66% for *marijuana*, 48% for *any illicit drug other than marijuana*, and 21% for *cocaine*. Put another way, even among young Americans who graduated from high school in 1996 and 1997—after the peak of the larger drug epidemic—less than a third (29%) have *never* tried an illegal drug.

The 2008 survey responses, *unadjusted* for previous answers, show somewhat lower lifetime prevalence for 29- to 30-year-olds: 66% for *any illicit drug*, 64% for *marijuana*, 39% for *any illicit drug other than marijuana*, and 19% for *cocaine*.

- Despite the higher lifetime prevalence rates among older age groups, these groups generally show *annual* or *30-day* prevalence rates that are no higher than those of today's 12th graders. Indeed, for a number of drugs, the levels reported by older respondents are lower—sometimes considerably lower—suggesting that the incidence of quitting more than offsets the incidence of initiating use of these drugs during the years after high school.

In analyses published elsewhere, we looked closely at patterns of change in drug use with age and identified post-high school experiences that contribute to declining levels of annual or current use of drugs as respondents grow older. For example, the likelihood of marriage increases with age, and we have found that marriage is consistently associated with declines in *alcohol* use, *heavy drinking*, *marijuana* use, and *cocaine* use, and most likely just about all of the other illicit drugs as well.³⁶

³⁶Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates; and Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Bryant, A. L., & Merline, A. C. (2002). *The decline of substance use in young adulthood: Changes in social activities, roles, and beliefs*. Mahwah, NJ: Lawrence Erlbaum Associates. See also Schulenberg, J. E., O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (2000). "Spread your wings and fly": The course of well-being and substance use during the transition to young adulthood. In L. J. Crockett & R. K. Silbereisen (Eds.), *Negotiating adolescence in times of social change* (pp. 224–255). New York: Cambridge University Press. And see O'Malley, P. M., Bachman, J. G., Johnston, L. D., & Schulenberg, J. E. (2004). Studying the transition from youth to adulthood: Impacts on substance use and abuse. In J. S. House, F. T. Juster, R. L. Kahn, H. Schuman, & E. Singer (Eds.), *A telescope on society: Survey research and social science at the University of Michigan and beyond* (pp. 305–329). Ann Arbor, MI: The University of Michigan Press.

- For use of **any illicit drug** (Figure 4-1), lifetime prevalence (unadjusted) is 66% among 29- to 30-year-olds versus 47% among 2008 twelfth graders. Annual prevalence, however, is highest among younger respondents (12th graders and 21- to 22-year-olds at 37%) with progressively lower rates among the older age groups, reaching 28% among 29- to 30-year-olds. Current (30-day) prevalence shows much the same pattern, with 12th graders having the highest rate (22%) and the rate generally declining with each age band, reaching 16% among 29- to 30-year-olds.
- Among 35-, 40-, 45-, and 50-year-olds, lifetime prevalence rates for **marijuana**, **any illicit drug**, and **any illicit drug other than marijuana** generally increase with each age level. (This is also true for many other illicit drugs.) However, annual and 30-day prevalence rates generally decline slightly across age levels from 18 to 35, and then remain fairly level through age 50 (the oldest age covered by MTF so far). Thus, it is clear that greater proportions of the older cohorts have discontinued use.
- Among young adults, a similar pattern exists for **marijuana**: a higher lifetime prevalence as a function of age, but considerably lower annual and 30-day prevalence rates through the late 20s (Figure 4-3). After age 35 there is little difference across age bands. Current **daily marijuana** use shows the least variation across age (as shown in Figure 5-3c). Still, in 2008 it ranges from 5.4% among 12th graders down to 4.1% among 19- to 20-year-olds, to 4.8% among 29- to 30-year-olds, then down to 2.0% among 50-year-olds. Clearly, most respondents who were daily users in their teenage years are no longer daily users. Note again the evidence of a cohort effect, in that the 50-year-olds had a higher rate of marijuana use when they were high school seniors (1976) than did those who are now 35 years old (class of 1991).
- Statistics on the use of **any illicit drug other than marijuana** (Figure 4-2) have a somewhat similar pattern. Like marijuana and the any-illicit-drug-use index, adjusted lifetime rates on this index also show an appreciable rise with age, reaching 48% among the 29- to 30-year-old age group and 73% among 50-year-olds. In other words, almost half of today's 30-year-olds have tried some illicit drug other than marijuana, and almost three quarters of all 50-year-olds have done so. Current use shows a slight decline across age bands, ranging from 9% among 12th graders to 4% among 40-year-olds. After ages 23–24, annual use is generally lower with increased age of the respondent. A number of the individual drugs that comprise this general category show lower rates of use at higher ages for annual prevalence, usually with the highest rate observed at ages 18–20. This is particularly true for **amphetamines**, **hallucinogens**, **LSD** specifically, **inhalants**, and **sedatives (barbiturates)**. The falloff with age is not as great nor as consistent for **cocaine**, **crack**, **other cocaine**, **crystal methamphetamine (ice)**, **heroin**, **narcotics other than heroin**, **tranquilizers**, and **ecstasy (MDMA)**, though in general, usage rates are somewhat lower among those in their 30s than among those in their early 20s. Several classes of drugs are discussed individually next.
- **Inhalants** show some very interesting differences across age strata (see Figure 4-13). There is little difference across age in contemporaneously reported lifetime prevalence, but a considerable difference in the lifetime prevalence figure *adjusted* for previous

reporting of use. The adjusted pattern—an increase with age—is the one we have come to expect, and we believe is the more accurate one. Annual prevalence rates drop off with age, while 30-day rates begin fairly low and can drop only a little. Clearly, use of inhalants is extremely low beyond about age 20, and we know from data presented in *Volume I* that much of the decline in use with age has already occurred by the time young people have reached 12th grade. Questions on inhalant use are not included in the surveys administered to respondents over the age of 30, given the extremely low rates of use reported by age 30.

- For ***amphetamines***, lifetime prevalence is again much higher among the older age groups—reflecting in part the addition of new users who initiate use in their 20s, but also reflecting some cohort differences (Figure 4-4). (There is also a considerable divergence between the corrected lifetime prevalence versus the contemporaneously reported lifetime prevalence, as is true for most psychotherapeutic drugs.) However, more recent use, as reflected in the annual prevalence figure, is considerably lower among the older age groups. This has not always been true; the present pattern reflects a sharper historic decline in use among older respondents than has occurred among 12th graders, as well as cohort differences in having ever used these drugs. These trends are discussed in the next chapter.
- ***Ritalin***, an amphetamine widely prescribed for the treatment of attention deficit hyperactivity disorder (ADHD), shows a drop-off in annual prevalence of use (outside of medical supervision) after ages 23–24 (2.9%) to ages 25–26 (2.4%) and 27–28 (0.7%), consistent with the interpretation that it is often used by college students and perhaps graduate students trying to stay awake and alert for studying and completing assignments (see Table 4-3).
- Questions on the use of ***methamphetamine*** are contained in two of the six questionnaire forms, making estimates less reliable than those based on all six forms. Annual prevalence of methamphetamine use does not decline with age until the late 20s, with annual prevalence at 1% from ages 18–26. Past 30-day use does decline from very low up to age 20, and is then negligible after that (Table 4-3 and Figure 4-5).
- ***Crystal methamphetamine (ice)*** is also included on only two questionnaire forms through age 30 (and is not asked of 35-, 40-, 45-, or 50-year-old respondents). Among the 19- to 30-year-old respondents *combined*, 0.7% reported some use in the prior year—lower than the 1.1% reported by 12th graders (see Table 4-3 and Figure 4-6). Generally, those above age 24 have lower annual prevalence rates than those who are younger.
- ***Sedatives (barbiturates)*** show lifetime prevalence rates that are fairly similar across the age band 23 through 40, but are appreciably higher among 45- and 50-year-olds. *Annual* use falls only very gradually with age from age 18 to 25–26, but then remains fairly constant at 3–4% through age 50 (Figure 4-14). At present, current usage rates are quite low at all ages (highest at age 18 at 2.8%). Because of the substantial long-term decline in

sedative (barbiturate) use over the life of MTF, the 50-year-olds have, by far, the highest adjusted lifetime prevalence rate (32%).³⁷

- Annual prevalence of *narcotics other than heroin* is highest in the 18-to-28 age range (at 8% to 10%), compared to 3% among 45- and 50-year-olds. Thirty-day prevalence shows a somewhat similar profile.
- *Tranquilizer* use shows an increase with age in lifetime prevalence through age 28, then some falloff through age 35, with the highest rates among those over 35—again, with the oldest cohorts having much higher levels of lifetime use. On the other hand, there is some modest decrease with age in annual prevalence after age 26. Thirty-day prevalence is 3% among 18- to 26-year-olds, and 1% to 2% thereafter (Figure 4-16).
- *Cocaine* had generally presented a unique case among the illicit drugs, in that lifetime, annual, and current prevalence rates have *all* tended to rise with age into the 20s. By 1994, however, 30-day cocaine use had reached such low levels that it varied rather little by age. Following the resurgence of cocaine use in the 1990s, some differences by age in annual prevalence emerged, though there are still rather few differences for current prevalence (Figure 4-7). Annual prevalence is now highest among respondents ages 19 through 28. The cohort differences in lifetime cocaine use are particularly vivid, with the 45- and 50-year-olds showing 44% and 46% adjusted lifetime prevalence rates, respectively, compared to 13% among 21- to 22-year-olds in 2008. Very few (around 1%) of the 40- to 50-year-olds today are current users of cocaine.
- In 2008, lifetime prevalence of *crack* use (Figure 4-8) rises fairly steadily between the age-19 stratum and the 28–35 strata (to 7–8%); it is higher among 45- and 50-year-olds (10–12%), reflecting something of a cohort effect due to the rather transient popularity of crack in the early to mid-1980s, and a brief resurgence in the mid-1990s. Current prevalence is 0.8% or below in all age groups. Annual prevalence is highest among 18-year-olds at 1.6% and lower among those above age 28. We believe that the omission of high school dropouts is likely to have a greater than average impact on the prevalence estimates for crack. It also seems likely that any panel respondents who become dependent on crack (or other illicit drugs like heroin) would be less likely than average to respond to the questionnaires; therefore, such extreme users are no doubt underrepresented among the panel respondents.
- *Ecstasy (MDMA)* was added to two of the six forms of the follow-up surveys in 1989 to assess how widespread its use had become among young adults. It was added to a third form in 2002. Questions about its use were not asked of high school students until 1996, primarily because of concern that its alluring name might stimulate interest in the drug.

³⁷Barbiturates were the dominant form of sedatives in use when these questions were first introduced. In the intervening years, a number of nonbarbiturate sedatives have entered the market and largely displaced barbiturates. We believe that a number of users of nonbarbiturate sedatives are reporting them in answer to this question, which also defines them in terms of the conditions for which they are prescribed. In recognition of this fact, we now label them as “sedatives (barbiturates).” The rewording of the question was made in half of the questionnaire forms in 2004 and in the other half in 2005.

That concern diminished considerably after the drug had become more widely known. (Ecstasy use is not asked of respondents 35 years of age and older.)

In 2008, among all 19- to 30-year-olds combined, 14% say they have tried *ecstasy*; among 12th graders, 6% say they have used it. The age differences in ecstasy use are quite dramatic, with lifetime prevalence increasing sharply with age and highest (21%) at ages 27–30 (see Figure 4-17). This very likely reflects the rapid rise in ecstasy use between 1997 and 2001, and the subsequent sharp decline. Annual prevalence is highest among 18- to 22-year-olds (at 4–5%), and lower among 23- to 30-year-olds (at 2–3%). Past-month ecstasy use is now at 1% or less for all age groups between 19 and 30.

- All *alcohol* prevalence rates are higher among young adults than among 12th graders, and they generally increase for the first three to five years after high school, through age 23 or 24 (Figures 4-20a and 4-20b). Prevalence rates vary only modestly among the older age groups. Lifetime prevalence changes very little after ages 23 to 24, due in large part to a “ceiling effect.” Current (30-day) alcohol use is considerably higher among those ages 21–22 (74%) than among 12th graders (43%); it drops some through the age strata, reaching 64% at age 50. Current *daily drinking* (Figure 4-20b) is also slightly higher and level among those 21 to 30 years old (5–6%) versus among those 18–20 (2–3%). The highest rate of daily drinking is among 50-year-olds (11%).
- Among the various measures of alcohol consumption, *occasional heavy drinking* (i.e., having five or more drinks in a row on at least one occasion in the two weeks prior to the survey) shows some differences by age (Figure 4-20b). There is a large difference between 18-year-olds (25%) and 21- to 24-year-olds, who have the highest prevalence of such heavy drinking (42%). Then there is a falloff at each subsequent age level above age 24, dropping to 20% by age 50. We have interpreted this curvilinear relationship as reflecting an age effect—not a cohort effect—because it seems to replicate across different graduating class cohorts, and also because it has been linked directly to age-related events such as leaving the parental home (which increases heavy drinking) and marriage (which decreases it), both of which are, in turn, related to attending college.³⁸ Among those at ages 40 to 50, about one fifth (20–22%) report such heavy drinking in the prior two-week interval.
- *Cigarette smoking* also shows an unusual pattern of age-related differences (Figure 4-21). Current (30-day) smoking rates used to be about the same among those in their early 20s and 12th graders, partly because most initiation of cigarette use happens in high school. Beginning in 2005, however, current smoking was somewhat lower among 12th graders, almost surely due to the sharp drop in smoking that has been occurring among secondary school students—a cohort effect. Smoking at heavier levels—such as smoking half a pack daily—is (and has been) higher among those in their 20s than among 12th graders, as many light or moderate smokers in high school move into a pattern of heavier

³⁸O’Malley, P. M., Bachman, J. G., & Johnston, L. D. (1988). Period, age, and cohort effects on substance use among young Americans: A decade of change, 1976–1986. *American Journal of Public Health*, 78, 1315–1321. See also Bachman, J. G., Wadsworth, K. N., O’Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates.

use after high school.³⁹ About a quarter (26%) of current 12th-grade smokers use at the rate of a half pack per day or more, compared to over one half (51%) of current smokers ages 29 to 30, and about three quarters (74%) of 50-year-old current smokers. Half-pack-per-day smoking rises to 12% by ages 23–24 and is virtually flat up to age 45. At age 50 it is slightly higher at 14%. But annual smoking is highest at ages 23–28 and then decreases with age, suggesting that many former occasional smokers quit as they get older.

- Questions on *steroid* use (Figure 4-18) were added to one form in 1989 and to an additional form in 1990, making it difficult to determine age-related differences with much accuracy due to limited sample sizes. (Steroid questions are not asked of 35- to 50-year-old respondents.) Overall, 1.9% of 19- to 30-year-olds in 2008 reported having used steroids in their lifetime. Annual and 30-day usage levels were very low, at 0.4% and 0.2%, respectively (see Tables 4-2 through 4-4). Rates among 12th graders tend to be higher than among older age groups, possibly reflecting both age and cohort effects. (As described in *Volume I*, the prevalence of steroid use among 12th graders rose sharply between 2000 and 2002 and then fell. At present, the highest annual use among young adults is among 19- to 20-year-olds, at 0.9%.)

In sum, lifetime prevalence rates in some of the older age groups studied here, who passed through adolescence in the heyday of the drug epidemic, show impressively high lifetime rates of illicit drug use—particularly when lifetime prevalence is corrected for the recanting of earlier reported use. However, *current* use of most illicit drugs is substantially lower among those over age 30 than among those in their late teens to early 20s. For the two licit drugs, alcohol and cigarettes, the picture is different, with less falloff in active use with age.

PREVALENCE COMPARISONS FOR SUBGROUPS OF YOUNG ADULTS

Subgroup differences for 19- to 30-year-olds are presented in Tables 4-1 through 4-5. While Table 4-1 provides only gender differences, the remaining tables have prevalence rates by gender, age, region of the country, and population density. Each of these subgroup dimensions is discussed separately below.

Gender Differences

In general, most of the gender differences in drug use that were observed in high school students may be found in the young adult (19- to 30-year-old) sample as well. See Table 4-1 for the full set of gender comparisons.

³⁹Because age is confounded with class cohort, and because we have established that cigarette smoking shows strong cohort effects (enduring differences among cohorts), one must be careful in interpreting age-related differences in a cross-sectional sample as if they were due only to age effects—that is, changes with age consistently observable across cohorts. However, multivariate analyses conducted on MTF panel data from multiple cohorts do show a consistent age effect of the type mentioned here (see O'Malley, Bachman, & Johnston, 1988, in previous footnote).

- Among young adults, more males than females report using **any illicit drug** during the prior year (37% vs. 30%). Males have higher annual prevalence rates for nearly all illicit drugs—with ratios of two or greater for **inhalants**, **LSD**, **hallucinogens other than LSD**, **PCP**, **heroin (with and without using a needle)**, **GHB**, **Ketamine**, and **steroids**. For example, among 19- to 30-year-olds, **LSD** was used by 1.8% of males versus 0.9% of females during the prior 12 months (see Table 4-1).
- All three measures of **cocaine** showed higher rates of use by males than females (19- to 30-year-olds). Annual **cocaine** use was reported by 8.0% of males and 4.4% of females; **other cocaine** use by 7.4% of males and 4.1% of females; and **crack** use by 1.0% of males and 0.8% of females.
- Other large gender differences among 19- to 30-year-olds are found in **daily marijuana** use (7.4% for males vs. 3.5% for females), **daily alcohol** use (7.5% vs. 3.7%), and occasions of drinking **five or more drinks in a row** in the prior two weeks (47% vs. 31%). This gender difference in occasions of heavy drinking is larger among young adults than among 12th graders, where it is 28% for males versus 21% for females.
- **Ecstasy (MDMA)** use is only modestly higher among males than among females in the young adult sample overall (annual prevalence of 3.8% vs. 2.6%, respectively).
- Annual prevalence of use of **narcotics other than heroin** outside of medical supervision is at 10.3% for males versus 7.7% for females. Use of **Vicodin**, one of the most widely used drugs in the class, is 11.1% versus 7.7%, respectively. There is a larger proportional gender difference for **OxyContin** (4.9% for males vs. 2.7% for females).
- The use of **amphetamines**, which is now about equivalent among males and females in high school, is also fairly similar for both genders in this post-high school period (annual prevalence of 5.5% vs. 4.4%, respectively).
- In the 1980s, there were few differences between males and females in rate of **cigarette** use. By the early 1990s, however, males had slightly higher rates of use. In 2008, among 19- to 30-year-olds, males are more likely to have smoked in the past month (29% vs. 22%), to have smoked daily (19% vs. 15%), and to have smoked half a pack or more per day (12.5% vs. 9.3%). This gender difference reflects a cohort effect in which the differences between the genders in high school during the 1990s are then carried up the age spectrum.
- **Steroid** use among young adults is much more prevalent among males than females, as is true for 12th graders. Among 12th graders, 2.5% of males reported steroid use in the past year versus 0.4% of females. These statistics are much lower among 19- to 30-year-olds, but use by males remains considerably higher (0.7% for males vs. 0.2% for females).

Regional Differences

Follow-up respondents are asked in what state they currently reside. States are then grouped into the same regions used in the analysis of high school data.⁴⁰ Tables 4-2 through 4-5 present regional differences in lifetime, annual, 30-day, and current daily prevalence for 19- to 30-year-olds combined.

- There exist some regional differences in the use of *marijuana*, with the Northeast and West somewhat higher than the South and Midwest. The Northeast and West are also slightly higher in the proportion using *any illicit drug* and *any illicit drug other than marijuana* (see Table 4-3).
- *Methamphetamine* use is highest in the West (annual prevalence of 1.5%), and there is not much difference among the other regions: the Midwest and the South are both at 0.8%, and the Northeast is at 0.6% (see Table 4-3).
- The use of *crystal methamphetamine (ice)* by 19- to 30-year-olds is at low levels in all regions, with the Northeast having 0.3% and all other regions at 0.8%.
- The West and Northeast continue to have higher rates than the other two regions for *hallucinogen* use, though the regional differences are not large.
- For *ecstasy (MDMA)*, annual prevalence is lower in the Northeast (1.7%) than in the other three regions (2.8% to 3.9%).
- *OxyContin* use is highest in the West (4.6%) and Midwest (4.3%) followed by the South (3.0%) and the Northeast (2.6%).
- Use of the club drugs *Rohypnol* and *GHB* appears to be higher in the West and next highest in the Midwest, though annual prevalence rates are very low in all regions (Table 4-3). *Ketamine* has a higher annual prevalence than the other two, and shows very little variation by region.
- The use of *steroids* is highest in the Northeast, though again the prevalence rates are very low and the differences between regions do not always replicate from year to year.
- For the remaining illicit drugs, regional differences are not substantial (see Tables 4-3 and 4-4).
- Prevalence rates for *alcohol* use are typically somewhat higher in the Northeast and Midwest regions than in the South and West; this pattern has generally been true among

⁴⁰States are grouped into regions as follows: *Northeast*—Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, and Pennsylvania; *Midwest*—Ohio, Indiana, Illinois, Michigan, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas; *South*—Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, Florida, Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, and Texas; *West*—Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada, Washington, Oregon, and California.

12th graders as well. For *binge drinking* among 19- to 30-year-olds, the Northeast and Midwest have a prevalence rate of 41%, whereas the West and South have rates of 34% and 33%. Self-reported *drunkenness* shows a similar pattern, as would be expected.

- As with alcohol, *cigarette smoking* among young adults is highest in the Midwest and Northeast, which have 30-day prevalence rates of 27% and 26%, respectively. It is lower in the South (24%) and particularly low in the West (20%). This difference is most pronounced at the current half-pack-a-day level (Table 4-5), where the rate in the West (6.0%) is about half the rate in the Midwest (12.9%). The South (11.1%) and Northeast (10.5%) are intermediate.

Population Density Differences

Population density is measured by asking respondents to select the response category that best describes the size and nature of the community where they lived during March of the year in which they were completing the follow-up questionnaire. The various categories are listed in Tables 4-2 through 4-5; the population sizes given to the respondent to help define each level are provided in a footnote to each table. An examination of the 1987 and 1988 drug use data for the two most urban strata revealed that the modest differences in prevalence rates between the suburbs and the corresponding cities were not worth the complexity of reporting them separately; accordingly, since then these categories have been merged to increase sample sizes. See Tables 4-2 through 4-5 for the tabular results discussed below.

- Differences in illicit drug use by population density tend to be very modest, perhaps more modest than is commonly supposed. Among the general population, use of most illicit drugs is broadly distributed among all areas from rural to urban. To the extent that there are variations, almost all of the associations are positive, with rural/country areas having the lowest levels of use, and small towns having the next lowest. Medium-sized cities, large cities, and very large cities tend to be higher, with only small variations among them. Positive associations with population density exist for annual prevalence of *any illicit drug*, *marijuana*, and *cocaine*. The association is strongest for cocaine, where the annual prevalence rate in the very large cities (8.1%) is almost twice that in small towns (4.4%).
- In 2008, annual prevalence rates of *methamphetamine* and *crystal methamphetamine (ice)* use are highest in small towns and very large cities (Table 4-3).
- *Amphetamine* use has a weak positive association with population density in 2008.
- Among young adults, the lifetime and annual *alcohol* use measures all show a slight positive association with population density, while 30-day use has a somewhat stronger positive association, with 56% of the farm/country stratum reporting use in the prior 30 days versus 78% of those in very large cities. *Occasions of heavy drinking* are positively associated with urbanicity as well (see Table 4-5). *Daily alcohol use* falls between 4.3% and 6.7% for all community size strata, again with a positive association with population density.

- Contrary to what we find for almost all other substances, there exists a negative association between population density and ***daily cigarette smoking***, which is highest in the rural/country stratum and lowest in the very large cities (daily prevalence rates of 21% and 13%, respectively). Smoking at the half-pack-a-day level is more than twice as high in the farm/country stratum (16%) as in very large cities (7%) (see Table 4-5).

TABLE 4-1
Prevalence of Use of Various Types of Drugs by Gender
among Respondents of Modal Ages 19–30, 2008
 (Entries are percentages.)

<i>Approximate Weighted N =</i>	<u>Males</u>	<u>Females</u>	<u>Total</u>
Any Illicit Drug ^a	2,400	3,500	5,900
Annual	36.8	30.3	32.9
30-Day	22.6	16.2	18.8
Any Illicit Drug ^a other than Marijuana			
Annual	21.0	16.9	18.5
30-Day	10.0	7.7	8.6
Marijuana			
Annual	32.5	24.3	27.6
30-Day	19.8	12.6	15.5
Daily ^p	7.4	3.5	5.0
Inhalants ^c			
Annual	1.9	0.8	1.3
30-Day	0.8	0.2	0.4
Hallucinogens			
Annual	5.6	2.4	3.7
30-Day	1.5	0.4	0.8
LSD			
Annual	1.8	0.9	1.3
30-Day	0.5	0.2	0.3
Hallucinogens other than LSD			
Annual	5.1	2.0	3.2
30-Day	1.3	0.2	0.7
PCP ^d			
Annual	0.5	0.2	0.3
30-Day	0.2	0.1	0.1
Ecstasy (MDMA) ^c			
Annual	3.8	2.6	3.1
30-Day	0.8	0.4	0.6
Cocaine			
Annual	8.0	4.4	5.8
30-Day	2.4	1.5	1.9
Crack ^e			
Annual	1.0	0.8	0.9
30-Day	0.4	0.4	0.4
Other Cocaine ^f			
Annual	7.4	4.1	5.4
30-Day	1.9	1.4	1.6
Heroin			
Annual	0.6	0.4	0.5
30-Day	0.1	0.1	0.1
With a Needle ^g			
Annual	0.2	0.1	0.1
30-Day	0.1	*	*
Without a Needle ^g			
Annual	0.7	0.1	0.4
30-Day	0.1	0.0	*

(Table continued on next page.)

TABLE 4-1 (cont.)
Prevalence of Use of Various Types of Drugs by Gender
among Respondents of Modal Ages 19–30, 2008
(Entries are percentages.)

	<u>Males</u>	<u>Females</u>	<u>Total</u>
<i>Approximate Weighted N =</i>	2,400	3,500	5,900
Narcotics other than Heroin ^h			
Annual	10.3	7.7	8.8
30-Day	4.0	3.2	3.5
Amphetamines, Adjusted ^{h,i}			
Annual	5.5	4.4	4.9
30-Day	2.4	1.8	2.0
Methamphetamine ^g			
Annual	0.6	1.1	0.9
30-Day	0.2	0.3	0.3
Crystal Methamphetamine (Ice) ^g			
Annual	0.7	0.7	0.7
30-Day	0.2	0.3	0.2
Sedatives (Barbiturates) ^h			
Annual	4.8	4.1	4.4
30-Day	2.0	1.7	1.8
Tranquilizers ^h			
Annual	7.6	6.2	6.7
30-Day	2.9	2.3	2.6
Alcohol			
Annual	84.3	84.3	84.3
30-Day	73.1	67.3	69.7
Daily ^p	7.5	3.7	5.2
5+ Drinks in a Row in Last 2 Weeks	46.5	30.5	37.0
10+ Drinks in a Row in Last 2 Weeks ^{d,j}	24.0	4.3	12.3
15+ Drinks in a Row in Last 2 Weeks ^{d,k}	9.1	1.1	4.4
Been Drunk ^c			
Annual	69.1	63.6	65.9
30-Day	45.8	36.0	40.0
Daily ^p	0.6	0.3	0.4
Flavored Alcoholic Beverages ^d			
Annual	48.5	62.4	56.9
30-Day	20.9	27.8	25.1
Cigarettes			
Annual	38.2	31.8	34.4
30-Day	28.5	21.8	24.5
Daily	19.2	15.3	16.9
1/2 Pack+/Day	12.5	9.3	10.6
Steroids ^g			
Annual	0.7	0.2	0.4
30-Day	0.3	0.1	0.2

Source. The Monitoring the Future study, the University of Michigan.

** indicates a prevalence rate of less than 0.05%.

(Table continued on next page.)

TABLE 4-1 (cont.)
Prevalence of Use of Various Types of Drugs by Gender
among Respondents of Modal Ages 19–30, 2008

^aUse of “any illicit drug” includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor’s orders.

^bDaily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes, for which actual daily use is measured, and for 5+ drinks, for which the prevalence of having five or more drinks in a row in the last two weeks is measured.

^cThis drug was asked about in three of the six questionnaire forms. Total *N* is approximately 3,000.

^dThis drug was asked about in one of the six questionnaire forms. Total *N* is approximately 1,000.

^eThis drug was asked about in five of the six questionnaire forms. Total *N* is approximately 5,000.

^fThis drug was asked about in four of the six questionnaire forms. Total *N* is approximately 3,900.

^gThis drug was asked about in two of the six questionnaire forms. Total *N* is approximately 2,000.

^hOnly drug use that was not under a doctor’s orders is included here.

ⁱBased on data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

^jErratum corrected on Nov. 10th, 2015. These are the correct values, rather than the 29.5%, 5.0%, and 14.7% previously contained here.

^kErratum corrected on Nov. 10th, 2015. These are the correct values, rather than the 11.2%, 1.3%, and 5.2% previously contained here.

TABLE 4-2
Lifetime Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2008

(Entries are percentages.)

	Approximate Weighted N	Any Illicit Drug ^a		Marijuana	Marijuana	Inhalants ^b	Hallucinogens		LSD	Hallucinogens other than LSD	PCP ^c	Ecstasy (MDMA) ^b		Cocaine	Crack ^d
		Any Illicit Drug ^a	other than Marijuana				Hallucinogens	LSD				MDMA	Cocaine		
Total	5,900	60.5	35.5	57.2	10.3	16.1	9.7	14.0	2.2	14.1	15.4	4.4			
Gender:															
Male	2,400	61.5	38.0	59.2	12.1	20.6	11.8	19.0	3.1	15.3	19.5	5.2			
Female	3,500	59.9	33.8	55.9	9.1	13.1	8.3	10.7	1.6	13.4	12.7	3.8			
Modal Age:															
19–20	1,100	49.2	24.8	45.9	6.6	8.0	3.8	7.5	1.7	7.4	8.5	2.3			
21–22	1,000	57.1	30.9	54.3	6.6	11.8	5.6	11.1	1.6	10.1	13.0	4.3			
23–24	1,000	62.8	37.9	58.8	12.0	15.4	7.3	13.9	1.9	12.9	15.2	4.3			
25–26	900	62.2	38.1	59.4	11.1	19.1	10.6	16.7	2.9	16.2	19.1	5.3			
27–28	900	66.7	43.2	62.4	11.4	20.6	14.0	16.6	2.9	19.6	18.8	5.5			
29–30	1,000	66.5	39.4	64.0	14.5	22.8	17.7	19.0	2.3	19.6	18.8	4.6			
Region:															
Northeast	1,100	64.5	36.8	62.6	10.5	17.3	10.4	15.1	3.3	16.0	16.4	3.5			
Midwest	1,600	60.3	34.7	56.9	10.2	15.7	9.5	13.5	1.8	11.9	14.7	4.8			
South	2,000	56.5	34.3	52.2	10.1	13.9	9.0	11.9	2.3	13.6	13.7	4.0			
West	1,200	63.4	37.4	60.7	10.3	19.7	10.9	17.4	1.4	16.5	18.7	5.2			
Population Density: ^e															
Farm/Country	700	53.1	32.4	49.0	10.9	13.0	8.7	10.5	2.4	9.6	13.5	5.2			
Small Town	1,500	56.4	33.3	53.1	9.9	14.7	9.0	13.2	3.1	13.0	14.2	6.0			
Medium City	1,400	61.7	34.8	58.6	10.7	14.8	9.2	13.0	2.3	11.4	13.5	3.6			
Large City	1,400	62.9	36.1	60.2	9.6	18.2	10.7	15.9	1.6	17.1	16.9	3.2			
Very Large City	900	67.1	40.9	63.8	10.9	19.8	11.2	17.1	1.4	20.8	19.2	3.3			

(Table continued on next page.)

TABLE 4-2 (cont.)
Lifetime Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2008

(Entries are percentages.)

	Approximate Weighted N	Other Cocaine ^f	Heroin	Heroin with a Needle ^g	Heroin without a Needle ^g	Narcotics other than Heroin ^h	Amphetamines ^{hi}	Methamphetamine ^g	Methamphetamine (Ice) ^g	Crystal Methamphetamine
Total	5,900	14.5	2.0	0.5	2.3	19.7	14.7	6.8	3.7	3.7
Gender:										
Male	2,400	18.1	2.8	0.8	3.6	22.4	15.9	8.0	4.0	4.0
Female	3,500	12.1	1.4	0.3	1.5	17.8	13.9	6.1	3.5	3.5
Modal Age:										
19–20	1,100	8.6	1.1	0.1	0.6	13.3	10.4	2.0	2.0	2.0
21–22	1,000	12.5	2.0	0.5	2.1	17.4	13.4	3.5	2.9	2.9
23–24	1,000	13.3	2.3	0.6	2.4	22.6	16.1	7.1	3.9	3.9
25–26	900	18.6	1.9	0.5	2.4	22.1	16.1	9.1	4.7	4.7
27–28	900	17.2	2.2	1.0	3.2	22.5	17.2	10.7	4.5	4.5
29–30	1,000	17.9	2.5	0.6	3.5	20.7	15.3	9.3	4.5	4.5
Region:										
Northeast	1,100	14.6	2.5	0.7	2.7	20.2	14.1	3.2	1.2	1.2
Midwest	1,600	13.8	1.5	0.4	1.8	20.0	15.1	7.5	3.7	3.7
South	2,000	13.3	2.0	0.3	2.5	18.3	15.0	5.7	4.0	4.0
West	1,200	17.8	2.3	0.8	2.5	20.3	14.2	11.0	6.0	6.0
Population Density: ^e										
Farm/Country	700	12.2	1.8	1.1	1.9	19.2	14.4	5.3	4.1	4.1
Small Town	1,500	13.9	2.5	0.2	2.7	19.0	14.7	7.7	4.9	4.9
Medium City	1,400	12.8	1.9	0.5	2.1	18.9	14.3	6.0	3.1	3.1
Large City	1,400	16.2	1.6	0.3	1.9	19.9	14.4	6.9	3.3	3.3
Very Large City	900	17.2	1.8	1.2	2.7	21.1	15.8	6.5	3.0	3.0

(Table continued on next page.)

TABLE 4-2 (cont.)
Lifetime Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2008

(Entries are percentages.)

	Approximate Weighted <i>N</i>	Sedatives (Barbiturates) ^b	Tranquilizers ^b	Alcohol	Been Drunk ^b	Flavored Alcoholic Beverages ^c	Cigarettes	Steroids ^d
Total	5,900	10.7	16.1	89.2	81.4	83.2	—	1.9
Gender:								
Male	2,400	12.1	17.8	88.1	81.3	77.7	—	4.2
Female	3,500	9.7	15.0	89.9	81.5	86.8	—	0.5
Modal Age:								
19–20	1,100	8.2	10.1	76.8	62.3	70.7	—	1.4
21–22	1,000	10.1	13.5	91.4	81.1	85.6	—	1.6
23–24	1,000	11.8	18.0	91.1	86.2	86.0	—	2.1
25–26	900	11.4	18.2	90.9	84.9	85.1	—	2.5
27–28	900	11.6	19.9	92.5	87.8	85.8	—	1.5
29–30	1,000	11.1	17.9	93.6	88.0	86.1	—	2.5
Region:								
Northeast	1,100	9.8	17.0	91.9	85.6	84.3	—	3.2
Midwest	1,600	10.3	13.9	91.5	85.7	88.3	—	1.9
South	2,000	11.4	18.2	87.3	77.5	80.7	—	1.5
West	1,200	10.6	14.8	86.0	78.4	78.5	—	1.5
Population Density: ^e								
Farm/Country	700	11.2	15.7	86.8	76.4	75.8	—	2.6
Small Town	1,500	10.7	15.6	86.7	80.0	83.2	—	1.4
Medium City	1,400	10.7	16.5	89.4	81.7	80.8	—	1.7
Large City	1,400	10.8	16.3	91.4	83.6	84.3	—	2.6
Very Large City	900	9.9	16.2	92.0	85.2	89.8	—	1.6

Source: The Monitoring the Future study, the University of Michigan.

“—” indicates data not available.

^aUse of “any illicit drug” includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor’s orders.

^bThis drug was asked about in three of the six questionnaire forms. Total *N* is approximately 3,000.

^cThis drug was asked about in one of the six questionnaire forms. Total *N* is approximately 1,000.

^dThis drug was asked about in five of the six questionnaire forms. Total *N* is approximately 5,000.

^eA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000–100,000; a large city as 100,000–500,000; and a very large city as having over 500,000. Within each level of population density, suburban and urban respondents are combined.

^fThis drug was asked about in four of the six questionnaire forms. Total *N* is approximately 3,900.

^gThis drug was asked about in two of the six questionnaire forms. Total *N* is approximately 2,000.

^hOnly drug use that was not under a doctor’s orders is included here.

ⁱBased on data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

TABLE 4-3
Annual Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2008

(Entries are percentages.)

	Approximate Weighted N	Any Illicit Drug ^a		Marijuana	Marijuana Inhalants ^b	Hallucinogens	LSD	Hallucinogens other than LSD	PCP ^c	Ecstasy (MDMA) ^b	Cocaine	Crack ^d
		Any Illicit Drug ^a	other than Marijuana									
Total	5,900	32.9	18.5	27.6	1.3	3.7	1.3	3.2	0.3	3.1	5.8	0.9
Gender:												
Male	2,400	36.8	21.0	32.5	1.9	5.6	1.8	5.1	0.5	3.8	8.0	1.0
Female	3,500	30.3	16.9	24.3	0.8	2.4	0.9	2.0	0.2	2.6	4.4	0.8
Modal Age:												
19–20	1,100	35.5	16.8	32.1	1.8	5.2	2.0	4.7	0.6	4.7	5.0	1.2
21–22	1,000	36.7	19.5	33.3	1.5	5.5	2.7	4.5	0.6	4.7	7.3	0.9
23–24	1,000	34.4	21.3	27.4	1.7	3.3	0.9	3.0	0.6	2.0	5.8	0.9
25–26	900	31.8	19.1	25.9	1.1	3.2	0.8	2.8	*	2.7	6.5	0.9
27–28	900	30.1	18.0	23.6	0.9	1.7	0.4	1.6	*	2.1	5.3	0.7
29–30	1,000	28.2	16.5	22.3	0.7	2.9	0.5	2.6	0.2	2.0	5.2	0.5
Region:												
Northeast	1,100	37.1	20.8	32.6	1.8	4.2	1.3	3.6	0.5	1.7	7.6	0.5
Midwest	1,600	30.5	17.3	25.7	0.9	3.0	1.0	2.6	0.4	2.8	5.1	0.7
South	2,000	30.6	18.2	24.4	1.3	3.4	1.2	3.0	0.4	3.4	4.8	1.0
West	1,200	35.6	18.6	30.6	1.2	4.4	1.7	4.0	*	3.9	6.8	0.9
Population Density: ^e												
Farm/Country	700	24.4	16.2	18.3	1.7	2.2	1.2	1.7	0.8	2.7	4.4	1.4
Small Town	1,500	29.6	17.0	25.0	1.3	4.1	1.3	3.6	0.1	2.8	5.3	1.2
Medium City	1,400	34.0	18.9	28.1	1.1	3.7	1.5	3.2	0.5	3.1	5.0	0.5
Large City	1,400	35.5	19.2	30.7	0.7	3.8	1.4	3.4	0.4	3.8	6.5	0.6
Very Large City	900	38.2	20.8	32.7	1.9	3.7	0.7	3.4	*	2.6	8.1	0.6

(Table continued on next page.)

TABLE 4-3 (cont.)
Annual Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2008

(Entries are percentages.)

	Approximate Weighted N	Heroin		Heroin with a Needle ^g		Heroin without a Needle ^g		Narcotics other than Heroin ^h		Amphetamines ^{hi}		Ritalin ^{gh}		Methamphetamine ^g	
		Other Cocaine ^f	Heroin	Heroin a Needle ^g	Heroin with a Needle ^g	Heroin without a Needle ^g	Heroin ^h	OxyContin ^{gh}	Vicodin ^{gh}	Amphetamines ^{hi}	Ritalin ^{gh}	Methamphetamine ^g			
Total	5,900	5.4	0.5	0.1	0.1	0.4	8.8	3.6	9.1	4.9	2.0	0.9			
Gender:															
Male	2,400	7.4	0.6	0.2	0.2	0.7	10.3	4.9	11.1	5.5	2.2	0.6			
Female	3,500	4.1	0.4	0.1	0.1	0.1	7.7	2.7	7.7	4.4	1.9	1.1			
Modal Age:															
19–20	1,100	5.0	0.6	0.1	0.1	0.2	8.6	4.8	8.9	5.9	3.0	0.9			
21–22	1,000	6.8	0.9	0.1	0.1	0.8	8.4	4.5	9.2	6.7	2.8	0.6			
23–24	1,000	4.3	0.6	0.3	0.3	0.7	10.5	3.4	9.8	5.7	2.9	1.2			
25–26	900	6.7	0.3	0.1	0.1	0.3	9.4	3.9	8.9	4.0	2.4	0.9			
27–28	900	5.0	0.2	*	*	*	8.6	2.6	8.8	4.1	0.7	1.5			
29–30	1,000	4.7	0.3	0.1	0.1	0.1	7.1	2.2	8.8	2.6	0.4	0.1			
Region:															
Northeast	1,100	6.2	0.7	0.2	0.2	0.3	9.4	2.6	6.7	5.0	2.7	0.6			
Midwest	1,600	4.6	0.3	*	*	*	8.6	4.3	11.8	5.0	2.2	0.8			
South	2,000	4.8	0.4	0.1	0.1	0.5	8.4	3.0	7.4	5.4	2.4	0.8			
West	1,200	6.8	0.7	0.4	0.4	0.7	9.1	4.6	11.2	3.6	0.6	1.5			
Population Density: ^e															
Farm/Country	700	3.8	0.5	*	*	*	9.8	3.9	11.0	4.4	1.9	1.0			
Small Town	1,500	5.1	0.7	*	*	0.4	8.6	3.3	9.3	4.6	2.2	1.3			
Medium City	1,400	4.5	0.6	0.2	0.2	0.8	8.2	2.4	8.3	4.8	2.3	0.4			
Large City	1,400	6.3	0.3	0.1	0.1	*	9.2	5.2	9.4	5.2	1.7	0.6			
Very Large City	900	7.0	0.3	0.3	0.3	0.3	8.7	2.7	8.3	5.3	2.0	1.3			

(Table continued on next page.)

TABLE 4-3 (cont.)
Annual Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2008

(Entries are percentages.)

	Approximate Weighted N	Crystal Methamphetamine (Ice) ^g	Sedatives (Barbiturates) ^h	Tranquilizers ^h	Rohypno ^g	GHB ^g	Ketamine ^g	Alcohol	Been Drunk ^k	Flavored Alcoholic Beverages ^c	Cigarettes	Steroids ^g
Total	5,900	0.7	4.4	6.7	0.2	0.2	0.5	84.3	65.9	56.9	34.4	0.4
Gender:												
Male	2,400	0.7	4.8	7.6	0.2	0.4	0.9	84.3	69.1	48.5	38.2	0.7
Female	3,500	0.7	4.1	6.2	0.2	0.1	0.2	84.3	63.6	62.4	31.8	0.2
Modal Age:												
19–20	1,100	0.7	5.4	7.1	*	0.3	*	72.3	54.4	55.8	33.6	0.9
21–22	1,000	0.2	4.9	6.3	0.4	0.1	0.7	88.6	71.3	63.1	35.6	0.1
23–24	1,000	1.4	5.0	8.1	0.1	0.5	0.4	86.6	68.7	64.7	37.1	0.3
25–26	900	0.8	4.0	6.7	0.1	0.4	0.7	86.4	71.4	52.7	34.5	0.3
27–28	900	0.7	4.1	5.7	0.5	0.1	0.2	84.7	65.1	55.2	34.2	0.3
29–30	1,000	0.3	2.9	6.5	*	*	0.8	87.8	64.9	49.4	31.3	0.3
Region:												
Northeast	1,100	0.3	3.8	7.2	0.1	0.1	0.6	88.1	70.9	55.7	34.3	0.9
Midwest	1,600	0.8	4.0	5.2	0.3	0.3	0.5	87.5	68.7	65.4	36.5	0.2
South	2,000	0.8	4.7	8.0	0.1	0.1	0.4	81.2	60.3	53.7	33.7	0.3
West	1,200	0.8	4.8	6.1	0.4	0.7	0.5	80.9	66.6	51.2	32.6	0.3
Population Density: ^e												
Farm/Country	700	0.6	4.6	6.6	0.6	*	0.2	76.3	51.1	53.6	33.8	0.2
Small Town	1,500	1.1	4.1	5.9	0.1	0.2	0.3	81.9	63.5	60.7	33.6	0.2
Medium City	1,400	0.1	4.5	6.9	0.3	0.1	0.5	83.9	68.2	54.6	35.6	0.2
Large City	1,400	0.5	5.0	7.9	0.1	0.2	1.0	88.7	71.0	58.0	34.6	1.0
Very Large City	900	1.3	3.3	5.8	0.1	0.8	0.2	88.9	70.2	53.6	33.2	0.3

Source: The Monitoring the Future study, the University of Michigan.

* indicates a prevalence rate of less than 0.05%.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in three of the six questionnaire forms. Total N is approximately 3,000.

^cThis drug was asked about in one of the six questionnaire forms. Total N is approximately 1,000.

^dThis drug was asked about in five of the six questionnaire forms. Total N is approximately 5,000.

^eA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000–100,000; a large city as 100,000–500,000; and a very large city as having over 500,000. Within each level of population density, suburban and urban respondents are combined.

^fThis drug was asked about in four of the six questionnaire forms. Total N is approximately 3,900.

^gThis drug was asked about in two of the six questionnaire forms. Total N is approximately 2,000.

^hOnly drug use that was not under a doctor's orders is included here.

ⁱBased on data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

TABLE 4-4
Thirty-Day Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2008
 (Entries are percentages.)

	Approximate Weighted N	Any Illicit Drug ^a		Marijuana	Marijuana	Inhalants ^b	Hallucinogens		Ecstasy (MDMA) ^b		Crack ^d	
		Any Illicit Drug ^a	other than Marijuana				LSD	Hallucinogens other than LSD	PCP ^c	Cocaine		
Total	5,900	18.8	8.6	15.5	0.4	0.8	0.3	0.7	0.1	0.6	1.9	0.4
Gender:												
Male	2,400	22.6	10.0	19.8	0.8	1.5	0.5	1.3	0.2	0.8	2.4	0.4
Female	3,500	16.2	7.7	12.6	0.2	0.4	0.2	0.2	0.1	0.4	1.5	0.4
Modal Age:												
19–20	1,100	20.2	8.3	17.9	0.6	1.5	0.7	1.2	*	1.1	1.7	0.5
21–22	1,000	20.8	10.0	17.8	0.3	1.3	0.5	1.1	*	1.0	2.0	0.2
23–24	1,000	20.7	9.9	16.2	0.8	0.6	0.3	0.5	0.6	0.5	2.1	0.5
25–26	900	17.3	8.5	13.3	0.3	0.5	0.3	0.3	*	0.2	2.3	0.5
27–28	900	17.2	7.5	14.2	0.2	0.2	*	0.2	*	0.3	1.6	0.3
29–30	1,000	16.3	7.5	12.9	0.2	0.6	0.1	0.6	0.2	0.5	1.8	0.2
Region:												
Northeast	1,100	21.6	9.5	18.9	0.4	0.9	0.2	0.8	*	0.4	3.0	0.3
Midwest	1,600	17.2	8.3	13.7	0.3	0.7	0.2	0.6	*	0.4	1.4	0.3
South	2,000	17.0	8.4	13.4	0.4	0.7	0.3	0.5	0.4	0.8	1.5	0.4
West	1,200	21.2	8.5	18.3	0.3	1.1	0.6	0.8	*	0.7	2.2	0.4
Population Density: ^e												
Farm/Country	700	15.1	8.2	11.2	0.9	0.7	0.4	0.6	0.8	0.4	1.8	0.8
Small Town	1,500	16.7	8.1	13.8	0.3	0.8	0.4	0.6	0.1	0.4	1.8	0.5
Medium City	1,400	19.4	8.8	15.9	0.3	1.0	0.3	0.9	*	1.0	1.4	0.3
Large City	1,400	20.4	8.6	17.5	0.3	0.9	0.3	0.7	*	0.7	1.9	0.2
Very Large City	900	21.5	9.6	17.7	0.4	0.5	0.2	0.4	*	0.4	2.9	0.1

(Table continued on next page.)

TABLE 4-4 (cont.)
Thirty-Day Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2008
 (Entries are percentages.)

	Approximate Weighted N	Other		Heroin		Heroin		Narcotics other than Heroin ^h		Amphetamines ^{h,i}		Methamphetamine ^g		Crystal Methamphetamine (Ice) ^g
		Cocaine ^f	Heroin	With Needle ^g	Without Needle ^g	Heroin	Heroin ^h	Amphetamines ^{h,i}	Methamphetamine ^g					
Total	5,900	1.6	0.1	*	*	*	*	3.5	2.0	0.3	0.2			
Gender:														
Male	2,400	1.9	0.1	0.1	0.1	0.1	0.1	4.0	2.4	0.2	0.2			
Female	3,500	1.4	0.1	*	*	*	*	3.2	1.8	0.3	0.3			
Modal Age:														
19–20	1,100	1.5	0.1	*	*	*	*	3.4	2.4	0.5	0.1			
21–22	1,000	2.2	0.2	0.1	0.1	0.1	0.1	3.7	3.1	*	*			
23–24	1,000	1.2	0.2	*	*	*	*	3.9	2.2	0.3	0.7			
25–26	900	2.2	*	*	*	*	*	3.4	1.3	0.1	0.1			
27–28	900	1.5	*	*	*	*	*	3.5	1.8	0.4	0.4			
29–30	1,000	0.9	*	0.1	*	*	*	3.4	1.3	0.1	0.1			
Region:														
Northeast	1,100	2.2	0.1	0.1	0.1	0.1	0.1	3.6	2.2	*	*			
Midwest	1,600	0.9	0.1	*	*	*	*	3.7	1.8	0.3	0.1			
South	2,000	1.5	0.1	0.1	0.1	*	*	3.4	2.6	0.3	0.4			
West	1,200	1.9	*	*	*	*	*	3.5	1.2	0.4	0.5			
Population Density: ^e														
Farm/Country	700	0.8	0.2	*	*	*	*	4.5	1.7	0.5	0.4			
Small Town	1,500	1.6	0.1	*	*	0.1	0.1	3.8	1.8	0.4	0.3			
Medium City	1,400	1.0	0.1	0.2	*	*	*	3.4	2.5	0.1	0.1			
Large City	1,400	1.8	0.1	*	*	*	*	3.0	1.9	*	*			
Very Large City	900	2.6	*	*	*	*	*	3.6	2.1	0.5	0.6			

(Table continued on next page.)

TABLE 4-4 (cont.)
Thirty-Day Prevalence of Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2008
 (Entries are percentages.)

	Approximate Weighted N	Sedatives (Barbiturates) ^b	Tranquilizers ^b	Alcohol	Been Drunk ^b	Flavored Alcoholic Beverages ^c	Cigarettes	Steroids ^d
Total	5,900	1.8	2.6	69.7	40.0	25.1	24.5	0.2
Gender:								
Male	2,400	2.0	2.9	73.1	45.8	20.9	28.5	0.3
Female	3,500	1.7	2.3	67.3	36.0	27.8	21.8	0.1
Modal Age:								
19–20	1,100	2.2	2.6	53.8	33.7	31.8	21.8	0.4
21–22	1,000	1.6	2.9	74.4	46.6	33.8	24.5	*
23–24	1,000	2.4	3.1	74.0	43.7	26.8	26.5	0.3
25–26	900	1.3	2.9	73.9	44.0	21.0	24.5	*
27–28	900	2.1	1.9	69.8	35.9	19.6	25.7	0.3
29–30	1,000	1.3	1.9	73.4	36.2	16.7	24.0	*
Region:								
Northeast	1,100	1.6	2.4	76.2	44.8	24.6	26.0	0.7
Midwest	1,600	1.7	2.0	72.4	45.0	29.1	26.8	*
South	2,000	2.1	3.1	65.6	34.4	23.8	24.4	0.1
West	1,200	1.9	2.5	65.7	37.9	23.5	19.7	*
Population Density: ^e								
Farm/Country	700	2.2	2.6	56.0	26.1	21.7	25.1	*
Small Town	1,500	1.5	2.6	66.1	38.0	24.4	24.5	0.1
Medium City	1,400	1.9	2.8	69.5	41.1	27.2	24.1	*
Large City	1,400	2.2	2.6	75.7	44.3	25.7	25.4	0.6
Very Large City	900	1.5	2.1	78.0	46.2	23.9	22.3	*

Source: The Monitoring the Future study, the University of Michigan.

*** indicates a prevalence rate of less than 0.05%.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders.

^bThis drug was asked about in three of the six questionnaire forms. Total N is approximately 3,000.

^cThis drug was asked about in one of the six questionnaire forms. Total N is approximately 1,000.

^dThis drug was asked about in five of the six questionnaire forms. Total N is approximately 5,000.

^eA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000–100,000; a large city as 100,000–500,000; and a very large city as having over 500,000. Within each level of population density, suburban and urban respondents are combined.

^fThis drug was asked about in four of the six questionnaire forms. Total N is approximately 3,900.

^gThis drug was asked about in two of the six questionnaire forms. Total N is approximately 2,000.

^hOnly drug use that was not under a doctor's orders is included here.

ⁱBased on data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

TABLE 4-5
Thirty-Day Prevalence of Daily Use of Various Types of Drugs by Subgroups
among Respondents of Modal Ages 19–30, 2008

(Entries are percentages.)


	<i>Approximate Weighted N</i>	Marijuana Daily ^a	Alcohol Daily ^a	Alcohol: 5+ Drinks in a Row in Last 2 Weeks	Cigarettes Daily	Cigarettes: 1/2 Pack+ per Day
Total	5,900	5.0	5.2	37.0	16.9	10.6
Gender:						
Male	2,400	7.4	7.5	46.5	19.2	12.5
Female	3,500	3.5	3.7	30.5	15.3	9.3
Modal Age:						
19–20	1,100	4.1	2.3	30.7	14.3	7.3
21–22	1,000	6.1	5.5	42.1	16.1	9.8
23–24	1,000	5.4	6.3	42.2	17.9	11.5
25–26	900	5.5	6.4	40.0	17.4	10.9
27–28	900	4.3	6.3	35.0	18.3	12.0
29–30	1,000	4.8	4.8	31.9	17.4	12.3
Region:						
Northeast	1,100	5.6	4.4	40.9	17.5	10.5
Midwest	1,600	4.5	4.8	40.8	19.5	12.9
South	2,000	4.8	5.6	33.2	17.6	11.1
West	1,200	5.6	6.0	33.7	11.1	6.0
Population Density: ^b						
Farm/Country	700	3.7	4.3	26.4	21.1	16.3
Small Town	1,500	5.3	4.6	33.3	17.4	11.6
Medium City	1,400	5.2	4.7	38.1	16.2	10.0
Large City	1,400	4.9	6.1	41.0	17.0	9.2
Very Large City	900	5.4	6.7	43.1	12.8	6.9

Source. The Monitoring the Future study, the University of Michigan.

^aDaily use is defined as use on 20 or more occasions in the last 30 days except for cigarettes, for which actual daily use is measured, and for 5+ drinks, for which the prevalence of having five or more drinks in a row in the last two weeks is measured.

^bA small town is defined as having fewer than 50,000 inhabitants; a medium city as 50,000–100,000; a large city as 100,000–500,000; and a very large city as having over 500,000. Within each level of population density, suburban and urban respondents are combined.


FIGURE 4-1
Any Illicit Drug: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.


FIGURE 4-2
Any Illicit Drug other than Marijuana: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.


FIGURE 4-3
Marijuana: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.


FIGURE 4-4
Amphetamines: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

FIGURE 4-5
Methamphetamine: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^a
by Age Group, 2008




Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

^aQuestions about the use of methamphetamines were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.

FIGURE 4-6
Crystal Methamphetamine (Ice): Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^a
by Age Group, 2008




Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

^aQuestions about the use of crystal methamphetamine were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.


FIGURE 4-7
Cocaine: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.


FIGURE 4-8
Crack Cocaine: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.


FIGURE 4-9
Other Cocaine: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

FIGURE 4-10
Hallucinogens:^a Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008




Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

^aUnadjusted for the possible underreporting of PCP.

FIGURE 4-11
LSD: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^a
by Age Group, 2008




Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

^aQuestions about the use of LSD were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.

FIGURE 4-12
Hallucinogens other than LSD:^a Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^b
by Age Group, 2008




Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

^aUnadjusted for the possible underreporting of PCP.

^bQuestions about the use of hallucinogens other than LSD were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.

FIGURE 4-13
Inhalants:^a Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^b
by Age Group, 2008




Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

^aUnadjusted for the possible underreporting of amyl and butyl nitrites.

^bQuestions about the use of inhalants were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.


FIGURE 4-14
Sedatives (Barbiturates): Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.


FIGURE 4-15
Narcotics other than Heroin: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.


FIGURE 4-16
Tranquilizers: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

FIGURE 4-17
Ecstasy (MDMA): Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^a
by Age Group, 2008




Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

^aQuestions about the use of ecstasy were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.

FIGURE 4-18
Steroids: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 30^a
by Age Group, 2008




Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

^aQuestions about the use of steroids were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.


FIGURE 4-19
Heroin: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.

FIGURE 4-20a
Alcohol: Lifetime, Annual, and 30-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008




Source. The Monitoring the Future study, the University of Michigan.

Note. Lifetime prevalence estimates were adjusted for inconsistency in self-reports of drug use over time. See text for discussion.


Some bars with the same number are not even due to rounding.

FIGURE 4-20b
Alcohol: 2-Week Prevalence of 5 or More Drinks in a Row
and 30-Day Prevalence of Daily Use among Respondents of
Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 4-21
Cigarettes: Annual, 30-Day, Daily, and Half-Pack-a-Day Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group, 2008



Source. The Monitoring the Future study, the University of Michigan.

Chapter 5

TRENDS IN DRUG USE IN EARLY AND MIDDLE ADULTHOOD

In this chapter we use data from the MTF longitudinal studies of many graduating cohorts to examine trends for the various age bands under study. Panel data are typically used to study changes in the same individuals over time. Here, however, we are looking across historical time at the behaviors and attitudes of particular age groups in the population. In other words, the data can be used much as we use the repeated cross-sectional surveys of secondary school students to track changes in behaviors over time for particular grade levels. Here we report historical trends in the use of various licit and illicit drugs by respondents of modal ages 19 through 50.

In the early 1990s, we began to document large and important increases among secondary school students in the use of several substances, particularly marijuana and cigarettes. The increases continued among 12th graders through 1997, as discussed in *Volume I*. One of the important issues addressed in this chapter is whether such increases occurred only among adolescents or whether those higher-using graduating classes have carried their higher levels of drug use with them as they moved into young adulthood. In other words, are they exhibiting lasting cohort effects?

Figures 5-1 through 5-19c present separate trend lines for two-year age strata (that is, respondents who are one to two years beyond high school, three to four years beyond high school, etc.); these two-year age strata are used to reduce the random fluctuations that would be seen with one-year strata.⁴¹ Each data point through age 32 in these figures is based on approximately 1,200 weighted cases drawn from two adjacent high school classes; actual (unweighted) numbers of cases are somewhat higher than those shown in the tables.⁴² Figures 5-1 through 5-19c also present some recent trend data from the age-35, age-40, age-45, and age-50 follow-ups. These older age strata are constituted in a slightly different way, in that the two half samples from a *single* graduating class (which through age 30 had been surveyed in alternating years) are now both surveyed in the same year.⁴³

Tables 5-1 through 5-5 are derived from the same data but presented in tabular form for 19- to 28-year-olds combined. Data are given for each year in which they are available for that full age

⁴¹Strictly speaking, these two-year strata are not age strata, because they are based on all respondents in the given year from two adjacent high school classes, and they do not take into account the minor differences in individual respondents' ages within each graduating class; however, they are close approximations to age strata, and we characterize them by the modal age of the respondents as ages 19 to 20, 21 to 22, and so on.

⁴²For the 2008 data, the 19- to 20-year-old stratum is composed of participating respondents from the high school graduating classes of 2007 and 2006, respectively; the 21- to 22-year-old stratum contains data from the classes of 2005 and 2004, respectively; and so on.

⁴³In 2008, the 35-year-olds are graduates from the high school class of 1991 (weighted $N = 870$), the 40-year-olds are graduates from the high school class of 1986 (weighted $N = 850$), the 45-year-olds are graduates from the high school class of 1981 (weighted $N = 840$), and the 50-year-olds are graduates from the high school class of 1976 (weighted $N = 870$). (Again, the unweighted or actual N s are somewhat higher.)

band (i.e., from 1986 onward). Respondents ages 29 and over are omitted because their inclusion would shorten the time period over which trends can be examined. However, the full data for them are contained in Figures 5-1 through 5-19c.

TRENDS IN PREVALENCE: EARLY AND MIDDLE ADULTHOOD

The trend results are as follows:

- Longer term declines among young adults in the annual prevalence of several drugs appeared to end in 1992 or 1993 (see Table 5-2). Among the 19- to 28-year-old young adult sample, this was true for the use of *any illicit drug, any illicit drug other than marijuana, marijuana, hallucinogens, narcotics other than heroin, crack, amphetamines, sedatives (barbiturates), and tranquilizers*. In 1994, annual prevalence for most drugs remained steady. *Cocaine other than crack* reached its low point in 1994 after a period of substantial decline. In 1995 there were modest increases (a percentage point or less) in the annual prevalence of almost all of the drug classes in Table 5-2, some of which were statistically significant.

Thus, it is clear that by 1992 or 1993 the downward secular trend (i.e., period effect) observable in all of these age strata (as well as among adolescents) had ended.⁴⁴ What has happened since then, however, is more of a cohort effect, reflecting an interaction between age and period such that only adolescents showed an increase in illicit drug use initially, and then they carried those new (higher) levels of drug use with them as they entered older age bands. Figure 5-1 shows the effects of generational replacement as the teens of the early 1990s reached their 20s. While all age groups generally moved in parallel through about 1992, the youngest age bands first showed signs of increase in their overall level of illicit drug use. The 18-year-olds shifted up first, followed by the 19- to 20-year-olds in 1994, the 21- to 22-year-olds in 1996, the 23- to 26-year-olds in 1999, the 29- to 30-year-olds in 2004, and the 35-year-olds in 2008. So far, the 40- and 45-year-olds have not shown much increase. (In fact, the 8th graders, who are not included in these graphs, actually showed an increase in use a year earlier than the 12th graders.)

To repeat, in the earlier decline phase of the drug epidemic, annual prevalence of use of *any illicit drug* moved in parallel for all age strata, as illustrated in Figure 5-1; this pattern reflects a secular trend, because a similar change is observed simultaneously across different age levels. After 1992—in what we have called the “relapse phase” of the popular drug epidemic that began in the 1960s—a quite different pattern emerged, with 8th graders increasing their drug use first, followed by 10th and 12th graders; then the next oldest age group, but with a little delay; the next oldest then following, but with a longer delay; and the oldest two groups (45- and 50-year-olds) not yet showing an increase. This pattern reflects a classic cohort effect, in which different age groups are not all moving in parallel; rather, different age groups show increases when the cohorts (i.e., high school classes) having heavier use at an earlier stage in development reach the relevant age level. Further, the slopes of the age bands are successively less steep in the

⁴⁴Actually, the downturn ended at least a year earlier among 8th graders, who showed the beginning of an increase in 1992 (see Table 2-2).

older age groups, suggesting that some of the cohort effect may be dissipating with maturation. But we think it unlikely that only cohort effects are occurring (in addition to the long-established age effects); period effects also likely play a role.

- Use of *marijuana*, the major component of the illicit drug use index, shows an almost identical pattern to the index (Figure 5-3a). After a long and steady decline from the late 1970s to the early 1990s, annual use leveled for a while among young adults before beginning a gradual increase. Virtually all of this increase was attributable to the two youngest age bands (18 and 19 to 20) until 1996, when the 21- to 22-year-olds began to show a rise. The older age bands then tended to show increases fairly sequentially, with 29- to 30- and 35-year-olds showing significant increases in 2008. The 18-year-olds' use of marijuana in the prior 12 months declined after 1997, and later several of the succeeding age bands through age 26 began to show declines in a pattern that again suggests lasting cohort differences.
- A similar pattern emerged for current *daily marijuana* use (Figure 5-3c). In the mid- to late 1990s, daily marijuana use among 35- and 40-year-olds was as high as, or higher than, use among some younger age groups, suggesting a lasting cohort effect on this behavior. However, in recent years, the 35-, 40-, and 45-year-olds have been similar to respondents ages 27 to 30, who have had among the lowest levels of daily use. An important finding shown in Figure 5-3c is that, although the various age groups had been moving in parallel for many years at fairly similar levels of prevalence, the trends diverged considerably in the 1990s in a staggered fashion, such that now 18- to 30-year-olds have distinctly higher levels of daily marijuana use than the older age groups, again reflecting stable cohort differences.
- The index of using *any illicit drug other than marijuana* has shown a similar transition in the pattern of change. Period effects seemed to predominate until about 1992, but a cohort-related pattern of change emerged thereafter (Figure 5-2). And, while annual use leveled by 1997 among 18-year-olds, it began rising during that interval among 25- to 26-year-olds and is now also rising among 27- to 30-year-olds and 35-year-olds. The primary difference from the picture for marijuana is that the increases were not as sharp in the 1990s for most of the age bands. (Compare Figure 5-2 with Figure 5-1 to see the difference.)
- In the 1980s and 1990s, *LSD* use also increased among those in their teens and early 20s more than among the older strata, as Figure 5-6 illustrates. Over the interval 1985 to 1996, there was a gradual but considerable increase in annual LSD use among respondents ages 18 to 24, which was sharpest among 12th graders and 19- to 20-year-olds. (In this case the increase did not seem to radiate up the age spectrum beyond age 26.) A turnaround began among 12th graders after 1995 and then among the older age groups in a somewhat staggered fashion, again indicative of a cohort effect. Declines in the years since have been greatest among 18- to 24-year-olds, who had attained the highest rates of LSD use. LSD use declined considerably from 2001 to 2003 in all age bands (including 8th and 10th graders), and then leveled through 2007 at historically low rates, suggesting that an important secular trend may have set in, quite possibly related to

decreased availability of the drug. The youngest groups—ages 18, 19–20, and 21–22—all showed increases in 2008, suggesting a possible comeback for LSD.

- Several drug classes actually exhibited a faster decline in use among the older age groups than among 12th graders during the earlier period of decline in the 1980s (see Figures 5-1 through 5-19c). These included *any illicit drug*, *any illicit drug other than marijuana*, *amphetamines*, *hallucinogens* (until 1987), *LSD* (through 1989), and *methaqualone*.
- In fact, there was a crossover for some drugs when 12th graders are compared to young adults. In earlier years 12th graders had lower usage levels, but more recently they have tended to have higher ones than post–high school respondents for use of *any illicit drug*, *marijuana*, *any illicit drug other than marijuana*, *hallucinogens*, *LSD* specifically, *crack cocaine*, *amphetamines*, *sedative (barbiturates)*, *narcotics other than heroin*, and *crystal methamphetamine (ice)*.
- *Cocaine* (Figure 5-9) gives a quite dramatic picture of change. Unlike most other drugs, active use of cocaine has generally tended to rise with age after high school, usually peaking approximately three to four years past graduation. This is a classic example of an age effect. Despite the large age differences in absolute prevalence, all age strata moved in a fairly parallel way through 1991, indicating that a secular trend was taking place in addition to the age effect. All began a sharp and sustained decline in use after 1986. The two youngest strata (12th graders and 19- to 20-year-olds) leveled by 1992, whereas use continued a decelerating decline for a few years beyond that in the older age groups, signaling the beginning of a cohort effect. From 1994 to 1999, cocaine use rose some in the five youngest strata (i.e., those younger than 27) on a somewhat staggered basis, with the three older groups still decreasing a bit more over that same period. This, to some degree, reversed the age differences that were so prominent in the 1970s and 1980s. Cohort-related change appears to have predominated in the 1990s, quite possibly as the result of “generational forgetting” of the cocaine-related casualties so evident in the early to mid-1980s. In other words, those in the older cohorts retained that learning, but those in the newer cohorts never had it. The fact that from 1994 to 1996 the 35-year-olds had higher lifetime prevalence levels of cocaine use than some of the younger age groups also suggests some lasting cohort-related differences established during the peak years of the cocaine epidemic.
- *Crack* use was added to the 12th graders’ questionnaires in 1986 and to the follow-up questionnaires in 1987. The decline in crack use, which began right after the introduction of these questions, ended in 1991 among 12th graders, and by 1994 it had ended among young adults (see Figure 5-10 and Table 5-2). Among 19- to 28-year-olds, the annual prevalence rate held at about 1%, which was down from the peak levels of just over 3% in 1986 through 1988. As was true for a number of other drugs, crack use began to rise after 1993 among 12th graders, but not in the older age strata until years later, when increases were observed in a somewhat staggered pattern going up the age scale. Again, a cohort effect due to generational replacement seems to have been occurring. Since 1994, 18-year-olds have had the highest reported rates of use, with 19- to 22-years-olds next highest. Importantly, all groups now have annual prevalence rates below 2.0%.

- With regard to *inhalants*, the large separation of trend lines for the younger age groups in Figure 5-4 shows that, across many cohorts, use has dropped consistently and sharply with age, particularly in the first few years after high school. In fact, of all the populations covered by MTF, the 8th graders (not shown in Figure 5-4) have had the highest rate of use, indicating that the decline in use with age starts at least as early as 8th or 9th grade. Like cocaine, inhalants have shown a strong age effect, but unlike cocaine, use of inhalants declines with age.

Figure 5-4 also shows that, until the mid-1990s, there was a long-term gradual increase in annual inhalant use (unadjusted for underreporting of nitrite inhalants), one which was greatest among 12th graders, next greatest among 19- to 20-year-olds, and next greatest among 21- to 22-year-olds. Respondents more than six years past high school, who historically have had a negligible rate of use, did not exhibit the increases in use seen among the younger respondents, which began at least as early as 1977 among 12th graders and in 1983 among 19- to 20-year-olds. There was some subsequent increase among 21- to 22-year-olds and later still an increase among 23- to 24-year-olds. After 1995, this long-term trend, reflecting a cohort effect, began to reverse in the two youngest age strata and subsequently among the next two age strata. The older age strata have generally shown negligible rates of inhalant use.

- The annual prevalence for *ecstasy (MDMA)* use among the entire young adult sample (ages 19 to 28) was at about 1.5% in 1989 and 1990 (Table 5-2 and Figure 5-8). After 1991 it dropped to around 0.8% for several years before starting to rise significantly in 1995. Then ecstasy use began to rise in all of the young adult age strata, most notably in the younger age bands (19 through 26) through 2001. Use among 12th graders, which was not measured until 1996, was by then the highest of any of the age groups at 4.6% annual prevalence. Twelfth graders' use slipped by a full percentage point through 1998 before jumping significantly—by two full percentage points—in 1999. (Use by 10th graders also jumped significantly in 1999.) Thus it appears that young people from their mid-teens to mid-20s discovered ecstasy after some years of low and relatively level use. In 2000 the sharp increase in use continued among ages 15 to 16 (10th graders) through age 26, and also showed up among 8th graders (13- to 14-year-olds) for the first time. By 2001 the increase had slowed and even begun to reverse among ages 18 to 26, even as the 31- to 32-year-olds showed their first appreciable increase in ecstasy use. We attributed the deceleration in 2001 to a fairly sharp increase in perceived risk that year and predicted a turnaround in use in 2002. In 2002, and again in 2003, perceived risk increased sharply and, as Figure 5-8 illustrates, all age bands showed a reversal, with a sharp decrease in use. Clearly, the decrease has been sharpest in the younger age bands, perhaps because a cohort effect is at work in the upper ages, helping to offset a downward secular trend.
- In the late 1970s, *amphetamine* use rose with age beyond high school; but after a long period of decline in use from 1981 to the early 1990s, this relationship had reversed (see Figure 5-13). The declines were greatest in the older strata and least among 12th graders, even though use decreased substantially in all groups. As was true for many illicit drugs,

amphetamine use began to rise among 12th graders after 1992, and eventually among the 19- to 24-year-olds; but there has only recently been a small increase among 25- to 30-year-old respondents. In other words, another cohort-related pattern of change seems to have emerged in the 1990s for amphetamines, though in this case it may be dissipating quickly after respondents reach their early 20s. In fact, some decline in amphetamine use is observable in the younger age strata since 2003, and among 19- to 20-year-olds since 2004. At present the age differences through age 50 are of considerable magnitude and mostly ordinal (with 18- to 22-year-olds showing the highest rates of use and those over 30 much lower rates). There has been little further change in the upper age bands in recent years.

- Since 1990, when it was first measured, use of *crystal methamphetamine (ice)* has remained at fairly low rates in this young adult population (Figure 5-14). However, among 19- to 28-year-olds combined, annual prevalence rose from 0.4% in 1992 to 1.6% by 2005, falling to 0.8% by 2008 (Table 5-2). Use had been rising among 12th graders and 19- to 20-year-olds specifically between 2000 and 2002, but since then their use has declined to low levels. General *methamphetamine* use has only been measured since 1999, and its use until 2005 was stable among 19- to 28-year-olds, with annual prevalence fluctuating between 2.4% and 2.8%. Use has declined to 1.0% by 2008 (Table 5-2). This recent decline may well reflect a cohort effect, as use has been declining steadily among 12th graders since 2002.
- Use of *heroin* increased appreciably in 1995 among 12th graders and young adults ages 19 to 24, but not among the older age bands (Figure 5-11). It has remained at this higher plateau in these younger age bands since then. Among young adults generally, annual use had previously been quite stable from at least as far back as 1986 through 1994 (Table 5-2), and it stabilized again at a higher level after 1994—a level roughly twice as high as the previous one. Heroin use among 12th graders and 19- to 20-year-olds has declined slightly since 2000 or 2001, respectively, but the older age groups have maintained a fairly stable, very low rate of use.
- Among 19- to 28-year-olds, use of *narcotics other than heroin* leveled after 1991, following a long period of slow, fairly steady decline (Figure 5-12). Twelfth graders showed an appreciable increase in use beginning in 1993, which continued into 2004, while 19- to 20-year-olds showed some increase after 1994, 21- to 22-year-olds after 1996, 23- to 24-year-olds after 1997, and the older age groups after 2000. Thus, cohort-related change appears to have been occurring during the 1990s and beyond for this class of drugs, following a long period of secular trends. In 2002, the question text was changed on three of the six questionnaire forms to update the list of examples of narcotics other than heroin. Talwin, laudanum, and paregoric, each of which had negligible rates of use by 2001, were replaced by Vicodin, OxyContin, and Percocet. As a consequence of this revision, reported use rates increased in 2002. Data presented here for 2002 are from three of the six questionnaire forms with the new wording (which showed higher prevalence rates than the older question did). All six questionnaire forms contained the new wording beginning in 2003, so the data presented for 2003 and after are based on all forms. Although the older version of the question showed no significant changes

occurring in 2002, there was a significant increase in narcotics use observed in 2003 (based on the new question in both 2002 and 2003). Among 19- to 28-year-olds, annual prevalence reached a peak level of 9.0% in 2004, about where it is in 2008 (9.1%). Some turnaround was observed in some of the younger age bands after 2004 in the use of this important class of drugs, but use continues to rise in most of the older age bands.

- The annual prevalence rates for *Vicodin* and *OxyContin*, which were first measured in 2002 (separately from the general question about narcotics other than heroin), were appreciable (8.2% and 1.9%, respectively) for 19- to 28-year-olds. Increases were observed for these two drugs in the subsequent years. Among 19- to 28-year-olds (see Table 5-2), the annual prevalence of OxyContin use rose from 1.9% in 2002 to 3.1% in 2004 through 2006—changes that were fairly parallel to those observed among 12th graders over the same interval (when their annual prevalence rose from 4.0% in 2002 to 5.5% in 2005). The 2002–2005 increases in OxyContin use were significant for both 12th graders and 19- to 28-year-olds. Annual prevalence was stable from 2004 to 2007 at about 3%, but increased to 3.9% in 2008. Vicodin use rose by less, but started from a higher base, with annual prevalence increasing slightly among 19- to 28-year-olds, from 8.2% in 2002 to 9.3% in 2005; it has been at about 9% since 2006.
- *Sedative (barbiturate)* use (Figure 5-15) showed a long-term parallel decline in all age groups covered through the late 1970s and 1980s, leveling by about 1988. While use remained low and quite level for most of the age bands for about five years, it began to rise by 1993 among 18-year-olds, by 1995 among 19- to 20-year-olds, by 1997 among 21- to 22-year-olds, by 1998 among 23- to 24-year-olds, by 2001 among 25- to 28-year-olds, and by 2005 among 29- to 30-year-olds. The same cohort-related pattern of change seen during the 1990s for many other drugs also exists for sedatives (barbiturates); like most other drugs, this pattern was preceded by a long period of secular change. While use has now leveled off among most younger age groups, the 35-, 40-, and 45-year-olds all showed increases in sedative (barbiturate) use between 2006 and 2008.
- *Tranquilizers* (Figure 5-16) follow a similar pattern to that just described for sedatives (barbiturates). One difference is that the 12th graders' annual prevalence rate has not always been the highest among the various age groups, as was the case for sedatives (barbiturates), although it was highest between 1994 and 2000 as a result of a greater increase in tranquilizer use among the 12th graders than in the young adult strata. In the last six years or so, however, as use rose and then leveled among those in their early 20s, the 12th graders no longer stand out as having the highest rate of tranquilizer use. In fact, it is the 21- to 22-year-olds that had the highest rate in 2005 through 2007, with 23- to 24-year-olds having the highest rate in 2008. This is another clear example of a cohort-related pattern of change. Now that tranquilizer use has leveled and begun to decline among 12th graders and 19- to 20-year-olds, it is likely that we will see that pattern echoed in the older age groups as a new cohort effect plays out.
- The use of *anabolic steroids* (Figure 5-17) is substantially lower after high school than during, and this has been true since measures of steroid use were first introduced into two of the follow-up questionnaires in 1991. The age-related differences are not consistent;

prevalence rates among the young adult strata are all quite low and do not appear to trend in any systematic way. In general, it seems that the rise in steroid use from 1999 to 2003 among 8th and 10th graders and from 2001 to 2004 among 12th graders was largely specific to those age groups. Annual prevalence rates are now very low (0.3% or less) for respondents ages 21 and older.

- **Alcohol** trends for the older age groups (see Figures 5-18a–d) have been somewhat different than for the younger age groups in some interesting ways. For **30-day prevalence** and **occasions of heavy drinking**, the declines for the two youngest age strata (12th graders and those one to two years past high school) during the 1980s were greater than for the older age groups. These differential trends were due in part to the effects of changes in minimum drinking age laws in many states, changes that would be expected to affect primarily the age groups under age 21. However, because similar (though weaker) trends were evident among 12th graders in states that maintained a constant minimum drinking age of 21, the changed laws cannot account for all the downward trends, suggesting that there was also a more general downward trend in alcohol consumption during the 1980s.⁴⁵ By 1994, these declines in 30-day prevalence had slowed or discontinued for virtually all age groups until 1997, when they began to turn downward again for 12th graders, and 1999, when they started down among 19- to 20-year-olds.

Those respondents three to four years past high school stand out for showing the smallest downward trend in **occasional heavy drinking** or **binge drinking** (i.e., having five or more drinks in a row at least once in the past two weeks) since the early 1980s (see Figure 5-18d). One important segment of that age stratum is composed of college students, who have shown very little decline in use over the past quarter century (see chapter 9).

The older age groups have generally shown only a modest long-term decline in annual prevalence rates and no recent decline in binge drinking or 30-day prevalence rates. Note that the binge drinking trend lines for different age groups (Figure 5-18d) are spread out on the vertical dimension, reflecting large and persisting age differentials (age effects) in this behavior. The relationship with age is curvilinear, however. In recent years the 21- to 22-year-olds have consistently shown the highest rates of binge drinking, while the two adjacent age bands have shown the next highest. Binge drinking appears to have been gradually increasing in recent years, particularly among 23- to 24-year-olds and 25- to 26-year-olds, perhaps reflecting increased college enrollment in these age groups. (College attendance is strongly correlated with higher rates of binge drinking.⁴⁶) In 2008 the binge drinking rate for 19- to 20-year-olds is considerably lower (31%) than for those just one to two years older (42%).

⁴⁵O'Malley, P. M., & Wagenaar, A. C. (1991). Effects of minimum drinking age laws on alcohol use, related behaviors, and traffic crash involvement among American youth: 1976–1987. *Journal of Studies on Alcohol*, 52, 478–491.

⁴⁶Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates.

From the early 1980s through the mid-1990s, rates of *daily drinking* (Figure 5-18c) fell by considerable proportions in all age strata for which we have data, reflecting an important change in drinking patterns in the culture. Among 19- to 28-year-olds combined, daily drinking declined from 1987 (6.6%) to 2000 (4.1%), and has increased since then, to 5.3% in 2008 (see Table 5-4). Daily drinking rates have generally been highest for 35-, 40-, and 45-year-olds in recent years.

It is worth noting that the 35-, 40-, 45-, and 50-year-olds have had among the lowest rates of binge drinking but among the highest rates of daily drinking in recent years for which we have data available. These patterns—particularly the high rate of daily drinking—likely reflect age effects as well as perhaps some enduring cohort differences (because these cohorts had considerably higher rates of daily drinking when they were in high school).

- The prevalence rates for *cigarette smoking* show more complex trends than most other substances, due to the long-term presence of both cohort and age effects, plus slightly different patterns of such effects on different measures of smoking in the past 30 days (one or more cigarettes per month, one or more cigarettes per day, and a half pack or more of cigarettes per day).

In the earlier years of MTF, the curves across time were of the same general shape for each age band (Figures 5-19a–c), but each of those curves tended to be displaced to the right of the immediately preceding age group, which was two years younger. The pattern is clearest in Figure 5-19c (half pack plus per day). This pattern is very similar to the one described in *Volume I* for lifetime smoking rates for various grade levels below senior year; it is the classic pattern exhibited by a cohort effect,⁴⁷ and we believe that the persisting cohort differences are due to the dependence-producing characteristics of cigarette smoking.

The declining levels of cigarette smoking across cohorts at age 18, which were observed when the classes of 1978 through 1981 became 12th graders, were later observable in the early-30s age band, as those same high school graduating classes grew older (see Figures 5-19b and c). This was true at least through about 1991. After that, there was a considerable convergence of rates across age groups, largely because there were few cohort differences among the senior classes who graduated from the early to mid-1980s through the early 1990s—a period of fairly level use.

In addition to these cohort differences, there are somewhat different age trends in which, as respondents grow older, the proportion smoking at all in the past 30 days declines some, while the proportion smoking a half pack per day actually increases. Put another

⁴⁷O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (1988). Period, age, and cohort effects on substance use among young Americans: A decade of change, 1976–1986. *American Journal of Public Health*, 78, 1315–1321.

way, many of the light smokers in high school either become heavy smokers or quit smoking.⁴⁸

The picture was further complicated in the 1990s, when it appears that a new cohort effect emerged, with smoking among adolescents rising sharply (beginning after 1991 for 8th and 10th graders and after 1992 for 12th graders). The 19- to 20-year-olds soon showed a rise at the beginning of the 1990s—perhaps responding to some of the same social forces as the adolescents (including the Joe Camel advertising campaign); but 21- to 24-year-olds did not show an increase until about 1995, and 25- to 26-year-olds until about 1996. Those young adults over age 26 have not yet shown much increase; it is possible that an upward cohort effect was offset by a downward secular trend during this period.

After about 1999, smoking rates among virtually all age groups leveled or declined, suggesting that societal forces may be affecting all age groups in a similar way, giving rise to a secular trend. Large increases in price and a great deal of adverse publicity for the tobacco industry are highly plausible candidates, as are an increase in state and national antismoking advertising, the demise of the Joe Camel campaign and billboard advertising, and the imposition of no-smoking regulations in many public and workplace settings by states and municipalities. From 2003 to 2007 thirty-day, daily, and half-pack smoking all declined among 35-, 40-, and 45-year-olds (Figures 5-19a through 5-19c). The age-35 group did not continue the decline in 2008, though both the 40- and 45-year-olds did.

- Apart from cigarettes, none of the other drugs included in the study showed a clear long-term pattern of enduring cohort differences in the earlier years of MTF (the 1970s and 1980s), despite wide variations in their use by different cohorts at a given age. There was one exception; a modest cohort effect was observable for *daily marijuana* use (Figure 5-3c) during the late 1970s and early 1980s.⁴⁹ (But as more recent classes leveled at lower rates of use, evidence for the cohort effect faded.) The emergence in the 1990s of a new epidemic of marijuana use among teens once again yielded a strong pattern of cohort effects. As can be seen in Figure 5-3c, daily use rose sharply among 12th graders and 19- to 20-year-olds after 1992, among 21- to 22-year-olds after 1993 with a sharp rise occurring in 1997, among 23- to 24-year-olds after 1998, among 25- to 26-year-olds after 2000, among 27- to 28-year-olds in 2003, among 29- to 30-year-olds in 2005, among 35- and 40-year-olds in 2006, and among 45-year-olds in 2007. This is not unlike the pattern of change for cigarette smoking that occurred in the 1990s (Figure 5-19a). The cohort effect for daily marijuana use may be attributable, in part, to the very strong association between that behavior and regular cigarette smoking. It is noteworthy that even among the 35-, 40-, and 45-year-olds in the study, 2.2–2.7% report that they currently smoke marijuana on a daily basis. That amounts to 1 in every 37 to 46 adults at those ages. And

⁴⁸ To illustrate, in the graduating class cohort of 1976, 39% were 30-day smokers in senior year, 39% by ages 19 to 20, but only 28% by ages 31 to 32—a net drop of 11 percentage points over the entire interval. By way of contrast, 19% of that class were half-pack-a-day smokers in senior year, 24% by ages 19 to 20, and 21% at ages 31 to 32—a net gain of five percentage points and two percentage points over the respective intervals.

⁴⁹ O'Malley, P. M., Bachman, J. G., & Johnston, L. D. (1988). Period, age, and cohort effects on substance use among young Americans: A decade of change, 1976–1986. *American Journal of Public Health*, 78, 1315–1321.

we know from results published in *Volume I* that many more have been daily marijuana users for at least a month at some time in their lives.

- In sum, except for *cigarettes* and *alcohol*, substance use prior to 1992 among 12th graders and young adults had shown longer term trends that were highly parallel across age groups, indicating that general secular trends predominated in that period. Since 1992, however, there has been considerable divergence in the trends for different age bands on a number of drugs as use among adolescents rose sharply, followed by subsequent rises among 19- to 20-year-olds, 21- to 22-year-olds, and so on. This divergence indicates a new cohort effect, quite possibly reflecting a generational forgetting of the dangers of drugs by the cohorts who reached senior year in the early to mid-1990s. Data discussed in chapter 6, “Attitudes and Beliefs about Drugs among Young Adults,” provide additional evidence for this interpretation.

TRENDS FOR IMPORTANT SUBGROUPS OF YOUNG ADULTS

Four-year age bands have been used here to examine subgroup trends in order to yield sufficiently large numbers of cases to permit reliable estimates for the various subgroups being examined. Subgroup data for respondents of each gender and for respondents from communities of different sizes are available for 19- to 22-year-olds since 1980, 23- to 26-year-olds since 1984, and 27- to 30-year-olds since 1988. Beginning in 1987, a question about state of residence was added to all follow-up questionnaires, permitting trend data to be calculated for the four regions of the country since then. These various subgroup data are not presented in tables or figures here because of the substantial amount of space they would require. Rather, a verbal synopsis of what they contain is presented.

Gender Differences in Trends

- Over the long term, gender differences narrowed for some drugs among young adults in each of these three age bands, primarily when a steeper decline in use among males (who generally had higher rates of use) occurred in the 1980s. The overall picture, though, is one of parallel trends, with use among males remaining higher for most drugs, including the indexes of *any illicit drug* use in the prior year and use of *any illicit drug other than marijuana* (see Table 5-5, for example).

The downward trend in *marijuana* use among 19- to 22-year-olds between 1980 and 1989 was also a bit sharper among males than females, narrowing the gap between the two groups. Annual prevalence fell by 22 percentage points (to 34%) among males, compared to a drop of 14 percentage points (to 31%) among females, leaving a difference of three percentage points. Since 1995, the gender gap has averaged about 5% to 7% in all three age groups.

Similarly, between 1980 and 1993, *daily marijuana* use for the 19–22 age group fell more steeply, from 12.9% to 2.9% among males, versus 6.1% to 1.7% among females, considerably narrowing the rather large gap. As use began to rise after 1993, the gap widened again. Among 23- to 26-year-olds, as daily use first began to increase in 1998

and 1999, the gap between the genders began to widen. In the oldest age group (ages 27–30), the difference had been fairly constant, with daily marijuana use among males generally being two to three times higher than among females. After 2001, however, use increased more among females, narrowing the gap somewhat, but an increase from 2004 to 2008 among males opened the largest gender gap seen for daily marijuana use in this age band.

- Males have shown slightly higher proportions using *any illicit drug other than marijuana* in all three age bands, a fact that has not changed appreciably over the years, though the differences tended to narrow some as use dropped, and to widen as use increased.
- For *LSD*, males have consistently had higher rates of use than females. Among 19- to 22-year-olds, the male–female differences tended to diminish as use declined (from 1980 to 1985 and again from 1999 to 2004) and expand as use increased (1986–1995). In the two older age bands there was less change in use, and differences had been relatively consistent (with males higher) since data have been available, beginning in 1984 for 23- to 26-year-olds and in 1988 for 27- to 30-year-olds. After 1999 and 2001 for the two groups, respectively, LSD use dropped, substantially narrowing the gender differences. Males began to show these declines first, and both genders have moved to almost no use since about 2003.
- *Ecstasy (MDMA)* exhibited little or no gender difference in any of the three age bands before use began to grow in the late 1990s. Even since then, among 19- to 22-year-olds there has been little gender difference, except that use among males started to decline one year ahead of use among females. But in the older age groups a gender difference did open up after 1997, with males fairly consistently having higher rates of use among both 23- to 26-year-olds and 27- to 30-year-olds.
- Males have had higher rates of *cocaine* use than females since MTF began. During the period of sharp decline from the peak levels in annual cocaine prevalence (1986–1993), use dropped more among males than females, narrowing the gender differences. In the 19- to 22-year-old age band, annual prevalence for males declined by 16 percentage points (to 4.5%) versus 13 percentage points among females (to 2.8%) in 1993. In the 23- to 26-year-old age band, there was also a narrowing of the gender difference between 1986 and 1993, with annual prevalence down 19 percentage points among males (to 6.9%) and 13 percentage points among females (to 4.2%). Use in the 27- to 30-year-old group also dropped faster among males between 1988 (when data were first available) and 1997—down 13 percentage points versus 7 among females. In sum, during the period of sharp decline in overall cocaine use, the gender differences—which had been fairly large—narrowed considerably in all three of these age bands. During the more recent resurgence in cocaine use, which has occurred on a somewhat staggered basis over the years, the gap between genders expanded only slightly.
- A similar thing happened with *crack* during the earlier period of decline, though the proportional difference between the two genders has consistently been higher than for

cocaine overall. With crack, though, there was some gender convergence (between 1992 and 1998) among 19- to 22-year-olds, as use among males declined slightly and use among females rose gradually. Since 1999, there has been no consistent change in differences between males and females. In the two older age bands, males consistently had slightly higher crack usage rates, at least until a decline among males in recent years diminished the difference.

- As *sedative (barbiturate)* use declined through the 1980s, the modest gender differences (males were higher) were virtually eliminated in all three age bands. Beginning in the early 1990s, there emerged a staggered increase in use by both genders across all three age groups, with males increasing more than females, thereby opening a small difference again in the late 1990s and 2000s.
- During the period from 1982 through about 1994, there was little gender difference in annual prevalence for *heroin*. After 1994, use increased some, particularly among the younger groups, with males generally attaining slightly higher rates than females. As of 2008, the gender differences in heroin rates are small for all groups, but males continue to remain slightly higher.
- Among 19- to 22-year-olds, both genders showed some decline in their use of *narcotics other than heroin* between 1980 and 1991, with a near elimination of previous gender differences (males had been higher). Beginning in 1994, use by males began to rise in this age group, while use by females began to rise a year later. Some gender differences have developed as use has increased, with males at 9.6% and females at 7.7% in 2008. The picture for 23- to 26-year-olds is very similar: the gender difference (males higher) had been eliminated by 1988, but re-emerged after 1995 as use increased more among males. Among 27- to 30-year-olds, there has been a smaller gender difference and the least increase in use in the 2000s. Still, use has increased for both genders since 1999, with males having higher rates of use.
- The use of *OxyContin* has generally been higher among males than females for all three age bands in the seven years for which we have data; and both genders have shown some increase in use over that seven-year interval. *Vicodin* use also has been consistently higher among males, and there has been an increase in use among males in the upper two age bands and among females in the oldest age band.
- In general there has been no appreciable gender difference in *amphetamine* use for some years in any of these three age bands. Between 1981 and 1991, rates of amphetamine use were similar for males and females and showed substantial and parallel downward trends for both genders. Among 19- to 22-year-olds, annual prevalence of use dropped 22 percentage points for males (to 5.2% in 1991) and 21 percentage points for females (to 4.7% in 1991). There were small increases in annual prevalence for both genders in the 19- to 22-year-old age group after 1991, in the 23- to 26-year-old age group after 1995, and in the 27- to 30-year-old age band after 2000, but the genders diverged only slightly (males higher), and use has leveled in all three age bands. Nonmedical use of *Ritalin*, a

prescription amphetamine added to MTF in 2002, has generally been higher among males than females, though there was little gender difference in 2008.

- **Crystal methamphetamine (ice)** was added in 1990. In the early 1990s, use was low and very similar for both genders in all three young adult age bands. More of the increase in use that occurred in the mid-1990s in the younger two age bands occurred among males—opening a gender gap. The gap then narrowed, though males on average were slightly more likely to report use of crystal methamphetamine until 2005. It should be noted that the estimates are a bit unstable for this drug due to limited sample sizes.
- For **tranquilizers**, both genders showed a long, gradual decline (and very similar rates of use) from 1980 through about 1993 in all three age bands. Beginning in 1995, use increased for both genders in the 19- to 22-year-old group, followed by an increase beginning after 1997 among 23- to 26-year-olds and after 1999 among 27- to 30-year-olds, again reflecting cohort effects driven by generational replacement. Some gender difference emerged in this period of increase (and subsequent decrease after 2002 and 2003, respectively, for the lower two age bands), with males reporting somewhat higher usage rates. Among 27- to 30-year-olds, males' use is higher (e.g., 7.0% annual prevalence vs. 5.4% for females in 2008).
- **Inhalant** use has generally been quite a bit higher among males than females in all three age groups. The 19- to 22-year-old group showed a gradual upward shift from 1980 to 1988, followed by a leveling for some years for both genders. In 1997, female inhalant use began to decline among 19- to 22-year-olds, followed by males in 2001; however, the gender gap did not diminish much with this decline until 2005, when there was a convergence. Among 23- to 26-year-olds there was a widening gender gap as use by males increased between 1992 and 1999, though a decline among males since then has narrowed the gap, and since 2005 the gap has been very small. In the oldest age stratum, use among males has consistently been slightly higher, though the prevalence of inhalant use is very low by this age.
- Use of three “club drugs”—**GHB**, **ketamine**, and **Rohypnol**—has tended to be a bit more concentrated among males in all three age strata. However, the estimates are not very stable because of the limited numbers of cases upon which they are based; prevalence rates are all very low.
- For **alcohol**, 30-day prevalence rates exhibited a gradual, parallel decline from 1981 through 1992 for both genders in the 19- to 22-year-old age group. Thirty-day prevalence fell from 83% to 72% among males and from 75% to 62% among females by 1992. In the two older age bands, there had also been a modest, parallel decline for both genders, from 1985 through 1992 in the case of 23- to 26-year-olds, and at least from 1988 (when data were first available) to 1991 or 1992 in the case of 27- to 30-year-olds. Since 1992, both genders in the older two age bands have shown fairly level rates of use, with males somewhat higher; however, among 19- to 22-year-olds there has been a convergence, because use by males has declined slightly while use by females has increased slightly.

The increasing proportion of women attending college may explain this convergence, at least in part.

There was also a general long-term decline in *daily drinking* from about 1981 or 1982 through about 1992, with daily use falling more among males, considerably reducing, but far from eliminating, what had been a large gender difference among 19- to 22-year-olds. To illustrate, in 1981, 11.8% of males reported daily use versus 4.0% of females; the comparable 1992 statistics were 5.3% and 2.7%. After 1995, daily drinking began to increase among 19- to 22-year-olds for both genders, but leveled a few years later. From 2002 to 2005 daily use was rising among males and falling among females, increasing their differences, but since 2005 there has been a considerable convergence with daily use among males falling and use among females increasing some. There was still a gender difference for daily drinking among the 19- to 22-year-old age group in 2008—5.0% for males versus 3.1% for females—but not nearly as large as it had been in 1981 (11.8% vs. 4.0%). The gender differences have been similar for the older age groups (in 2008, for example, 9.3% for males vs. 4.4% for females among 23- to 26-year-olds), and there has been little evidence of any convergence or divergence in the two upper age bands.

There are also long-established and large gender differences in all age groups in the prevalence of *occasional heavy drinking*. Males in the 19- to 22-year-old band showed some longer term decline in this statistic, from 54% in 1986 to 45% in 1995, as female use remained steady, thus narrowing the gender gap (from 24 percentage points in 1986 to 17 percentage points in 1995). Since 1995, binge drinking among 19- to 22-year-old females has been drifting upward from 28% to 32% by 2008, whereas among males it has held fairly steady, narrowing the difference some. In the two older age bands (23- to 26-year-olds and 27- to 30-year-olds), the sizable gender differences have been mostly stable as the binge drinking rates have drifted steadily upward in both genders. To illustrate, among 23- to 26-year-olds, the binge drinking rate among females has risen from 22% in 1995 to 34% in 2008, while among males it has risen from 44% in 1994 to 53% in 2008. (The increases observed among 27- to 30-year-olds have not been as steep.) *This sustained increase for 23- to 26-year-olds is striking, and may be due at least in part to the increased rate of college attendance and decreased rate of marriage among this age group.*

- Most striking for *cigarette smoking* are the similarities between the genders in both absolute levels and trends. All three age groups showed a long-term decline in *daily smoking* rates for both males and females after data were first available for each: 19- to 22-year-olds from 1980 to 1990, 23- to 26-year-olds from 1984 to 1992, and 27- to 30-year-olds from 1988 to 1999—again reflecting a cohort effect moving up the age scale. Male and female *30-day smoking* rates have also been very close over most of the time for which data are available, particularly in the two older age groups. But among 19- to 22-year-olds there was a crossover after 1993—before that point, females had slightly higher 30-day prevalence rates, but after that males did from 1994 on. Both genders have shown a considerable decline since 1999. Among 23- to 26-year-olds, the genders had very similar smoking rates until males started reporting higher 30-day smoking rates from 1998 on. Males have shown almost no decline since 1998, but females in this age band

have shown some. In the oldest age band, the two genders were quite close until males first reported higher rates in 2002, and their rate has generally remained somewhat higher since; in 2008 it was 29% among males and 22% among females.

There were some increases in the last decade and a half in 30-day smoking rates among the two younger groups and especially among males. For example, from 1993 to 1999, 19- to 22-year-old males increased from 29% to 37%, while females increased from 29% to 34%. Because smoking rates in high school graduating classes after 1992 had been on the rise, and because we know that class cohorts tend to maintain their relative differences over time, we had predicted the increase in smoking among 19- to 22-year-olds and eventually in the older age bands as the heavier smoking high school class cohorts grew older. Beginning in 1996, smoking began to rise among 23- to 26-year-olds, before leveling after 1998. Again, it rose more among males, opening a small gender gap.

Regional Differences in Trends

The respondent's current state of residence was first asked in the 1987 follow-up survey; thus trend data by region exist only for the interval since then. In this case, changes have been examined for all 19- to 28-year-olds combined to increase estimate reliability. Because gender, for example, crosscut all regions, it has less sampling error than when the sample is divided into four separate regions. (Each region is represented by between 1,000 and 2,200 cases in all years.) By combining across all ages, we lose the ability to see the cohort effects that have occurred with many drugs, but we are able to see whether overall trends are similar across regions. In general, the changes that have occurred since 1987 have been fairly consistent across regions, particularly in terms of the direction of change. The four regions of the country—Northeast, Midwest, South, and West—have generally moved in parallel. There were substantial drops among young adults in all four regions between 1987 (the initial measurement point) and 1991 for *any illicit drug, marijuana, any illicit drug other than marijuana, cocaine, crack, and amphetamines*. Since 1991 in most or all regions, there has been some increase and then a leveling in the use of these drugs (except cocaine, which continued to decline through the mid-1990s, inched up thereafter, and then leveled since 2004).

- The proportion of 19- to 28-year-olds using *any illicit drug* has been consistently lowest in the South and highest in the West and Northeast. For *marijuana* use, the South has usually been lowest, and for the most part the Midwest has been second lowest. Generally, the other two regions have been fairly close to one another. For the use of *any illicit drug other than marijuana*, the West stood out as consistently highest, with the other three regions being very similar, at least until 2000; since 2001, use in the Northeast has been about as high as in the West, though the regional differences are not large.
- From 1991 through 1995, the West had slightly higher annual prevalence rates of *LSD* use than the other three regions among young adults (use dropped in 1995 in the West). Otherwise the usage rates have been quite similar in all four regions; all showed sharp declines in LSD use after 2001, though use had been declining in the Northeast for several years prior to that.

- Questions about *ecstasy (MDMA)* were added to the follow-up surveys of young adults in 1989. Through 1993, rates were highest in the West and South and lower in the Northeast and Midwest regions. Subsequently, use in the Northeast began to increase (as was true among 12th graders), exceeding levels of use found in the South and West from 1999 to 2001. The Midwest has consistently had a much lower level of ecstasy use than the other three regions. In 2000 all four regions showed a sharp and fairly parallel increase in ecstasy use; the rise decelerated in 2001 and began to decline thereafter in all regions. As discussed elsewhere, we believe that this decrease may be caused by growing concern about the hazards of ecstasy use. By 2003, little regional difference remained in annual prevalence, largely because the declines in use were most pronounced in the Northeast and West. By 2007, use was down a little more in all regions; but in 2008 ecstasy use increased some in three of the four regions.
- The considerable declines in *cocaine* use, observed in all regions between 1987 and 1991, were greatest in the two regions that had attained the highest levels of use by the mid-1980s—the West and Northeast. Thus, regional differences had diminished considerably by 1992. Similar to the finding for 12th graders, in 1992 these declines stalled in all regions except the Northeast. A gradual further decline then occurred in all regions through 1996 (1997 for the West) before a slight rise began to occur, likely reflecting the effects of generational replacement. Regional variability in annual cocaine prevalence has been minimal since the mid-1990s, though use in the West and Northeast has been a little higher than the Midwest and South.
- All four regions also exhibited an appreciable drop in *crack* use between 1987 and 1991, again with the greatest declines in the West and Northeast, where prevalence had been the highest. Use then generally leveled in all regions except the South, where it continued a gradual decline through 1997. As was true for cocaine generally, annual prevalence rates among the regions have converged; they now stand between 0.7% in the South and West, and 1.1% in the Midwest and Northeast. (Lifetime use of crack has been highest in the West since crack use was first measured in 1987, as is true for cocaine in general.)
- From 1987 (when data were first available) through 1994, rates of *inhalant* use remained relatively stable, quite low, and about equal in all four regions among 19- to 28-year-olds. Annual use then rose in the Northeast in 1995 and 1996 and remained higher than in the other regions through 2000, when it dropped back to rates comparable to the other three regions. Except for that divergence, the regions have moved very much in parallel for this class of drugs. Annual prevalence in 2008 is at low levels in this age group, between 0.9% and 1.7%.
- The regions have trended fairly similarly in their prevalence of *amphetamine* use by young adults. The only modest exception was that use declined more in the Northeast (which started out lowest) in the period 1987 to 1992, giving it a substantially lower rate than the other three regions; it remained lowest until 1998. The West has fairly consistently had the highest rate, but not by much. By the late 1990s, the Northeast had caught up to the Midwest and South, making the regional differences very small; there have been essentially no regional differences since 2000.

- **Methamphetamine** use in general has been measured only since 1999 (though crystal methamphetamine, discussed next, has been in the study for a longer time). It shows some differences in rates among the regions and some differential trending, with a gradual decline in annual prevalence in the Northeast (where use has generally been lowest) and a gradual increase in the West (where use has usually been highest) in 2000–2004. Use in the other two regions remained fairly flat until 2006, when both showed some decline. Use in the West has fallen since 2006, leaving very little variability among regions by 2008. Lifetime prevalence has been particularly high in the West, ranging between 11% and 16% since 1999, though it has been declining.
- The West has consistently had the highest rates for **crystal methamphetamine (ice)** use, and the regional differences have been very substantial, particularly in terms of lifetime use. The Northeast has generally had the lowest rates. When data were first available on crystal methamphetamine in 1990, the West had a lifetime prevalence of 5.1% versus a range of 1.7% to 2.3% in the other three regions. By 2006, the lifetime prevalence rate in the West had increased to 8.8%, and lifetime prevalence in the Midwest and South grew quite steadily over that interval. This strongly suggests that crystal methamphetamine use among young adults diffused from the West primarily to the South and Midwest regions, but diffused much less to the Northeast, which has had the lowest prevalence since 2001. The annual prevalence figures tell a similar story, but also show that there was a spike in past-year use in the West from 1991 to 1995 before use declined and then stabilized at around 2% from 1997 through 2001. Rates then rose again in the West between 2001 and 2003 and stabilized at a higher level around 2.7%. In 2007 and 2008, use in the West declined, narrowing the differences with the other regions.
- The annual prevalence for **sedatives (barbiturates)** remained flat, and at about equivalent levels, in all four regions of the country from 1987 through 1994. Rates then rose gradually in all regions for a number of years, followed by some leveling; regional differences have been consistently small.
- The picture for **tranquilizers** is similar to that for sedatives (barbiturates). Use generally held fairly steady in all regions from 1987 through 1993. Since then there has been some increase in all regions, with the South experiencing the most increase through 2004, after which all regions showed a leveling in use. The regional differences have been small, though the South tends to have a slightly higher rate—this difference grew a bit larger during the period of increasing use in the late 1990s.
- Levels and trends in **heroin** use have been quite comparable across the four regions since 1987. All regions had low and stable rates through the early 1990s. A gradual increase was observed from about 1993 through 2000, and prevalence has been stable since then. On average the Northeast has tended to have the highest rate of heroin use, but the annual prevalence rates have remained below 1% in all regions since 1987.
- Trends in prevalence of the use of **narcotics other than heroin** have also been quite parallel for the four regions. A gradual and long-term increase occurred from the mid-

1990s through 2003 or 2004, depending on the region, with little systematic change thereafter. The South has tended to have the lowest rate in recent years.

- The annual prevalence of *OxyContin* use was highest in the Northeast and lowest in the West in 2002, when it was first measured. In 2008, that pattern has reversed, with the West being highest (5.3%) and the Northeast lowest (2.9%). Use has risen some in all regions since 2002 except in the Northeast, and the rise has been most sharp in the West. Annual prevalence of use for *Vicodin* has shown some variation among the regions. The West and Midwest have generally had the highest rates, with the South the lowest and the Northeast in between. There has been little net change between 2002 and 2008 in the annual prevalence in each region, but use patterns in the interim have been different for each region. (It should be noted that the sample sizes are more limited than usual for these two drugs because questions about them occur on only two of the six questionnaire forms.)
- When two club drugs, *ketamine* and *GHB*, were first measured in 2002, the Northeast stood out as having a higher rate of annual use; but use in the Northeast has dropped since then, bringing that region's usage rates down to the same very low levels as the other three regions. *Rohypnol* use has remained very low in all four regions over the same interval, not reaching 1% in any region.
- With respect to *alcohol* use, there were modest declines in 30-day prevalence in all four regions between 1987 (when the first measurement was available for 19- to 28-year-olds) and 1992. The rates for 30-day prevalence then leveled in all regions. The West and South have consistently had lower rates of 30-day use than the Northeast and Midwest (as has generally been true among 12th graders).

Current *daily use* of alcohol also showed a decline from the first (1987) data collection through about 1994 or 1995 in all regions. The proportional declines were substantial—on the order of 40–50%. (This decline corresponds to a period of appreciable decline in daily drinking among 12th graders, though we can tell from their longer term data that their decline started in 1980; thus the decline may well have started earlier among 19- to 28-year-olds as well.) Since the mid-1990s there has been some upward trending in daily prevalence in all regions; the rates are all between 4.7% and 6.3% in 2008, with the Northeast having the highest average rate over the past five years and the South the lowest.

Occasional heavy drinking (or “binge drinking”) was fairly level in all regions between 1987 and the late 1990s or early 2000s. There have been some modest increases in recent years, particularly in the Northeast and West. The rates have consistently been appreciably higher in the Northeast (43% in 2008) and Midwest (42%) than in the West (35%) and South (34%).

- There have been highly consistent regional differences among young adults in *cigarette smoking* since data were first available in 1987—they exist for monthly, daily, and half-pack-daily prevalence rates. The West has consistently had the lowest rates (e.g., 11%

daily prevalence in 2008, compared to 17–19% in the other three regions) and the Northeast and Midwest the highest. The South has generally fallen in between. After some slight decline in 30-day prevalence in all regions between 1987 and 1989, rates leveled off for about five years (roughly through 1994). There then followed a very gradual increase of a few percentage points through 1998, followed by a gradual decrease. Daily use showed a very similar pattern. For half-pack-a-day smoking, the decline phase was longer (from 1987 through about 1992 or 1993), likely reflecting the lag between smoking initiation and regular heavy smoking. Since 1998, half-pack smoking rates have gradually declined in all four regions. In general, all of these measures have shown parallel movements across regions, suggesting that the forces accounting for whatever changes there have been are nationwide in scope. It should be remembered that, as illustrated earlier in this chapter, there are strong cohort effects in smoking which are weakened to some degree when we combine age groups across a 10-year age span, as we have done in the present analyses.

Population Density Differences in Trends

The analyses presented here for population density return to the use of four-year age groupings, which allows a longer time interval to be examined for the younger strata and for cross-age comparisons of the trends. Among young adults, five levels of population density are distinguished based on the respondent's answer to the question, "In March of this year did you live mostly . . .?"; answer alternatives were in a very large city (over 500,000 people), in a large city (100,000 to 500,000), in a medium-sized city (50,000 to 100,000), in a small city or town (under 50,000), or on a farm/in the country? Suburbs of cities of each size were combined with the city.

- The proportions of young adults using *any illicit drug* have moved in parallel among the various community-size strata. In general, the farm/country stratum has tended to have the lowest use. The other four community-size strata have tended to differ little from one another, though the very large cities have generally ranked at the top. In 2008, the proportions of 19- to 22-year-olds reporting use of an illicit drug in the past year were 26% for the farm/country stratum, 31% for small towns, 39% for medium-sized cities, 40% for large-sized cities, and 41% for very large cities.
- Trends in the use of *any illicit drug other than marijuana* tell a similar story. There was a long period of fairly parallel decline along with some convergence of usage rates among the community-size strata at all three age levels (among 19- to 22-year-olds it was between 1981 and 1992) followed by an increase in use and then a leveling. In general, medium, large, and very large cities have all tended to have about the same rates, and the farm/country stratum has tended to have the lowest rates, particularly prior to 1990; the differences by population density have been quite small since about 2000, ranging from 15% to 20% across the different community-size strata among 19- to 22-year-olds in 2008, for example. These trends have been somewhat staggered across the different age strata, indicating some cohort effects.
- *Marijuana* use has moved pretty much in parallel among the various community-size strata over the time intervals for which data exist. Among 19- to 22-year-olds, the rates

have been quite close among all communities, except for the farm/country stratum. The most rural stratum has consistently had the lowest rate of marijuana use, and it fell less in the decline period and rose more slowly in the subsequent increase than in the other community-size strata. Use has also tended to be lower in the more rural areas in the older two age bands. *Daily marijuana use* has also moved in parallel among the five population-density strata within each age band.

- In general, there have not been large differences in *LSD* use among adults as a function of community size since 1983. Among 19- to 22-year-olds (the young adult age group with by far the highest rates of LSD use prior to 2003), use in communities of all sizes declined appreciably in the early to mid-1980s, particularly in the urban strata, eliminating modest prior differences by 1984. From around 1989 through 1996, there was some increase in LSD use in all population-density strata among 19- to 22-year-olds, with the most rural region generally continuing to have the lowest prevalence. After 1997, there was some decline in LSD use in all community-size strata among 19- to 22-year-olds, followed by a sharp decline occurring from 2001 to 2003. The 23- to 26-year-old respondents had some modest increases after 1989 in all community-size strata, though the increases had virtually ended by 1995; since about 1999, there have been declines in all strata, with the largest decline in 2001–2003 in most regions. In *Volume I* in this series, we discussed how a sharp decline in supply may be responsible for the sizable decline in use among all ages after 2001. In the oldest age group, LSD use has remained very low and for the most part quite stable, with a small decline in the last few years.
- The use of *hallucinogens other than LSD*, taken as a class, has also shown considerably higher rates in the youngest age band compared to the two older ones, suggesting a sharp falloff in use with age. (The drug most often reported in this general class has been psilocybin or “shrooms,” as is true among 12th graders as well.) Use of this general class of drugs fell in communities of all sizes among young adults between 1980 and about 1988. Then there was a leveling of use for a few years, followed by an extended increase in use among all community-size strata in the 19- to 22-year-old age band. By 2003 the rates attained by each stratum exceeded those originally observed in 1980, but there have been declines since then, with very large cities being the last to show the decline. The 23- to 26-year-old group showed slightly higher rates of use from 1998 to 2004, followed by a leveling. The sharpest increase occurred in the very large cities in 1999 and 2000, possibly as a result of growing ecstasy use. The 27- to 30-year-olds have generally had low rates of use, and the trend lines have been very flat with only minor community-size stratum differences, until 2001 when all strata, especially the very large cities, began to increase before leveling after 2005. All three age groups have shown higher rates of use than previously observed—since 1994 for 19- to 21-year-olds, since about 2000 for 22- to 26-year-olds, and since about 2002 for 27- to 30-year-olds—in part reflecting a cohort effect in the use of these drugs, but also reflecting the change in question wording to include “shrooms” as an example.
- *Ecstasy (MDMA)* use was first measured in 1989, and since then has shown the largest increase among young adults of any drug. Use in 1989 was highest among 19- to 22-year-olds in the very large cities (5% annual prevalence); but prevalence declined in all

population-density strata between 1989 and 1994 (to 1.6% or less). By 1998, use had begun to increase in all community-size strata within this age band, except in the farm/country stratum. The farm/country stratum moved up sharply in 1999, but then the three most urban strata jumped sharply in 2000, opening a fair gap in use, with large and very large cities having rates nearly twice as high as any of the other strata in 2002. All community-size strata showed large declines in ecstasy use since 2000 or 2001, which lasted through 2004, narrowing the differences among them. Among 23- to 26-year-olds, use began to increase a little later, and again the most urban stratum showed the most increase, particularly in 2000; but use began to decline after 2001 (in the urban areas) or 2002 (in the rural and small town strata), which had the effect of narrowing the differences among strata. All population-density strata have continued to decline, or at least remain level, since 2003 among the 23- to 26-year-olds, but there has been a gradual increase in ecstasy use among 19- to 22-year-olds in all population strata in the past year or so. Considerably less increase in ecstasy use occurred among 27- to 30-year-olds, though there was some increase in the largest cities starting after 1996, and in the large and medium-sized cities after 1999. Since 1997 the very large cities have stood out as having the highest rates of ecstasy use.

Ecstasy use trends in the past several years tell an interesting story. In the very large cities, where use had spiked early, use peaked in all three age bands in 2000 and then began to decline. The medium-sized cities were beginning to level or decline in all three age bands. The small town and farm/country strata peaked in 2001 in all age groups. These data support our belief, based on school-level analyses of secondary schools, that the presence of this drug was still diffusing geographically—in this case, from more urban to more rural areas—and, were it not for this continued diffusion, ecstasy use would actually have declined nationally a year earlier. The data from 12th graders on perceived risk provide the clue as to the most likely cause of this turnaround; they showed a large jump in the level of perceived risk associated with ecstasy use from 2000 through 2003. Unlike most other drugs discussed here, the pattern of change since the mid-1990s appears to reflect secular trends, with all age groups moving pretty much in parallel, more than cohort effects.

- In the early 1980s, *cocaine* use was positively correlated with population density, with the highest use in the very large cities. The important drop in cocaine use that began after 1986 slowed considerably after 1992 or 1993 in all three age strata and in communities of all sizes, by which time the positive association with population density had been virtually eliminated. Among 19- to 22-year-olds there has been a sustained increase in cocaine use among all community-size strata after about 1993 or 1994, and among 23- to 26-year-olds after about 1998. There has been some decline in the mid-2000s in all strata except large cities. As just stated, usage rates among the population-density strata tended to converge considerably during the period of decline, and this convergence remains, with the very large cities showing rates of cocaine use only slightly higher than the less densely populated areas. In the 27- to 30-year-old age group, a gradual increase in use has emerged in nearly all population-density strata since 2000, no doubt reflecting a cohort effect working its way up the age spectrum. By 2004 all of these strata in the

oldest age band leveled or declined from their peak rates, the single exception being within very large cities, where use has remained relatively high.

- **Crack** use among all age groups peaked in 1987 or 1988 (strongly suggesting a secular trend at work at that time) and then, after declining appreciably, bottomed out in all population-density strata for several years. Use reported in these young adult samples at all three age levels has borne practically no systematic association with community size, and for the most part the strata have all tended to move in parallel.
- **Amphetamine** use shows virtually no differences associated with urbanicity in any of the three age groups, and this has generally been true since 1983. The trend curves are highly similar for all levels of population density within each age stratum.
- Use of **crystal methamphetamine (ice)**, first measured in 1990, has shown a modest increase since the early 1990s among the two youngest age strata, and since the late 1990s among the oldest stratum. This has been observable in most population-density groupings. It is difficult to see any sustained differences in use as a function of population density, but the trend lines across time are quite erratic due to the small numbers of cases.
- The use of **methamphetamine** in any form has been measured only since 1999. In general, the farm/country stratum has had higher than average rates of use in the two youngest age groups: otherwise there has been little systematic difference. Among 19- to 22- year-olds, all community-size strata have shown a decline in use since 2003 or 2004, reaching very low levels by 2008. Use has held quite steady over the same interval among 23- to 26-year-olds, while it has generally declined in all population-density strata since 2002 among 27- to 30-year-olds.
- **Methaqualone** use, which in 1981 was strongly positively associated with population density, dropped to annual prevalence rates of 0.8% or below in all community-size strata for all three age bands by 1989. For that reason, its use is no longer measured in MTF.
- Unlike methaqualone, **sedatives (barbiturates)** have never shown much correlation with urbanicity, at least as far back as 1980. This remains true in all three age bands, and the trends have been quite parallel within each age band, showing a long period of increasing use from about 1993 for 19- to 22-year-olds, 1996 for 23- to 36-year-olds, and about 1999 for 27- to 30-year-olds. These increases leveled off sequentially after about 2003 to 2005.
- **Tranquilizer** use among young adults has also had little or no association with population density over this time interval. Like sedatives (barbiturates), there was an earlier period of decline, staggered inflection points, a long period of gradual increase, and then a leveling staggered up the age band from about 2003–2005.
- From 1980 to 1995, annual **heroin** prevalence was less than 1.0%—usually much less—in all population-density strata for all three age bands. After 1994, use among 19- to 22-year-olds in all community-size strata rose and reached 1.0% in the three most urban

strata by 1998. In fact, in the very large cities, it reached 2.1% in 2000 (vs. 0.3–0.6% in the other strata). Use levels have been lower among 23- to 26-year-olds and lower still among 27- to 30-year-olds, making it difficult to discern systematic differences among the population-density strata in those age bands. In 2008 the annual prevalence of heroin is 1.0% or lower in all community-size strata in all three young adult age bands.

- The annual use of *narcotics other than heroin* had some positive association with population density among 19- to 22-year-olds through the early 1990s; however, it has shown rather little association since then. Use of narcotics other than heroin has increased substantially in all community-size strata since 1993 in the case of 19- to 22-year-olds, since about 1996 in the case of 23- to 26-year-olds, and since about 1998 in the case of 27- to 30-year-olds; however, no systematic differentiation by community size is evident. Clearly a cohort effect is at work, and the increasing use of these drugs was quite widespread. Use tended to level off since about 2003 or 2004 in all three age bands.
- Unfortunately, sample sizes for two of the narcotic drugs of particular interest, *OxyContin* and *Vicodin*, are not sufficient to estimate population-density differences or trends with a reasonable degree of accuracy.
- The absolute levels of *inhalant* use have remained low in these age groups, particularly above age 22. However, during the mid- to late 1980s, there was a gradual increase in use among 19- to 22-year-olds in all community-size strata. No strong or consistent association with population density has appeared, though the very large cities have generally tended to have higher rates than the other areas among 19- to 22-year-olds, particularly in the period 1998 through 2000. Among 19- to 22-year-olds, there has been some falloff in use since the late 1990s in all population-density strata.
- There have been few differences as a function of population density in the 30-day prevalence of drinking *alcohol* among 19- to 22-year-olds, since data were first available in 1980, except that the farm/country stratum has tended to have lower-than-average use. In the two older age bands, however, there has been a fairly consistent positive correlation between population density and use of alcohol in the past 30 days. So, for example, in 2008 59% of 27- to 30-year-olds in the farm/country stratum have had alcohol in the prior 30 days, compared to 79% of those in very large cities. But there have been no consistent differences in current *daily drinking* associated with urbanicity in any of the three age bands. For *occasional heavy drinking*, all community-size strata have been fairly close across time at all three age levels, again with the exception that the farm/country stratum has fairly consistently shown a slightly lower rate of binge drinking in the youngest two age bands.
- *Cigarette smoking* has generally been negatively associated with urbanicity in all three age strata, without much evidence of differential trends related to degree of urbanicity, with one exception. Among 19- to 22-year-olds, all smoking prevalence measures rose from 1997 through 1999 in the farm/country and small town strata, while most other strata remained level. The differences in 1999 were most striking for half-pack-a-day smoking among the 19- to 22-year-olds—24% for farm/country, 19% for small town,

15% for both medium-sized and large cities, and 10% for very large cities—compared with 1985, when there was virtually no difference in half-pack-a-day smoking rates among these strata (all were at 18% or 19%). Thus, smoking among those in their early 20s has become more concentrated in the nonurban populations. Among 19- to 22-year-olds, there has been a decline in 30-day prevalence in most population-density strata since about 2000. Continuing declines in smoking among 12th graders would lead us to expect still further declines in the young adults, as well. Among the two older age groups, the farm/country stratum has been highest more often than not, but in general the rates are not widely dispersed among them.

TABLE 5-1
Trends in Lifetime Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28

(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change	
<i>Approximate</i>																									
<i>Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900		
Any Illicit Drug ^a	70.5	69.9	67.9	66.4	64.5	62.2	60.2	59.6	57.5	57.4	56.4	56.7	57.0	57.4	58.2	58.1	59.0	60.2	60.5	60.4	59.7	59.8	59.3	-0.4	
Any Illicit Drug ^a other than Marijuana	48.4	47.0	44.6	42.7	40.8	37.8	37.0	34.6	33.4	32.8	31.0	30.5	29.9	30.2	31.3	31.6	32.8	33.9	35.2	34.0	34.8	34.2	34.7	+0.5	
Marijuana	66.5	66.0	63.8	62.8	60.2	58.6	56.4	55.9	53.7	53.6	53.5	53.8	54.4	54.6	55.1	55.7	56.8	57.2	57.4	57.0	56.7	56.7	55.9	-0.8	
Inhalants ^b	12.3	12.7	12.6	13.2	12.5	13.4	13.5	14.1	13.2	14.5	14.1	14.1	14.2	14.2	14.3	12.8	12.4	12.2	11.6	10.3	10.9	9.1	9.5	+0.3	
Nitrites ^c	2.6	6.9	6.2	—	1.9	1.4	1.2	1.3	1.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Hallucinogens ^d	18.5	17.1	17.0	15.9	16.1	15.7	15.7	15.4	15.4	16.1	16.4	16.7	17.4	18.0	18.4	18.3	19.6	19.7	19.3	17.6	17.2	16.0	14.8	-1.3	
LSD	14.6	13.7	13.8	12.7	13.5	13.5	13.8	13.6	13.8	14.5	15.0	15.0	15.7	16.2	16.4	16.0	15.1	14.6	13.4	11.2	10.1	9.6	8.1	-1.5 ss	
Hallucinogens other than LSD ^d	12.6	11.4	10.6	9.4	9.1	8.4	8.0	7.6	7.4	7.8	7.9	8.4	9.4	9.3	9.9	12.0	15.0	16.4	15.6	15.4	14.9	14.1	13.0	-1.1	
PCP ^e	8.4	4.8	5.0	—	2.5	3.1	2.0	1.9	2.0	2.2	1.9	2.4	2.7	2.3	2.3	3.1	2.5	3.0	2.7	2.0	2.4	2.1	2.2	+0.1	
Ecstasy (MDMA) ^f	—	—	—	3.3	3.7	3.2	3.9	3.8	3.8	4.5	5.2	5.1	7.2	7.1	11.6	13.0	14.6	15.3	16.0	14.9	14.4	13.1	13.1	-0.1	
Cocaine	32.0	29.3	28.2	25.8	23.7	21.0	19.5	16.9	15.2	13.7	12.9	12.0	12.3	12.8	12.7	13.1	13.5	14.7	15.2	14.3	15.2	14.7	14.8	0.0	
Crack ^g	—	6.3	6.9	6.1	5.1	4.8	5.1	4.3	4.4	3.8	3.9	3.6	3.8	4.3	4.6	4.7	4.3	4.7	4.2	4.1	4.4	3.9	4.3	+0.5	
Other Cocaine ^h	—	28.2	25.2	25.4	22.1	19.8	18.4	15.1	13.9	12.4	11.9	11.3	11.5	11.8	11.7	12.1	12.8	13.5	14.4	13.3	14.4	14.0	13.9	-0.1	
Heroin	1.3	1.3	1.1	1.0	0.9	0.9	0.9	0.8	1.1	1.3	1.3	1.3	1.6	1.7	1.8	2.0	1.8	1.9	1.9	1.7	1.9	1.6	1.9	+0.3	
With a Needle ⁱ	—	—	—	—	—	—	—	—	—	0.4	0.4	0.3	0.4	0.6	0.4	0.6	0.4	0.5	0.4	0.6	0.6	0.5	0.5	+0.0	
Without a Needle ⁱ	—	—	—	—	—	—	—	—	—	1.0	1.4	1.5	1.7	1.9	2.1	2.1	1.8	2.2	2.1	1.8	2.4	1.9	2.1	+0.2	
Narcotics																									
other than Heroin ^k	10.7	10.6	9.8	9.6	9.4	9.3	8.9	8.1	8.2	9.0	8.3	9.2	9.1	9.5	10.0	11.5	13.9	16.8	17.6	17.8	18.7	18.8	19.5	+0.6	
Amphetamines, Adj. ^{l,j}	32.3	30.8	28.8	25.3	24.4	22.4	20.2	18.7	17.1	16.6	15.3	14.6	14.3	14.1	15.0	15.0	14.8	15.2	15.9	14.6	15.6	15.3	14.6	-0.7	
Methamphetamine ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	-0.4	
Crystal Meth. (Ice) ^l	—	—	—	—	2.5	2.9	2.2	2.7	2.5	2.1	3.1	2.5	3.4	3.3	3.9	4.0	4.1	4.7	4.7	4.4	4.7	4.4	3.7	3.6	-0.1

(Table continued on next page.)

TABLE 5-1 (cont.)
Trends in Lifetime Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28

(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change	
<i>Approximate</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900		
<i>Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900		
Sedatives																									
(Barbiturates) ^j	11.1	9.7	8.9	7.9	8.7	8.2	7.4	6.5	6.4	6.7	6.6	6.5	6.9	7.4	8.1	7.8	8.0	8.7	9.7	10.0	9.5	9.8	10.6	+0.8	
Sedatives, Adj. ^{j,m}	16.7	15.0	13.2	12.1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Methaqualone ^l	13.1	11.6	9.7	8.7	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Tranquilizers ^{d,j}	17.6	16.5	15.1	13.5	12.9	11.8	11.3	10.5	9.9	9.7	9.3	8.6	9.6	9.6	10.5	11.9	13.4	13.8	14.9	14.5	15.0	14.5	15.8	+1.3	
Alcohol ^l	94.8	94.9	94.8	94.5	94.3	94.1	93.4	92.1	91.2	91.6	91.2	90.7	90.6	90.2	90.7	89.9	90.2	89.3	89.4	89.1	88.9	87.9	88.4	+0.5	
Been Drunk ^k	—	—	—	—	—	82.9	81.1	81.4	80.7	82.1	80.7	81.4	79.8	81.6	80.4	81.1	81.2	80.9	80.1	79.9	80.9	80.1	80.1	80.1	0.0
Flavored Alcoholic																									
Beverage ^p	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	83.2	84.6	84.4	84.0	82.6	-1.4	
Cigarettes	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Steroids ^q	—	—	—	1.1	1.2	1.7	1.9	1.5	1.3	1.5	1.5	1.4	1.4	1.9	1.4	1.4	1.6	1.8	1.9	1.8	1.8	1.7	1.8	+0.1	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

"—" indicates data not available.

See footnotes on next page.

Footnotes for Tables 5-1 through 5-4

^aUse of “any illicit drug” includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), methaqualone (until 1990), or tranquilizers not under a doctor’s orders.

^bThis drug was asked about in four of the five questionnaire forms in 1986–1989; *N* is four fifths of *N* indicated. Data was based on five of the six questionnaire forms in 1990–1998; *N* is five sixths of *N* indicated. Data was based on three of six questionnaire forms in 1999–2008; *N* is three sixths of *N* indicated.

^cThis drug was asked about in one questionnaire form. *N* is one fifth of *N* indicated in 1986–1988 and one sixth of *N* indicated in 1990–1994.

^dIn 2001 the question text was changed on three of the six questionnaire forms. “Other psychedelics” was changed to “other hallucinogens,” and “shrooms” was added to the list of examples. For tranquilizers, Miltown was replaced with Xanax. Beginning in 2002 the remaining forms were changed to the new wording.

^eThis drug was asked about in one of the five questionnaire forms in 1986–1988; *N* is one fifth of *N* indicated. Data was based on one of six questionnaire forms in 1990–2008; *N* is one sixth of *N* indicated.

^fThis drug was asked about in two of the five questionnaire forms in 1989; *N* is two fifths of *N* indicated. Data was based on two of the six questionnaire forms in 1990–2001; *N* is two sixths of *N* indicated. Data was based on three of the six questionnaire forms in 2002–2008; *N* is three sixths of *N* indicated.

^gThis drug was asked about in two of the five questionnaire forms in 1987–1989; *N* is two fifths of *N* indicated. Data based on all six questionnaire forms in 1990–2001. Data based on five of six questionnaire forms in 2002–2008; *N* is five sixths of *N* indicated.

^hThis drug was asked about in one of the five questionnaire forms in 1987–1989; *N* is one fifth of *N* indicated. Data based on four of the six questionnaire forms in 1990–2008; *N* is four sixths of *N* indicated.

ⁱThis drug was asked about in two of the six questionnaire forms; *N* is two sixths of *N* indicated.

^jOnly drug use that was not under a doctor’s orders is included here.

^kIn 2002 the question text was changed on three of the six questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only; *N* is three sixths of *N* indicated. In 2003 the remaining forms were changed to the new wording. The data are based on all forms in 2003 and beyond.

^lBased on the data from the revised question, which attempts to exclude the inappropriate reporting of nonprescription amphetamines.

^m“Sedatives, adjusted” data are a combination of barbiturate and methaqualone data.

ⁿIn 1993 and 1994, the question text was changed slightly in three of the six questionnaire forms to indicate that a “drink” meant “more than just a few sips.” Because this revision resulted in rather little change in reported prevalence in the surveys of high school graduates, the data for all forms combined are used in order to provide the most reliable estimate of change. After 1994 the new question text was used in all six of the questionnaire forms.

^oThis drug was asked about in three of the six questionnaire forms; *N* is three sixths of *N* indicated.

^pThis drug was asked about in one of the six questionnaire forms; *N* is one sixth of *N* indicated.

^qThis drug was asked about in one of the five questionnaire forms in 1989; *N* is one fifth of *N* indicated. Data based on two of the six questionnaire forms in 1990–2008; *N* is two sixths of *N* indicated.

^rDaily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes, for which actual daily use is measured, and for 5+ drinks, for which the prevalence of having five or more drinks in a row in the last two weeks is measured.

TABLE 5-2
Trends in Annual Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28
 (Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change	
Approximate Weighted N =	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900		
Any Illicit Drug ^a	41.9	39.3	36.3	32.8	30.7	27.0	28.3	28.4	28.4	29.8	29.2	29.2	29.9	30.3	30.8	32.1	32.4	33.0	33.7	32.8	32.1	32.5	33.8	+1.2	
Any Illicit Drug ^a other than Marijuana	27.0	23.9	21.3	18.3	16.7	14.3	14.1	13.0	13.0	13.8	13.2	13.6	13.2	13.7	14.9	15.4	16.3	18.1	18.8	18.5	18.4	18.1	18.9	+0.8	
Marijuana	36.5	34.8	31.8	29.0	26.1	23.8	25.2	25.1	25.5	26.5	27.0	26.8	27.4	27.6	27.9	29.2	29.3	29.0	29.2	28.2	27.7	28.5	28.6	+0.1	
Inhalants ^b	1.9	2.1	1.8	1.9	1.9	2.0	1.9	2.1	2.1	2.4	2.2	2.3	2.1	2.3	2.1	1.7	1.6	1.4	1.7	1.3	1.3	0.8	1.4	+0.6	
Nitrites ^c	2.0	1.3	1.0	—	0.4	0.2	0.1	0.4	0.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Hallucinogens ^d	4.5	4.0	3.9	3.6	4.1	4.5	5.0	4.5	4.8	5.6	5.6	5.8	5.2	5.4	5.4	4.7	4.7	5.2	4.7	4.5	4.1	3.8	3.8	0.0	
LSD	3.0	2.9	2.9	2.7	3.3	3.8	4.3	3.8	4.0	4.6	4.5	4.4	3.5	4.0	3.7	3.4	1.8	1.2	0.9	0.8	1.2	1.1	1.4	+0.3	
Hallucinogens other than LSD ^d	2.5	2.1	1.9	1.8	1.7	1.7	1.9	1.9	2.0	2.5	2.8	3.1	3.0	3.0	3.4	3.5	4.0	4.9	4.5	4.2	3.8	3.6	3.4	-0.2	
PCP ^e	0.8	0.4	0.4	—	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.5	0.6	0.6	0.3	0.6	0.3	0.3	0.1	0.6	0.2	0.3	0.4	+0.1	
Ecstasy (MDMA) ^f	—	—	—	1.4	1.5	0.8	1.0	0.8	0.7	1.6	1.7	2.1	2.9	3.6	7.2	7.5	6.2	4.5	3.5	3.0	3.0	2.5	3.3	+0.8	
Cocaine	19.7	15.7	13.8	10.8	8.6	6.2	5.7	4.7	4.3	4.4	4.1	4.6	4.9	5.4	5.4	5.8	5.8	6.6	7.1	6.9	6.6	6.2	6.0	-0.3	
Crack ^g	3.2	3.1	3.1	2.5	1.6	1.2	1.4	1.3	1.1	1.1	1.1	1.0	1.1	1.4	1.2	1.3	1.0	1.0	1.3	1.2	1.1	1.0	0.9	-0.1	
Other Cocaine ^h	—	13.6	11.9	10.3	8.1	5.4	5.1	3.9	3.6	3.9	3.8	4.3	4.5	4.8	4.8	5.3	5.6	6.1	6.4	6.3	5.9	5.6	5.5	-0.1	
Heroin	0.2	0.2	0.2	0.2	0.1	0.1	0.2	0.2	0.1	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.2	0.4	0.3	0.4	0.4	0.3	0.5	+0.2	
With a Needle ⁱ	—	—	—	—	—	—	—	—	—	0.1	0.1	0.1	0.1	0.1	*	0.3	*	*	0.1	0.2	0.3	0.1	0.1	0.0	
Without a Needle ⁱ	—	—	—	—	—	—	—	—	—	0.3	0.4	0.4	0.6	0.6	0.5	0.9	0.2	0.4	0.3	0.4	0.5	0.3	0.4	+0.1	
Narcotics other than Heroin ^{ik}	3.1	3.1	2.7	2.8	2.7	2.5	2.5	2.2	2.5	3.0	2.9	3.3	3.4	3.8	4.1	5.0	7.1	8.5	9.0	8.7	9.1	8.7	9.1	+0.4	
OxyContin ^{jl}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.9	2.6	3.1	3.1	3.1	2.9	3.9	+0.9	
Vicodin ^{jl}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	8.2	8.6	8.9	9.3	9.1	8.9	9.1	+0.2	
Amphetamines, Adj. ^{jl}	10.6	8.7	7.3	5.8	5.2	4.3	4.1	4.0	4.5	4.6	4.2	4.6	4.5	4.7	5.4	5.8	5.9	5.8	6.2	5.1	5.6	5.6	5.3	-0.3	
Ritalin ^{jl}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.9	2.9	2.7	2.5	2.6	2.4	2.4	0.0	
Methamphetamine ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.8	2.5	2.8	2.5	2.7	2.8	2.4	1.9	1.5	1.0	-0.5
Crystal Meth. (Ice) ⁱ	—	—	—	—	0.4	0.3	0.4	0.8	0.9	1.2	0.9	0.9	1.1	0.9	1.2	1.1	1.4	1.3	1.5	1.6	1.1	1.1	0.8	-0.3	

(Table continued on next page.)

TABLE 5-2 (cont.)
Trends in Annual Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28

(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change	
<i>Approximate</i>																									
<i>Weighted N =</i> 6,900 6,800 6,700 6,600 6,700 6,800 6,800 6,700 6,500 6,400 6,300 6,400 6,400 6,200 6,000 5,700 5,800 5,300 5,300 5,700 5,400 5,100 4,800 4,900																									
Sedatives																									
(Barbiturates) ^j	2.3	2.1	1.8	1.7	1.9	1.8	1.6	1.9	1.8	2.1	2.2	2.4	2.5	2.8	3.4	3.7	3.9	3.9	4.4	4.2	3.9	4.2	4.7	4.7	+0.5
Sedatives, Adj. ^{i,m}	3.0	2.5	2.1	1.8	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Methaqualone ^l	1.3	0.9	0.5	0.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers ^{d,i}	5.4	5.1	4.2	3.7	3.7	3.5	3.4	3.1	2.9	3.4	3.2	3.1	3.8	3.7	4.6	5.5	7.0	6.8	7.4	6.7	6.5	7.1	6.8	6.8	-0.4
Rohypnol ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
GHB ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Ketamine ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Alcohol ⁿ	88.6	89.4	88.6	88.1	87.4	86.9	86.2	85.3	83.7	84.7	84.0	84.3	84.0	84.1	84.0	84.3	84.9	83.3	84.4	83.8	84.4	84.0	84.0	83.6	-0.4
Been Drunk ^o	—	—	—	—	—	62.0	60.9	61.1	58.8	61.6	59.9	63.2	59.6	63.2	60.6	63.1	61.8	62.9	63.8	63.5	65.7	65.8	66.0	66.0	+0.3
Flavored Alcoholic																									
Beverage ^p	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cigarettes	40.1	40.3	37.7	38.0	37.1	37.7	37.9	37.8	38.3	38.8	40.3	41.8	41.6	41.1	40.9	41.1	39.1	38.6	39.0	39.1	36.9	36.2	35.0	35.0	-1.2
Steroids ^q	—	—	—	0.5	0.3	0.5	0.4	0.3	0.4	0.5	0.3	0.5	0.4	0.6	0.4	0.4	0.4	0.5	0.5	0.5	0.3	0.7	0.4	0.4	-0.3

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: $s = .05$, $ss = .01$, $sss = .001$.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

“—” indicates data not available.

“***” indicates a prevalence rate of less than 0.05%.

See relevant footnotes at the end of Table 5-1.

TABLE 5-3
Trends in 30-Day Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28

(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
<i>Approximate</i>																								
<i>Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	
Any Illicit Drug ^a	25.8	23.4	20.5	17.7	15.9	15.1	14.8	14.9	15.3	15.8	15.8	16.4	16.1	17.1	18.1	18.8	18.9	19.9	19.1	18.6	18.5	18.9	19.3	+0.4
Any Illicit Drug ^a other than Marijuana	13.0	10.7	9.5	7.5	6.0	5.4	5.5	4.9	5.3	5.7	4.7	5.5	5.5	6.0	6.4	7.0	7.7	8.3	8.5	8.2	8.1	8.6	8.9	+0.2
Marijuana	22.0	20.7	17.9	15.5	13.9	13.5	13.3	13.4	14.1	14.0	15.1	15.0	14.9	15.6	16.1	16.7	16.9	17.3	16.5	15.8	15.7	16.0	16.0	0.0
Inhalants ^b	0.4	0.6	0.6	0.5	0.6	0.5	0.6	0.7	0.5	0.7	0.5	0.5	0.7	0.8	0.5	0.4	0.5	0.3	0.3	0.2	0.3	0.2	0.4	+0.3
Nitrites ^c	0.5	0.5	0.4	—	0.1	*	0.1	0.2	0.1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Hallucinogens ^d	1.3	1.2	1.1	1.1	0.9	1.1	1.5	1.2	1.4	1.7	1.2	1.5	1.4	1.3	1.2	1.2	0.9	1.2	0.9	0.8	0.7	0.9	0.9	0.0
LSD	0.9	0.8	0.8	0.8	0.6	0.8	1.1	0.8	1.1	1.3	0.7	0.9	1.0	0.8	0.8	0.7	0.3	0.2	0.1	0.1	0.2	0.2	0.4	+0.2 s
Hallucinogens other than LSD ^d	0.6	0.6	0.4	0.5	0.4	0.3	0.5	0.6	0.6	0.6	0.6	0.7	0.5	0.6	0.7	0.6	0.8	1.2	0.9	0.8	0.6	0.8	0.7	-0.1
PCP ^e	0.2	0.1	0.3	—	0.2	0.1	0.2	0.2	0.1	*	0.1	0.1	0.2	0.2	*	*	0.1	0.1	0.1	*	*	*	0.1	+0.1
Ecstasy (MDMA) ^f	—	—	—	0.4	0.2	0.1	0.3	0.3	0.2	0.4	0.3	0.6	0.8	1.3	1.9	1.8	1.3	0.8	0.6	0.6	0.7	0.5	0.6	+0.1
Cocaine	8.2	6.0	5.7	3.8	2.4	2.0	1.8	1.4	1.3	1.5	1.2	1.5	1.7	1.9	1.7	2.2	2.2	2.4	2.2	2.2	2.3	2.1	1.9	-0.2
Crack ^g	—	1.0	1.2	0.7	0.4	0.4	0.4	0.4	0.3	0.2	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.4	+0.1
Other Cocaine ^h	—	4.8	4.8	3.4	2.1	1.8	1.7	1.1	1.0	1.3	1.1	1.5	1.5	1.6	1.5	1.8	2.0	2.1	2.1	1.9	1.9	2.0	1.7	-0.3
Heroin	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.3	*	0.1	0.1	0.1	0.2	0.1	0.1	0.0
Narcotics other than Heroin ^{i,k}	0.9	0.9	0.7	0.7	0.7	0.6	0.7	0.7	0.6	0.9	0.7	0.9	0.9	1.2	1.4	1.7	2.9	2.9	3.0	3.5	3.2	3.4	3.6	+0.2
Amphetamines, Adj. ^{j,l}	4.0	3.2	2.7	2.1	1.9	1.5	1.5	1.5	1.7	1.7	1.5	1.7	1.7	1.9	2.3	2.4	2.5	2.5	2.4	2.1	2.2	2.3	2.2	-0.2
Methamphetamine ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	0.7	1.0	0.7	0.6	0.7	0.5	0.6	0.3	-0.3
Crystal Meth. (Ice) ^j	—	—	—	—	—	*	0.1	0.3	0.5	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.5	0.4	0.4	0.6	0.3	0.3	0.3	0.0

(Table continued on next page.)

TABLE 5-3 (cont.)
Trends in 30-Day Prevalence of Various Types of Drugs
among Respondents of Modal Ages 19–28

(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
<i>Approximate</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	4,900
<i>Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	4,900
Sedatives																								
(Barbiturates) ^l	0.7	0.7	0.7	0.5	0.6	0.5	0.5	0.6	0.6	0.8	0.8	0.9	0.9	1.1	1.3	1.7	1.5	1.5	1.8	1.7	1.5	1.6	1.9	+0.3
Sedatives, Adj. ^{l,m}	0.9	0.8	0.7	0.5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Methaqualone ^l	0.3	0.2	0.1	*	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers ^{di}	1.8	1.6	1.4	1.2	1.1	0.9	1.0	1.0	0.8	1.1	0.7	1.1	1.2	1.3	1.8	2.1	2.8	2.4	2.7	2.6	2.3	2.8	2.7	-0.2
Alcohol ⁿ	75.1	75.4	74.0	72.4	71.2	70.6	69.0	68.3	67.7	68.1	66.7	67.5	66.9	68.2	66.8	67.2	68.3	67.0	68.4	68.6	68.7	69.5	68.9	-0.5
Been Drunk ^e	—	—	—	—	—	35.4	35.6	34.2	34.3	33.0	33.2	35.6	34.2	37.7	35.7	36.8	37.1	37.8	39.0	39.0	42.1	41.4	40.7	-0.7
Flavored																								
Alcoholic Beverage ^p	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	29.5	27.6	24.9	25.9	26.7	+0.9
Cigarettes	31.1	30.9	28.9	28.6	27.7	28.2	28.3	28.0	28.0	29.2	30.1	29.9	30.9	30.3	30.1	30.2	29.2	28.4	29.2	28.6	27.0	26.2	24.6	-1.6
Steroids ^q	—	—	—	0.2	0.1	0.2	0.1	*	0.1	0.2	0.2	0.2	0.2	0.3	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.4	0.2	-0.2

Source: The Monitoring the Future study, the University of Michigan.

Notes: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

"—" indicates data not available.

"**" indicates a prevalence rate of less than 0.05%.

See relevant footnotes at the end of Table 5-1.

TABLE 5-4
Trends in 30-Day Prevalence of Daily Use of Various Types of Drugs
among Respondents of Modal Ages 19–28

(Entries are percentages.)

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
<i>Approximate</i>																								
<i>Weighted N =</i>	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	
Marijuana ^f	4.1	4.2	3.3	3.2	2.5	2.3	2.3	2.4	2.8	3.3	3.3	3.8	3.7	4.4	4.2	5.0	4.5	5.3	5.0	4.9	5.0	5.0	5.1	+0.1
Cocaine ^f	0.2	0.1	0.2	0.1	*	0.1	*	0.1	*	0.1	*	*	*	0.1	*	0.1	*	*	0.1	0.1	0.1	*	*	0.0
Amphetamines, Adjusted ^{g,h,r}	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.2	0.2	0.3	0.2	0.1	0.2	0.1	0.1	0.0
Alcohol																								
Daily ^{h,r}	6.1	6.6	6.1	5.5	4.7	4.9	4.5	4.5	3.9	3.9	4.0	4.6	4.0	4.8	4.1	4.4	4.7	5.1	4.5	5.2	5.4	5.6	5.3	-0.3
Been Drunk ^{o,r}	—	—	—	—	—	0.5	0.4	0.4	0.5	0.3	0.4	0.9	0.5	0.9	0.5	0.4	0.6	0.8	0.7	0.5	0.6	0.6	0.5	-0.1
5+ Drinks in a Row in Last 2 Weeks	36.1	36.2	35.2	34.8	34.3	34.7	34.2	34.4	33.7	32.6	33.6	34.4	34.1	35.8	34.7	35.9	35.9	35.8	37.1	37.0	37.6	37.8	37.9	+0.1
Cigarettes																								
Daily	25.2	24.8	22.7	22.4	21.3	21.7	20.9	20.8	20.7	21.2	21.8	20.6	21.9	21.5	21.8	21.2	21.2	20.3	20.8	19.6	18.6	17.3	16.7	-0.6
1/2 Pack+/Day	20.2	19.8	17.7	17.3	16.7	16.0	15.7	15.5	15.3	15.7	15.3	14.6	15.6	15.1	15.1	14.6	14.2	13.9	13.5	12.5	11.9	11.1	10.2	-0.9

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

The illicit drugs not listed here show a daily prevalence of 0.2% or less in all years.

*** indicates a prevalence rate of less than 0.05%.

“—” indicates data not available.

See relevant footnotes at the end of Table 5-1.

TABLE 5-5
Trends in Annual and 30-Day Prevalence of an Illicit Drug Use Index^a
among Respondents of Modal Ages 19–28
Total and by Gender

	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007– 2008 change
	Percentage who used in last 12 months																							
Any Illicit Drug	41.9	39.3	36.3	32.8	30.7	27.0	28.3	28.4	28.4	29.8	29.2	29.2	29.9	30.3	30.8	32.1	32.4	33.0	33.7	32.8	32.1	32.5	33.8	+1.2
Males	45.3	42.6	39.5	35.7	33.6	30.0	31.4	31.1	32.3	32.1	31.6	31.9	33.6	33.9	34.4	34.9	35.6	36.0	37.0	35.3	35.9	35.4	37.4	+2.1
Females	39.0	36.5	33.6	30.5	28.3	24.5	25.8	26.1	25.3	28.1	27.3	27.1	27.1	27.6	28.2	30.1	30.2	31.0	31.4	31.1	29.5	30.7	31.4	+0.7
Any Illicit Drug other than Marijuana	27.0	23.9	21.3	18.3	16.7	14.3	14.1	13.0	13.0	13.8	13.2	13.6	13.2	13.7	14.9	15.4	16.3	18.1	18.8	18.5	18.4	18.1	18.9	+0.8
Males	30.4	26.5	23.8	21.0	19.1	16.4	16.3	14.7	16.2	16.2	15.4	15.6	16.2	16.7	17.8	17.2	18.9	19.8	21.3	20.4	21.8	20.3	21.1	+0.8
Females	24.0	21.6	19.4	16.2	14.7	12.5	12.2	11.6	10.5	12.0	11.4	12.0	11.0	11.5	12.9	14.1	14.6	17.0	17.1	17.3	16.0	16.7	17.5	+0.8
	Percentage who used in past 30 days																							
Any Illicit Drug	25.8	23.4	20.5	17.7	15.9	15.1	14.8	14.9	15.3	15.8	15.8	16.4	16.1	17.1	18.1	18.8	18.9	19.9	19.1	18.6	18.5	18.9	19.3	+0.4
Males	29.9	27.1	23.7	21.1	18.8	18.3	17.9	17.4	19.5	18.6	19.0	19.8	20.1	20.0	21.5	21.9	22.8	22.4	23.1	22.0	22.5	22.7	22.8	+0.1
Females	22.2	20.2	17.8	15.0	13.5	12.5	12.4	12.9	12.1	13.5	13.3	13.8	13.2	15.0	15.6	16.6	16.3	18.3	16.3	16.4	15.7	16.4	16.9	+0.5
Any Illicit Drug other than Marijuana	13.0	10.7	9.5	7.5	6.0	5.4	5.5	4.9	5.3	5.7	4.7	5.5	5.5	6.0	6.4	7.0	7.7	8.3	8.5	8.2	8.1	8.6	8.9	+0.2
Males	15.2	12.3	10.6	9.1	6.8	6.6	6.5	5.9	7.1	6.8	5.7	6.8	7.1	7.3	7.8	8.1	8.5	9.2	10.6	9.2	10.2	10.0	10.0	0.0
Females	11.0	9.4	8.7	6.2	5.3	4.4	4.7	4.0	3.9	4.8	4.0	4.5	4.4	5.1	5.4	6.3	7.1	7.7	7.1	7.6	6.8	7.7	8.1	+0.4
	Approximate Weighted N																							
All Respondents	6,900	6,800	6,700	6,600	6,700	6,600	6,800	6,700	6,500	6,400	6,300	6,400	6,200	6,000	5,700	5,800	5,300	5,300	5,700	5,400	5,100	4,800	4,900	
Males	3,200	3,100	3,000	2,900	3,000	3,000	3,000	3,000	2,900	2,800	2,700	2,800	2,700	2,600	2,400	2,400	2,200	2,200	2,300	2,200	2,100	1,900	2,000	
Females	3,700	3,700	3,700	3,700	3,600	3,600	3,700	3,600	3,600	3,600	3,600	3,600	3,500	3,400	3,300	3,400	3,100	3,100	3,400	3,200	3,000	2,900	2,900	


Source: The Monitoring the Future study, the University of Michigan.

Notes: Level of significance of difference between the two most recent years: $s = .05$, $ss = .01$, $sss = .001$.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aUse of "any illicit drug" includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), methaqualone (until 1990), or tranquilizers not under a doctor's orders.

FIGURE 5-1
Any Illicit Drug: Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group




Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05	'06	'07	'08	
18	48.1	51.1	53.8	54.2	53.1	49.4	47.4	45.8	46.3	44.3	41.7	38.5	35.4	32.5	29.4	27.1	31.0	35.8	39.0	40.2	42.4	41.4	42.1	40.9	41.4	41.0	39.3	38.8	38.4	36.5	35.9	36.6		
19-20			55.8	54.5	53.4	50.2	47.4	45.9	45.7	42.6	39.5	39.4	35.7	32.3	28.1	29.1	30.5	32.2	35.6	36.1	36.7	40.6	40.4	39.3	38.4	39.4	38.1	38.0	38.9	36.3	35.2	35.5		
21-22				55.3	55.4	51.2	49.9	47.3	46.3	45.8	42.3	38.2	35.0	32.7	29.9	30.0	30.2	31.6	31.9	33.0	33.5	34.1	33.3	36.9	40.2	36.7	38.3	36.5	36.4	36.0	35.0	36.7		
23-24					51.7	48.9	44.0	47.8	42.8	37.9	36.6	31.4	30.7	27.0	29.2	29.8	27.3	28.5	27.6	27.3	27.4	31.1	29.6	31.1	35.2	34.6	34.5	31.9	32.7	34.1	34.4			
25-26						44.0	45.2	39.3	40.1	34.4	30.5	29.6	25.2	26.4	25.6	25.5	27.3	23.4	25.4	23.9	24.5	25.5	27.4	27.6	27.5	31.6	32.0	28.6	29.3	31.8				
27-28							38.4	36.2	32.5	30.9	27.4	23.9	24.6	23.6	23.9	23.7	20.7	22.0	20.8	21.4	22.9	22.9	26.3	26.8	24.3	25.7	28.5	30.1						
29-30								30.5	28.9	23.0	24.5	23.1	21.7	22.4	21.3	22.2	22.2	19.6	19.0	20.3	21.1	20.9	20.6	22.0	25.2	25.9	22.7	28.2						
31-32 ^a									23.7	23.8	21.9	22.3	22.4	19.8	21.7	21.2	19.3	17.7	17.6	20.2														
35										19.5	21.6	21.2	20.3	18.1	17.7	19.1	17.8	18.1	17.9	18.5	18.2	17.5	17.5	22.1										
40											20.3	16.7	17.2	15.8	18.2	15.8	17.5	19.1	16.2	17.4	17.5													
45												17.8	15.8	15.3	17.2	18.3	17.3																	
50																																		17.9

Source: The Monitoring the Future study, the University of Michigan.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).


FIGURE 5-2
Any Illicit Drug other than Marijuana: Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).


FIGURE 5-3a
Marijuana: Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).


FIGURE 5-3b
Marijuana: Trends in 30-Day Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.


^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-3c
Marijuana: Trends in 30-Day Prevalence of Daily Use
among Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.
^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-4
Inhalants:^a Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 32^{b,c}
by Age Group



Source: The Monitoring the Future study, the University of Michigan.


* indicates a percentage of less than 0.05%.

^aUnadjusted for the possible underreporting of amyl and butyl nitrites. Chapter 5, *Volume 1*, shows that such an adjustment would flatten the trend for seniors considerably because the line was adjusted up more in the earlier years, when nitrite use was more prevalent. Questions about nitrite use were dropped from the follow-up questionnaires beginning in 1995.

^bQuestions about the use of inhalants were not included in the questionnaires for 35-, 40-, 45-, and 50-year-olds.

^cBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-5
Hallucinogens:^a Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group




Source: The Monitoring the Future study, the University of Michigan.

^aUnadjusted for the possible underreporting of PCP.

^bBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-6
LSD: Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 35^a
by Age Group




Source: The Monitoring the Future study, the University of Michigan.

^a*** indicates a percentage of less than 0.05%.

^bQuestions about LSD use were not included in the questionnaires administered to the 40-, 45-, and 50-year-olds, or the 35-year-olds after 2006.

^cBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-7
Hallucinogens other than LSD:^a Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 35^b
by Age Group




^aUnadjusted for the possible underreporting of PCP.

^bQuestions about the use of hallucinogens other than LSD were not included in the questionnaires administered to the 40-, 45-, and 50-year-olds, or the 35-year-olds after 2006.

^cBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-8
Ecstasy (MDMA): Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 32^a
by Age Group




Source: The Monitoring the Future study, the University of Michigan.

* indicates a percentage of less than 0.05%.

^aQuestions about use of ecstasy (MDMA) were not included in the questionnaires administered to the 35-, 40-, 45-, and 50-year-olds.

^bBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).


FIGURE 5-9
Cocaine: Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.


^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-10
Crack Cocaine: Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.
^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-11
Heroin: Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group




Source: The Monitoring the Future study, the University of Michigan.

* indicates a percentage of less than 0.05%.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-12
Narcotics other than Heroin:^a Trends in Annual Prevalence
among Respondents of Modal Ages 18 through 50
by Age Group




Source: The Monitoring the Future study, the University of Michigan.

^aIn 2002 the question text was changed on half of the questionnaire forms for 18-30-year-olds. The list of examples of narcotics other than heroin was updated. Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. The 2001 data presented here are based on all forms. The 2002 data are based on the changed forms or in 2003 the remaining forms were changed to the new wording. The data are based on all forms in 2003. Beginning in 2002 data were based on the changed question text for 35- and 40-year-olds.


^bBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-13
Amphetamines: Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.
^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-14
Crystal Methamphetamine (Ice): Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 32^a
by Age Group




Source: The Monitoring the Future study, the University of Michigan.

* indicates a percentage of less than 0.05%.

^aQuestions about the use of crystal methamphetamine were not included in the questionnaires administered to the 35-, 40-, 45-, and 50-year-olds.

^bBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).


FIGURE 5-15
Sedatives (Barbiturates): Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).


FIGURE 5-16
Tranquilizers: Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-17
Steroids: Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 32^a
by Age Group




Source: The Monitoring the Future study, the University of Michigan.

* indicates a percentage of less than 0.05%.

^aQuestions about the use of steroids were not included in the questionnaires administered to the 35-, 40-, 45-, and 50-year-olds.

^bBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).


FIGURE 5-18a
Alcohol: Trends in Annual Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).


FIGURE 5-18b
Alcohol: Trends in 30-Day Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).


FIGURE 5-18c
Alcohol: Trends in 30-Day Prevalence of Daily Use among
Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-18d
Alcohol: Trends in 2-Week Prevalence of Having 5 or More Drinks in a Row
among Respondents of Modal Ages 18 through 50
by Age Group




Source: The Monitoring the Future study, the University of Michigan.

^aDue to a coding error for 18-year-olds, previous versions of this figure contained a value that was slightly off for the measure of five or more drinks in a row for 2005. This has been corrected here.

^bBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).


FIGURE 5-19a
Cigarettes: Trends in 30-Day Prevalence among
Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-19b
Cigarettes: Trends in 30-Day Prevalence of Daily Use among
Respondents of Modal Ages 18 through 50
by Age Group




Age	'76	'77	'78	'79	'80	'81	'82	'83	'84	'85	'86	'87	'88	'89	'90	'91	'92	'93	'94	'95	'96	'97	'98	'99	'00	'01	'02	'03	'04	'05	'06	'07	'08		
18	28.8	28.8	27.5	25.4	21.3	20.3	21.1	21.2	18.7	19.5	18.7	18.7	18.1	18.9	19.1	18.5	17.2	19.0	19.4	21.6	22.2	24.6	22.4	23.1	20.6	19.0	16.9	15.8	15.6	13.6	12.2	12.3	11.4		
19-20	31.0	31.2	29.3	26.0	23.9	24.4	24.1	23.2	21.9	22.5	19.5	18.9	19.2	19.4	20.5	21.1	21.9	22.2	22.5	22.7	23.8	25.6	22.7	21.9	20.6	18.8	18.2	17.6	14.4	12.9	14.3				
21-22			31.1	31.4	28.6	26.0	25.3	24.4	24.2	22.3	22.5	20.2	20.6	21.2	20.5	21.1	24.0	22.8	21.4	22.8	24.2	25.1	23.6	23.9	20.8	21.5	19.2	17.7	18.3	16.1					
23-24					30.1	30.6	27.8	25.1	25.2	26.0	24.0	23.3	22.2	22.5	20.9	20.1	19.9	20.0	22.8	21.5	21.2	21.4	21.2	22.4	23.5	21.5	23.3	20.4	19.5	17.5	17.9				
25-26							28.7	30.4	27.3	23.7	22.9	25.0	23.3	22.8	20.3	21.9	19.8	19.2	21.1	19.2	21.9	19.6	20.1	20.9	19.8	20.4	22.7	22.5	22.0	19.2	17.4				
27-28									27.6	27.9	25.0	22.9	22.2	23.9	21.8	20.1	20.5	20.9	19.4	17.6	19.5	16.0	19.7	17.2	18.1	19.8	18.2	18.6	20.2	19.3	18.3				
29-30												25.4	26.4	24.2	21.0	20.3	21.7	20.9	20.1	18.6	19.7	17.2	17.2	15.8	14.4	17.4	16.4	16.7	18.9	18.3	16.8	17.4			
31-32 ^a													23.9	24.9	22.8	21.4	20.9	21.2	19.8	19.1	17.9	18.9	18.1	16.1	—	—	—	—	—	—	—	—	—		
35																	22.5	23.0	22.1	18.3	20.4	19.7	20.1	16.4	18.2	16.3	14.8	14.5	13.5	13.9	16.5				
40																																			
45																																			
50																																			

Source: The Monitoring the Future study, the University of Michigan.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

FIGURE 5-19c
Cigarettes: Trends in 30-Day Prevalence of Smoking a Half Pack or More per Day
among Respondents of Modal Ages 18 through 50
by Age Group



Source: The Monitoring the Future study, the University of Michigan.

^aBeginning in 2002, respondents were followed through age 30 (instead of 32, as in past years).

Chapter 6

ATTITUDES AND BELIEFS ABOUT DRUGS AMONG YOUNG ADULTS

One of the most important theoretical and practical contributions of MTF has been to demonstrate the extent to which attitudes and beliefs about drugs determine use. Earlier volumes in this monograph series, as well as other publications from the study, have demonstrated that shifts in certain attitudes and beliefs—in particular the degree of risk of harm perceived to be associated with use of a particular drug—are important in explaining changes in actual drug-using behavior. Indeed, on a number of occasions we have accurately predicted such changes.⁵⁰ In this chapter, we review trends since 1980 in the attitudes and beliefs among young adults.

PERCEIVED HARMFULNESS OF DRUGS

Table 6-1 provides trends in the percentages of young adults who perceive a “great risk” of harm associated with differing usage levels of various licit and illicit drugs. These questions are contained in one questionnaire form only, limiting the numbers of follow-up cases; accordingly, we use four-year age bands in order to increase the available sample size (to about 300–600 weighted cases per year for each age band) and, thus, to improve the reliability of the estimates. (The numbers of weighted cases are given at the end of Table 6-1. The actual numbers of respondents are somewhat larger.) Still, these are relatively small sample sizes compared to those available for 8th, 10th, and 12th graders, and thus the change estimates may be relatively less stable. Because of the nature of the MTF design, trend data are available for a longer period for 19- to 22-year-olds (since 1980) than for 23- to 26-year-olds (since 1984) or 27- to 30-year-olds (since 1988). Also displayed in this table are comparison data for 12th graders, shown here as 18-year-olds, from 1980 onward. (See also Table 8-3 in *Volume I* for the longer term trends in 12th graders’ levels of perceived risk.) Questions about these attitudes and beliefs are not included in the questionnaires for 35-, 40-, 45-, and 50-year-old respondents.

- Table 6-1 illustrates considerable differences in the degree of risk young adults associate with various drugs. In general, the results closely parallel the distinctions made by 12th graders.
- **Marijuana** is seen as the least risky of the illicit drugs, although sharp distinctions are made between different levels of marijuana use. In 2008, experimental use of marijuana

⁵⁰Bachman, J. G., Johnston, L. D., O’Malley, P. M., & Humphrey, R. H. (1988). Explaining the recent decline in marijuana use: Differentiating the effects of perceived risks, disapproval, and general lifestyle factors. *Journal of Health and Social Behavior*, 29, 92–112; Bachman, J. G., Johnston, L. D., & O’Malley, P. M. (1990). Explaining the recent decline in cocaine use among young adults: Further evidence that perceived risks and disapproval lead to reduced drug use. *Journal of Health and Social Behavior*, 31, 173–184; Bachman, J. G., Johnston, L. D., & O’Malley, P. M. (1998). Explaining recent increases in students’ marijuana use: Impacts of perceived risks and disapproval, 1976 through 1996. *American Journal of Public Health*, 88, 887–892; Johnston, L. D. (1981). Characteristics of the daily marijuana user. In R. de Silva, R. L. DuPont, & G. K. Russell (Eds.), *Treating the marijuana-dependent person* (pp. 12–15). New York: The American Council on Marijuana; Johnston, L. D. (1985). The etiology and prevention of substance use: What can we learn from recent historical changes? In C. L. Jones & R. J. Battjes (Eds.), *Etiology of drug abuse: Implications for prevention* (NIDA Research Monograph No. 56, pp. 155–177). (DHHS Publication No. (ADM) 85-1335). Rockville, MD: National Institute on Drug Abuse.

is perceived as being of great risk by only 11–17% of all high school graduates ages 19–30, whereas regular use is perceived to carry great risk by about half (51–52%).

In the mid-1980s and early 1990s, fewer of the older age groups attached great risk to marijuana use than did the younger respondents. Indeed, there was a regular negative ordinal relationship between age and perceived risk for some years after 1980, when the first comparisons were available. Although this may have first looked like an age effect, the MTF design allows us to recognize it as a cohort effect: the younger cohorts initially perceived marijuana as more dangerous than the older cohorts and persisted in such beliefs as they grew older. Newer cohorts, however, have shown lower levels of perceived risk. Twelfth graders from the class of 2008 are much less likely to perceive regular marijuana use as dangerous than 12th-grade cohorts in the late 1980s and early 1990s. This reflects what we have interpreted as generational forgetting, a phenomenon wherein younger replacement cohorts no longer carry the knowledge—or had the direct or vicarious experience on which the knowledge is based—that the older cohorts had at that age. (The implications of this concept for prevention are discussed in the last section of this chapter.)

The decline in perceived risk that began in the 1990s was greater in the younger age bands, including grades 8 and 10, and least among the 27- to 30-year-olds. We believe that much of the decline in perceived risk that occurred in the older age bands resulted directly from generational replacement of earlier cohorts by later, less concerned ones. The credibility of this view is strengthened by the 1993–1995 reversal of the relationship between age and perceived risk of regular use. This reversal is consistent with an underlying cohort effect and could not simply reflect an association between age and a regular change in these attitudes. The decline in perceived risk for regular marijuana use ended in a somewhat staggered fashion—among 12th graders in 1999, among 19- to 22-year-olds in 2001, among 23- to 26-year-olds in 2002, and among 27- to 30-year-olds in 2004. In 2007 all four age strata showed declines of three to four percentage points in perceived risk for regular marijuana use, though no one of the declines was statistically significant. The decline continued in 2008 among 12th graders, and the two-year decline was statistically significant.

- Young adults view experimental use of any of the other illicit drugs as distinctly more risky than use of marijuana. About 28–32% of young adults ages 19 to 30 think trying *sedatives (barbiturates)* involves great risk; the corresponding figures are 36–42% for *amphetamines*, 42–51% for *LSD* or *ecstasy (MDMA)*, 44–48% for *cocaine powder*, 51–56% for *crack*, and 68–70% for *heroin*. (Note that the two classes of prescription drugs have the lowest levels of perceived risk among this set.)
- In the past, the older age groups were more likely to see *LSD* as dangerous. The age distinctions for LSD use became sharper through about 2001 as perceived risk declined more in the younger age groups—again indicating some important cohort changes in these attitudes, quite likely as a result of generational forgetting of the dangers of LSD. In recent years, there has been a decline in perceived risk in the older age groups, again

likely resulting from generational replacement as newer cohorts move up the age spectrum, which has begun to diminish the age differences.

- Young adults have generally reported somewhat higher perceived risk with respect to *cocaine* use than 12th graders, who have had less experience with this drug. Among 12th graders and the young adult age groups, the danger associated with using cocaine on a regular basis grew considerably (by 13 and 17 percentage points, respectively) between 1980 and 1986. However, these changed beliefs did not translate into changed behavior until the perceived risk associated with *experimental* and *occasional* use began to rise sharply after 1986. When these two measures rose, a sharp decline in actual use occurred.

We hypothesized that respondents saw only these lower levels of use as relevant to them and, therefore, saw themselves as vulnerable only to the dangers of such use. (No one starts out planning to be a heavy user; further, in the early 1980s, cocaine was not believed to be addictive.) Based on this hypothesis, we included the additional question about occasional use in 1986, just in time to capture a sharp increase in perceived risk later that year. This increase occurred largely in response to the growing media frenzy about cocaine—and crack cocaine, in particular—and to the widely publicized, cocaine-related deaths of several public figures (most notably Len Bias, a collegiate basketball star and high National Basketball Association draft pick). After stabilizing for a few years, perceived risk began to fall off after about 1991 among 12th graders, but not among the older age groups—again suggesting that lasting cohort differences were emerging, quite possibly as a result of generational forgetting of the dangers of cocaine in the younger age groups. A decline in perceived risk began among 19- to 22-year-olds starting after 1994, among 23- to 26-year-olds after 1997, and among 27- to 30-year-olds after 2001. There has been rather little systematic change in perceived risk for cocaine in recent years.

- A similar situation now exists for *crack*, for which perceived risk is highest in the oldest age band and lowest among 12th graders. Trend data (available since 1987) on the risks perceived to be associated with crack use show increases in 1987–1990 for all age groups, followed by relatively little change in the older age strata. Since 1990, twelfth graders have shown decreases in the perceived risk of experimental use of crack—perhaps reflecting the onset of generational forgetting of its dangers—leaving them as perceiving considerably less risk than the older groups. The young adult age groups have shown a staggered decline in this measure, with 19- to 22-year-olds showing a decline after 1994, 23- to 26-year-olds since 1996, and 27- to 30-year-olds after 2001. As a result, the different ages spread out more in their levels of perceived risk of crack use, until recent declines in the older age groups after about 2002.
- Measures of perceived risk of *crystal methamphetamine (ice)* use were introduced in 1990, and the results show what may be an important reason for its lack of rapid spread. More than half of all 12th graders and young adults perceived it as quite dangerous even to try, perhaps because it was likened to crack in many media accounts. (Both drugs come in crystal form, both are burned and the fumes inhaled, both are stimulants, and both can produce a strong dependence.) There was rather little age-related difference in

perceived risk associated with use of crystal methamphetamine in 1990 and 1991 (although the two youngest age groups were somewhat higher). But as perceived risk fell considerably among 12th graders (and eventually among 19- to 22-year-olds) and held steady or rose in the oldest two age groups, an age-related difference emerged. Now perceived risk has risen some among 19- to 22-year-olds, narrowing the age-related differences that had emerged for a few years. Still, 12th graders are less likely to see risk attached to trying crystal methamphetamine (62%) than are 19- to 22-year-olds (70%), again very likely due to generational forgetting.

- Questions about perceived risk of *ecstasy* (*MDMA*) were introduced in the follow-up surveys in 1989, but were not asked of 12th graders until 1997. At the beginning of the 1990s, all young adult age groups viewed ecstasy as a fairly dangerous drug, even for experimentation. But, again, the different age bands had diverging trends during the 1990s, with the oldest two age bands continuing to see ecstasy as quite dangerous, but the 19- to 22-year-olds (and very likely the 12th graders, for whom we did not have data until 1997) coming to see it as less so. In 2000, 38% of 12th graders saw great risk in trying ecstasy versus 49% of 27- to 30-year-olds; in 2001, the corresponding figures were 46% and 54%. In fact, three of the four age groups showed appreciable increases in perceived risk for ecstasy in 2001, which led us to predict a decline in use. The increase in perceived risk continued in 2002 in the two youngest age strata, and their use of ecstasy did, indeed, begin to decline—and decline sharply (see chapter 5). Perceived risk continued to rise through 2005 for the two youngest age groups, with the increase extending through 2007 among 23- to 26-year-olds. In 2008 there were substantial declines among the two oldest age bands in the perceived risk of trying ecstasy, but the younger groups were still more concerned than the older age group about the dangers of ecstasy use—a reversal of the situation that occurred in the late 1990s and early 2000s.
- Young adults have been more cautious about *heroin* use than 12th graders. In general, there has been relatively little change over the years in the proportions of all age groups seeing *regular* heroin use as dangerous; the great majority of each group (over 86%) consistently held this viewpoint. With regard to heroin experimentation, from 1975 to 1986 there had been a downward shift among 12th graders in the proportion seeing great risk associated with trying heroin. Following this decline (although their data do not extend back as far), young adults showed an increased caution about heroin use in the latter half of the 1980s—possibly due to the association of heroin injection with the spread of HIV—followed by a leveling through most of the 1990s. In 1996 and 1997, young adults' perceived risk increased some, as happened among 12th graders (as well as among 8th and 10th graders). These various trends may reflect, respectively, (a) the lesser attention paid to heroin by the media during the late 1970s and early 1980s as cocaine took center stage; (b) the subsequent great increase in attention paid to intravenous heroin use in the latter half of the 1980s due to the recognition of its important role in the spread of HIV/AIDS; (c) the emergence in the 1990s of heroin so pure that people no longer needed to use a needle to administer it, resulting in lower perceived risk; and (d) the subsequent increased attention given to heroin by the media (partly as a result of some overdose deaths by public figures and partly prompted by the emergence of “heroin chic” in the design industry), as well as through an anti-heroin media campaign launched by the

Partnership for a Drug-Free America in June 1996. At present, young adults still see heroin use as more dangerous than do 12th graders (Table 6-1); the differences have generally been largest for experimental and occasional use.

- A large minority of young adults see *heavy drinking on weekends* as dangerous (37–42%), as do 12th graders (46%). The belief that heavy drinking carries great risk has increased over the years in these age groups, rising among 12th graders from 36% in 1980 to 49% in 1992. Among 19- to 22-year-olds, it rose from a low of 30% in 1981 to 42% in 1992; the increases among the older groups were smaller. The increase in this belief may well help to explain the important decline in actual heavy drinking, and may in turn be explained by the media campaigns against drunk driving and the increase in the drinking age in a number of states.⁵¹ Following a staggered pattern, perceived risk peaked among 18-year-olds in 1992, among 19- to 22-year-olds in 1993, among 23- to 26-year-olds in 1994, and among 27- to 30-year-olds in 1995, suggesting a cohort effect. Since then, perceived risk of heavy drinking eased back some in all age strata.
- Between 1980 and 1991, a gradually increasing proportion of all four age groups viewed drinking *one or two drinks per day* as dangerous; but then they all showed a parallel decrease in perceived risk for this behavior through at least 2000. It seems likely that the earlier increase was due to the general rising concern about the consequences of alcohol use, particularly drunk driving, and that the subsequent decline was due to increasing reports of cardiovascular health benefits of light-to-moderate daily alcohol consumption. In recent years there has been little systematic change in this belief in any of the age strata, and there has been little difference by age.
- More than four fifths (80–87%) of young adults now perceive regular *pack-a-day cigarette smoking* as entailing high risk. In recent years, 18-year-olds have consistently shown lower perceived risk than young adults, while 10th graders have been still lower and 8th graders lowest. Clearly, there is an age effect in young people coming to understand the dangers of smoking. Unfortunately, it appears that much of the learning about the risks of smoking happens after a great deal of smoking initiation has occurred and many young people have already become addicted. These beliefs about smoking risks have strengthened very gradually in all age groups from senior year forward during the years we have monitored them (see Table 6-1). The parallel changes in these beliefs across the different age groups indicate a period effect, suggesting that all of these age groups were responding to common influences in the larger culture. These influences are discussed at some length in the chapter on attitudes and beliefs in *Volume I*. The rise in perceived risk has stalled some in recent years; only the oldest group (ages 27–30) sees a higher risk in 2008 than in 2002.
- The regular use of *smokeless tobacco* is seen as dangerous by 49–62% of young adults and 43% of 12th graders. These beliefs have also gradually strengthened in all age groups

⁵¹See O'Malley, P. M., & Johnston, L. D. (1999). Drinking and driving among U.S. high school seniors: 1984–1997. *American Journal of Public Health*, 89, 678–684. See also O'Malley, P. M., & Johnston, L. D. (2003). Unsafe driving by high school seniors: National trends from 1976 to 2001 in tickets and accidents after use of alcohol, marijuana and other illegal drugs. *Journal of Studies on Alcohol*, 64, 305–312.

over the intervals covered (Table 6-1). As with cigarettes, the change appears to reflect a secular trend (period effect), because it has been occurring in parallel for all age groups. Also like cigarettes, smokeless tobacco has seen the degree of perceived risk level off in the past several years—this time in all strata, including 12th graders.

PERSONAL DISAPPROVAL OF DRUG USE

The questions asked of 12th graders concerning the extent to which they personally disapprove of various drug-using behaviors among “people (who are 18 or older)” are also asked of follow-up respondents in one of the six questionnaire forms. Trends in the answers of young adults in the age bands of 19 to 22, 23 to 26, and 27 to 30 are contained in Table 6-2. Comparison data for 12th graders are also provided for 1980 onward. (See Table 8-6 in *Volume I* for the longer term trends in 12th graders’ levels of disapproval associated with using the various drugs.)

- In general, disapproval levels of adult use of the various drugs rank similarly across substances for both 12th graders and young adults. The great majority of young adults disapprove of using, or even experimenting with, all of the *illicit drugs other than marijuana*. For example, 95% or more of young adults in 2008 disapprove of regular use of each of the following drugs: *LSD*, *cocaine*, *amphetamines*, *sedatives (barbiturates)*, and *heroin*. Fully 78% to 96% of young adults disapprove of experimentation with each of these same drugs. Many of these attitudes differ rather little as a function of age at present; when there is a difference, the younger age groups are usually the least disapproving.
- Even for *marijuana*, about half of young adults now disapprove of experimentation (46–53%), nearly two thirds (63–65%) disapprove of occasional use, and the great majority (84–86%) disapprove of regular use.

Marijuana use shows the widest fluctuations in disapproval over time—generally, fluctuations that parallel the changes in perceived risk (though sometimes with a one-year lag, with the change in perceived risk coming first). The most fluctuation has occurred among the younger age groups (Table 6-2). Among 12th graders, disapproval of regular marijuana use increased substantially in the 1980s, peaked in the early 1990s, declined through much of the 1990s, and then leveled around 1998 with little change since then. The 19- to 22-year-olds had a quite similar pattern, though the decline continued a year longer, likely due to generational replacement. Among 23- to 26-year-olds, there were some declines starting later in the 1990s, but the declines have been very modest. In 2008 there was a decline among 12th graders in their disapproval of marijuana use (significant in the case of regular marijuana use). Whether this signals the beginning of a longer trend, and possibly cohort effect, should become clearer in the next survey.

- The great majority of all age groups have disapproved of even experimenting with *LSD* since 1980, when these data were first available. Beginning around 1990, all age groups decreased some in their disapproval of trying LSD (from high levels of disapproval, all at 90–91%). The decline was steepest among 12th graders, but there was a reversal in their

disapproval in 1997, and then an increase through 2006. Disapproval in the older age groups declined less, and in a staggered fashion; this trend has shown some evidence of a reversal among 19- to 22-year-olds and 23- to 26-year-olds since 2001 and 2002, respectively. This pattern again suggests some lasting cohort differences in these attitudes. All four age bands showed some decline in their disapproval of LSD use in 2008, while two of the four showed a parallel decline in perceived risk for this drug. It appears that during this recent period of very low rates of use, attitudes about this drug are softening.

- Since 2001, when it was first measured among young adults, disapproval of *ecstasy* use has risen in all age groups, and now all have very high levels. Experimenting with ecstasy is disapproved of by 88% of 12th graders in 2008, declining only slightly with age to 83% among 27- to 30-year-olds. This decline with age may reflect cohort differences in past experience with ecstasy. At the beginning of the decade the age differences were in the opposite direction.
- Disapproval of all three levels of *heroin* use has remained very high and stable since MTF began. There was one minor exception, however: a little slippage in disapproval of experimental use among 12th graders from 1991 through 1996 (from 96% to 92%)—a period during which heroin usage rates began to rise.
- Disapproval of regular *cocaine* use rose gradually among 19- to 22-year-olds, from 89% in 1981 to 99% in 1990, with little change thereafter (it was 98% in 2008). All three young adult age bands are now near 100%. Disapproval of experimental cocaine use increased during the 1980s, peaking first among 12th graders at 94% in 1991. It then peaked among 19- to 22-year-olds (at 94%) and 23- to 26-year-olds (at 92%) in 1995. Finally, it peaked among 27- to 30-year-olds at 90% in 1999. All age groups have had some modest falloff in disapproval since then. Again, the lag in inflection points between the successive age groups suggests some lasting cohort differences in these attitudes. For the last few years, all age groups' disapproval of experimental cocaine use has hovered around 85–90%.
- Disapproval of experimental use has moved very much in parallel for *amphetamines* and *sedatives (barbiturates)*. Disapproval of both drugs increased significantly during the 1980s, accompanied by declining use. Trying *amphetamines* once or twice was disapproved of by 73–74% of 19- to 26-year-olds in 1984, compared to 84% by 1990. The corresponding figures for disapproval of trying *sedatives* were 84–85% in 1984 compared to 89–91% by 1990. Disapproval of amphetamine and sedative use slipped some among 12th graders after 1992 and among 19- to 22-year-olds after 1994, with the 23- to 26-year-olds following suit after 1996, and the 27- to 30-year-old stratum in 2004. This pattern of staggered change again suggests cohort effects, reflecting lasting cohort differences in these attitudes. In recent years there has been a staggered increase in disapproval of both amphetamines and sedatives that has shown up in all age groups.
- The story for *alcohol* is quite an interesting one, in that changes in the minimum drinking age seem to have led to modest changes in norms for the affected cohorts. Between 1980

and 1992, an increasing proportion of 12th graders favored total abstinence; the percent who disapproved of drinking even just once or twice rose from 16% in 1980 to 33% in 1992. (This figure has fallen back slightly, and stands at 30% in 2008.) Among 19- to 22-year-olds, there was a modest increase from 15% to 22% disapproving of any use between 1985 and 1989, where it held for some years; this figure is 21% in 2008. For the two oldest age groups, there has been rather little change in these attitudes so far. These differing trends may reflect the fact that during the 1980s, the drinking age was raised in a number of states so that by 1987 it was 21 in all states; this change would have had the greatest effect on 12th graders, who may have incorporated the legal restrictions into their normative structure and, as they entered young adulthood, brought these new norms with them. But the changes may be exhibited only among respondents in the age groups directly impacted by the law.

Disapproval of *daily drinking* (one or two drinks) has not shown any such cohort effects, because all age groups have moved in parallel, at similar levels of disapproval. The three youngest age bands (which include 12th graders through 26-year-olds) showed an increase in disapproval of daily drinking up until about 1990 (there was little data yet available on the oldest age group), but disapproval has declined a fair amount in all of the age groups since then. A bit of a gap between 12th graders and young adults opened up between 2004 and 2008, when 12th graders increased their disapproval of daily drinking while young adults did not. This pattern of cross-time change closely parallels what was observed for the perceived risk of light daily drinking, discussed previously; and the later decline in both variables may well be due to widely publicized reports that some cardiovascular benefits result from having one or two drinks per day.

There was a considerable increase in disapproval of *heavy drinking on weekends* from the early 1980s for the two youngest age groups, and this continued through 1992 for 12th graders (who then showed some drop-off) and through 1996 among 19- to 22-year-olds (who also then showed some drop-off). As Figure 5-18d illustrates, the prevalence of occasional heavy drinking declined substantially among 12th graders and 19- to 22-year-olds between 1981 and the early 1990s, as norms became more restrictive. There was little or no change in the older age strata, either in their levels of disapproval or in their rates of occasional heavy drinking, until the early 2000s, when disapproval began to drop some in both strata.

At present, 12th graders are most likely to disapprove of trying alcoholic beverages (as has been the case for some years), but are the least disapproving of heavy daily drinking. Weekend binge drinking is generally least disapproved of by 19- to 22-year-olds, who also report the highest levels of engaging in such behavior.

- Some fluctuations in the disapproval of *cigarette smoking* have occurred over the intervals covered by MTF. Twelfth graders showed some increase in disapproval of pack-a-day-or-more smoking between 1982 (69%) and 1992 (74%). Their disapproval then fell through 1997 (to 67%) as their smoking increased; disapproval then increased for several years (to 82% in 2006) before leveling as smoking declined. The 19- to 22-year-olds

showed a similar increase in disapproval from 66% in 1982 to 81% in 2006, and are at 80% in 2008.

All four age strata showed some upward drift in their level of disapproval of smoking since about 1999, suggesting a secular change in these attitudes during this historical period.

COHORT DIFFERENCES AND THEIR IMPLICATIONS FOR PREVENTION AND THEORY

An important theoretical point to be made—based on the strong evidence reported here for cohort effects in perceived risk and disapproval of many of the drugs under study—is that one cause for cohort effects in actual use is lasting cohort differences in these critical attitudes and beliefs. *The attitudes and beliefs brought into adulthood from adolescence tend to persevere.*

A second point has to do with the causes of these attitudinal cohort effects. We noted earlier that the older respondents are more likely than the younger ones to see the use of *marijuana*, *LSD*, *heroin*, *amphetamines*, *ecstasy*, *crystal methamphetamine*, *cocaine*, *crack*, and *sedatives (barbiturates)* as dangerous. We have offered the framework for a theory of drug epidemics in which direct learning (from personal use) and vicarious learning (from observing use by others in both the immediate and mass media environments) play important roles in changing these key attitudes.⁵² To the extent that the data on perceived risk represent cohort effects (enduring differences between class cohorts), these findings would be consistent with this theoretical perspective. Clearly, use of these particular drugs was greater when the older cohorts were growing up, and public attention and concern regarding the consequences of these drugs were greatest in the 1970s and early to mid-1980s. In the early 1970s, LSD was alleged to cause brain and chromosomal damage, as well as bad trips, flashbacks, and behavior that could prove dangerous. Methamphetamine use was discouraged with the slogan “speed kills.” In addition, there was a serious epidemic of heroin use in the early 1970s. More recent cohorts (through the mid-1990s, at least) were not exposed to those experiences. While there may have been a secular trend toward greater perceived risk for drugs in general, in the case of LSD there may also have been an operating cohort effect (with younger cohorts seeing less danger) offsetting the secular trend among 12th graders; the net effect was a decrease in 12th graders’ perceived risk of LSD use after 1980.

This vicarious learning process has a very practical application for national strategy for preventing future epidemics. Because fewer in their immediate social circles and fewer public role models may be using these drugs and exhibiting the adverse consequences of use, future cohorts of youth may have less opportunity to learn about the adverse consequences of these drugs in the normal course of growing up. Unless those hazards are convincingly communicated to them in *other ways*—for example, through school prevention programs, by their parents, and

⁵²Johnston, L. D. (1991). Toward a theory of drug epidemics. In L. Donohew, H. E. Sypher, & W. J. Bukoski (Eds.), *Persuasive communication and drug abuse prevention* (pp. 93–131). Hillsdale, NJ: Lawrence Erlbaum. Available at <http://monitoringthefuture.org/pubs.html#chapters>.

through the mass media, including public service advertising—they will become more susceptible to a new epidemic of use of the same or similar drugs.

Volume I, the companion to this volume, reported an increase in use of several drugs in 8th, 10th, and 12th grades in 1994 through 1997. This increase suggests that this form of generational forgetting may well have been taking place during those years. For the cohorts that follow such a rise in use, there is once again an increased opportunity for vicarious learning from the adverse experiences of those around them, but by that time, members of affected cohorts have had to learn the hard way what consequences await those who become involved with the various drugs.

TABLE 6-1
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. How much do you think people risk harming themselves (physically or in other ways), if they . . .

	Age Group	Percentage saying "great risk" ^a														
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
Try marijuana once or twice	18	10.0	13.0	11.5	12.7	14.7	14.8	15.1	18.4	19.0	23.6	23.1	27.1	24.5	21.9	19.5
	19–22	8.3	7.8	9.7	9.7	12.8	11.2	13.0	12.9	16.8	16.9	17.8	19.1	19.7	19.4	18.8
	23–26	—	—	—	—	9.6	10.0	12.4	14.5	16.0	14.0	17.7	14.0	15.0	13.0	15.0
	27–30	—	—	—	—	—	—	—	—	14.6	16.0	17.0	15.7	15.1	14.0	14.8
Smoke marijuana occasionally	18	14.7	19.1	18.3	20.6	22.6	24.5	25.0	30.4	31.7	36.5	36.9	40.6	39.6	35.6	30.1
	19–22	13.9	14.2	16.9	16.7	21.7	20.6	22.4	23.0	28.7	29.1	30.1	30.2	29.5	30.3	31.3
	23–26	—	—	—	—	15.8	16.3	20.9	20.8	26.8	25.3	30.4	26.2	27.4	24.0	25.5
	27–30	—	—	—	—	—	—	—	—	24.2	25.7	28.7	27.4	27.5	26.8	28.1
Smoke marijuana regularly	18	50.4	57.6	60.4	62.8	66.9	70.4	71.3	73.5	77.0	77.5	77.8	78.6	76.5	72.5	65.0
	19–22	43.9	47.8	52.4	58.4	62.2	66.8	67.6	69.4	72.4	74.9	73.0	75.0	69.3	69.2	65.0
	23–26	—	—	—	—	52.9	57.5	59.4	65.3	68.3	72.1	71.0	70.9	67.3	64.1	63.2
	27–30	—	—	—	—	—	—	—	—	67.5	69.1	69.2	67.5	68.8	69.4	65.6
Try LSD once or twice	18	43.9	45.5	44.9	44.7	45.4	43.5	42.0	44.9	45.7	46.0	44.7	46.6	42.3	39.5	38.8
	19–22	44.8	44.4	45.0	44.7	46.0	44.3	47.6	49.4	49.2	49.5	49.3	48.0	45.6	42.4	42.3
	23–26	—	—	—	—	48.3	46.9	47.9	51.5	53.7	50.7	52.0	50.1	49.7	49.0	46.8
	27–30	—	—	—	—	—	—	—	—	53.3	55.6	54.6	52.5	53.0	51.5	53.5
Take LSD regularly	18	83.0	83.5	83.5	83.2	83.8	82.9	82.6	83.8	84.2	84.3	84.5	84.3	81.8	79.4	79.1
	19–22	83.4	85.3	86.2	86.0	84.5	86.4	87.1	85.6	85.4	85.5	85.8	86.6	87.0	81.3	81.0
	23–26	—	—	—	—	89.0	86.6	88.7	90.0	89.2	89.0	88.2	89.1	87.3	85.3	87.5
	27–30	—	—	—	—	—	—	—	—	89.1	91.2	92.0	87.1	88.5	89.0	89.2
Try PCP once or twice	18	—	—	—	—	—	—	—	55.6	58.8	56.6	55.2	51.7	54.8	50.8	51.5
	19–22	—	—	—	—	—	—	—	63.6	63.8	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	64.8	63.2	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	65.9	—	—	—	—	—	—
Try ecstasy (MDMA) once or twice	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	19–22	—	—	—	—	—	—	—	—	—	45.2	47.1	48.8	46.4	45.0	51.1
	23–26	—	—	—	—	—	—	—	—	—	49.5	47.2	47.4	45.5	41.9	50.6
	27–30	—	—	—	—	—	—	—	—	—	44.9	48.7	47.7	44.2	51.7	47.3
Take ecstasy (MDMA) occasionally	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try cocaine once or twice	18	31.3	32.1	32.8	33.0	35.7	34.0	33.5	47.9	51.2	54.9	59.4	59.4	56.8	57.6	57.2
	19–22	31.4	30.4	33.3	28.7	33.1	33.2	35.5	45.9	51.9	51.5	58.1	58.7	56.1	60.5	63.8
	23–26	—	—	—	—	31.3	31.1	35.9	48.0	47.1	51.3	51.5	50.5	53.5	54.1	56.0
	27–30	—	—	—	—	—	—	—	—	45.3	53.0	51.6	52.6	51.8	54.7	53.5
Take cocaine occasionally	18	—	—	—	—	—	—	54.2	66.8	69.2	71.8	73.9	75.5	75.1	73.3	73.7
	19–22	—	—	—	—	—	—	53.8	61.3	67.1	72.6	74.6	72.6	74.9	75.4	78.0
	23–26	—	—	—	—	—	—	50.9	62.6	63.2	69.9	69.9	70.3	69.9	72.8	70.3
	27–30	—	—	—	—	—	—	—	—	62.6	66.6	66.6	69.1	69.9	69.1	69.9
Take cocaine regularly	18	69.2	71.2	73.0	74.3	78.8	79.0	82.2	88.5	89.2	90.2	91.1	90.4	90.2	90.1	89.3
	19–22	65.2	69.3	71.5	75.2	75.1	82.9	82.0	88.0	90.3	89.1	93.9	93.5	92.9	91.7	92.2
	23–26	—	—	—	—	75.6	76.9	83.0	88.9	90.9	91.2	91.2	92.7	89.9	91.9	92.6
	27–30	—	—	—	—	—	—	—	—	88.9	92.0	91.4	90.9	92.0	91.6	92.1
Try crack once or twice	18	—	—	—	—	—	—	—	57.0	62.1	62.9	64.3	60.6	62.4	57.6	58.4
	19–22	—	—	—	—	—	—	—	59.4	67.3	68.5	69.4	66.9	65.4	63.5	70.1
	23–26	—	—	—	—	—	—	—	59.1	63.5	69.8	67.3	66.9	67.1	64.2	69.3
	27–30	—	—	—	—	—	—	—	—	66.5	64.9	68.7	66.8	64.3	68.8	65.6
Take crack occasionally	18	—	—	—	—	—	—	—	70.4	73.2	75.3	80.4	76.5	76.3	73.9	73.8
	19–22	—	—	—	—	—	—	—	75.0	77.3	81.8	82.3	82.7	81.9	83.6	84.3
	23–26	—	—	—	—	—	—	—	70.3	74.0	79.9	81.1	83.9	84.4	81.6	83.2
	27–30	—	—	—	—	—	—	—	—	76.4	76.7	82.6	81.8	79.1	83.6	78.6

(Table continued on next page.)

TABLE 6-1 (cont.)
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. How much do you think people risk harming themselves (physically or in other ways), if they . . .

	Age Group	Percentage saying "great risk" ^a														2007–2008 change
		1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Try marijuana once or twice	18	16.3	15.6	14.9	16.7	15.7	13.7	15.3	16.1	16.1	15.9	16.1	17.8	18.6	17.4	-1.2
	19–22	13.3	16.9	14.8	13.4	12.5	14.3	11.9	13.3	17.1	15.3	15.6	14.4	10.8	17.4	+6.6 ss
	23–26	15.8	18.5	15.1	16.7	16.4	13.1	13.0	15.1	15.3	13.6	13.0	13.9	13.0	12.5	-0.5
	27–30	16.1	16.2	16.1	16.4	16.1	14.4	17.3	16.2	18.0	13.8	14.5	14.5	16.6	11.4	-5.2 s
Smoke marijuana occasionally	18	25.6	25.9	24.7	24.4	23.9	23.4	23.5	23.2	26.6	25.4	25.8	25.9	27.1	25.8	-1.4
	19–22	25.5	25.6	22.0	22.0	19.8	25.8	18.0	21.0	24.1	23.2	24.3	22.1	22.3	23.6	+1.3
	23–26	27.7	27.3	26.4	26.8	26.4	24.9	20.5	24.5	22.2	22.7	21.6	22.3	20.2	18.5	-1.7
	27–30	28.3	28.1	26.0	25.8	25.3	25.8	25.0	30.2	27.9	25.1	24.8	21.8	25.6	21.6	-4.0
Smoke marijuana regularly	18	60.8	59.9	58.1	58.5	57.4	58.3	57.4	53.0	54.9	54.6	58.0	57.9	54.8	51.7	-3.1
	19–22	62.1	61.3	60.7	53.4	55.2	58.0	49.6	56.7	57.8	57.2	55.3	54.5	50.4	51.6	+1.3
	23–26	64.2	62.7	64.1	62.7	60.1	60.3	55.1	53.7	56.7	54.2	53.6	55.9	52.5	52.4	-0.2
	27–30	69.2	67.3	65.0	63.6	66.1	64.0	61.7	63.5	64.7	59.3	57.0	54.9	51.5	51.2	-0.3
Try LSD once or twice	18	36.4	36.2	34.7	37.4	34.9	34.3	33.2	36.7	36.2	36.2	36.5	36.1	37.0	33.9	-3.1
	19–22	40.3	44.4	40.1	38.7	38.1	37.9	37.5	35.3	39.7	39.2	38.7	43.5	40.9	46.5	+5.6
	23–26	45.8	46.1	46.6	45.7	49.3	44.9	48.5	45.7	43.8	40.7	39.9	38.1	42.8	43.8	+1.0
	27–30	52.5	50.1	52.0	52.0	49.9	46.4	46.7	44.9	47.5	47.2	47.9	44.9	44.6	42.4	-2.1
Take LSD regularly	18	78.1	77.8	76.6	76.5	76.1	75.9	74.1	73.9	72.3	70.2	69.9	69.3	67.3	63.6	-3.7 s
	19–22	80.5	82.4	83.6	78.6	82.2	81.6	79.2	81.1	78.6	78.4	77.8	78.9	77.5	73.9	-3.5
	23–26	86.3	84.7	85.6	82.1	85.4	84.1	86.0	85.3	84.3	83.5	80.8	82.0	80.3	80.2	0.0
	27–30	88.4	87.0	87.2	90.5	87.8	85.3	86.9	85.3	87.5	83.9	87.9	82.2	85.7	82.9	-2.8
Try PCP once or twice	18	49.1	51.0	48.8	46.8	44.8	45.0	46.2	48.3	45.2	47.1	46.6	47.0	48.0	47.4	-0.7
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try ecstasy (MDMA) once or twice	18	—	—	33.8	34.5	35.0	37.9	45.7	52.2	56.3	57.7	60.1	59.3	58.1	57.0	-1.1
	19–22	48.3	46.7	45.5	42.7	37.6	37.9	40.5	46.8	50.1	52.3	53.8	51.0	50.3	51.4	+1.1
	23–26	49.3	50.4	50.5	47.7	50.0	46.7	45.7	45.6	45.9	44.9	51.2	46.4	51.4	46.3	-5.1
	27–30	50.0	50.6	48.8	50.4	50.9	48.9	53.6	52.0	58.8	49.1	50.2	46.5	51.9	43.5	-8.4 s
Take ecstasy (MDMA) occasionally	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	19–22	—	—	—	—	—	—	72.5	77.8	81.7	78.3	80.0	82.5	79.3	81.9	+2.6
	23–26	—	—	—	—	—	—	72.5	71.9	73.6	77.4	77.2	77.0	78.7	78.6	0.0
	27–30	—	—	—	—	—	—	75.2	76.5	79.9	76.9	74.7	70.4	72.0	71.3	-0.7
Try cocaine once or twice	18	53.7	54.2	53.6	54.6	52.1	51.1	50.7	51.2	51.0	50.7	50.5	52.5	51.3	50.3	-1.0
	19–22	57.7	61.9	55.5	55.4	52.8	56.7	48.9	55.5	55.0	55.5	55.6	54.0	55.8	56.7	+0.9
	23–26	58.7	57.2	63.1	60.2	62.6	63.1	62.4	61.0	55.4	52.1	53.0	52.5	56.9	55.0	-1.9
	27–30	56.4	53.6	54.6	60.5	61.7	59.9	60.9	58.8	56.4	61.4	56.5	58.1	54.8	56.1	+1.3
Take cocaine occasionally	18	70.8	72.1	72.4	70.1	70.1	69.5	69.9	68.3	69.1	67.2	66.7	69.8	68.8	67.1	-1.7
	19–22	73.4	76.6	76.1	71.2	68.0	72.4	70.0	69.9	70.3	70.2	72.1	71.0	71.5	72.4	+0.9
	23–26	76.0	71.3	76.5	74.2	77.8	76.2	74.2	75.4	68.3	74.1	70.4	68.5	70.9	67.2	-3.7
	27–30	70.0	67.8	73.8	73.2	75.4	76.5	78.1	74.3	72.6	75.3	76.2	74.6	72.1	73.9	+1.8
Take cocaine regularly	18	87.9	88.3	87.1	86.3	85.8	86.2	84.1	84.5	83.0	82.2	82.8	84.6	83.3	80.7	-2.6
	19–22	91.5	92.2	91.6	88.7	88.5	90.7	85.1	88.3	87.4	87.1	89.2	86.2	86.7	87.0	+0.3
	23–26	93.3	90.6	93.2	92.9	92.7	92.9	91.1	91.5	88.5	91.5	88.0	90.9	88.0	86.5	-1.5
	27–30	91.3	91.6	92.7	93.0	92.4	92.3	94.5	91.2	92.9	91.3	94.0	90.0	89.9	91.1	+1.3
Try crack once or twice	18	54.6	56.0	54.0	52.2	48.2	48.4	49.4	50.8	47.3	47.8	48.4	47.8	47.3	47.5	+0.2
	19–22	61.9	65.2	62.0	59.3	56.1	52.9	54.1	54.1	55.1	56.8	56.6	55.3	51.9	54.9	+2.9
	23–26	64.8	68.6	64.7	67.3	64.6	63.2	59.8	60.9	58.5	56.4	60.6	54.7	58.4	50.5	-7.8 s
	27–30	66.4	66.7	68.5	66.5	65.0	62.9	69.3	67.4	66.0	62.6	61.9	56.8	64.1	56.2	-7.9 s
Take crack occasionally	18	72.8	71.4	70.3	68.7	67.3	65.8	65.4	65.6	64.0	64.5	63.8	64.8	63.6	65.2	+1.6
	19–22	78.8	83.5	79.1	79.1	75.5	74.9	72.3	75.3	75.3	76.0	75.0	72.8	77.7	75.7	-2.0
	23–26	81.4	85.9	80.8	84.2	81.6	84.0	80.1	82.2	77.1	76.4	78.6	76.8	79.8	75.2	-4.6
	27–30	81.1	81.3	85.3	81.7	79.8	81.6	84.4	81.5	81.9	82.1	79.5	82.8	79.1	77.3	-1.8

(List of drugs continued.)

TABLE 6-1 (cont.)
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. How much do you think people risk harming themselves (physically or in other ways), if they . . .

	Age Group	Percentage saying "great risk" ^a															
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	cont.
Take crack regularly	18	—	—	—	—	—	—	—	84.6	84.8	85.6	91.6	90.1	89.3	87.5	89.6	
	19–22	—	—	—	—	—	—	—	89.6	91.1	94.1	94.9	95.6	93.4	96.2	96.0	
	23–26	—	—	—	—	—	—	—	88.0	89.2	91.5	94.2	95.4	94.1	93.4	94.9	
	27–30	—	—	—	—	—	—	—	—	89.6	89.5	95.3	94.4	93.3	93.5	93.0	
Try cocaine powder once or twice	18	—	—	—	—	—	—	—	45.3	51.7	53.8	53.9	53.6	57.1	53.2	55.4	
	19–22	—	—	—	—	—	—	—	44.0	48.6	51.1	54.5	52.7	56.2	49.7	62.0	
	23–26	—	—	—	—	—	—	—	41.0	43.6	48.4	48.9	47.4	45.9	45.6	52.5	
	27–30	—	—	—	—	—	—	—	—	42.0	45.1	46.2	43.3	42.3	49.9	47.1	
Take cocaine powder occasionally	18	—	—	—	—	—	—	—	56.8	61.9	65.8	71.1	69.8	70.8	68.6	70.6	
	19–22	—	—	—	—	—	—	—	58.0	59.0	63.2	70.0	69.9	72.6	70.6	75.4	
	23–26	—	—	—	—	—	—	—	50.0	53.2	62.2	63.3	67.0	65.8	64.0	68.8	
	27–30	—	—	—	—	—	—	—	—	53.6	52.7	60.9	59.2	61.2	64.3	61.0	
Take cocaine powder regularly	18	—	—	—	—	—	—	—	81.4	82.9	83.9	90.2	88.9	88.4	87.0	88.6	
	19–22	—	—	—	—	—	—	—	86.6	87.6	91.3	92.5	93.8	92.1	94.0	94.9	
	23–26	—	—	—	—	—	—	—	82.9	84.1	88.5	92.4	93.8	91.3	92.4	92.8	
	27–30	—	—	—	—	—	—	—	—	85.1	86.7	92.7	91.1	91.5	92.5	90.7	
Try heroin once or twice	18	52.1	52.9	51.1	50.8	49.8	47.3	45.8	53.6	54.0	53.8	55.4	55.2	50.9	50.7	52.8	
	19–22	57.8	56.8	54.4	52.5	58.7	51.0	55.5	57.9	58.9	59.6	58.3	59.9	59.8	58.9	60.8	
	23–26	—	—	—	—	58.2	59.2	60.8	66.6	65.4	62.3	64.1	62.4	63.7	65.0	63.3	
	27–30	—	—	—	—	—	—	—	—	66.0	69.7	67.5	66.1	66.5	69.3	69.6	
Take heroin occasionally	18	70.9	72.2	69.8	71.8	70.7	69.8	68.2	74.6	73.8	75.5	76.6	74.9	74.2	72.0	72.1	
	19–22	77.5	77.8	73.6	74.5	74.9	73.6	77.2	77.6	77.5	79.8	80.8	80.2	81.6	78.8	79.0	
	23–26	—	—	—	—	81.2	80.7	78.9	84.5	82.4	80.8	83.4	84.4	81.5	82.1	80.8	
	27–30	—	—	—	—	—	—	—	—	86.0	86.8	85.3	84.3	84.9	86.2	86.8	
Take heroin regularly	18	86.2	87.5	86.0	86.1	87.2	86.0	87.1	88.7	88.8	89.5	90.2	89.6	89.2	88.3	88.0	
	19–22	87.2	89.9	87.5	88.6	86.8	90.2	90.7	90.2	89.6	90.8	91.2	91.5	92.2	89.2	91.2	
	23–26	—	—	—	—	92.0	90.1	90.6	92.8	91.5	91.3	91.0	92.6	91.3	91.6	93.0	
	27–30	—	—	—	—	—	—	—	—	92.7	93.5	93.0	90.7	91.3	92.6	93.8	
Try amphetamines once or twice	18	29.7	26.4	25.3	24.7	25.4	25.2	25.1	29.1	29.6	32.8	32.2	36.3	32.6	31.3	31.4	
	19–22	24.6	24.6	27.8	24.8	26.9	23.9	27.1	27.4	31.7	28.9	35.6	32.8	34.5	33.3	36.3	
	23–26	—	—	—	—	29.6	29.4	29.4	34.1	33.2	32.5	35.3	31.0	32.7	32.6	32.9	
	27–30	—	—	—	—	—	—	—	—	35.2	37.5	36.9	36.5	36.2	34.0	37.5	
Take amphetamines regularly	18	69.1	66.1	64.7	64.8	67.1	67.2	67.3	69.4	69.8	71.2	71.2	74.1	72.4	69.9	67.0	
	19–22	71.9	69.9	68.3	69.9	68.4	68.5	72.3	72.0	73.9	71.3	74.0	77.1	73.5	73.5	71.6	
	23–26	—	—	—	—	75.8	77.2	75.6	78.2	77.4	76.7	77.8	79.4	76.4	76.2	73.6	
	27–30	—	—	—	—	—	—	—	—	80.6	82.9	83.3	79.4	80.3	79.8	78.4	
Try crystal meth. (ice)	18	—	—	—	—	—	—	—	—	—	—	—	61.6	61.9	57.5	58.3	
	19–22	—	—	—	—	—	—	—	—	—	—	—	57.8	58.6	57.7	57.5	
	23–26	—	—	—	—	—	—	—	—	—	—	—	56.5	56.0	55.6	52.0	
	27–30	—	—	—	—	—	—	—	—	—	—	—	59.6	57.2	52.7	60.3	
Try sedatives/ barbiturates^b once or twice	18	30.9	28.4	27.5	27.0	27.4	26.1	25.4	30.9	29.7	32.2	32.4	35.1	32.2	29.2	29.9	
	19–22	27.6	26.4	30.5	25.4	29.9	25.0	30.7	29.6	32.7	30.5	36.4	33.5	33.5	33.4	35.0	
	23–26	—	—	—	—	32.2	29.9	30.2	35.5	35.8	32.9	37.9	31.8	33.5	32.8	34.0	
	27–30	—	—	—	—	—	—	—	—	37.2	38.7	39.0	37.0	38.2	36.5	40.5	
Take sedatives/ barbiturates^b regularly	18	72.2	69.9	67.6	67.7	68.5	68.3	67.2	69.4	69.6	70.5	70.2	70.5	70.2	66.1	63.3	
	19–22	74.0	73.3	72.7	71.3	71.6	71.7	74.5	73.0	74.0	71.7	75.5	75.5	73.6	71.1	69.4	
	23–26	—	—	—	—	77.4	77.0	74.9	79.9	79.8	76.6	80.5	77.7	76.3	75.0	74.3	
	27–30	—	—	—	—	—	—	—	—	81.5	83.7	84.0	79.6	78.6	80.2	78.3	

(Table continued on next page.)

TABLE 6-1 (cont.)
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. How much do you think people risk harming themselves (physically or in other ways), if they . . .

	Age Group	Percentage saying "great risk" ^a														2007–2008 change
		1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Take crack regularly	18	88.6	88.0	86.2	85.3	85.4	85.3	85.8	84.1	83.2	83.5	83.3	82.8	82.6	83.4	+0.8
	19–22	94.2	94.7	93.3	92.8	92.3	91.1	89.6	91.1	93.8	93.3	92.5	90.3	90.3	93.6	+3.4
	23–26	95.5	96.1	91.4	95.6	94.4	95.6	93.4	94.7	92.2	92.5	93.1	93.3	93.1	91.8	-1.3
	27–30	94.0	94.3	96.0	94.3	95.2	93.5	96.8	94.2	94.4	94.0	95.2	94.1	93.6	93.1	-0.5
Try cocaine powder once or twice	18	52.0	53.2	51.4	48.5	46.1	47.0	49.0	49.5	46.2	45.4	46.2	45.8	45.1	45.1	0.0
	19–22	55.8	57.1	53.8	53.0	47.9	48.0	47.1	47.9	49.4	48.7	50.2	48.7	46.8	48.3	+1.5
	23–26	48.9	57.2	53.6	54.1	53.8	53.2	53.9	52.5	50.8	46.0	53.3	45.8	48.1	44.1	-3.9
	27–30	48.2	48.9	49.1	49.8	49.7	52.2	53.3	54.4	56.6	52.5	52.9	49.0	53.6	47.2	-6.4
Take cocaine powder occasionally	18	69.1	68.8	67.7	65.4	64.2	64.7	63.2	64.4	61.4	61.6	60.8	61.9	59.9	61.6	+1.7
	19–22	73.0	77.4	70.7	73.0	69.3	69.3	64.4	68.9	69.3	68.6	68.1	66.4	67.1	68.5	+1.3
	23–26	68.8	76.1	72.8	77.0	70.8	76.0	70.5	73.7	67.9	64.6	69.9	66.7	69.9	64.5	-5.4
	27–30	65.9	68.2	69.7	68.5	70.1	71.3	73.5	71.9	71.7	71.5	71.7	73.1	69.3	64.9	-4.3
Take cocaine powder regularly	18	87.8	86.8	86.0	84.1	84.6	85.5	84.4	84.2	82.3	81.7	82.7	82.1	81.5	82.5	+1.1
	19–22	93.5	93.8	92.8	91.5	92.4	90.7	89.8	91.0	92.0	91.6	90.7	89.1	89.5	92.3	+2.8
	23–26	92.1	94.8	90.8	93.7	93.6	94.2	92.2	93.4	89.1	89.4	91.2	92.9	92.3	90.5	-1.9
	27–30	92.7	91.7	93.0	92.3	93.1	91.5	94.0	93.3	94.1	93.1	93.9	92.4	92.5	90.1	-2.3
Try heroin once or twice	18	50.9	52.5	56.7	57.8	56.0	54.2	55.6	56.0	58.0	56.6	55.2	59.1	58.4	55.5	-2.9
	19–22	58.9	61.0	63.9	60.7	63.5	63.2	64.0	63.1	64.6	67.3	66.5	65.0	69.6	67.7	-1.9
	23–26	64.1	63.5	67.3	67.3	68.0	70.7	71.9	69.8	70.6	67.5	69.2	67.0	68.3	70.1	+1.7
	27–30	66.4	66.4	67.9	69.7	70.1	67.4	68.2	70.9	72.3	68.4	74.4	70.8	70.2	70.2	0.0
Take heroin occasionally	18	71.0	74.8	76.3	76.9	77.3	74.6	75.9	76.6	78.5	75.7	76.0	79.1	76.2	75.3	-1.0
	19–22	77.9	82.1	84.7	80.4	82.5	82.0	83.6	82.2	84.9	85.1	83.8	84.3	85.4	84.5	-0.9
	23–26	85.3	82.4	86.5	83.9	88.5	86.6	88.4	90.0	88.3	86.7	87.5	85.2	86.5	88.0	+1.5
	27–30	83.1	83.8	85.8	86.6	87.1	86.5	86.4	87.9	87.4	88.6	91.2	88.3	88.5	87.7	-0.8
Take heroin regularly	18	87.2	89.5	88.9	89.1	89.9	89.2	88.3	88.5	89.3	86.8	87.5	89.7	87.8	86.4	-1.4
	19–22	89.9	94.0	93.7	92.4	92.8	94.0	91.3	92.6	93.9	94.3	94.9	94.2	93.6	92.3	-1.3
	23–26	93.5	92.7	94.4	93.4	93.7	94.8	95.9	96.3	96.5	96.0	94.8	95.8	93.1	95.7	+2.6
	27–30	92.4	92.1	93.8	95.0	93.7	94.2	94.5	95.9	94.9	95.0	97.3	95.3	94.8	95.4	+0.6
Try amphetamines once or twice	18	28.8	30.8	31.0	35.3	32.2	32.6	34.7	34.4	36.8	35.7	37.7	39.5	41.3	39.2	-2.2
	19–22	32.9	36.8	30.1	31.7	33.7	35.0	34.2	38.1	40.2	36.8	38.3	40.0	38.4	42.1	+3.7
	23–26	34.3	34.9	37.8	40.9	41.8	39.9	41.6	38.0	38.3	33.2	39.1	37.0	38.0	40.8	+2.8
	27–30	36.0	36.2	34.5	37.6	36.3	39.4	38.5	39.0	40.5	39.2	38.2	39.7	37.4	36.5	-0.9
Take amphetamines regularly	18	65.9	66.8	66.0	67.7	66.4	66.3	67.1	64.8	65.6	63.9	67.1	68.1	68.1	65.4	-2.6
	19–22	72.2	75.8	72.3	71.9	72.4	73.4	71.1	72.7	75.0	72.4	74.1	72.1	73.8	74.2	+0.4
	23–26	80.5	78.5	79.1	77.5	78.7	79.0	77.7	77.9	80.1	75.1	80.1	78.3	77.0	76.5	-0.5
	27–30	77.7	75.6	77.4	81.1	82.6	80.8	79.9	79.8	81.5	77.6	78.9	78.9	77.6	78.9	+1.3
Try crystal meth. (ice)	18	54.4	55.3	54.4	52.7	51.2	51.3	52.7	53.8	51.2	52.4	54.6	59.1	60.2	62.2	+2.0
	19–22	58.9	61.1	56.4	55.8	50.6	49.2	52.5	56.5	60.0	60.3	63.1	63.5	65.0	70.0	+5.0
	23–26	57.8	64.1	60.7	58.2	61.3	60.1	59.2	57.7	58.6	55.9	63.9	63.9	66.6	65.6	-1.0
	27–30	58.5	59.1	59.8	59.9	61.0	59.7	66.4	62.5	66.6	62.8	62.6	64.9	67.9	62.0	-5.9
Try sedatives/ barbiturates^b once or twice	18	26.3	29.1	26.9	29.0	26.1	25.0	25.7	26.2	27.9	24.9	24.7	28.0	27.9	25.9	-2.0
	19–22	30.5	34.1	31.4	27.7	28.5	30.3	30.0	30.7	32.7	26.7	26.9	28.9	28.1	31.9	+3.7
	23–26	34.8	35.8	37.3	40.3	39.4	37.0	38.5	34.7	36.5	22.2	29.8	26.3	25.9	28.4	+2.5
	27–30	36.6	37.2	35.7	36.7	35.2	36.3	40.9	37.3	38.6	31.4	31.7	28.8	28.0	27.8	-0.1
Take sedatives/ barbiturates^b regularly	18	61.6	60.4	56.8	56.3	54.1	52.3	50.3	49.3	49.6	54.0	54.1	56.8	55.1	50.2	-4.9 ss
	19–22	66.4	70.7	69.5	65.1	64.7	64.6	61.8	64.5	63.8	60.2	64.4	61.3	63.2	64.0	+0.8
	23–26	77.6	77.1	75.2	73.9	75.1	73.8	73.1	73.1	72.8	63.9	67.0	67.6	64.8	66.8	+2.1
	27–30	77.7	74.1	77.1	79.9	80.7	75.5	78.2	75.4	79.0	70.1	75.2	68.0	70.0	70.4	+0.4

↓
(List of drugs continued.)

TABLE 6-1 (cont.)
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. How much do you think people risk harming themselves (physically or in other ways), if they . . .

Percentage saying "great risk"^a

	Age Group	Percentage saying "great risk" ^a														
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
Try one or two drinks of an alcoholic beverage (beer, wine, liquor)	18	3.8	4.6	3.5	4.2	4.6	5.0	4.6	6.2	6.0	6.0	8.3	9.1	8.6	8.2	7.6
	19–22	3.0	3.4	3.1	2.3	4.7	3.1	5.4	3.5	3.9	5.9	6.1	5.4	5.8	6.6	6.5
	23–26	—	—	—	—	5.5	3.0	6.5	6.6	4.2	5.1	5.7	4.4	5.6	3.2	4.5
	27–30	—	—	—	—	—	—	—	—	5.0	6.3	4.4	6.6	5.6	4.7	4.1
Take one or two drinks nearly every day	18	20.3	21.6	21.6	21.6	23.0	24.4	25.1	26.2	27.3	28.5	31.3	32.7	30.6	28.2	27.0
	19–22	22.7	22.9	23.2	23.2	25.0	26.3	27.3	26.1	26.5	28.1	30.1	29.1	30.2	28.0	27.5
	23–26	—	—	—	—	27.8	27.4	26.9	30.2	29.1	27.8	31.1	30.4	31.6	25.9	26.2
	27–30	—	—	—	—	—	—	—	—	27.4	31.7	32.2	31.7	30.9	28.0	27.4
Take four or five drinks nearly every day	18	65.7	64.5	65.5	66.8	68.4	69.8	66.5	69.7	68.5	69.8	70.9	69.5	70.5	67.8	66.2
	19–22	71.2	72.7	73.3	72.7	76.2	74.1	74.0	76.4	72.8	75.7	76.1	75.5	71.8	72.1	70.3
	23–26	—	—	—	—	76.7	77.9	80.1	77.2	81.8	76.9	79.7	80.2	78.0	76.7	77.5
	27–30	—	—	—	—	—	—	—	—	79.3	81.7	84.7	79.1	79.9	79.1	76.6
Have five or more drinks once or twice each weekend	18	35.9	36.3	36.0	38.6	41.7	43.0	39.1	41.9	42.6	44.0	47.1	48.6	49.0	48.3	46.5
	19–22	34.2	30.1	33.5	36.6	37.9	40.2	34.6	36.7	36.9	42.4	40.6	40.8	41.8	42.4	41.9
	23–26	—	—	—	—	38.4	39.7	39.1	39.8	35.8	37.7	40.2	39.3	37.6	36.2	40.2
	27–30	—	—	—	—	—	—	—	—	41.0	42.3	44.1	42.2	45.1	42.9	43.2
Smoke one or more packs of cigarettes per day	18	63.7	63.3	60.5	61.2	63.8	66.5	66.0	68.6	68.0	67.2	68.2	69.4	69.2	69.5	67.6
	19–22	66.5	61.7	64.0	62.1	69.1	71.4	70.4	70.6	71.0	73.4	72.5	77.9	72.6	76.0	71.2
	23–26	—	—	—	—	71.1	70.1	75.7	73.6	75.5	71.4	78.5	75.3	76.3	78.4	76.4
	27–30	—	—	—	—	—	—	—	—	72.8	75.2	77.8	75.4	77.6	75.0	75.3
Use smokeless tobacco regularly	18	—	—	—	—	—	—	25.8	30.0	33.2	32.9	34.2	37.4	35.5	38.9	36.6
	19–22	—	—	—	—	—	—	29.7	34.1	31.1	37.1	33.5	38.9	40.1	43.3	37.6
	23–26	—	—	—	—	—	—	37.0	38.5	35.8	37.9	40.1	38.9	41.6	44.6	42.9
	27–30	—	—	—	—	—	—	—	—	42.8	42.8	43.8	44.3	44.1	47.3	46.3
<i>Approximate Weighted N =</i>	18	3,234	3,604	3,557	3,305	3,262	3,250	3,020	3,315	3,276	2,796	2,553	2,549	2,684	2,759	2,591
	19–22	590	585	583	585	579	547	581	570	551	565	552	533	527	480	490
	23–26	—	—	—	—	540	512	545	531	527	498	511	505	518	503	465
	27–30	—	—	—	—	—	—	—	—	513	587	490	486	482	473	443

cont. →

(Table continued on next page.)

TABLE 6-1 (cont.)
Trends in Harmfulness as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. How much do you think people risk harming

themselves (physically or in other ways), if they . . .

	Age Group	Percentage saying "great risk" ^a														2007–2008 change
		1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Try one or two drinks of an alcoholic beverage (beer, wine, liquor)	18	5.9	7.3	6.7	8.0	8.3	6.4	8.7	7.6	8.4	8.6	8.5	9.3	10.5	10.0	-0.5
	19–22	4.5	3.3	3.2	4.2	5.7	5.4	4.8	6.6	7.5	5.1	3.8	7.7	5.1	7.9	+2.8
	23–26	4.3	4.8	4.4	4.4	6.6	3.5	5.5	5.1	5.7	4.7	5.3	5.1	4.8	6.5	+1.7
	27–30	6.7	4.7	4.0	6.2	5.9	4.7	5.5	3.1	6.9	4.6	7.3	4.2	6.2	3.4	-2.8
Take one or two drinks nearly every day	18	24.8	25.1	24.8	24.3	21.8	21.7	23.4	21.0	20.1	23.0	23.7	25.3	25.1	24.2	-0.9
	19–22	24.0	23.0	24.2	22.1	23.9	22.1	19.6	22.7	19.8	21.3	22.1	22.0	19.0	24.4	+5.4
	23–26	26.1	22.0	20.2	21.0	26.0	21.7	23.5	23.4	19.1	22.9	19.9	22.5	21.2	21.0	-0.2
	27–30	27.2	24.0	24.8	20.8	25.3	22.0	22.7	21.7	21.4	21.8	23.7	20.2	21.5	21.5	0.0
Take four or five drinks nearly every day	18	62.8	65.6	63.0	62.1	61.1	59.9	60.7	58.8	57.8	59.2	61.8	63.4	61.8	60.8	-1.0
	19–22	72.5	68.5	71.4	70.4	69.9	69.9	64.5	71.1	66.4	65.3	63.0	66.6	68.8	68.5	-0.3
	23–26	75.2	72.0	75.1	69.3	72.8	71.7	75.8	74.9	71.1	74.2	71.2	72.4	70.2	70.0	-0.2
	27–30	82.2	76.1	79.3	75.7	75.1	77.4	72.8	76.2	70.6	72.1	77.5	73.0	76.5	77.1	+0.6
Have five or more drinks once or twice each weekend	18	45.2	49.5	43.0	42.8	43.1	42.7	43.6	42.2	43.5	43.6	45.0	47.6	45.8	46.3	+0.6
	19–22	39.9	40.7	36.6	42.0	37.2	38.9	37.2	37.8	40.4	38.1	37.5	37.2	43.4	41.7	-1.7
	23–26	37.9	39.1	37.4	41.1	40.2	34.9	39.0	36.8	36.3	37.9	36.8	38.4	39.7	37.0	-2.7
	27–30	44.6	41.5	40.0	40.2	41.9	37.9	41.6	40.6	42.5	40.5	44.0	39.1	40.4	40.4	0.0
Smoke one or more packs of cigarettes per day	18	65.6	68.2	68.7	70.8	70.8	73.1	73.3	74.2	72.1	74.0	76.5	77.6	77.3	74.0	-3.3 s
	19–22	71.6	73.8	76.3	77.2	75.7	77.1	76.6	80.6	77.8	81.1	80.5	80.8	79.3	79.5	+0.2
	23–26	76.0	76.0	77.6	76.5	80.9	79.7	83.9	85.1	83.6	84.1	81.6	86.4	80.7	83.6	+3.0
	27–30	75.6	73.0	80.3	80.9	80.7	78.4	82.7	80.6	82.0	81.7	84.1	83.8	84.3	86.6	+2.2
Use smokeless tobacco regularly	18	33.2	37.4	38.6	40.9	41.1	42.2	45.4	42.6	43.3	45.0	43.6	45.9	44.0	42.9	-1.1
	19–22	42.3	40.9	46.5	47.4	47.0	52.0	48.4	53.6	50.8	49.9	47.6	46.4	48.9	48.7	-0.2
	23–26	46.6	47.2	46.2	48.4	53.1	49.8	59.8	61.4	58.9	57.8	55.8	59.1	55.3	51.0	-4.3
	27–30	44.2	43.6	50.2	52.6	53.6	49.9	53.2	56.7	58.2	55.7	58.9	57.5	61.4	61.7	+0.3
<i>Approximate Weighted N =</i>	18	2,603	2,449	2,579	2,564	2,306	2,130	2,173	2,198	2,466	2,491	2,512	2,407	2,450	2,389	
	19–22	500	469	464	431	447	424	430	395	402	447	412	411	375	377	
	23–26	446	438	420	413	418	400	392	382	401	426	408	361	351	375	
	27–30	450	422	434	416	400	377	384	369	380	388	374	358	344	350	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

"—" indicates data not available.

^aAnswer alternatives were: (1) No risk, (2) Slight risk, (3) Moderate risk, (4) Great risk, and (5) Can't say, drug unfamiliar.

^bIn 2004 the question text was changed from "barbiturates" to "sedatives/barbiturates" and the list of examples was changed from "downers, goofballs, reds, yellows, etc." to just "downers." These changes likely explain the discontinuity in the 2003 and 2004 results.

TABLE 6-2
Trends in Proportions Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

		Percentage disapproving ^a															
		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
<i>Q. Do you disapprove of people (who are 18 or older) doing each of the following?</i>																	
Trying marijuana once or twice	18	39.0	40.0	45.5	46.3	49.3	51.4	54.6	56.6	60.8	64.6	67.8	68.7	69.9	63.3	57.6	
	19–22	38.2	36.1	37.0	42.0	44.1	46.6	51.6	52.8	55.8	62.4	59.6	60.4	57.8	60.6	63.5	
	23–26	—	—	—	—	41.2	38.6	42.6	49.1	48.7	52.5	57.5	58.8	55.0	54.6	52.3	
	27–30	—	—	—	—	—	—	—	—	49.0	50.9	53.8	54.6	51.9	56.8	55.7	
Smoking marijuana occasionally	18	49.7	52.6	59.1	60.7	63.5	65.8	69.0	71.6	74.0	77.2	80.5	79.4	79.7	75.5	68.9	
	19–22	49.6	49.1	51.3	56.0	60.4	62.6	66.7	67.2	69.5	77.3	76.3	77.0	74.8	75.8	76.9	
	23–26	—	—	—	—	54.8	52.8	57.0	64.9	63.4	69.4	73.7	73.3	74.0	71.9	70.9	
	27–30	—	—	—	—	—	—	—	—	65.3	67.1	68.9	73.0	67.2	72.2	69.4	
Smoking marijuana regularly	18	74.6	77.4	80.6	82.5	84.7	85.5	86.6	89.2	89.3	89.8	91.0	89.3	90.1	87.6	82.3	
	19–22	74.3	77.2	80.0	81.8	84.9	86.7	89.2	88.7	89.1	91.2	93.1	91.3	89.5	90.2	90.1	
	23–26	—	—	—	—	80.6	81.3	83.3	87.4	86.9	90.4	91.0	89.6	90.2	92.1	90.3	
	27–30	—	—	—	—	—	—	—	—	87.6	87.5	89.7	89.6	87.2	89.4	88.7	
Trying LSD once or twice	18	87.3	86.4	88.8	89.1	88.9	89.5	89.2	91.6	89.8	89.7	89.8	90.1	88.1	85.9	82.5	
	19–22	87.4	84.8	85.9	88.4	88.1	89.1	90.4	90.0	90.9	89.3	90.5	88.4	84.6	88.5	86.8	
	23–26	—	—	—	—	87.3	87.1	88.0	89.9	91.4	91.0	90.7	89.1	88.8	86.9	87.3	
	27–30	—	—	—	—	—	—	—	—	91.0	87.2	89.7	87.9	85.6	88.8	88.2	
Taking LSD regularly	18	96.7	96.8	96.7	97.0	96.8	97.0	96.6	97.8	96.4	96.4	96.3	96.4	95.5	95.8	94.3	
	19–22	98.2	97.4	97.7	97.6	97.6	98.8	98.5	98.0	98.1	97.5	99.1	97.5	97.0	97.8	97.7	
	23–26	—	—	—	—	99.2	98.0	98.5	99.0	98.0	98.4	98.3	98.4	98.3	98.1	97.7	
	27–30	—	—	—	—	—	—	—	—	98.8	97.1	98.9	98.9	97.5	98.5	98.7	
Trying ecstasy (MDMA) once or twice	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Taking ecstasy (MDMA) occasionally	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Trying cocaine once or twice	18	76.3	74.6	76.6	77.0	79.7	79.3	80.2	87.3	89.1	90.5	91.5	93.6	93.0	92.7	91.6	
	19–22	73.0	69.3	69.9	74.1	72.5	77.6	78.9	82.3	85.3	88.8	90.1	91.2	90.6	92.7	93.9	
	23–26	—	—	—	—	70.2	70.5	72.1	80.0	82.9	85.5	88.3	88.0	87.3	89.2	89.2	
	27–30	—	—	—	—	—	—	—	—	82.1	81.0	85.5	86.9	83.9	85.7	86.6	
Taking cocaine regularly	18	91.1	90.7	91.5	93.2	94.5	93.8	94.3	96.7	96.2	96.4	96.7	97.3	96.9	97.5	96.6	
	19–22	91.6	89.3	91.9	94.6	95.0	96.3	97.0	97.2	97.9	97.4	98.9	97.9	98.4	97.8	98.8	
	23–26	—	—	—	—	95.7	95.3	97.3	98.1	97.6	98.3	98.4	98.5	98.7	98.4	98.8	
	27–30	—	—	—	—	—	—	—	—	98.1	97.0	99.3	99.0	97.2	98.7	99.0	
Trying heroin once or twice	18	93.5	93.5	94.6	94.3	94.0	94.0	93.3	96.2	95.0	95.4	95.1	96.0	94.9	94.4	93.2	
	19–22	96.3	95.4	95.6	95.2	95.1	96.2	96.8	96.3	97.1	96.4	98.3	95.9	95.9	96.3	96.6	
	23–26	—	—	—	—	96.7	94.9	96.4	97.1	97.4	96.7	96.8	96.9	96.3	95.4	96.5	
	27–30	—	—	—	—	—	—	—	—	97.9	95.8	97.5	96.6	94.8	97.3	94.7	
Taking heroin occasionally	18	96.7	97.2	96.9	96.9	97.1	96.8	96.6	97.9	96.9	97.2	96.7	97.3	96.8	97.0	96.2	
	19–22	98.6	97.8	98.3	98.3	98.6	98.7	98.3	98.3	98.3	97.9	99.2	98.2	98.1	98.1	98.3	
	23–26	—	—	—	—	99.2	98.2	98.8	99.1	98.4	98.3	98.1	99.0	98.7	98.4	98.6	
	27–30	—	—	—	—	—	—	—	—	99.2	97.3	99.0	98.9	97.0	98.9	98.7	
Taking heroin regularly	18	97.6	97.8	97.5	97.7	98.0	97.6	97.6	98.1	97.2	97.4	97.5	97.8	97.2	97.5	97.1	
	19–22	99.2	98.5	98.6	98.7	98.7	99.1	98.9	98.6	98.4	98.3	99.5	98.5	98.3	98.4	98.8	
	23–26	—	—	—	—	99.4	98.8	99.1	99.4	98.7	98.7	98.5	99.3	99.2	98.9	98.8	
	27–30	—	—	—	—	—	—	—	—	99.4	97.6	99.4	99.0	97.8	99.0	99.4	

(Table continued on next page.)

TABLE 6-2 (cont.)
Trends in Proportions Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. Do you disapprove of people (who are 18 or older) doing each of the following?	Age Group	Percentage disapproving ^a														2007–2008 change
		1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Trying marijuana once or twice	18	56.7	52.5	51.0	51.6	48.8	52.5	49.1	51.6	53.4	52.7	55.0	55.6	58.6	55.5	-3.0
	19–22	57.1	55.4	56.2	55.9	54.0	55.2	49.3	48.7	54.2	48.3	50.3	51.2	47.6	52.7	+5.1
	23–26	51.9	56.3	54.5	55.3	55.7	54.8	51.2	52.4	47.8	53.4	47.7	47.5	54.6	46.2	-8.4 s
	27–30	57.5	54.1	59.0	55.7	52.6	58.0	54.4	56.9	54.9	55.4	52.1	52.0	50.9	49.3	-1.6
Smoking marijuana occasionally	18	66.7	62.9	63.2	64.4	62.5	65.8	63.2	63.4	64.2	65.4	67.8	69.3	70.2	67.3	-2.9
	19–22	70.4	68.9	70.2	67.8	66.4	70.7	64.6	62.3	68.0	64.3	67.9	62.6	64.1	63.3	-0.9
	23–26	68.1	72.5	69.2	70.4	71.1	68.6	67.4	64.0	63.8	69.3	65.6	62.2	68.0	64.5	-3.5
	27–30	72.5	70.5	74.5	72.4	71.5	72.2	70.9	69.1	71.2	69.1	68.2	68.7	67.5	63.7	-3.9
Smoking marijuana regularly	18	81.9	80.0	78.8	81.2	78.6	79.7	79.3	78.3	78.7	80.7	82.0	82.2	83.3	79.6	-3.7 ss
	19–22	86.8	87.7	88.1	85.3	84.5	86.6	84.5	82.8	84.8	82.7	84.4	82.5	83.7	83.6	-0.1
	23–26	90.1	88.9	88.1	87.5	86.1	83.9	86.4	81.7	82.3	87.4	84.3	81.9	85.3	84.3	-1.0
	27–30	91.9	89.9	92.1	89.2	90.0	89.5	89.3	88.8	87.7	88.6	86.3	86.4	86.8	86.0	-0.8
Trying LSD once or twice	18	81.1	79.6	80.5	82.1	83.0	82.4	81.8	84.6	85.5	87.9	87.9	88.0	87.8	85.5	-2.4
	19–22	84.2	83.0	83.1	80.8	83.2	82.3	81.4	83.7	86.2	85.0	87.6	85.4	88.5	86.5	-2.0
	23–26	87.1	86.7	87.9	84.1	84.8	80.3	83.0	79.2	80.1	84.0	84.0	84.5	87.6	81.8	-5.8 s
	27–30	87.4	88.7	88.7	87.3	86.6	87.2	85.7	82.7	85.6	82.5	82.2	82.0	84.1	82.7	-1.4
Taking LSD regularly	18	92.5	93.2	92.9	93.5	94.3	94.2	94.0	94.0	94.4	94.6	95.6	95.9	94.9	93.5	-1.5
	19–22	96.8	97.0	97.4	96.3	97.0	96.8	96.5	96.9	98.4	97.3	98.9	97.8	97.7	96.8	-0.9
	23–26	96.7	97.7	96.1	97.6	98.0	97.0	97.1	97.9	96.9	97.1	98.7	97.0	98.4	97.4	-1.0
	27–30	98.6	98.1	97.5	97.4	97.9	98.6	98.2	98.0	98.2	98.2	97.2	96.7	97.2	97.1	-0.1
Trying ecstasy (MDMA) once or twice	18	—	—	82.2	82.5	82.1	81.0	79.5	83.6	84.7	87.7	88.4	89.0	87.8	88.2	+0.4
	19–22	—	—	—	—	—	—	81.5	80.3	87.2	83.5	90.3	87.5	88.5	89.5	+1.0
	23–26	—	—	—	—	—	—	80.6	80.6	80.2	83.1	83.9	83.9	87.4	83.9	-3.5
	27–30	—	—	—	—	—	—	84.2	84.0	86.3	83.2	82.4	82.2	81.8	82.7	+0.9
Taking ecstasy (MDMA) occasionally	18	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	19–22	—	—	—	—	—	—	92.8	91.8	95.6	93.8	96.7	94.0	95.3	94.8	-0.5
	23–26	—	—	—	—	—	—	90.5	91.8	92.1	93.3	94.4	93.7	94.3	94.0	-0.3
	27–30	—	—	—	—	—	—	91.7	93.0	94.3	91.0	92.1	93.4	92.8	94.1	+1.3
Trying cocaine once or twice	18	90.3	90.0	88.0	89.5	89.1	88.2	88.1	89.0	89.3	88.6	88.9	89.1	89.6	89.2	-0.5
	19–22	94.2	92.0	91.7	89.9	90.9	89.9	87.7	87.9	89.3	87.7	92.3	88.2	89.2	85.8	-3.4
	23–26	91.8	90.7	91.5	89.0	91.3	87.1	90.1	85.8	86.4	87.4	88.3	84.4	87.6	84.5	-3.1
	27–30	86.6	88.3	89.2	90.3	90.4	89.4	90.3	88.5	91.5	88.0	87.0	85.8	87.7	87.4	-0.3
Taking cocaine regularly	18	96.1	95.6	96.0	95.6	94.9	95.5	94.9	95.0	95.8	95.4	96.0	96.1	96.2	94.8	-1.4
	19–22	98.2	97.9	98.0	97.8	97.6	98.0	97.2	97.0	98.2	98.5	98.7	98.9	99.0	97.6	-1.4
	23–26	97.7	97.8	96.9	98.5	98.3	97.8	97.5	97.5	97.6	98.1	98.9	97.3	98.1	98.0	-0.2
	27–30	98.9	98.5	97.9	97.8	98.8	98.7	98.4	97.8	98.8	98.8	97.8	97.2	97.9	97.3	-0.6
Trying heroin once or twice	18	92.8	92.1	92.3	93.7	93.5	93.0	93.1	94.1	94.1	94.2	94.3	93.8	94.8	93.3	-1.5
	19–22	95.6	95.2	95.6	95.1	95.5	94.1	94.2	95.0	96.4	95.9	98.8	95.6	97.6	95.7	-2.0
	23–26	95.9	96.1	95.2	94.6	96.3	93.1	95.0	94.8	95.0	95.0	96.1	93.7	97.2	95.6	-1.6
	27–30	96.3	96.0	96.9	95.9	96.7	95.9	96.4	94.4	97.6	94.9	95.6	93.9	96.4	96.2	-0.2
Taking heroin occasionally	18	95.7	95.0	95.4	96.1	95.7	96.0	95.4	95.6	95.9	96.4	96.3	96.2	96.8	95.3	-1.4 s
	19–22	97.7	97.9	97.8	98.2	97.2	98.0	97.9	97.9	98.3	98.9	99.4	98.2	98.8	97.3	-1.5
	23–26	97.7	98.7	97.4	97.5	98.5	98.2	97.8	97.5	97.2	98.5	98.3	97.7	98.8	98.3	-0.4
	27–30	98.9	98.0	98.7	97.6	98.8	98.6	98.4	98.6	98.7	98.1	97.7	97.1	98.1	98.2	+0.1
Taking heroin regularly	18	96.4	96.3	96.4	96.6	96.4	96.6	96.2	96.2	97.1	97.1	96.7	96.9	97.1	95.9	-1.2
	19–22	98.4	98.3	98.1	98.3	98.2	98.5	98.2	98.3	98.8	99.0	99.2	98.9	99.1	98.3	-0.9
	23–26	98.7	98.9	97.6	98.5	98.7	98.8	98.4	98.3	98.6	98.9	98.9	98.0	99.0	99.1	+0.1
	27–30	99.1	98.6	98.4	98.1	98.8	98.7	98.7	98.4	99.3	98.8	99.1	97.5	98.2	98.4	+0.2

(List of drugs continued.)

TABLE 6-2 (cont.)
Trends in Proportions Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

		Percentage disapproving ^a																
		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	cont.
<i>Q. Do you disapprove of people (who are 18 or older) doing each of the following?</i>																		
Trying amphetamines once or twice	18	75.4	71.1	72.6	72.3	72.8	74.9	76.5	80.7	82.5	83.3	85.3	86.5	86.9	84.2	81.3		
	19–22	74.5	70.5	68.9	74.0	73.0	75.6	78.9	79.9	81.8	85.3	84.4	83.9	83.8	87.2	88.3		
	23–26	—	—	—	—	74.2	74.2	74.6	80.3	83.5	83.3	84.1	84.8	83.4	84.8	82.7		
	27–30	—	—	—	—	—	—	—	—	83.5	81.0	84.3	83.7	80.9	83.5	82.0		
Taking amphetamines regularly	18	93.0	91.7	92.0	92.6	93.6	93.3	93.5	95.4	94.2	94.2	95.5	96.0	95.6	96.0	94.1		
	19–22	94.8	93.3	94.3	93.4	94.9	96.6	96.9	95.1	97.5	96.8	97.5	97.7	96.7	97.3	97.9		
	23–26	—	—	—	—	96.6	95.9	96.6	97.0	97.2	98.1	97.9	97.9	97.7	98.4	97.7		
	27–30	—	—	—	—	—	—	—	—	98.1	96.5	98.6	97.8	96.8	97.7	99.0		
Trying sedatives/barbiturates^b once or twice	18	83.9	82.4	84.4	83.1	84.1	84.9	86.8	89.6	89.4	89.3	90.5	90.6	90.3	89.7	87.5		
	19–22	83.5	82.3	83.8	85.1	85.2	86.1	88.3	87.5	90.1	92.0	91.1	90.4	88.8	90.7	91.1		
	23–26	—	—	—	—	84.0	84.5	84.4	89.8	90.7	89.4	88.8	87.9	88.8	88.5	88.0		
	27–30	—	—	—	—	—	—	—	—	90.5	88.3	88.4	88.8	86.6	88.9	87.6		
Taking sedatives/barbiturates^b regularly	18	95.4	94.2	94.4	95.1	95.1	95.5	94.9	96.4	95.3	95.3	96.4	97.1	96.5	97.0	96.1		
	19–22	96.6	95.6	97.3	96.5	96.6	98.1	98.0	97.0	97.9	97.7	98.7	98.0	97.9	98.2	98.7		
	23–26	—	—	—	—	98.4	98.5	97.7	98.6	98.3	98.3	98.5	98.5	98.6	98.5	98.5		
	27–30	—	—	—	—	—	—	—	—	98.4	97.1	99.1	98.5	97.7	98.4	99.1		
Trying one or two drinks of an alcoholic beverage (beer, wine, liquor)	18	16.0	17.2	18.2	18.4	17.4	20.3	20.9	21.4	22.6	27.3	29.4	29.8	33.0	30.1	28.4		
	19–22	14.8	14.5	13.9	15.5	15.3	15.4	16.9	16.0	18.4	22.4	17.6	22.2	16.9	20.8	22.2		
	23–26	—	—	—	—	17.4	16.1	13.2	17.7	13.7	17.5	18.6	19.5	17.4	18.1	17.6		
	27–30	—	—	—	—	—	—	—	—	19.5	19.1	18.7	18.8	17.9	19.5	18.6		
Taking one or two drinks nearly every day	18	69.0	69.1	69.9	68.9	72.9	70.9	72.8	74.2	75.0	76.5	77.9	76.5	75.9	77.8	73.1		
	19–22	67.8	69.7	71.3	73.3	74.3	71.3	77.4	75.3	76.5	80.0	79.7	77.1	76.0	75.0	78.0		
	23–26	—	—	—	—	71.4	73.7	71.6	72.7	74.6	74.4	77.6	76.9	75.5	74.2	73.3		
	27–30	—	—	—	—	—	—	—	—	76.0	73.9	73.3	76.1	69.5	73.5	72.4		
Taking four or five drinks nearly every day	18	90.8	91.8	90.9	90.0	91.0	92.0	91.4	92.2	92.8	91.6	91.9	90.6	90.8	90.6	89.8		
	19–22	95.2	93.4	94.6	94.6	94.6	94.8	94.9	95.7	94.8	96.1	95.8	96.4	95.5	95.1	96.2		
	23–26	—	—	—	—	96.2	95.0	95.5	96.9	94.3	95.9	96.9	96.1	95.7	95.7	95.7		
	27–30	—	—	—	—	—	—	—	—	97.4	94.6	96.1	95.3	94.8	94.8	96.4		
Having five or more drinks once or twice each weekend	18	55.6	55.5	58.8	56.6	59.6	60.4	62.4	62.0	65.3	66.5	68.9	67.4	70.7	70.1	65.1		
	19–22	57.1	56.1	58.2	61.0	59.7	59.4	60.3	61.6	64.1	66.3	67.1	62.4	65.6	63.5	68.1		
	23–26	—	—	—	—	66.2	68.3	66.5	67.5	65.2	63.2	66.9	64.6	69.6	66.8	66.9		
	27–30	—	—	—	—	—	—	—	—	73.9	71.4	73.1	72.1	68.4	73.4	73.5		
Smoking one or more packs of cigarettes per day	18	70.8	69.9	69.4	70.8	73.0	72.3	75.4	74.3	73.1	72.4	72.8	71.4	73.5	70.6	69.8		
	19–22	68.7	68.1	66.3	71.6	69.0	70.5	71.4	72.7	73.8	75.6	73.7	73.2	72.6	72.8	75.3		
	23–26	—	—	—	—	69.9	68.7	67.5	69.7	66.4	71.1	71.5	77.2	73.6	72.9	70.3		
	27–30	—	—	—	—	—	—	—	—	72.8	69.4	73.5	71.2	70.7	73.8	72.3		
<i>Approximate Weighted N =</i>	18	3,261	3,610	3,651	3,341	3,254	3,265	3,113	3,302	3,311	2,799	2,566	2,547	2,645	2,723	2,588		
	19–22	588	573	605	579	586	551	605	587	560	567	569	533	530	489	474		
	23–26					542	535	560	532	538	516	524	495	538	514	475		
	27–30									526	509	513	485	512	462	442		

(Table continued on next page.)

TABLE 6-2 (cont.)
Trends in Proportions Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. Do you disapprove of people (who are 18 or older) doing each of the following?	Age Group	Percentage disapproving ^a														2007–2008 change
		1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Trying	18	82.2	79.9	81.3	82.5	81.9	82.1	82.3	83.8	85.8	84.1	86.1	86.3	87.3	87.2	-0.1
 amphetamines	19–22	85.0	84.4	83.3	84.6	84.9	83.8	82.1	81.4	86.3	82.1	88.2	84.9	84.8	86.7	+1.9
 once or twice	23–26	86.0	86.4	85.7	83.5	84.5	82.4	83.9	83.5	79.9	81.6	81.3	79.0	85.8	79.7	-6.1 s
	27–30	83.1	85.8	86.3	85.9	86.4	84.5	86.0	86.4	84.9	82.4	81.3	81.1	84.5	83.7	-0.9
Taking	18	94.3	93.5	94.3	94.0	93.7	94.1	93.4	93.5	94.0	93.9	94.8	95.3	95.4	94.2	-1.3
 amphetamines	19–22	96.8	97.2	97.8	96.7	97.5	96.1	97.3	96.4	97.1	97.1	98.4	97.5	98.6	96.2	-2.4 s
 regularly	23–26	97.0	97.9	97.0	98.0	97.0	97.6	96.8	96.3	97.2	95.9	98.3	96.2	97.6	97.3	-0.3
	27–30	98.9	98.2	98.1	97.7	98.2	98.5	97.6	97.4	98.1	98.0	97.6	96.4	98.4	97.2	-1.2
Trying sedatives/ barbiturates^b	18	87.3	84.9	86.4	86.0	86.6	85.9	85.9	86.6	87.8	83.7	85.4	85.3	86.5	86.1	-0.4
 once or twice	19–22	90.5	89.1	86.6	85.8	86.6	84.2	85.2	84.2	87.7	81.8	86.6	83.4	82.7	82.1	-0.7
	23–26	89.3	88.3	88.3	87.4	87.3	85.2	86.9	86.8	81.8	80.3	81.6	80.5	84.3	77.7	-6.6 s
	27–30	88.0	89.4	88.8	88.4	87.6	87.3	88.5	86.9	89.2	81.8	78.7	80.1	83.5	80.5	-3.0
Taking sedatives/ barbiturates^b	18	95.2	94.8	95.3	94.6	94.7	95.2	94.5	94.7	94.4	94.2	95.2	95.1	94.6	94.3	-0.3
 regularly	19–22	97.7	97.9	97.7	97.7	97.3	97.4	96.9	97.8	98.5	96.6	98.3	98.1	98.3	96.7	-1.6
	23–26	97.4	98.4	97.4	98.5	97.6	97.4	97.0	97.1	97.1	96.1	98.0	96.3	97.8	96.7	-1.1
	27–30	99.0	98.5	97.9	97.7	98.5	98.1	98.4	97.2	98.4	98.1	96.5	95.6	97.4	97.4	0.0
Trying one or two drinks of an alcoholic beverage (beer, wine, liquor)	18	27.3	26.5	26.1	24.5	24.6	25.2	26.6	26.3	27.2	26.0	26.4	29.0	31.0	29.8	-1.2
	19–22	22.0	22.0	18.3	21.5	18.3	18.4	16.3	18.3	20.1	20.7	22.3	17.8	17.3	20.5	+3.2
	23–26	16.5	18.0	15.8	18.6	19.1	19.9	15.9	18.1	13.0	16.3	13.5	14.7	14.9	12.5	-2.4
	27–30	18.2	16.1	17.4	15.2	15.9	14.8	15.9	18.4	15.4	18.8	16.1	15.0	14.2	11.9	-2.3
Taking one or two drinks nearly every day	18	73.3	70.8	70.0	69.4	67.2	70.0	69.2	69.1	68.9	69.5	70.8	72.8	73.3	74.5	+1.2
	19–22	74.7	73.5	73.2	70.3	67.3	66.7	68.3	63.9	66.9	68.1	64.6	68.2	65.1	65.2	+0.1
	23–26	69.7	70.6	68.4	70.2	73.4	66.3	66.5	62.7	65.0	61.7	64.4	62.0	62.4	66.4	+4.1
	27–30	71.8	71.4	71.8	69.8	67.9	65.9	68.9	70.9	63.1	66.7	60.5	62.0	65.8	59.5	-6.3
Taking four or five drinks nearly every day	18	88.8	89.4	88.6	86.7	86.9	88.4	86.4	87.5	86.3	87.8	89.4	90.6	90.5	89.8	-0.7
	19–22	95.5	94.2	93.9	92.4	92.4	92.8	94.2	92.6	92.5	92.2	93.2	92.9	92.9	94.0	+1.1
	23–26	95.2	96.5	93.8	96.1	95.1	94.3	93.5	93.7	92.6	93.1	94.8	92.9	95.6	94.9	-0.7
	27–30	96.7	96.4	96.2	95.0	97.2	95.3	96.1	95.4	95.6	96.0	92.8	92.7	95.0	93.9	-1.2
Having five or more drinks once or twice each weekend	18	66.7	64.7	65.0	63.8	62.7	65.2	62.9	64.7	64.2	65.7	66.5	68.5	68.8	68.9	+0.2
	19–22	66.0	69.2	66.5	63.2	63.5	65.1	58.3	57.5	61.9	59.4	60.1	59.3	59.1	63.4	+4.2
	23–26	65.3	70.9	66.6	69.5	68.1	66.2	66.0	61.2	65.5	60.9	64.5	59.7	62.4	63.0	+0.6
	27–30	73.7	72.4	73.0	71.1	73.1	73.1	73.0	70.9	71.5	73.8	67.5	67.3	71.5	66.4	-5.2
Smoking one or more packs of cigarettes per day	18	68.2	67.2	67.1	68.8	69.5	70.1	71.6	73.6	74.8	76.2	79.8	81.5	80.7	80.5	-0.1
	19–22	69.8	72.2	74.3	72.3	70.1	73.1	73.2	73.4	73.4	74.8	81.5	77.2	81.0	80.4	-0.6
	23–26	72.2	73.0	71.7	73.9	73.8	72.7	77.3	74.8	75.7	76.2	74.8	74.1	76.2	77.9	+1.6
	27–30	73.9	72.7	74.3	71.7	71.0	78.6	75.2	78.8	76.2	77.6	77.3	73.9	81.1	74.5	-6.7 s
<i>Approximate</i>	18	2,603	2,399	2,601	2,545	2,310	2,150	2,144	2,160	2,442	2,455	2,460	2,377	2,450	2,314	
<i>Weighted N =</i>	19–22	465	480	470	446	449	416	413	402	396	431	378	378	333	365	
	23–26	466	449	423	401	397	389	404	346	385	403	374	364	325	335	
	27–30	450	430	453	449	429	395	368	359	346	370	367	330	355	339	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

“—” indicates data not available.

^aAnswer alternatives were: (1) Don't disapprove, (2) Disapprove, and (3) Strongly disapprove.

Percentages are shown for categories (2) and (3) combined.

^bIn 2004 the question text was changed from “barbiturates” to “sedatives/barbiturates” and the list of examples was changed from “downers, goofballs, redds, yellows, etc.” to just “downers.” These changes likely explain the discontinuity in the 2003 and 2004 results.

Chapter 7

THE SOCIAL CONTEXT

The social contexts in which individuals place and otherwise find themselves can influence their likelihood of using drugs in a number of ways. The environment can bring about an awareness of new drugs (knowledge of their existence and potential for altering mood and consciousness) and provide social norms regarding drug use. It can also provide social modeling of either use or abstention from use, and influence the availability of drugs (through friends and friends' contacts). Since its inception, MTF has measured three important features of the social environment: (1) peer groups' norms about drug use, (2) amount of direct exposure to drug use through friends and others, and (3) perceived availability of drugs. All three factors are measured by self-reports and are, therefore, measures of the perceived environment, though evidence suggests that they bear a strong correlation with the actual environment. We believe that these three factors exert important influences on substance use, at both the individual (micro) and the aggregate (macro) level.

In *Volume I*, we examined these factors among secondary school students. In this chapter, we do the same for the young (and sometimes middle) adult population, whose social contexts typically differ considerably from what they were in high school. Most high school graduates today enter college, many get civilian jobs, and some enter military service. These transitions almost always change the institutional environments experienced by young adults (e.g., colleges, work organizations, military services, etc.) and therefore the circles of people to whom they are exposed and with whom they develop friendships.

Each of the question sets discussed here is contained in only one of the six questionnaire forms, so the case counts are lower than those presented in most chapters in this volume. (Also, in comparison to the secondary school samples covered in *Volume I*, follow-up samples are necessarily much smaller.) Therefore, the prevalence and trend estimates are more subject to fluctuation.

PEER NORMS AMONG ADULTS

Table 7-1 provides current levels and trends in perceived friends' disapproval of drug use as reported by 12th graders, 19- to 22-year-olds, 23- to 26-year-olds, and 27- to 30-year-olds. (These are the same age groupings used in chapter 6.) Trend data are available since 1980, 1984, and 1988, respectively, for these three 4-year age groupings of young adults.

The questions regarding friends' disapproval include the same answer scale (stated in terms of strength of disapproval associated with different use levels of the various drugs) as the questions on the respondents' own attitudes about those behaviors (discussed in chapter 6). The list of drug-using behaviors is shorter here, and the questions appear on a different questionnaire form and, therefore, have a different set of respondents. However, because the questionnaire forms are distributed randomly in senior year, there should be no systematic sample differences across forms. Furthermore, the results for perceived peer norms are generally quite consistent with

those for personal *disapproval* in the aggregate.⁵³ Exceptions are *trying marijuana once or twice* and *smoking one or more packs of cigarettes per day*, to which respondents have consistently reported their friends' attitudes as *more* disapproving than their own attitudes (especially in the oldest age band), and *heavy weekend drinking*, to which friends' attitudes are seen as *less* disapproving than their own.

Current Perceptions of Friends' Attitudes

Table 7-1 provides trends for each age band in the proportions of respondents indicating how their close friends would feel about the respondent engaging in various drug-using behaviors. For purposes of simplification, we begin by addressing results across the entire 19- to 30-year age band (tabular data for the entire age band are not presented). Then we distinguish among the three young adult age bands: 19–22, 23–26, and 27–30, along with 18-year-olds.

- We find that the peer norms reported by young adults 1 to 12 years past high school are quite similar to those reported by 12th graders. That is, for each of the *illicit drugs other than marijuana*, the great majority of young adults think that their close friends would disapprove of their even *trying* such drugs once or twice (90% for *cocaine*, 88% for *LSD*, and 88% for *amphetamines*).
- Well over half of young adults (61%) now think their friends would disapprove of their even trying *marijuana*, while two thirds (68%) think their friends would disapprove of occasional use, and about 84% think they would disapprove of regular use. So, clearly the norms differ as a function of level of marijuana use, but for all levels of use they tend to be restrictive for the majority of young adults.
- Nearly two thirds (64%) of young adults say their friends would disapprove if they were *daily drinkers*, and 9 out of 10 (91%) think friends would disapprove if they had four or five drinks nearly every day.
- Friends' disapproval of *heavy drinking on weekends* is distinctly lower. Only 55–57% of any of the young adult age groups think that their friends would disapprove of their having five or more drinks once or twice each weekend. The 19- to 22-year-olds and the 23- to 26-year-olds—among whom this kind of drinking is most prevalent—have slightly lower levels of disapproval (both 55%) than 18-year-olds (62%) or 27- to 30-year-olds (57%).
- Peer disapproval of *cigarette* smoking is high in all four age bands: 83% of 12th graders say their friends would disapprove of pack-a-day smoking, as do 84–88% of 19- to 30-year-olds.

⁵³The question reads, "How do you think your close friends feel (or would feel) about you... [smoking marijuana once or twice]?" The answer categories are "don't disapprove," "disapprove," and "strongly disapprove." Percentages discussed are for the last two categories combined.

Trends in Peer Norms

- Important changes in the social acceptability of drug-using behaviors among both 12th graders' and young adults' peers have occurred since MTF began (see Table 7-1). Among 12th graders, the proportion saying their friends would disapprove of their trying *marijuana* rose from 41% in 1979 to 73% in 1992—a period of substantial decline in use. Friends' disapproval also grew substantially stronger in all of the young adult age bands in the years for which data are available. For example, among 19- to 22-year-olds, the proportion thinking their friends would disapprove if they even tried marijuana rose from 41% in 1981 to 65% in 1992. A similar peak in disapproval occurred for 23- to 26-year-olds in 1992 and 1993, and among 27- to 30-year-olds in 1994 and 1995—66% for both age bands. In all age groups, disapproval subsequently declined, though the declines in peer disapproval were earliest and greatest among 12th graders. The decline ended in 1997 for 12th graders and began to reverse, but continued through 2002 among 19- to 26-year-olds. There has been little systematic change in the past several years.

Friends' disapproval of more frequent marijuana use also rose until the early 1990s among 18-year-olds, and then declined between 1992 and 1997. It declined through 1999 among 19- to 22-year-olds and continued to decline among 23- to 30-year-olds through 2003. In essence, peer norms have moved in a way consistent with the existence of some lasting cohort differences in use.

- There was a more gradual increase in peer disapproval levels of trying an *amphetamine* for all age groups through 1991, followed by a small decline evident among 12th graders through 1997. Since 1997, levels of disapproval among 18- to 30-year-olds have increased to some extent, though not dramatically. In 2008, disapproval levels ranged from 85% to 90%.
- Through 1991, peer disapproval of trying *LSD* showed very little change in any of the age bands, but it fell some in the 1990s, especially among 18-year-olds and subsequently among 19- to 22-year-olds. These declines bottomed out in a staggered fashion, beginning with the 12th graders in 1997, who have since shown a seven-percentage-point increase in peer disapproval. There has been an eight-percentage-point increase among 19- to 22-year-olds (since 2000), and a six-percentage-point increase among 23- to 26-year-olds (since 2001). There is now almost no difference among the age groups, with 85–90% of all of them saying their friends would disapprove of their trying LSD.
- Perceived peer norms regarding *cocaine* use were first measured in 1986. During the next eight years, self-reported cocaine use declined substantially as peer norms in all age bands shifted considerably toward disapproval. For example, by 1994, 95% of the 19- to 22-year-olds thought their friends would disapprove of their even trying cocaine. After 1994, peer norms against use continued to strengthen a bit in the upper age bands, perhaps through generational replacement, but weakened slightly in the younger age bands, likely reflecting a new cohort effect. In recent years there has been little difference by age in peer norms against cocaine use, with 88–91% saying their friends would disapprove of their even trying it. By way of contrast, in 1986 that statistic ranged between 71% and 80% among 18- to 26-year-olds.

- Peer norms regarding *heavy drinking on weekends* (five or more drinks once or twice each weekend) among the three young adult age groups have tended to be weakest among the 19- to 22-year-old age stratum (where such behavior is most common) and strongest among the 27- to 30-year-old stratum. Since 2002, however, the 23- to 26-year-olds were also low relative to the other two age bands in their disapproval of weekend binge drinking. Among 12th graders, friends' attitudes became somewhat more restrictive between 1981 and 1992 (and binge drinking declined during that interval), but attitudes have been fairly level since then. There was a similar upward trend in disapproval among the various young adult age bands that followed a staggered pattern, likely reflecting a cohort effect. However, between 1997 and 2000 the 19- to 22-year-old age group became somewhat less disapproving of occasional binge drinking; this was followed by a decline in peer disapproval between 2001 and 2004 among 23- to 26-year-olds, but little net change by 2008 among 27- to 30-year-olds. Despite some increases in peer disapproval over the years, this rather extreme form of drinking has the least restrictive perceived peer norms of all of the substance-using behaviors measured in MTF.
- Peer norms regarding *cigarette smoking* (at the rate of one or more packs per day) became somewhat more restrictive among 12th graders in the early years of MTF; peer disapproval rose from 64% in 1975 to 73% in 1979. There was little further net change for 13 years through 1992, when friends' disapproval stood at 76%. However, peer disapproval of smoking slipped some in the 1990s. Between 1992 or 1993 and 1997 or 1998, all age groups showed a decrease in perceived peer disapproval of smoking. Since then, the two younger strata have shown a considerable strengthening of norms against smoking. Between 1998 and 2008, the proportion saying that their close friends would disapprove of their smoking a pack or more of cigarettes per day rose from 69% to 83% among 18-year-olds and from 69% to 86% among 19- to 22-year-olds. The two older strata did not see a comparable change until 2006, when there was a significant increase observed among 23- to 26-year-olds, who have seen peer disapproval rise from 77% in 2005 to 88% in 2008. The change has not yet manifested itself among the 27- to 30-year-olds; their rates have stayed between 81% and 86% during the 2000s. In 2008, the norms across the age bands range between 83% and 88% reporting their friends would disapprove of pack-a-day smoking.

ADULTS' EXPOSURE TO DRUG USE THROUGH FRIENDS AND OTHERS

Exposure to drug use is important because it provides both the modeling of the behavior by peers (possibly including direct encouragement to use) and immediate access. Exposure is measured by two sets of questions, each appearing on a (different) single questionnaire form. The first set asks the respondent to estimate what proportion of his or her friends use each drug, while the second asks, "During the LAST TWELVE MONTHS how often were you around people who were using each of the following to get high or for 'kicks'?" The same questions are asked of 12th graders, and their results are included for comparison purposes in Tables 7-2 and 7-3. (Questions about direct exposure to drug use were not included in the questionnaires for 35–50-year-olds.) We continue to deal with four-year age bands for the friends' use measures in order to increase the reliability of the estimates. (Ages 35, 40, 45, and 50 are included as one-year age

bands, with both half samples from those cohorts surveyed; those years have larger numbers of cases than single years at the earlier ages because only one questionnaire form is used in the surveys of those age groups, instead of six.) At the end of each table is a summary of the weighted number of cases upon which each annual estimate is based. (The actual numbers of cases are somewhat higher.)

Exposure to Drug Use

- Relatively high proportions of young adults in all of these age bands have at least *some* friends who use *some illicit drug* (Table 7-2). In 2008, illicit drug use by at least some friends is reported by 80% of 12th graders, 67% of 27- to 30-year-olds, 45% of 35-year-olds, 36% of 45-year-olds, and 39% of 50-year-olds. The proportions who say that *most or all* of their friends use one or more of the illicit drugs are much lower: 21% for 12th graders, falling to 14% of 19- to 22-year-olds, 9% of 23- to 26-year-olds, 6% of 27- to 30-year-olds, and between 1.3% and 2.2% for the 35-, 40-, 45-, and 50-year-olds—quite a dramatic difference across ages, and one that is consistent with the large differences in their own self-reported current use.
- With regard to *illicit drugs other than marijuana*, taken as a whole, considerably fewer report that *any* of their friends use compared to what is true for marijuana use (see below): 49% for 12th graders, 46% for 19- to 22-year-olds, 36% for 27- to 30-year-olds, 24% for 35-year-olds, 21% for 45-year-olds, and 25% for 50-year-olds. The proportions saying that *most or all* of their friends use *illicit drugs other than marijuana* in 2008 are 5%, 3%, and 2%, respectively, for the three youngest age bands, with fewer than 1% of respondents over the age of 30 reporting such high proportions of their friends using other illicit drugs. Thus, relatively few in these age groups appear to be deeply immersed in a drug culture involving illicit drugs beyond marijuana.
- With respect to individual illicit drugs, exposure among all of the age groups is greatest for *marijuana*, with 78% of 12th graders, 75% of 19- to 22-year-olds, 64% of 27- to 30-year-olds, and 29–42% of 35- through 50-year-olds reporting that at least some of their friends use the drug. The next highest exposures are for *hallucinogens other than LSD* (29% among 12th graders, 22% among 19- to 22-year-olds, 21% among 23- to 26-year-olds, and 17% among 27- to 30-year-olds), followed by *cocaine*, *amphetamines*, and *ecstasy (MDMA)*. Because of the dramatic increase in its use during the 1990s and early 2000s, ecstasy surpassed a number of the more traditional drugs, though its use has declined sharply in recent years. (It should be noted that use of several illicit drugs was not asked of the age groups above 30 due to space limitations in the single questionnaires used at those ages.)

For the remaining illicit drugs, the proportion of young adults reporting that some friends use a given drug is 10% or higher in at least one of the young adult age groups for the following drugs: *LSD* (9–13%), *narcotics other than heroin* (14–17%), *sedatives (barbiturates)* (13–18%), *steroids* (7–15%), *crack cocaine* (6–12%), and *tranquilizers* (12–13%). See Table 7-2 for specifics.

- For most illicit drugs, the proportion of young adults having any friends who use them decreases with age, consistent with the age differentials in self-reported use. The steepest declines occur with *inhalants* (18% of 18-year-olds down to 2.7% of 27- to 30-year-olds). (Inhalant use is not asked of the age groups above 30, precisely because of this sharp decline in use with age.)
- For some years, *cocaine* showed significantly higher rates of active use among adults compared to 12th graders. That is no longer true, although there is rather little drop-off with age in early adulthood; consequently, there is not a great difference associated with age in having friends who use cocaine (21–25% for all four of the younger age groups). (The 35-, 40-, 45-, and 50-year-olds are asked separately about cocaine powder and crack use; far fewer, but still a fair proportion, report having friends who use cocaine powder—11% for age 35 and between 6% to 7% in 2008 for the three older groups.)
- For *crack*, however, the story is different. Reported friends' use of crack now descends sharply with age, although this was not true in the mid-1980s, when measures of crack use were first included in the surveys. In 2008, 19% of 12th graders report having any friends using crack, versus 12% of 19- to 22-year-olds, 6% of 27- to 30-year-olds, and 2–3% of 35- to 50-year-olds.
- The proportion reporting that they have any friends who use *heroin* also decreases sharply with age, from 11% among 12th graders to 2% among 27- to 30-year-olds. *Narcotics other than heroin* also shows a less sharp decline with age, from 21% among 18-year-olds, to 17% among 19- to 22-year-olds, to 14% among 29- to 30-year-olds. (Older respondents are not asked these questions.)
- In general, it appears that some respondents who report that their friends use *illicit drugs* are not directly exposed to that use themselves, judging by the differences in proportions saying they have some friends who use (Table 7-2) and the proportions who say they have been around people who were using during the prior year (Table 7-3).
- With respect to *alcohol* use, the great majority of young adults have at least *some* friends who *get drunk at least once a week*, although this differs by age: 76% of 12th graders, 81% of 19- to 22-year-olds, 80% of 23- to 26-year-olds, 72% of 27- to 30-year-olds, 56% of 35-year-olds, 48% of 40-year-olds, 46% of 45-year-olds, and 40% of 50-year-olds. Given the potential serious consequences of this behavior, these rates are impressively high across a wide age range. The proportions who say *most or all* of their friends get drunk once a week differ more substantially by age: 25% of 12th graders and 29% of 19- to 22-year-olds, declining sharply to 11% of 27- to 30-year-olds and 3% of 50-year-olds. Note in particular how high these rates are among the high school and college-age populations. In terms of direct exposure during the past year to people who were drinking alcohol “to get high or for ‘kicks,’” having some such exposure is almost universal in the three 4-year age groups of young adults: 91%, 92%, and 94%, respectively (see Table 7-3).

- From ages 18 through 30, about four fifths of respondents (78–87%) have at least a few friends who *smoke cigarettes*, with considerable falloff by age 35. In fact, 13–14% of the 12th graders and 19- to 22-year-olds state that *most or all* of their friends smoke. Above those ages, the proportions decline to 11% of 27- to 30-year-olds and 4–6% for those 35 years of age and older. This increase in the segregation of smokers from nonsmokers likely reflects the stratification of young people after high school as a function of educational attainment, which is highly correlated with cigarette smoking. Also, it can be seen in Table 7-2 that there was much less age-related difference in the late 1980s, suggesting that the sharp rise in smoking among high school students during much of the 1990s, followed by a sharp decline in the years since, accentuated the age differentials, and that those differentials remain, reflecting lasting cohort effects.

Trends in Exposure to Drug Use

Tables 7-2 and 7-3 also provide *trend* data on the proportions of respondents' friends using drugs and the proportion of respondents directly exposed to drug use by others. Both of these measures of exposure to use will be discussed in this section. Once again, trends are available for 19- to 22-year-olds since 1980, for 23- to 26-year-olds since 1984, and for 27- to 30-year-olds since 1988. Data for 35-, 40-, 45-, and 50-year-olds are available on friends' use since 1994, 1998, 2003, and 2008, respectively. (Questions about frequency of being around drug users were not included in the questionnaires administered to respondents age 35 and older, so those age bands are not included in Table 7-3. However, they were asked about the proportions of their friends using.) Twelfth-grade data have also been included in these tables for comparison purposes.

- An examination of Table 7-3 shows that exposure to illicit drug use (in the 12 months preceding the survey) declines with age for *any illicit drug, marijuana, and any illicit drug other than marijuana*, as well as for nearly all of the specific illicit drugs. In general, these differences replicate across different historical periods, with the exception of *cocaine*, which did not show a decline in exposure with increasing age until after 1996. These declines reflect age effects (changes with age observed across multiple cohorts) in both exposure to use and in personal use of most drugs.
- Until 1992, young adults' trends in exposure to use tended to parallel those observed for 12th graders. From 1980 to 1992, that meant a decreasing number of respondents were exposed to *any illicit drug* use (Table 7-3) or reported any such use in their own friendship circle (Table 7-2). After 1992, however, an important *divergence* in trends emerged: 12th graders showed a substantial increase in both friends' use and exposure to use (as well as self-reported use); 19- to 22-year-olds showed a similar rise, but lagged by a few years; 23- to 26-year-olds subsequently showed some rise; while the 27- to 30-year-old age band did not show a rise until 2002. As discussed in earlier chapters, this pattern no doubt reflects the emergence of lasting cohort differences driven by generational replacement.
- *Marijuana* showed a very similar pattern of change. It is particularly noteworthy that, while 34% of 19- to 22-year-olds in 1980 said *most or all* of their friends used marijuana, only 9% said the same in 1993. Clearly, the number of friendship groupings in which marijuana use was widespread dropped dramatically over that earlier interval. Friends'

use increased to 19% by 1999, during the relapse phase in the larger epidemic, where it remained for a couple of years, before falling to 12% by 2008. Self-reported use and friends' use both increased significantly among 18-year-olds in 2008. Whether this reflects a turnaround in the marijuana situation may become evident with another year or two of data.

- The proportion reporting having any friends who use *any illicit drugs other than marijuana* began to decline after 1982. By 1991 or 1992 there had been a considerable drop in all four age groups. (This drop appears to be due particularly to decreases in friends' use of *cocaine* and *amphetamines*, although there were decreases for *sedatives (barbiturates)* and *tranquilizers* as well.) The levels then began to rise in the two youngest age bands in the early 1990s, while at the same time declining in the two oldest age bands, opening up a large age-related difference in exposure to use. Since 2001 there has been some decline in friends' use in the two youngest age strata, narrowing the age differences.
- Between 1987 and about 1992, there was a considerable drop in all four age groups in the proportion of respondents with friends who used *crack*. (Self-reported use declined in the same period.) After that decline, the rates of friends' use increased some in the two youngest age bands and decreased some in the four oldest ones, resulting in a large age differential that is still evident.
- There were substantial increases between the early 1990s and about 2000 in the proportion of 18- and 19- to 22-year-olds reporting that they have friends using *narcotics other than heroin*, and smaller increases among 23- to 30-year-olds, resulting in some fair age-related differences. By 2008, the 18- and 19- to 22-year-olds had declined some, while the 23- to 30-year-olds had increased some, narrowing the age differences.
- The proportions saying that any of their friends use *ecstasy (MDMA)* increased sharply in all age groups for which data are available, though in a staggered fashion. Twelfth graders showed the first sharp increase beginning after 1992, 19- to 22-year-olds after 1994, 23- to 26-year-olds after 1996, and 27- to 30-year-olds after 1997. These sharp increases ended among 12th graders in 2001 and among 19- to 30-year-olds a year later. Since those peak levels, the proportions saying that they had any friends using ecstasy have generally declined, corresponding with a decline in self-reported use. In all four age groups, 18–25% now report that any of their friends use ecstasy.
- For all four age groups, there were modest declines between 1987 and 1992 in the proportion saying that most or all of their friends drink *alcohol*. After 1992, there was very little change in the four youngest age bands until 2002, when 12th graders' friends' use dropped while continuing to climb in the other age bands. In the years for which data are available, the older three age bands have shown an increase in the proportions saying that most or all of their friends drink alcohol—an increase not seen in the younger three age bands. The age groups above age 30 have consistently been much less likely to report that most or all of their friends *get drunk* at least once a week, compared with those ages 18 to 26. The rates for the three younger groups are very high—between 23% and 29%.

- Among 12th graders, the proportion who said most or all of their friends smoked *cigarettes* declined appreciably between 1975 and 1981, the same period in which self-reported use declined. After that, neither measure showed much change until about 1992. Thereafter, substantial increases in both measures occurred. By 1997, fully one third (34%) of 12th graders reported that most or all of their friends smoked cigarettes (up from 21% in 1992); since then, that statistic has declined (to 14% in 2008) along with self-reported use. Among 19- to 22-year-olds, a decline in friends' use occurred between 1980 (or possibly earlier) and 1985, followed by a leveling through 1994. The percentage saying most friends smoke increased from 22% in 1994 to 29% in 2000, before beginning to decline. Among 23- to 26-year-olds, a downturn was evident between at least 1984 (the first year for which data are available) and 1988, and then reported friends' use leveled. After 2002, some slight increase occurred, but a reversal has been occurring since 2004. These staggered changes until about 1998 illustrate that cohort effects were moving up the age spectrum. Since 1998 (or the earliest year available, for the age bands above 30), the proportion saying that any of their friends smoke has shown some decline among those above age 30, but little or no change among 23- to 30-year-olds (which now contain the heavier smoking senior classes of the mid-1990s) until about 2006.

Nearly all of these changes in exposure to drug use parallel changes in self-reported use by these age groups. This pattern reinforces the validity of self-report data, because there would presumably be less motivation to distort answers about the proportion of an unnamed set of friends who use a drug than about one's own use. Also reassuring is the systematic nature of the patterns of change across age strata (whether in terms of parallel changes consistent with a secular trend, or staggered ones consistent with a cohort-related trend).

PERCEIVED AVAILABILITY OF DRUGS AMONG ADULTS

Adults participating in the follow-up surveys receive survey questions identical to those asked of 12th graders regarding how difficult they think it would be to get each of the various drugs if they wanted them. The questions are contained in only one of the six questionnaire forms. Data for the young adult follow-up samples, which are grouped into the same four-year age bands used above, are presented in Table 7-4, along with data for 12th graders and 35-, 40-, 45-, and 50-year-olds. Sample sizes are presented at the bottom of the table.

Perceived Availability

Much like 12th graders, substantial proportions of the American adult population have access to various illicit drugs. (We do not ask about access to alcohol and cigarettes because we assume these are readily available to all adults.)

- *Marijuana* is the most available illicit drug in 2008, with 85–88% of the young adult age strata saying it would be “fairly easy” or “very easy” to get. Access decreases steadily with age after age 30; but even at age 50, 64% of respondents say they can get it fairly or very easily.

Monitoring the Future

- Though less available than marijuana, **amphetamines** are fairly available, with 45–50% of young adults and 33–41% of 35-, 40-, 45-, and 50-year-olds saying they would be easy to get.
- **Cocaine** is reported as readily available by a significant proportion of young adults, with 44–51% saying it would be easy to get. **Powdered cocaine** availability does not differ much by age (39–48%) until after age 30, when it drops to 33% among 50-year-olds. **Crack** is available to somewhat smaller proportions than powdered cocaine—31–36% for all three post-high school young adult age strata and 33–35% for 35-, 40-, 45-, and 50-year-olds.
- **Sedatives (barbiturates)** are a bit less available than amphetamines to these age groups, with little variation across age up to age 30 (43–45%). More than a third of young adults (38–41%) say they can get **narcotics other than heroin** easily, as do 35% of 12th graders. **Tranquilizers** are reported as available by considerably fewer (23–33%), which historically was not always the case. None of these three drugs are included in follow-up surveys of respondents over age 30.
- About a quarter (23–26%) of young adults and 12th graders say that they could get **heroin** fairly or very easily, though far fewer report having used heroin. (The question is not asked of respondents above age 30.)
- **Ecstasy (MDMA)** is readily available to 35–42% of young adults and 12th graders. (The question is not asked of respondents above age 30.)
- **Hallucinogens other than LSD** are reported as available by 43% of 12th graders and 35–39% of the three young adult strata. (The question is not asked of respondents above age 30.)
- **LSD** is the least available drug, declining from 29% among 12th graders to 19% among 50-year-olds.
- **Crystal methamphetamine (ice)** is perceived to be available by between a quarter and a third of each young adult age group (24–31%). (The question is not asked of respondents above age 30.)
- **Steroids** show some declines in perceived availability with increasing age, as has generally been the case, ranging from 35% among 12th graders, 34% among 19- to 22-year-olds, down to 30–31% among 23- to 30-year-olds. (The question is not asked of respondents above age 30.)

Trends in Perceived Availability

- **Marijuana** has been almost universally available to the adolescent and young adult age groups throughout the historical periods covered by the data (since 1975 in the case of 12th graders). There was a slight decrease from 1979, the peak year, through 1991 among

12th graders, and a slightly larger decrease from 1980 through 1991 among 19- to 22-year-olds. Availability rose by a few percentage points in nearly all strata between about 1993 and 2001, and since then has slipped back a few percentage points in the two younger strata. Perceived availability is now a bit higher for the younger age groups (84–88% for 12th graders through 27- to 30-year-olds versus 64–76% for those ages 35 to 50). In general, though, their trends in availability have been quite parallel since the late 1990s, suggesting secular trends in prevailing conditions that affect availability.

- **Cocaine** availability increased among all three younger age strata in 1984–1988, reaching historic highs in 1988 and 1989. (Twelfth graders showed a rise in availability in earlier years—from 1975 to 1980—followed by a leveling between 1980 and 1984. Availability was also level during the latter period among 19- to 22-year-olds.) From a policy perspective, it is worth noting that in all three age bands for which we have data, the perceived availability of cocaine *increased* in 1987—the same year that use actually dropped sharply. Between 1988 and 1989, in the two younger age strata (ages 18–22), the proportions reporting that they could get cocaine fairly easily were still increasing, whereas in the older age strata the proportions were beginning to decrease. In 1990 and 1991, all four groups reported decreased availability—quite parallel to the number who had friends who were users and to personal use, both of which dropped substantially in these years and then leveled in 1992. Perceived availability of cocaine dropped to between 49% and 57% for all four age groups by 1993, with the absolute declines ranging from four to seven percentage points. After that there was some falloff in perceived availability in all age strata through age 30—particularly among those ages 23 through 30—and an increasing convergence. Since 2000, there does not appear to have been any substantial systematic change.
- **Crack** availability peaked in 1988–1989 for all age groups (it was first assessed in 1987) and declined through 1992, with little further change until 1995. Since 1995, crack availability has declined some in all of the lower five age strata. Data on 40- and 45-year-olds is available for a shorter interval, but also shows declines from peak levels.
- The trends in **LSD** availability among young adults have some parallels to those for 12th graders. Among 12th graders, there was a drop of about 10 percentage points in the mid-1970s, and a later drop from 1980 to 1986. The latter drop, at least, was paralleled in the data from 19- to 22-year-olds. After 1986, LSD availability increased considerably in all age bands, reaching its peak levels (the highest we have recorded since these measures were introduced) by 1995, with a considerable age-related difference developing (with availability lower in the older age groups). Since 1995, availability has fallen substantially in all age bands (with the exception of the 45-year olds, for whom there is limited trend data available), but particularly in the youngest two age strata, narrowing the differences among the age groups. Indeed, the drop-off in availability of LSD to 12th graders and 19- to 22-year-olds was quite sharp in 2002, possibly contributing to the steep decline in use that year. Availability among 50-year-olds is the lowest. In 2008, between 19% and 29% of each age group say they could get LSD fairly easily, which contrasts quite dramatically with the mid-1990s, when over 50% of those in the younger age strata said they could get it.

- In the early 1980s, there was a fair decline among all age groups in the availability of **hallucinogens other than LSD**; there was little additional change through 1992. From 1992 to 1995, the three youngest age groups all showed an increase in availability, with 12th graders showing the largest increase. From 1996 to 2000, availability was fairly steady. (This question is not asked of respondents over age 30.) All age groups showed substantial increases in 2001, undoubtedly due to the changed question wording in which, among other things, “shrooms” was added to the examples of hallucinogens. Since 2001 the general pattern has been one of stability, but with the levels of availability more differentiated by age. Generally, the lower the age stratum, the higher the reported availability. So, it appears that the inclusion of “shrooms,” or psilocybin mushrooms, introduced a greater variability with age in the availability of hallucinogens other than LSD, taken as a class.
- The availability of **ecstasy (MDMA)** rose very substantially in all of these age groups during the 1990s and early 2000s. (The questions were first introduced in 1989 and 1990, and are not asked of those over age 30.) Among 12th graders, reported availability nearly tripled, from 22% in 1989 to 62% in 2001—the peak year for 12th graders. All four age groupings showed sharp increases in 2000 and 2001, with the older age groups continuing to increase through 2002—their peak year. The three youngest strata have shown considerable declines since 2002, and reported availability of ecstasy now varies little by age, ranging between 35% and 42% among all the age strata covered in 2008.
- **Heroin** availability varied within a fairly narrow range from 1980 to 1986, but then showed a fair-sized increase among both 12th graders and young adults through 1990. It then rose further among 12th graders and 19- to 22-year-olds through 1995 before declining a bit. In the older two age groups, heroin availability remained fairly flat from 1990 to 1995, increased some through 1999, and declined thereafter. It is clear that heroin was much more available to all of these age groups in the 1990s than it was in the 1980s. This increase in the availability—and purity—of heroin most likely led to the emergence of noninjection forms of heroin administration during this period. All age groups have shown some decline in heroin availability since 1997 or 1998, during which interval there has been only modest variability in heroin availability across the 18-to-30 age range. (The question is not asked of respondents over age 30.)
- The availability of **narcotics other than heroin** slowly rose among all age groups between 1980 and 1989, followed by a period of considerable stability from 1989 through 1994. After 1994, availability increased modestly, accompanied by steadily rising use. But in recent years, there has been some modest falloff in availability among all age strata except the 27- to 30-year-olds. For the most part, there has not been a consistent difference by age in the availability of narcotics other than heroin. (Respondents over age 30 are not asked this question.)
- The reported availability of **amphetamines** peaked in 1982 for both 12th graders and 19- to 22-year-olds, after which it fell until 1991, by 14 percentage points among 12th graders and 15 percentage points among 19- to 22-year-olds. Between 1984 (when data were first available) and 2005, there was a decline of 14 percentage points among 23- to

26-year-olds as well. For 27- to 30-year-olds, reported availability decreased by nine percentage points between 1988 (the first measurement point) and 2005. There were decreases among 35-year-olds as well for awhile in the 2000s, but some reversal in recent years. In general, the age groups above age 30 have reported somewhat lower availability than the younger strata, but not dramatically lower. These differential rates of availability across the age groups emerged after 1992, when prevalence of use began to rise among 12th graders.

- By way of contrast, *crystal methamphetamine* or “ice” exhibited an increase in availability in the 1990s, rising for all four age strata from 1991 to 1998 or 1999, before stabilizing with similar rates of availability from ages 18 to 30. (This question was not asked of those over 30.)
- *Sedatives (barbiturates)* exhibited a long-term decline in availability over more than two decades, from about 1981 or 1982 through 2003 in the two younger groups—a 20-percentage-point drop among 12th graders and a 23-percentage-point drop among 19- to 22-year-olds. All groups showed an increase in 2004—no doubt due primarily to a change in the question wording—and little further systematic change since, except for a gradual decline among 12th graders.
- *Tranquilizer* availability has declined long term by more than two thirds among 12th graders, from 72% in 1975 to 22% in 2008. Since 1980, when data were first collected for 19- to 22-year-olds, tranquilizer availability declined sharply by more than half (from 67% in 1980 to 23% in 2008), such that previous differences in availability between these two groups were eliminated by 1992. The older age groups also showed a considerable decline in the availability of tranquilizers through 2008 among 23- to 26-year-olds and through 2006 among 27- to 30-year-olds. For the most part, trend lines for the different age groups have been quite parallel, as was true for sedatives (barbiturates). Indeed, this class of drugs has shown the most consistent pattern of change in perceived availability since MTF began.
- Data on *steroid* availability were first gathered in 1990. There appears to have been some decline in availability in all age groups since about 2000, including a substantial decline in the youngest three age strata in 2008. (This question was not asked of those over 30.)

TABLE 7-1
Trends in Proportions of Friends Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

		Percentage saying friends disapprove ^a																		
<i>Q. How do you think your close friends feel (or would feel) about you. . .</i>		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	cont. →
Trying marijuana once or twice	18	42.6	46.4	50.3	52.0	54.1	54.7	56.7	58.0	62.9	63.7	70.3	69.7	73.1	66.6	62.7	58.1	55.8		
	19–22	41.0	40.6	46.9	47.1	51.6	54.5	55.2	54.7	58.7	63.0	63.6	64.7	64.7	63.4	63.7	58.5	64.3		
	23–26	—	—	—	—	47.7	47.0	49.1	53.9	58.2	62.6	61.3	64.5	65.6	65.5	63.2	63.8	61.2		
	27–30	—	—	—	—	—	—	—	—	58.6	58.7	61.4	64.6	63.5	64.4	66.3	66.1	65.8		
Smoking marijuana occasionally	18	50.6	55.9	57.4	59.9	62.9	64.2	64.4	67.0	72.1	71.1	76.4	75.8	79.2	73.8	69.1	65.4	63.1		
	19–22	50.9	49.2	54.0	57.9	59.4	64.6	64.4	65.1	69.8	71.5	74.1	73.9	74.3	73.1	73.0	66.6	71.3		
	23–26	—	—	—	—	54.3	56.4	57.1	63.1	68.1	73.2	71.8	72.5	75.3	73.5	72.2	70.7	70.8		
	27–30	—	—	—	—	—	—	—	—	67.8	69.4	71.9	73.7	76.0	75.1	76.4	73.8	75.6		
Smoking marijuana regularly	18	72.0	75.0	74.7	77.6	79.2	81.0	82.3	82.9	85.5	84.9	86.7	85.9	88.0	83.5	80.6	78.9	76.1		
	19–22	70.3	75.2	75.7	79.5	80.0	82.7	83.5	84.8	86.9	87.5	89.1	88.4	89.1	87.6	85.9	83.9	84.5		
	23–26	—	—	—	—	77.8	78.4	80.9	82.0	85.8	89.2	88.1	87.9	90.3	89.1	88.8	84.9	89.5		
	27–30	—	—	—	—	—	—	—	—	85.4	86.0	88.4	89.2	88.7	88.2	88.9	89.7	89.6		
Trying LSD once or twice	18	87.4	86.5	87.8	87.8	87.6	88.6	89.0	87.9	89.5	88.4	87.9	87.9	87.3	83.5	83.4	82.6	80.8		
	19–22	87.4	90.5	88.0	89.3	89.3	91.1	90.5	91.8	90.8	91.2	89.1	89.9	87.2	87.7	87.9	84.6	85.3		
	23–26	—	—	—	—	87.4	90.8	88.6	89.8	88.9	91.0	90.1	92.4	88.9	87.7	86.3	85.3	88.5		
	27–30	—	—	—	—	—	—	—	—	88.8	89.7	92.3	91.1	91.4	89.9	91.2	89.7	89.3		
Trying cocaine once or twice	18	—	—	—	—	—	—	79.6	83.9	88.1	88.9	90.5	91.8	92.2	91.1	91.4	91.1	89.2		
	19–22	—	—	—	—	—	—	76.4	—	84.8	87.7	89.2	92.3	91.9	92.4	94.7	91.7	91.5		
	23–26	—	—	—	—	—	—	70.8	—	81.4	84.5	84.1	86.7	87.4	87.7	87.9	90.4	90.0		
	27–30	—	—	—	—	—	—	—	—	81.8	81.1	83.7	83.5	84.4	86.1	87.8	87.5	88.7		
Taking cocaine occasionally	18	—	—	—	—	—	—	87.3	89.7	92.1	92.1	94.2	94.7	94.4	93.7	93.9	93.8	92.5		
	19–22	—	—	—	—	—	—	84.9	—	91.0	93.8	94.2	95.6	95.9	95.6	97.5	95.6	95.7		
	23–26	—	—	—	—	—	—	81.7	—	88.2	91.5	92.4	94.1	93.8	93.5	94.3	94.6	95.4		
	27–30	—	—	—	—	—	—	—	—	87.7	89.5	90.0	92.2	92.3	92.8	94.6	94.1	94.6		
Trying an amphetamine once or twice	18	78.9	74.4	75.7	76.8	77.0	77.0	79.4	80.0	82.3	84.1	84.2	85.3	85.7	83.2	84.5	81.9	80.6		
	19–22	75.8	76.7	75.3	74.3	77.0	79.7	81.5	81.3	83.0	83.5	84.5	86.5	83.8	85.0	87.2	83.1	86.0		
	23–26	—	—	—	—	78.4	79.1	76.7	81.7	83.0	85.6	84.3	85.0	83.6	84.2	84.7	87.6	86.5		
	27–30	—	—	—	—	—	—	—	—	82.7	84.1	84.9	84.6	84.7	84.1	85.9	85.5	85.6		
Taking one or two drinks nearly every day	18	70.5	69.5	71.9	71.7	73.6	75.4	75.9	71.8	74.9	76.4	79.0	76.6	77.9	76.8	75.8	72.6	72.9		
	19–22	71.9	72.1	68.6	73.5	71.6	72.2	72.7	70.2	73.9	77.1	73.3	73.7	74.0	71.2	73.0	68.3	68.9		
	23–26	—	—	—	—	63.6	66.8	67.7	68.3	69.2	70.8	72.7	72.5	72.1	67.6	71.5	68.2	72.8		
	27–30	—	—	—	—	—	—	—	—	71.0	68.0	70.4	71.9	68.8	73.2	70.9	68.8	65.7		
Taking four or five drinks nearly every day	18	87.9	86.4	86.6	86.0	86.1	88.2	87.4	85.6	87.1	87.2	88.2	86.4	87.4	87.2	85.2	84.1	82.6		
	19–22	93.7	91.7	89.9	91.9	91.7	92.5	91.5	90.8	90.4	92.5	89.9	91.7	92.6	89.6	90.1	88.8	88.1		
	23–26	—	—	—	—	90.8	90.2	92.5	92.8	93.7	92.1	92.1	92.4	91.1	93.1	92.1	92.2	92.6		
	27–30	—	—	—	—	—	—	—	—	92.8	92.0	92.9	92.7	92.7	93.9	94.0	92.9	91.9		
Having five or more drinks once or twice each weekend	18	50.6	50.3	51.2	50.6	51.3	55.9	54.9	52.4	54.0	56.4	59.0	58.1	60.8	58.5	59.1	58.0	57.8		
	19–22	53.5	51.7	51.7	53.3	50.8	53.3	47.0	49.4	50.5	56.8	53.1	51.4	53.6	51.9	54.4	55.5	52.1		
	23–26	—	—	—	—	53.8	57.3	61.0	57.2	58.8	57.5	55.1	56.8	58.4	57.6	61.4	58.9	58.4		
	27–30	—	—	—	—	—	—	—	—	61.9	65.1	66.3	68.2	66.2	66.7	63.7	64.6	61.6		
Smoking one or more packs of cigarettes per day	18	74.4	73.8	70.3	72.2	73.9	73.7	76.2	74.2	76.4	74.4	75.3	74.0	76.2	71.8	72.4	69.2	69.3		
	19–22	75.6	75.1	75.4	78.5	76.2	79.7	77.7	78.6	80.2	78.4	77.5	78.3	79.0	76.0	73.8	70.9	73.9		
	23–26	—	—	—	—	73.9	77.3	80.3	80.5	79.5	80.5	78.5	83.3	82.3	77.4	80.1	78.8	78.3		
	27–30	—	—	—	—	—	—	—	—	81.2	80.9	82.9	84.5	83.1	86.8	82.5	83.4	81.9		
<i>Approximate Weighted N =</i>	18	2,766	3,120	3,024	2,722	2,721	2,688	2,639	2,815	2,778	2,400	2,184	2,160	2,229	2,220	2,149	2,177	2,030		
	19–22	569	597	580	577	582	556	577	595	584	555	559	537	520	510	470	480	471		
	23–26	—	—	—	—	510	548	549	540	510	513	516	507	481	463	445	436	—		
	27–30	—	—	—	—	—	—	—	—	483	518	479	480	451	451	457	439	439		

(Table continued on next page.)

TABLE 7-1 (cont.)
Trends in Proportions of Friends Disapproving of Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. How do you think your close friends feel (or would feel) about you. . .	Age Group	Percentage saying friends disapprove ^a											2007–2008 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007		2008
Trying marijuana once or twice	18	53.0	53.8	55.1	58.1	57.6	54.1	58.4	59.5	60.9	62.3	60.4	60.8	+0.4
	19–22	58.4	57.0	56.5	56.0	54.2	53.4	56.5	61.0	57.9	60.5	58.4	62.4	+4.0
	23–26	59.3	66.5	62.6	64.6	55.2	53.8	51.4	57.7	55.9	60.7	55.8	62.1	+6.3
	27–30	65.0	65.4	61.8	63.9	64.9	67.1	61.9	67.2	61.2	64.1	58.2	57.1	-1.1
Smoking marijuana occasionally	18	59.9	60.4	61.6	63.9	64.3	60.3	64.2	65.0	67.6	68.1	65.8	66.3	+0.6
	19–22	65.1	65.1	64.6	61.8	61.0	62.6	63.3	70.1	67.2	68.8	70.6	67.5	-3.1
	23–26	68.5	73.6	70.2	70.9	63.9	64.5	61.6	63.5	65.5	71.3	63.8	70.1	+6.3
	27–30	72.4	74.9	74.5	75.0	74.2	72.9	71.4	76.9	70.4	74.9	66.4	67.0	+0.6
Smoking marijuana regularly	18	74.1	74.7	74.5	76.1	77.8	75.3	77.0	77.3	79.5	79.8	78.3	78.0	-0.4
	19–22	83.3	81.1	78.2	78.5	80.0	80.5	79.1	84.4	82.2	84.1	83.7	81.4	-2.3
	23–26	85.6	87.1	86.8	86.9	83.7	82.8	80.0	79.2	82.7	83.7	81.9	87.1	+5.2
	27–30	87.8	90.8	89.2	91.6	90.1	87.9	87.2	88.0	87.7	88.2	84.3	84.5	+0.2
Trying LSD once or twice	18	79.3	81.7	83.2	84.7	85.5	84.9	87.5	87.3	88.4	89.5	88.4	86.3	-2.1
	19–22	83.6	81.7	82.0	82.1	85.2	86.9	86.9	88.6	90.5	90.4	90.0	90.0	0.0
	23–26	85.4	87.6	84.5	85.3	82.8	83.6	79.3	82.4	85.6	89.3	90.4	88.4	-2.0
	27–30	88.5	88.7	88.4	85.6	87.4	86.3	87.1	87.7	86.9	88.5	83.5	85.3	+1.9
Trying cocaine once or twice	18	87.3	88.8	88.7	90.2	89.3	89.1	91.2	87.9	89.0	88.7	89.6	88.7	-0.9
	19–22	91.8	90.0	91.2	89.4	89.1	91.7	90.6	90.3	90.3	91.2	93.3	90.2	-3.1
	23–26	91.1	92.0	89.6	90.5	88.0	88.5	83.6	84.2	84.6	88.7	91.7	91.0	-0.7
	27–30	89.4	89.3	90.5	90.4	89.3	88.8	89.9	91.8	89.5	92.0	86.4	88.0	+1.6
Taking cocaine occasionally	18	90.8	92.2	91.8	92.8	92.2	92.2	93.0	91.0	92.3	92.4	93.1	92.0	-1.1
	19–22	96.6	93.1	95.7	94.7	94.5	95.6	95.1	96.0	95.3	96.1	97.1	95.5	-1.6
	23–26	95.1	95.2	95.2	96.7	94.7	93.2	91.2	90.1	93.0	94.9	95.9	96.6	+0.7
	27–30	94.2	96.1	95.4	95.9	94.2	94.0	95.1	96.3	94.5	95.4	93.2	94.3	+1.2
Trying an amphetamine once or twice	18	80.4	82.6	83.0	84.1	83.8	83.3	85.9	84.7	86.1	86.7	87.3	87.1	-0.2
	19–22	84.5	84.0	85.8	81.6	84.5	87.6	87.6	89.4	88.9	89.4	89.1	90.2	+1.1
	23–26	83.3	87.0	85.9	85.1	83.1	83.9	81.5	82.7	86.2	89.9	89.3	89.6	+0.3
	27–30	85.9	85.8	87.2	87.8	86.4	86.0	87.9	88.9	87.5	88.5	82.9	85.3	+2.4
Taking one or two drinks nearly every day	18	71.5	72.3	71.7	71.6	73.4	71.6	74.7	72.8	74.0	73.2	74.5	75.2	+0.7
	19–22	73.5	67.3	68.6	66.6	64.9	68.5	64.4	72.4	68.3	68.7	68.4	69.5	+1.2
	23–26	68.1	66.9	66.1	65.4	64.4	61.6	62.1	61.8	62.3	66.1	62.5	63.4	+0.9
	27–30	67.3	66.7	64.3	67.3	67.1	64.0	64.5	65.0	62.8	64.9	59.4	58.9	-0.5
Taking four or five drinks nearly every day	18	82.5	82.8	82.2	82.8	84.4	80.1	83.1	82.9	82.7	83.3	84.8	84.7	-0.1
	19–22	90.0	85.9	87.9	86.6	84.6	87.7	86.8	89.8	86.8	89.0	90.7	88.8	-1.8
	23–26	90.7	93.7	89.9	92.5	91.1	88.1	89.3	87.8	89.1	90.8	87.8	93.8	+6.0 ss
	27–30	93.8	92.1	95.3	92.4	91.2	92.7	92.6	92.5	93.4	92.3	91.3	89.0	-2.3
Having five or more drinks once or twice each weekend	18	56.4	55.5	57.6	57.7	57.8	55.6	60.3	59.4	59.9	60.6	60.0	62.1	+2.1
	19–22	56.4	52.8	51.8	45.2	47.4	50.4	47.9	52.4	53.2	54.8	54.4	55.2	+0.8
	23–26	55.6	60.0	54.5	56.6	56.9	52.9	49.5	49.5	51.9	56.0	51.3	55.3	+4.0
	27–30	64.0	63.0	57.7	65.8	58.8	63.3	59.6	64.6	56.9	62.7	56.3	57.3	+1.0
Smoking one or more packs of cigarettes per day	18	68.5	69.0	71.2	72.6	74.5	75.7	79.2	78.6	81.1	81.2	81.4	82.5	+1.1
	19–22	76.5	69.2	73.9	71.1	74.3	77.3	78.3	82.1	82.7	84.8	87.0	85.5	-1.6
	23–26	75.8	76.5	78.0	79.9	77.0	75.4	78.3	77.6	77.4	84.4	82.6	88.2	+5.5 s
	27–30	80.5	81.9	82.6	84.0	83.6	86.1	84.0	84.6	82.2	84.1	81.3	83.9	+2.5
<i>Approximate</i>	18	2,095	2,037	1,945	1,775	1,862	1,820	2,133	2,208	2,183	2,183	2,161	2,090	
<i>Weighted N =</i>	19–22	466	436	430	379	402	361	399	427	395	395	361	370	
	23–26	419	425	394	398	378	366	363	377	361	344	349	336	
	27–30	422	440	397	394	374	364	346	408	362	327	330	318	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

“—” indicates data not available.

^aAnswer alternatives were: (1) Don't disapprove, (2) Disapprove, and (3) Strongly disapprove. Percentages are shown for categories (2) and (3) combined.

TABLE 7-2
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19-22, 23-26, 27-30, 35, 40, 45, and 50

Q. How many of your friends would you estimate...	Age Group	Percentage saying friends use ^a																	cont. →
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	
Take any illicit drug^b % saying any	18	87.5	85.4	86.3	82.6	81.0	82.4	82.2	81.7	79.1	76.9	71.0	69.1	67.3	71.0	78.3	78.6	80.6	
	19-22	90.2	88.0	86.8	85.0	82.3	82.9	80.5	76.7	77.2	78.4	72.7	71.5	66.8	71.7	71.6	71.6	76.2	
	23-26	—	—	—	—	83.6	82.7	80.3	80.9	74.4	73.8	65.8	63.0	67.3	64.6	66.7	65.3	64.6	
	27-30	—	—	—	—	—	—	—	—	74.8	72.9	69.6	67.1	61.5	60.2	57.1	58.5	59.1	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	38.1	37.4	39.7	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	% saying most or all	18	32.5	29.8	26.5	23.8	20.9	22.7	21.5	18.6	15.8	15.7	11.6	11.7	12.0	15.5	20.3	21.7	23.8
		19-22	34.9	32.8	28.1	22.4	21.9	18.2	16.2	14.0	13.5	10.9	10.5	8.8	9.0	10.4	14.9	13.1	17.3
		23-26	—	—	—	—	19.6	15.4	16.2	11.7	9.5	9.7	9.5	7.4	6.2	6.4	8.7	7.6	8.8
		27-30	—	—	—	—	—	—	—	—	8.6	6.4	5.9	2.9	5.8	5.0	5.6	6.1	3.6
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.1	1.9	2.0
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		Take any illicit drug^b other than marijuana % saying any	18	62.4	63.3	64.7	61.2	61.3	61.8	63.3	62.4	56.5	56.2	50.1	46.3	47.1	48.7	53.7	53.7
19-22			67.9	67.8	66.7	65.2	60.8	62.1	61.0	57.3	53.5	60.8	53.4	51.5	45.3	51.4	46.3	46.4	46.5
23-26			—	—	—	—	63.7	64.0	59.0	61.1	55.1	54.2	47.8	41.8	46.1	42.3	39.4	40.3	32.8
27-30			—	—	—	—	—	—	—	—	55.9	55.0	49.7	47.2	37.7	38.5	33.9	37.7	36.4
35			—	—	—	—	—	—	—	—	—	—	—	—	—	—	21.4	21.6	22.1
40			—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
45			—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
50			—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all			18	11.1	11.9	10.9	11.0	10.3	10.4	10.3	9.2	6.9	7.7	5.1	4.6	5.3	7.1	7.1	7.7
	19-22		9.8	12.9	11.8	9.8	9.3	8.6	7.6	5.0	5.3	4.0	3.2	2.6	3.3	4.0	4.4	3.5	6.2
	23-26		—	—	—	—	10.6	6.6	8.6	5.2	3.9	4.2	3.4	1.6	1.8	2.8	2.5	1.9	1.9
	27-30		—	—	—	—	—	—	—	—	4.6	3.0	2.8	1.0	1.4	1.5	1.5	1.5	0.9
	35		—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	0.5	0.7
	40		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	Smoke marijuana % saying any		18	86.4	83.0	84.4	80.3	77.7	79.5	79.2	78.4	75.3	72.5	68.3	65.8	63.1	67.4	75.6	76.1
		19-22	88.8	86.4	85.2	83.8	81.6	81.1	78.5	75.3	75.1	73.8	67.6	68.0	63.5	67.6	67.4	68.8	74.9
		23-26	—	—	—	—	82.0	80.8	77.7	79.4	71.6	69.8	61.8	59.6	61.3	61.2	62.6	63.2	62.6
		27-30	—	—	—	—	—	—	—	—	71.8	68.2	65.1	62.6	58.0	57.4	52.3	55.7	55.1
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	36.9	36.3	36.3
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		% saying most or all	18	31.3	27.7	23.8	21.7	18.3	19.8	18.2	15.8	13.6	13.4	10.1	10.0	10.3	13.9	18.9	20.7
19-22			34.1	30.6	25.6	20.6	19.4	16.0	13.3	12.5	12.2	9.0	9.2	8.3	8.2	8.5	13.0	12.5	16.3
23-26			—	—	—	—	17.0	14.3	13.7	10.4	7.8	8.6	8.3	6.9	5.6	5.6	7.5	6.6	8.2
27-30			—	—	—	—	—	—	—	—	6.8	4.4	4.0	2.8	5.1	5.2	5.0	5.6	3.5
35			—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.0	2.5	2.9
40			—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
45			—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
50			—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How many of your friends would you estimate . . .	Age Group	Percentage saying friends use ^a											2007–2008 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007		2008
Take any illicit drug^b														
% saying any	18	83.4	84.6	82.0	82.0	82.8	81.8	80.7	81.2	79.8	78.8	77.7	80.1	+2.4
	19–22	77.2	79.8	77.3	83.1	81.1	78.3	79.4	78.1	78.6	74.7	79.8	77.4	-2.4
	23–26	67.0	67.6	67.9	67.8	66.9	73.4	70.8	70.8	74.2	72.2	71.3	72.2	+0.9
	27–30	60.9	58.3	59.6	55.6	57.2	61.8	58.6	63.1	63.7	62.3	62.7	66.7	+4.0
	35	39.2	38.4	36.3	37.7	39.1	40.9	37.5	37.9	40.0	40.4	42.1	44.9	+2.8
	40	—	39.2	38.2	38.0	38.4	36.2	36.5	34.6	36.2	35.4	34.6	35.9	+1.3
	45	—	—	—	—	—	—	37.8	38.3	34.3	36.7	38.5	35.9	-2.6
	50	—	—	—	—	—	—	—	—	—	—	—	39.3	—
% saying most or all	18	23.7	25.9	25.5	24.5	25.2	23.1	23.5	23.0	20.2	20.9	21.7	21.3	-0.4
	19–22	16.2	16.8	20.6	18.9	20.3	20.2	17.3	14.7	15.8	16.8	14.5	13.7	-0.8
	23–26	10.5	9.6	8.4	9.7	10.4	10.3	10.3	11.7	9.7	11.1	8.1	8.9	+0.8
	27–30	4.5	5.3	5.7	5.3	7.1	6.9	6.9	3.9	4.7	5.4	6.5	6.3	-0.2
	35	3.0	3.1	2.8	3.1	3.2	2.9	3.2	2.8	2.5	2.1	2.2	2.2	0.0
	40	—	2.3	2.0	2.0	1.6	2.2	1.6	1.6	2.1	2.5	2.0	1.3	-0.7
	45	—	—	—	—	—	—	2.2	1.5	1.4	1.7	1.3	1.3	0.0
	50	—	—	—	—	—	—	—	—	—	—	—	1.4	—
Take any illicit drug^b other than marijuana														
% saying any	18	55.1	55.6	51.2	52.5	55.0	54.3	50.0	51.4	51.3	51.0	50.0	49.3	-0.7
	19–22	49.7	53.3	54.8	56.1	60.0	57.2	50.8	53.4	54.9	49.5	52.5	46.4	-6.1
	23–26	35.1	35.4	41.1	42.5	42.6	49.4	42.3	47.1	46.6	45.6	42.6	45.9	+3.2
	27–30	33.9	34.1	35.2	31.7	33.5	36.0	34.7	35.8	33.1	36.2	34.2	36.4	+2.2
	35	19.2	19.3	19.0	17.9	18.7	20.4	18.5	20.2	18.5	18.1	20.7	23.7	+3.0
	40	—	20.9	21.0	21.9	21.4	21.0	20.2	18.5	21.0	20.3	20.3	19.8	-0.5
	45	—	—	—	—	—	—	23.4	25.1	20.8	22.7	25.0	21.2	-3.8 s
	50	—	—	—	—	—	—	—	—	—	—	—	24.5	—
% saying most or all	18	7.0	8.9	7.4	7.4	7.0	6.1	6.7	7.3	6.7	5.3	6.5	5.3	-1.2
	19–22	4.1	4.3	5.1	7.7	8.0	5.7	5.1	3.5	4.8	4.2	3.9	3.4	-0.4
	23–26	2.6	2.8	2.2	3.8	3.7	2.8	3.7	3.1	3.2	3.2	1.0	1.5	+0.5
	27–30	1.2	0.9	1.3	1.5	2.6	2.3	0.7	0.8	0.9	1.4	2.2	2.5	+0.3
	35	0.5	0.7	0.9	1.0	0.9	0.6	0.6	0.4	0.5	0.5	0.6	0.5	-0.1
	40	—	0.4	0.8	0.7	0.5	0.3	0.3	0.2	0.5	0.7	0.6	0.7	0.0
	45	—	—	—	—	—	—	0.7	0.7	0.4	0.9	0.5	0.5	+0.1
	50	—	—	—	—	—	—	—	—	—	—	—	0.5	—
Smoke marijuana														
% saying any	18	81.4	83.2	80.7	80.5	81.2	79.4	78.9	79.5	77.4	76.4	74.8	78.2	+3.4 s
	19–22	74.7	77.2	73.9	81.2	78.4	77.2	76.5	75.6	75.8	72.0	76.6	74.7	-1.9
	23–26	63.5	65.0	64.4	64.8	64.5	68.8	67.7	68.4	70.7	67.6	69.0	67.7	-1.3
	27–30	58.3	55.5	57.0	51.7	56.5	59.0	55.8	60.4	60.8	61.0	60.2	64.2	+4.0
	35	35.0	34.6	33.3	34.9	35.6	37.4	32.9	34.7	37.2	37.3	38.6	42.1	+3.4
	40	—	34.6	32.5	32.3	31.8	31.4	30.7	29.9	30.4	29.4	29.2	29.6	+0.4
	45	—	—	—	—	—	—	31.1	29.4	26.3	28.4	30.0	28.6	-1.4
	50	—	—	—	—	—	—	—	—	—	—	—	30.1	—
% saying most or all	18	22.5	23.8	24.2	23.2	24.0	21.4	21.7	21.1	17.9	19.6	19.2	19.9	+0.7
	19–22	16.2	16.4	19.4	16.6	18.5	18.6	16.0	15.0	13.4	15.7	13.4	11.5	-1.8
	23–26	9.8	9.0	8.5	8.2	9.0	8.7	9.3	9.8	8.0	10.1	7.9	8.5	+0.6
	27–30	3.9	4.8	5.5	4.9	6.3	6.2	6.7	3.5	4.3	5.0	6.6	5.0	-1.5
	35	2.9	2.8	2.6	2.8	2.6	2.7	3.1	2.7	2.3	2.0	2.1	1.9	-0.2
	40	—	2.1	1.4	1.9	1.2	2.0	1.4	1.6	1.8	2.1	1.6	0.9	-0.7
	45	—	—	—	—	—	—	1.9	0.9	1.3	1.1	1.0	1.0	0.0
	50	—	—	—	—	—	—	—	—	—	—	—	1.2	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19-22, 23-26, 27-30, 35, 40, 45, and 50

		Percentage saying friends use ^a																	
Q. How many of your friends would you estimate. . .		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Use inhalants																			
% saying any		18	17.8	16.5	18.4	16.1	19.3	21.2	22.4	24.7	20.8	22.1	20.0	19.2	22.2	23.7	26.5	27.5	27.2
		19-22	11.9	13.2	13.8	12.3	11.7	9.6	10.9	12.7	10.9	11.7	13.0	12.2	12.6	13.8	14.0	14.2	16.2
		23-26	—	—	—	—	7.7	6.7	7.2	6.1	6.2	5.9	6.1	4.4	5.1	6.3	7.0	9.3	5.6
		27-30	—	—	—	—	—	—	—	—	4.6	3.5	2.9	2.5	3.3	2.9	3.5	4.0	4.1
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	1.2	0.9	1.3	1.1	1.1	1.5	2.0	1.9	1.2	1.9	1.0	0.7	1.8	1.8	2.0	2.0	2.4
		19-22	0.5	0.4	0.7	0.3	0.5	0.6	0.7	0.7	0.7	0.4	0.6	0.2	0.8	0.7	0.7	0.6	1.1
		23-26	—	—	—	—	0.6	0.2	0.6	0.1	0.2	0.4	0.4	0.1	*	0.1	0.2	0.7	0.5
		27-30	—	—	—	—	—	—	—	—	0.3	*	0.2	0.2	*	0.2	*	*	*
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Use nitrites																			
% saying any		18	19.0	17.4	17.5	14.5	15.0	15.6	18.0	18.3	13.6	13.3	10.4	8.9	9.0	10.7	10.0	10.7	11.2
		19-22	18.4	16.0	14.2	13.8	8.9	9.9	11.7	13.2	10.2	—	—	—	—	—	—	—	—
		23-26	—	—	—	—	10.8	7.8	8.0	7.9	5.2	—	—	—	—	—	—	—	—
		27-30	—	—	—	—	—	—	—	—	6.6	—	—	—	—	—	—	—	—
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	1.3	1.2	0.9	0.7	1.2	1.0	1.2	1.3	0.7	0.9	0.6	0.4	0.7	0.7	0.8	0.8	0.8
		19-22	0.3	0.4	0.9	0.6	0.6	0.6	0.4	0.4	0.2	—	—	—	—	—	—	—	—
		23-26	—	—	—	—	0.8	0.3	0.4	0.3	0.1	—	—	—	—	—	—	—	—
		27-30	—	—	—	—	—	—	—	—	0.5	—	—	—	—	—	—	—	—
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take LSD																			
% saying any		18	28.1	28.5	27.8	24.0	23.9	24.4	24.5	25.3	24.1	25.2	25.0	23.4	28.1	31.3	34.1	36.9	37.9
		19-22	30.9	25.9	26.5	22.6	21.6	18.8	18.7	18.2	19.0	20.1	20.1	22.0	22.2	28.8	23.8	26.9	28.6
		23-26	—	—	—	—	21.5	17.2	15.4	15.9	13.3	14.1	12.3	12.5	15.0	17.2	17.3	21.5	15.3
		27-30	—	—	—	—	—	—	—	—	10.4	7.7	9.1	8.6	10.9	8.7	8.1	12.0	11.6
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	1.8	2.2	2.4	1.4	2.0	1.5	1.8	1.6	1.5	2.4	1.9	1.7	2.4	3.8	4.2	4.8	5.0
		19-22	1.2	0.8	0.9	1.0	0.6	0.8	0.9	0.6	1.3	0.4	1.2	1.4	1.9	2.1	2.5	2.3	3.8
		23-26	—	—	—	—	0.8	0.5	1.0	0.2	0.6	0.5	0.6	0.2	0.4	0.7	1.1	0.7	0.7
		27-30	—	—	—	—	—	—	—	—	0.3	0.2	0.3	0.3	*	0.3	0.4	0.3	0.4
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How many of your friends would you estimate...	Age Group	Percentage saying friends use ^a											2007–2008 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007		2008
Use inhalants														
% saying any	18	27.4	25.9	21.6	23.5	22.2	21.0	17.5	17.9	18.1	18.9	17.9	18.0	+0.1
	19–22	13.7	16.2	16.3	13.7	13.7	10.4	10.0	9.5	11.1	11.0	9.6	7.4	-2.1
	23–26	7.5	6.2	7.9	6.9	7.5	7.4	7.9	6.2	5.8	5.2	3.7	6.1	+2.4
	27–30	3.6	3.8	4.2	3.6	6.0	4.5	3.2	2.6	3.2	3.3	2.8	2.7	-0.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	1.9	2.7	1.8	1.4	1.4	1.2	1.1	1.2	2.0	1.2	1.6	1.1	-0.6
	19–22	0.7	1.3	0.8	0.6	1.2	0.4	0.6	*	0.1	0.5	0.3	0.6	+0.3
	23–26	0.8	*	0.1	0.7	0.1	0.4	0.3	*	0.1	0.3	*	*	0.0
	27–30	*	*	*	*	0.3	0.3	*	*	*	*	*	0.3	+0.3
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Use nitrites														
% saying any	18	11.9	12.9	10.9	11.0	11.9	11.2	8.5	9.4	9.1	8.1	7.7	7.3	-0.4
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	0.7	1.0	0.7	1.0	0.6	0.8	1.0	1.2	1.0	0.5	0.7	0.5	-0.2
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Take LSD														
% saying any	18	36.5	36.8	32.2	31.9	32.2	28.6	21.9	23.5	19.5	18.7	18.3	20.9	+2.7
	19–22	24.7	29.4	28.2	27.8	28.4	24.0	15.4	15.9	13.9	14.2	15.1	12.5	-2.6
	23–26	18.2	15.2	18.1	19.3	16.8	15.8	16.1	14.4	12.0	11.7	11.2	9.2	-1.9
	27–30	12.3	12.6	13.4	11.8	12.5	13.1	11.4	8.9	6.6	9.1	7.6	8.8	+1.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	3.7	4.7	3.9	3.1	2.9	1.7	1.9	1.5	1.5	0.8	1.2	1.1	0.0
	19–22	1.4	2.5	1.8	2.1	2.7	1.6	0.8	0.3	0.3	0.2	0.8	0.2	-0.5
	23–26	0.6	1.0	1.5	0.9	0.3	0.4	0.8	0.4	0.2	0.3	*	0.1	+0.1
	27–30	0.4	0.1	0.6	0.4	0.4	0.3	0.1	*	0.3	0.4	0.4	*	-0.4
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19-22, 23-26, 27-30, 35, 40, 45, and 50

		Percentage saying friends use ^a																		
Q. How many of your friends would you estimate. . .		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	cont.
Take other hallucinogens^c																				
% saying any		18	28.2	26.3	25.6	22.1	21.3	22.0	22.3	21.7	17.8	18.1	15.9	15.1	17.0	19.3	21.4	23.8	26.4	
		19-22	33.4	25.5	25.1	21.0	20.2	16.6	15.8	15.0	16.1	13.9	15.3	14.2	12.0	15.0	13.8	14.9	17.2	
		23-26	—	—	—	—	20.0	16.7	13.2	13.2	11.7	9.6	8.7	8.5	9.8	9.4	10.3	11.7	10.4	
		27-30	—	—	—	—	—	—	—	—	10.6	7.4	7.1	6.8	7.9	7.1	6.6	7.9	7.5	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	2.2	2.1	1.9	1.6	1.9	1.4	1.3	1.2	0.9	1.4	1.0	0.8	1.0	1.7	2.2	2.2	2.3	
		19-22	1.5	0.9	1.1	1.2	0.7	1.0	0.7	0.6	0.9	0.2	0.5	0.8	0.7	0.9	1.6	1.5	1.0	
		23-26	—	—	—	—	0.8	0.3	0.5	0.3	0.2	0.3	0.8	0.1	0.4	0.7	0.6	0.8	0.1	
		27-30	—	—	—	—	—	—	—	—	0.2	0.1	0.3	0.2	*	0.2	0.3	0.1	0.2	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take PCP																				
% saying any		18	22.2	17.2	17.3	14.2	14.2	15.9	16.1	15.5	13.5	14.7	13.0	12.0	12.7	15.6	15.5	18.3	20.3	
		19-22	24.1	15.3	15.3	12.6	9.5	8.9	10.1	9.7	10.1	—	—	—	—	—	—	—	—	
		23-26	—	—	—	—	11.6	6.8	7.4	6.9	5.1	—	—	—	—	—	—	—	—	
		27-30	—	—	—	—	—	—	—	—	6.7	—	—	—	—	—	—	—	—	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	1.6	0.9	0.9	1.1	1.1	1.2	1.2	1.1	0.8	1.2	0.5	0.5	0.9	1.9	1.2	1.2	1.3	
		19-22	0.5	0.3	0.3	0.5	0.7	0.7	0.2	0.1	0.3	—	—	—	—	—	—	—	—	
		23-26	—	—	—	—	0.6	*	0.4	*	0.2	—	—	—	—	—	—	—	—	
		27-30	—	—	—	—	—	—	—	—	0.4	—	—	—	—	—	—	—	—	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Take ecstasy (MDMA)																				
% saying any		18	—	—	—	—	—	—	—	—	—	—	12.4	11.9	10.7	12.8	15.9	20.7	24.2	
		19-22	—	—	—	—	—	—	—	—	—	16.3	14.3	12.0	12.9	13.7	11.3	17.2	20.7	
		23-26	—	—	—	—	—	—	—	—	—	7.6	9.0	9.5	11.0	9.8	11.4	11.2	11.3	
		27-30	—	—	—	—	—	—	—	—	—	5.6	6.3	5.4	4.6	6.6	5.8	6.9	10.1	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all		18	—	—	—	—	—	—	—	—	—	—	2.2	1.7	2.1	1.2	1.7	2.8	3.0	
		19-22	—	—	—	—	—	—	—	—	—	0.4	0.7	0.2	0.7	0.7	0.5	0.5	0.8	
		23-26	—	—	—	—	—	—	—	—	—	0.5	0.2	0.1	0.1	0.5	0.1	0.4	0.1	
		27-30	—	—	—	—	—	—	—	—	—	0.5	0.3	*	0.1	0.3	0.2	0.5	0.1	
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How many of your friends would you estimate. . .	Age Group	Percentage saying friends use ^a											2007–2008 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007		2008
Take other hallucinogens^c														
% saying any	18	26.3	27.4	22.5	24.0	35.4	33.6	30.1	31.9	31.0	30.1	30.1	29.4	-0.8
	19–22	17.2	19.1	18.9	20.9	33.6	33.5	24.8	26.8	25.1	27.8	26.7	21.9	-4.8
	23–26	13.0	11.7	9.6	11.3	18.6	22.4	20.2	24.5	18.5	18.9	15.9	21.1	+5.1
	27–30	6.8	7.8	9.4	8.0	14.6	14.9	13.5	12.4	9.4	14.9	10.6	16.9	+6.3 s
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	2.6	3.1	2.4	2.4	2.9	2.3	2.4	2.6	2.2	1.7	1.7	1.8	+0.1
	19–22	1.1	1.7	0.8	2.0	2.3	2.2	1.5	1.1	0.6	0.9	1.0	1.3	+0.3
	23–26	0.8	0.7	0.8	0.3	0.6	0.7	1.0	0.8	0.7	1.0	0.4	0.4	0.0
	27–30	0.3	0.2	0.2	0.4	0.6	1.0	0.1	*	0.4	0.4	0.4	0.4	0.0
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Take PCP														
% saying any	18	19.7	20.2	16.8	17.5	19.1	17.2	13.6	11.8	10.1	10.6	9.4	9.4	0.0
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	1.4	1.6	1.5	1.7	1.3	1.0	1.5	1.1	1.0	0.5	0.8	0.5	-0.2
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Take ecstasy (MDMA)														
% saying any	18	27.7	24.5	26.7	37.3	41.9	38.0	34.2	28.9	23.1	23.0	23.6	24.7	+1.1
	19–22	21.4	26.0	30.7	42.4	43.3	43.4	31.3	27.6	28.3	25.2	21.6	19.3	-2.3
	23–26	15.1	13.7	15.2	25.9	29.4	36.8	27.0	31.2	25.3	23.4	16.5	20.8	+4.3
	27–30	7.4	8.5	12.4	13.1	17.8	20.6	19.4	20.6	15.6	22.6	15.9	17.8	+1.9
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	2.6	2.5	2.7	4.8	5.2	3.7	2.7	3.2	2.5	1.9	2.1	2.4	+0.3
	19–22	1.7	2.0	2.9	4.9	5.8	2.7	1.9	1.9	1.1	0.7	0.8	0.9	+0.1
	23–26	0.8	0.8	0.4	2.9	1.7	1.2	2.0	1.1	1.3	0.9	0.3	0.4	+0.1
	27–30	0.3	*	0.8	0.4	0.3	0.9	0.5	0.6	0.2	*	0.1	0.9	+0.7
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19-22, 23-26, 27-30, 35, 40, 45, and 50

Q. How many of your friends would you estimate...	Age Group	Percentage saying friends use ^a																
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Take cocaine																		
% saying any	18	41.6	40.1	40.7	37.6	38.9	43.8	45.6	43.7	37.7	37.4	31.7	26.8	26.3	24.5	26.1	24.8	28.1
	19-22	51.0	48.9	49.8	46.5	47.6	45.9	48.3	45.7	42.0	42.7	33.2	29.7	22.8	24.3	21.5	22.0	19.4
	23-26	—	—	—	—	52.4	53.2	51.6	50.7	47.1	40.8	34.8	29.0	28.8	27.1	22.3	24.4	18.1
	27-30	—	—	—	—	—	—	—	—	47.9	43.3	38.3	35.7	29.9	27.6	22.6	26.2	20.8
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	6.1	6.3	4.9	5.1	5.1	5.8	6.2	5.1	3.4	3.7	2.1	1.5	1.5	2.1	1.5	2.0	2.2
	19-22	7.0	8.6	7.8	6.1	6.3	6.1	6.1	3.3	3.5	2.1	1.2	1.1	1.0	0.5	1.5	0.9	1.0
	23-26	—	—	—	—	9.1	5.3	7.0	4.1	3.1	2.7	2.1	0.6	0.9	0.8	1.0	0.3	0.4
	27-30	—	—	—	—	—	—	—	—	3.8	2.0	2.3	0.9	1.2	0.8	0.8	0.4	0.4
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take crack																		
% saying any	18	—	—	—	—	—	—	—	27.4	25.4	26.1	19.2	17.6	17.8	17.9	20.0	19.2	21.6
	19-22	—	—	—	—	—	—	—	23.8	21.8	20.6	14.6	14.3	11.8	13.6	13.8	14.0	9.4
	23-26	—	—	—	—	—	—	—	26.4	22.4	19.8	14.4	10.8	10.8	8.8	8.8	11.1	8.2
	27-30	—	—	—	—	—	—	—	22.1	18.4	16.6	11.6	10.3	10.2	10.4	10.3	8.6	8.6
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4.5	5.1	4.4
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	—	—	—	—	—	—	—	2.2	1.1	2.1	0.6	0.6	0.7	0.9	1.0	1.1	0.9
	19-22	—	—	—	—	—	—	—	0.7	0.8	1.0	0.6	0.2	0.1	0.3	0.4	0.3	0.5
	23-26	—	—	—	—	—	—	—	0.8	0.9	0.8	0.5	0.1	0.1	0.5	0.2	*	0.3
	27-30	—	—	—	—	—	—	—	1.2	0.9	0.9	0.3	0.3	*	0.6	0.3	0.1	0.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.6	0.3	0.4
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take cocaine powder																		
% saying any	18	—	—	—	—	—	—	—	—	—	25.3	24.6	19.8	19.7	18.1	20.7	19.2	22.8
	19-22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	23-26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	27-30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	14.2	12.9	15.4
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	—	—	—	—	—	—	—	—	—	2.3	2.5	1.8	2.0	1.6	1.9	1.7	1.9
	19-22	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	23-26	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	27-30	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	0.3	0.6
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How many of your friends would you estimate. . .	Age Group	Percentage saying friends use ^a											2007–2008 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007		2008
Take cocaine														
% saying any	18	28.2	31.2	27.8	27.2	27.1	26.8	23.8	29.3	28.1	29.7	29.7	25.2	-4.5 ss
	19–22	22.2	26.8	25.7	24.8	27.4	28.2	25.5	26.2	27.2	26.6	29.4	21.8	-7.6 s
	23–26	19.7	18.7	20.1	20.3	19.4	23.7	21.9	27.4	25.6	24.6	23.1	23.1	0.0
	27–30	21.5	18.6	20.7	16.5	19.7	16.0	17.0	17.0	17.9	19.5	18.6	20.7	+2.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	2.0	3.2	2.9	2.0	1.7	1.7	2.4	2.3	2.3	1.9	2.1	1.2	-0.9
	19–22	0.8	1.5	1.1	1.0	1.8	1.0	1.4	0.7	2.1	1.0	1.1	1.3	+0.2
	23–26	1.1	0.9	0.5	0.8	1.6	1.0	1.6	1.0	1.5	1.4	0.8	0.6	-0.2
	27–30	0.6	0.1	0.4	0.4	0.5	0.6	0.3	0.3	*	*	1.4	0.9	-0.5
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Take crack														
% saying any	18	22.2	24.4	19.0	21.4	23.4	21.5	18.7	22.5	22.9	22.3	21.8	19.1	-2.8
	19–22	13.1	16.4	15.7	16.5	17.4	18.0	11.8	16.0	14.9	14.5	16.0	12.2	-3.8
	23–26	8.3	8.3	8.8	7.9	8.6	10.1	10.4	10.8	10.8	10.0	8.7	9.8	+1.1
	27–30	6.3	6.4	8.7	6.0	7.1	6.4	6.5	5.2	8.5	9.1	6.9	5.8	-1.1
	35	3.1	2.8	3.2	3.9	2.8	3.2	2.8	3.1	2.6	2.8	3.0	2.8	-0.1
	40	—	3.8	3.0	2.9	3.5	2.6	2.7	2.6	2.8	2.3	3.1	1.9	-1.2
	45	—	—	—	—	—	—	3.7	3.3	2.4	3.0	2.9	1.8	-1.1
	50	—	—	—	—	—	—	—	—	—	—	—	2.0	—
% saying most or all	18	1.1	1.7	1.5	1.4	0.8	0.8	1.4	1.6	1.6	1.0	1.3	1.1	-0.2
	19–22	0.3	0.9	0.9	0.5	0.3	0.2	0.4	0.1	1.0	0.8	0.3	0.4	+0.1
	23–26	0.5	0.4	*	0.5	0.3	*	0.3	0.5	0.2	0.7	0.1	0.3	+0.2
	27–30	0.2	0.1	*	*	*	0.3	0.1	*	0.1	*	0.3	0.6	+0.3
	35	*	0.1	0.3	0.5	0.2	0.3	0.3	0.1	0.2	0.2	0.2	*	-0.2
	40	—	*	0.2	0.2	0.1	*	*	*	0.1	0.2	0.1	*	0.0
	45	—	—	—	—	—	—	0.4	0.3	0.2	0.2	*	*	0.0
	50	—	—	—	—	—	—	—	—	—	—	—	0.2	—
Take cocaine powder														
% saying any	18	24.8	22.9	22.0	21.3	20.1	22.4	23.2	25.4	23.2	22.8	22.3	22.6	+0.3
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	11.1	10.4	10.0	10.3	9.4	9.4	8.2	9.2	8.3	8.4	9.1	11.4	+2.3
	40	—	10.8	8.9	8.8	8.8	8.5	7.6	7.6	8.9	7.3	6.7	6.2	-0.5
	45	—	—	—	—	—	—	8.3	8.0	7.0	7.4	8.0	6.7	-1.2
	50	—	—	—	—	—	—	—	—	—	—	—	6.0	—
% saying most or all	18	2.0	1.9	1.9	1.8	1.5	1.9	1.9	3.3	1.7	1.7	1.8	1.5	-0.3
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	0.4	0.4	0.6	0.7	0.4	0.4	0.4	0.2	0.2	0.2	0.4	0.2	-0.2
	40	—	0.2	0.2	*	0.2	0.1	0.1	0.1	0.2	0.5	0.2	0.2	0.0
	45	—	—	—	—	—	—	0.5	0.5	0.2	0.4	0.1	0.1	0.0
	50	—	—	—	—	—	—	—	—	—	—	—	0.2	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19-22, 23-26, 27-30, 35, 40, 45, and 50

		Percentage saying friends use ^a																	
Q. How many of your friends would you estimate. . .		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Take heroin																			
% saying any		18	13.0	12.5	13.2	12.0	13.0	14.5	15.3	13.9	12.4	14.0	11.4	11.4	13.2	13.3	14.3	14.5	15.6
		19-22	11.0	8.1	9.4	7.5	7.1	6.5	8.5	8.5	7.8	6.8	6.5	6.1	4.7	7.0	8.1	10.4	6.7
		23-26	—	—	—	—	6.1	4.4	4.3	6.5	3.6	5.2	4.2	3.6	3.8	4.5	4.9	5.8	4.0
		27-30	—	—	—	—	—	—	—	—	3.8	2.8	4.5	2.7	3.1	3.6	4.2	3.6	4.4
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	1.0	0.5	0.7	0.8	0.8	0.9	1.1	0.9	0.7	1.1	0.4	0.4	0.7	1.1	1.0	1.1	0.9
		19-22	0.3	0.5	0.1	0.2	0.4	0.6	0.2	0.3	0.2	0.2	0.3	0.2	0.1	0.2	0.4	0.4	0.4
		23-26	—	—	—	—	0.4	0.2	0.2	*	0.2	0.4	0.2	0.3	0.4	0.1	0.2	0.2	*
		27-30	—	—	—	—	—	—	—	—	0.2	0.1	0.2	0.2	*	0.2	0.3	*	*
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take other narcotics																			
% saying any		18	22.4	23.1	23.9	20.8	21.4	22.8	21.8	23.2	19.2	19.2	17.2	13.7	14.9	16.1	18.5	19.5	21.8
		19-22	22.8	20.4	21.9	17.9	17.4	16.9	14.6	15.4	14.1	15.0	12.9	14.1	10.8	13.2	10.5	15.9	13.4
		23-26	—	—	—	—	16.0	14.9	14.0	13.0	10.6	10.8	10.5	8.5	8.4	8.7	8.0	10.5	8.9
		27-30	—	—	—	—	—	—	—	—	12.1	8.6	9.1	9.3	7.5	8.2	8.0	7.7	9.5
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	1.7	1.5	1.4	1.4	1.6	1.4	1.8	1.4	1.2	1.4	0.9	0.5	1.1	1.2	1.0	1.6	1.5
		19-22	0.9	0.7	0.6	0.5	0.8	1.0	0.5	0.4	0.9	0.1	0.6	0.4	0.5	0.6	0.6	0.6	0.4
		23-26	—	—	—	—	0.4	0.3	0.7	*	0.3	0.2	0.2	*	*	*	0.3	0.2	*
		27-30	—	—	—	—	—	—	—	—	0.3	*	0.2	0.2	0.1	0.2	0.2	*	0.2
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take amphetamines																			
% saying any		18	43.9	48.8	50.6	46.1	45.1	43.3	41.8	39.5	33.4	33.5	28.7	24.3	24.3	27.5	28.1	30.3	32.2
		19-22	54.1	52.2	51.3	49.7	46.1	42.1	38.5	34.5	26.8	29.6	23.3	26.2	19.5	21.0	20.9	21.7	21.6
		23-26	—	—	—	—	45.6	40.1	33.5	32.1	28.4	23.1	20.6	17.1	15.1	16.8	16.2	18.2	12.5
		27-30	—	—	—	—	—	—	—	—	26.1	21.6	19.3	17.0	15.3	14.0	13.1	13.7	15.5
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all		18	4.8	6.4	5.4	5.1	4.5	3.4	3.4	2.6	1.9	2.6	1.9	1.3	1.3	2.0	1.8	2.0	2.8
		19-22	3.8	5.7	4.6	3.8	3.3	2.9	1.3	1.9	1.4	0.7	1.0	0.6	0.9	0.2	1.1	1.2	0.7
		23-26	—	—	—	—	1.9	1.8	1.7	1.2	0.3	0.6	0.7	0.8	0.4	1.5	0.9	0.5	0.2
		27-30	—	—	—	—	—	—	—	—	0.6	0.4	0.5	0.5	0.1	0.5	0.5	0.3	0.3
		35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How many of your friends would you estimate. . .	Age Group	Percentage saying friends use ^a											2007–2008 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007		2008
Take heroin														
% saying any	18	15.6	16.5	12.7	14.9	13.1	12.9	10.3	12.7	13.1	12.7	12.9	11.2	-1.7
	19–22	7.4	9.4	9.7	7.7	8.7	8.9	5.3	7.0	6.4	7.5	9.0	6.4	-2.6
	23–26	6.2	5.8	4.8	4.7	5.0	5.2	6.1	2.9	5.1	3.5	4.3	3.1	-1.3
	27–30	4.2	3.5	3.8	2.8	4.3	3.9	3.4	3.0	3.8	2.5	3.0	2.1	-1.0
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	0.8	1.3	1.0	1.1	0.9	0.7	0.9	0.9	1.1	0.8	1.4	0.7	-0.8
	19–22	0.2	0.5	0.1	0.3	0.6	*	0.3	*	0.3	0.4	0.3	0.6	+0.3
	23–26	0.7	*	*	0.3	*	0.1	*	*	0.3	0.3	*	*	0.0
	27–30	*	0.1	*	*	*	0.3	*	*	*	*	*	*	0.0
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Take other narcotics														
% saying any	18	22.2	24.8	22.9	23.1	24.0	27.5	21.6	24.6	21.4	23.0	20.7	20.6	-0.1
	19–22	13.2	15.2	19.8	23.2	23.0	21.8	21.9	22.6	19.9	17.6	23.7	16.8	-6.9 s
	23–26	9.9	9.4	10.4	11.2	13.5	14.6	18.4	16.8	18.3	17.6	14.2	16.0	+1.9
	27–30	7.9	8.3	7.2	8.4	11.2	11.8	11.0	12.0	12.5	13.1	10.6	14.3	+3.7
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	1.4	2.9	1.8	2.0	2.0	2.1	2.4	2.4	1.9	1.8	2.6	1.3	-1.3 s
	19–22	0.4	0.8	0.4	1.2	1.8	1.3	1.0	0.5	1.0	0.9	0.3	1.3	+1.0
	23–26	0.6	0.3	*	0.4	0.5	0.6	0.6	0.5	0.8	0.5	0.3	0.1	-0.2
	27–30	*	*	0.2	*	*	0.3	0.1	*	*	0.6	*	0.6	+0.6
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Take amphetamines														
% saying any	18	32.7	33.8	30.8	32.9	33.2	34.4	28.1	31.4	28.8	29.0	27.4	27.3	-0.1
	19–22	21.1	24.4	25.5	28.4	28.0	28.6	24.0	23.5	25.9	25.4	26.9	19.9	-7.0 s
	23–26	14.4	14.1	14.2	14.5	17.5	18.4	18.0	18.8	18.4	19.7	17.6	17.9	+0.3
	27–30	12.9	11.0	11.8	11.9	12.9	12.3	12.0	13.5	11.8	12.5	10.0	12.8	+2.8
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	2.4	3.4	2.8	3.1	2.2	2.4	2.1	2.9	2.2	2.0	2.4	1.8	-0.6
	19–22	0.7	1.2	0.7	1.7	1.6	1.3	1.2	0.5	0.7	1.1	0.4	1.3	+0.8
	23–26	0.8	0.5	0.6	0.3	0.5	0.3	0.7	0.1	0.3	0.7	*	0.1	+0.1
	27–30	0.1	0.3	0.6	0.1	0.5	0.9	0.1	*	0.4	0.4	0.4	0.3	-0.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19-22, 23-26, 27-30, 35, 40, 45, and 50

Q. How many of your friends would you estimate...	Age Group	Percentage saying friends use ^a																
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Take sedatives/barbiturates^d																		
% saying any	18	30.5	31.1	31.3	28.3	26.6	27.1	25.6	24.3	19.7	20.3	17.4	14.8	16.4	17.8	18.2	17.8	21.6
	19-22	33.2	27.9	27.7	23.6	22.0	17.2	18.8	15.5	14.0	14.1	11.9	12.8	10.7	11.7	9.7	13.3	11.6
	23-26	—	—	—	—	22.2	18.7	16.3	14.1	11.2	10.4	8.9	8.3	8.7	8.2	7.6	9.6	6.9
	27-30	—	—	—	—	—	—	—	—	12.0	8.5	8.8	7.1	6.6	6.7	7.4	7.2	6.7
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	2.6	2.1	1.8	1.7	1.7	1.6	1.4	1.1	1.1	1.4	0.6	0.5	0.6	1.0	1.1	1.4	1.6
	19-22	1.1	1.3	1.0	0.8	0.8	0.5	0.3	0.4	0.8	0.1	0.2	0.3	0.1	0.1	0.3	0.8	0.2
	23-26	—	—	—	—	0.4	0.3	0.3	0.3	0.1	0.2	0.2	0.1	0.1	0.3	0.2	*	*
	27-30	—	—	—	—	—	—	—	—	0.2	*	0.4	0.2	0.2	0.2	*	*	0.3
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take quaaludes																		
% saying any	18	32.5	35.0	35.5	29.7	26.1	26.0	23.5	22.0	17.1	16.6	14.3	12.0	13.1	14.2	14.2	15.5	18.1
	19-22	38.3	36.2	35.4	30.5	24.6	19.9	20.3	16.9	12.5	10.9	10.0	10.6	9.2	10.0	7.8	11.5	10.1
	23-26	—	—	—	—	25.7	21.0	17.4	15.0	12.1	10.3	8.6	5.9	6.4	7.6	7.7	9.0	6.3
	27-30	—	—	—	—	—	—	—	—	11.8	7.9	8.2	7.0	7.1	6.5	6.6	4.5	6.9
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	3.6	3.6	2.6	2.6	1.7	1.3	1.6	1.0	1.0	1.3	0.8	0.5	0.8	1.1	1.1	1.3	1.7
	19-22	1.9	2.7	1.2	1.3	1.2	0.6	0.2	0.4	0.4	0.2	0.6	0.2	0.1	0.1	0.2	0.7	0.1
	23-26	—	—	—	—	0.6	0.3	0.7	0.2	0.2	0.4	0.2	0.1	0.2	0.6	0.2	0.2	*
	27-30	—	—	—	—	—	—	—	—	0.5	0.2	0.2	0.2	*	0.2	*	*	0.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take tranquilizers																		
% saying any	18	29.7	29.5	29.9	26.7	26.6	25.8	24.2	23.3	19.9	18.0	14.9	13.5	14.6	15.5	16.5	15.8	18.1
	19-22	37.5	33.9	28.7	22.9	22.0	19.7	20.6	18.0	16.4	14.8	13.4	13.0	11.3	11.9	9.5	13.6	10.5
	23-26	—	—	—	—	29.3	26.3	22.3	20.8	15.5	13.1	14.8	12.1	12.5	11.0	13.4	10.4	10.7
	27-30	—	—	—	—	—	—	—	—	20.1	16.6	16.9	14.9	12.0	12.5	13.9	11.9	11.0
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	14.3	12.2	13.1
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	1.9	1.4	1.1	1.2	1.5	1.2	1.3	1.0	0.7	1.5	0.5	0.4	0.7	0.9	0.9	1.1	1.4
	19-22	0.7	0.9	0.5	0.8	0.3	0.7	0.3	0.6	0.4	0.1	0.4	0.5	0.1	0.1	0.2	0.7	0.7
	23-26	—	—	—	—	0.4	0.3	0.5	*	0.3	0.4	0.2	0.3	0.1	0.4	0.2	*	*
	27-30	—	—	—	—	—	—	—	—	0.5	0.3	0.4	0.2	0.1	0.2	0.4	*	0.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.5	0.3	0.3
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How many of your friends would you estimate. . .	Age Group	Percentage saying friends use ^a												2007–2008 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Take sedatives/barbiturates^d														
% saying any	18	20.4	22.8	20.9	21.6	22.1	25.3	18.1	25.2	22.3	22.5	20.8	19.8	-1.0
	19–22	12.1	14.8	16.0	15.2	18.6	17.1	14.4	18.8	19.6	18.7	20.1	17.8	-2.2
	23–26	8.4	7.9	8.3	6.6	11.1	10.9	12.9	16.7	15.7	16.2	16.5	13.4	-3.1
	27–30	6.5	6.1	5.7	6.4	7.9	7.4	7.3	11.5	10.5	13.5	12.5	15.2	+2.7
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	1.1	2.5	1.4	1.7	1.1	1.7	1.9	2.0	1.8	1.3	1.6	1.3	-0.4
	19–22	0.7	0.4	0.4	1.0	0.9	0.8	0.7	0.3	0.3	0.2	0.6	0.5	-0.1
	23–26	0.8	*	*	0.4	0.4	*	0.2	0.4	0.2	0.5	*	0.4	+0.4
	27–30	*	*	0.2	*	0.3	0.6	0.1	*	0.5	0.4	0.6	0.1	-0.5
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Take quaaludes														
% saying any	18	16.1	17.4	15.5	16.2	17.8	18.0	14.2	16.6	13.6	13.4	13.6	11.2	-2.4
	19–22	9.3	10.6	11.4	13.1	14.6	13.0	10.3	8.3	8.2	8.6	8.8	5.9	-2.9
	23–26	6.5	6.6	6.4	4.9	7.7	8.5	8.9	6.5	7.7	5.6	5.6	4.1	-1.5
	27–30	4.9	4.1	5.1	5.0	4.9	6.6	4.3	4.4	3.6	4.9	4.3	5.8	+1.5
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	1.1	2.0	1.4	1.4	1.2	1.2	1.2	1.6	1.3	1.3	1.6	0.8	-0.9 s
	19–22	0.6	0.5	0.4	0.9	0.8	0.1	0.4	*	0.4	0.2	*	0.2	+0.2
	23–26	0.8	*	0.2	0.3	0.3	0.1	0.2	0.1	0.2	0.3	0.3	*	-0.3
	27–30	*	*	0.2	0.3	*	0.3	*	*	0.3	0.7	*	0.3	+0.3
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Take tranquilizers														
% saying any	18	17.9	19.7	16.4	19.4	18.6	21.2	17.2	18.3	16.9	15.3	15.5	15.0	-0.5
	19–22	11.7	13.7	16.2	16.7	21.3	18.1	14.5	12.3	11.5	13.0	17.2	11.6	-5.7 s
	23–26	9.6	8.5	9.8	11.2	12.4	14.9	12.9	15.1	13.1	10.7	12.3	12.6	+0.4
	27–30	10.8	12.6	10.4	10.6	9.6	10.6	10.4	9.9	9.7	8.5	9.1	12.3	+3.2
	35	10.8	10.7	11.4	10.8	12.2	12.5	11.4	12.7	12.4	12.2	14.7	16.1	+1.5
	40	—	13.7	14.8	15.2	15.1	15.6	15.0	13.6	14.1	16.1	16.0	15.0	-1.0
	45	—	—	—	—	—	—	17.3	19.8	15.4	18.3	20.7	17.3	-3.4
	50	—	—	—	—	—	—	—	—	—	—	—	19.7	—
% saying most or all	18	0.8	2.3	1.3	2.1	1.3	1.6	1.5	1.7	1.6	1.2	1.8	1.2	-0.6
	19–22	0.8	0.6	0.3	0.6	0.9	0.4	0.4	0.3	0.3	0.3	0.3	0.1	-0.2
	23–26	1.1	0.1	*	0.5	0.8	0.1	*	0.5	0.7	0.4	*	0.1	+0.1
	27–30	*	*	0.4	*	0.4	0.6	0.1	*	0.2	0.2	*	0.1	+0.1
	35	0.1	0.2	0.6	0.6	0.2	0.1	0.2	0.3	0.5	0.3	0.5	0.3	-0.2
	40	—	*	0.4	0.1	0.3	0.2	*	0.2	0.2	0.3	0.3	0.3	0.0
	45	—	—	—	—	—	—	0.3	0.2	0.2	0.1	0.3	*	-0.3
	50	—	—	—	—	—	—	—	—	—	—	—	0.3	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How many of your friends would you estimate...	Age Group	Percentage saying friends use ^a																
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Drink alcoholic beverages																		
% saying any	18	96.1	94.7	95.7	95.5	94.6	94.6	95.6	95.4	95.7	95.1	92.0	91.2	90.5	88.9	90.1	90.9	89.6
	19–22	96.3	96.7	96.6	97.3	96.8	95.8	96.9	95.6	97.0	97.6	96.1	95.2	93.1	95.1	92.5	94.8	93.7
	23–26	—	—	—	—	96.8	96.8	96.2	95.9	95.3	95.4	94.7	93.9	95.1	94.4	94.0	94.1	92.7
	27–30	—	—	—	—	—	—	—	—	96.1	96.0	95.2	94.4	95.6	93.4	93.3	93.3	93.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	89.6	89.9	90.3
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	68.9	67.7	69.7	69.0	66.6	66.0	68.0	71.8	68.1	67.1	60.5	58.6	56.9	57.0	59.6	56.4	56.4
	19–22	76.6	77.6	75.2	75.1	74.9	71.9	74.2	71.3	73.4	74.1	70.0	71.4	67.4	66.5	68.7	63.9	67.0
	23–26	—	—	—	—	73.2	74.4	69.5	74.9	68.9	69.8	67.1	69.3	68.8	68.7	70.7	67.0	68.9
	27–30	—	—	—	—	—	—	—	—	66.7	67.8	62.0	62.7	63.3	61.3	63.2	62.6	64.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	43.8	45.1	49.5
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Get drunk at least once a week																		
% saying any	18	83.1	81.8	83.1	83.9	81.5	82.5	84.7	85.6	84.4	82.8	79.2	79.8	79.9	79.2	81.4	78.9	78.5
	19–22	80.9	79.9	80.0	80.4	79.8	76.7	82.0	81.1	80.6	80.4	80.1	80.8	76.5	81.1	79.6	83.2	80.9
	23–26	—	—	—	—	73.1	72.7	73.5	73.7	72.1	73.1	72.2	74.0	73.1	74.3	72.1	73.1	74.5
	27–30	—	—	—	—	—	—	—	—	66.3	61.8	65.4	65.2	65.5	64.5	62.7	67.1	66.7
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	44.3	43.2	44.9
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	30.1	29.4	29.9	31.0	29.6	29.9	31.8	31.3	29.6	31.1	27.5	29.7	28.6	27.6	28.4	27.4	29.0
	19–22	21.9	23.3	22.0	20.2	22.7	21.7	20.8	21.3	24.0	22.6	23.6	24.9	22.6	28.8	26.3	28.2	26.0
	23–26	—	—	—	—	11.4	11.6	12.5	11.9	12.8	12.0	13.9	11.6	14.6	13.2	15.2	15.2	14.0
	27–30	—	—	—	—	—	—	—	—	5.2	6.3	6.7	6.6	5.9	6.7	6.4	7.9	8.6
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.6	3.6	5.4
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Smoke cigarettes																		
% saying any	18	90.6	88.5	88.3	87.0	86.0	87.0	87.8	88.3	87.7	86.5	84.9	85.7	84.4	84.8	88.1	87.9	88.3
	19–22	94.4	94.3	93.4	93.1	91.9	91.6	91.1	90.3	89.3	90.0	86.1	86.1	86.7	86.7	86.1	88.8	89.2
	23–26	—	—	—	—	93.9	95.0	91.6	92.1	89.8	90.1	88.7	89.6	85.6	88.3	86.4	86.8	85.3
	27–30	—	—	—	—	—	—	—	—	92.6	89.8	90.7	90.4	88.0	85.8	84.8	84.9	85.4
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	72.7	71.7	71.7
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	23.3	22.4	24.1	22.4	19.2	22.8	21.5	21.0	20.2	23.1	21.4	21.8	21.4	25.0	25.3	27.5	30.4
	19–22	31.8	27.6	25.6	25.2	25.6	22.7	21.9	22.5	19.3	19.9	19.2	20.2	20.3	22.2	21.7	28.4	24.0
	23–26	—	—	—	—	25.6	22.7	19.7	18.5	16.5	20.5	16.9	18.1	16.0	15.5	16.6	13.9	17.6
	27–30	—	—	—	—	—	—	—	—	15.8	14.2	11.6	12.9	11.9	14.3	10.9	12.3	10.4
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	7.9	7.2	9.3
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How many of your friends would you estimate . . .	Age Group	Percentage saying friends use ^a											2007–2008 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007		2008
Drink alcoholic beverages														
% saying any	18	90.7	91.2	90.2	89.8	89.2	88.0	87.9	87.8	87.2	86.0	85.1	85.2	+0.1
	19–22	94.5	94.5	92.8	95.2	93.4	94.5	92.5	90.4	95.0	91.9	94.0	92.6	-1.4
	23–26	95.4	95.5	93.3	94.5	93.1	95.3	92.8	94.9	91.6	93.6	94.7	93.3	-1.4
	27–30	95.1	93.1	94.4	92.7	91.4	92.8	90.5	94.4	93.7	95.6	92.4	91.7	-0.7
	35	89.5	88.1	88.7	89.6	89.3	90.1	87.4	93.4	91.3	90.6	90.5	91.0	+0.5
	40	—	88.4	88.9	90.7	89.6	90.5	89.2	90.5	92.1	90.8	93.0	89.3	-3.7 ss
	45	—	—	—	—	—	—	87.9	90.3	89.8	90.1	89.8	90.5	+0.7
	50	—	—	—	—	—	—	—	—	—	—	—	88.9	—
% saying most or all	18	60.9	61.0	58.2	57.2	59.2	53.7	53.1	53.9	55.3	52.4	52.0	51.6	-0.4
	19–22	63.8	69.4	67.8	70.1	65.4	68.8	63.9	66.4	71.8	65.4	71.1	64.4	-6.7
	23–26	66.6	67.4	63.6	70.8	65.7	73.4	66.0	71.3	69.3	69.2	70.2	76.3	+6.1
	27–30	66.6	62.9	64.4	64.8	64.9	66.3	61.5	69.0	66.2	70.7	65.6	67.1	+1.5
	35	46.6	47.1	46.0	49.1	48.4	52.9	51.6	53.7	55.5	55.2	56.1	55.7	-0.4
	40	—	37.7	41.4	42.5	44.7	44.8	47.2	43.3	47.2	45.9	50.3	48.9	-1.5
	45	—	—	—	—	—	—	38.9	41.7	42.4	45.1	46.6	47.0	+0.4
	50	—	—	—	—	—	—	—	—	—	—	—	37.7	—
Get drunk at least once a week														
% saying any	18	82.4	81.1	81.5	79.5	79.6	78.3	77.3	79.0	78.7	77.4	75.5	76.2	+0.7
	19–22	79.2	82.3	82.8	82.2	81.9	81.5	81.5	80.5	85.1	81.7	84.4	81.3	-3.1
	23–26	71.9	74.1	71.0	76.5	74.7	81.0	76.4	75.8	80.7	80.9	80.4	79.5	-0.9
	27–30	65.4	65.5	65.9	64.3	64.7	68.9	66.5	73.8	72.4	74.6	72.0	71.7	-0.3
	35	42.9	46.1	44.5	46.9	47.6	48.3	47.9	52.0	50.7	52.6	55.0	56.0	+1.1
	40	—	41.6	40.6	42.2	41.3	42.6	42.9	43.2	48.4	47.2	46.3	48.2	+1.9
	45	—	—	—	—	—	—	41.6	42.2	41.6	40.0	42.7	45.7	+3.0
	50	—	—	—	—	—	—	—	—	—	—	—	40.0	—
% saying most or all	18	30.9	31.7	30.1	32.4	32.7	28.3	27.1	27.6	28.5	27.7	27.0	25.2	-1.8
	19–22	26.6	29.8	29.3	28.1	30.2	31.0	29.6	29.0	31.2	32.9	32.0	28.9	-3.1
	23–26	17.0	16.0	16.8	17.4	19.1	19.2	18.3	24.0	24.0	20.3	22.8	23.1	+0.3
	27–30	7.7	9.3	12.1	9.8	11.7	8.9	13.0	9.4	11.2	13.5	12.2	10.9	-1.3
	35	3.2	4.4	4.9	4.6	4.8	4.5	5.2	5.3	5.3	5.6	6.1	7.3	+1.2
	40	—	2.8	3.0	2.5	2.9	3.8	3.9	3.0	3.6	4.0	3.4	4.8	+1.4
	45	—	—	—	—	—	—	3.6	2.7	2.7	3.1	3.7	4.1	+0.4
	50	—	—	—	—	—	—	—	—	—	—	—	3.2	—
Smoke cigarettes														
% saying any	18	89.9	89.5	89.3	87.2	86.8	85.4	83.3	83.7	81.8	81.4	77.1	78.4	+1.3
	19–22	91.3	92.6	91.0	90.9	90.9	89.7	86.5	89.7	89.3	85.8	86.8	84.4	-2.3
	23–26	85.4	88.7	84.1	86.5	86.7	86.4	86.5	87.0	87.3	85.4	84.1	86.8	+2.7
	27–30	84.1	81.1	86.3	85.1	84.9	87.0	82.8	83.5	81.0	84.4	81.7	82.1	+0.4
	35	72.4	71.8	69.9	70.8	69.2	66.6	67.0	67.7	65.5	67.0	64.8	67.6	+2.8
	40	—	70.2	70.0	67.8	64.3	65.5	65.1	62.4	63.8	64.6	59.2	59.7	+0.6
	45	—	—	—	—	—	—	66.1	67.0	62.9	60.9	58.5	56.1	-2.4
	50	—	—	—	—	—	—	—	—	—	—	—	62.1	—
% saying most or all	18	34.4	33.9	31.1	28.2	25.0	23.0	19.6	20.6	16.7	15.8	16.4	13.9	-2.5
	19–22	25.1	28.8	26.8	29.4	27.0	25.7	20.2	20.7	20.4	15.2	17.9	12.9	-5.0
	23–26	17.0	16.8	17.5	17.0	15.5	15.1	18.3	19.8	19.6	13.9	14.7	15.0	+0.4
	27–30	12.1	12.3	13.4	11.7	10.2	12.9	12.2	9.2	12.6	12.6	12.7	10.8	-1.9
	35	7.2	8.0	9.0	6.7	8.8	6.6	6.3	6.9	6.0	6.8	5.7	5.9	+0.2
	40	—	8.1	7.4	6.8	5.7	5.8	5.9	6.0	7.0	5.1	4.7	4.5	-0.2
	45	—	—	—	—	—	—	5.7	5.9	6.1	5.4	4.5	3.7	-0.8
	50	—	—	—	—	—	—	—	—	—	—	—	4.0	—

↓
(List of drugs continued.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19-22, 23-26, 27-30, 35, 40, 45, and 50

Q. How many of your friends would you estimate. . .	Age Group	Percentage saying friends use ^a																
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Take steroids																		
% saying any	18	—	—	—	—	—	—	—	—	—	25.9	24.7	21.5	19.0	18.1	19.5	17.9	
	19-22	—	—	—	—	—	—	—	—	23.4	21.5	22.2	19.7	20.7	16.8	16.6	16.1	
	23-26	—	—	—	—	—	—	—	—	15.3	15.0	12.3	14.5	11.1	10.5	12.4	7.3	
	27-30	—	—	—	—	—	—	—	—	9.9	10.5	7.5	8.0	8.0	8.0	8.0	10.2	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
% saying most or all	18	—	—	—	—	—	—	—	—	—	1.8	1.0	1.7	0.9	1.2	1.3	0.8	
	19-22	—	—	—	—	—	—	—	—	0.2	0.6	*	0.1	0.4	0.2	0.1	*	
	23-26	—	—	—	—	—	—	—	—	0.4	*	*	0.2	0.1	0.1	*	*	
	27-30	—	—	—	—	—	—	—	—	0.5	*	*	*	0.2	0.1	*	*	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
<i>Approximate</i>	18	2,987	3,307	3,303	3,095	2,945	2,971	2,798	2,948	2,961	2,587	2,361	2,339	2,373	2,410	2,337	2,379	2,156
<i>Weighted N =</i>	19-22	576	592	564	579	543	554	579	572	562	579	556	526	510	468	435	470	469
	23-26					527	534	546	528	528	506	510	507	516	495	449	456	416
	27-30									516	507	499	476	478	461	419	450	464
	35														1,200	1,187	1,187	
	40																	
	45																	
	50																	

(Table continued on next page.)

TABLE 7-2 (cont.)
Trends in Friends' Use of Drugs as Estimated by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How many of your friends would you estimate . . .	Age Group	Percentage saying friends use ^a											2007–2008 change	
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007		2008
Take steroids														
% saying any	18	18.9	18.3	20.0	19.8	21.7	21.6	21.1	22.8	19.1	19.8	20.1	19.4	-0.7
	19–22	16.8	20.0	20.6	18.9	20.0	19.3	17.1	21.4	20.1	21.0	18.3	14.8	-3.5
	23–26	13.0	9.2	15.0	12.2	13.6	14.3	12.9	12.4	11.6	13.4	13.8	13.3	-0.5
	27–30	9.1	7.0	11.2	9.3	10.7	6.4	11.6	10.1	7.4	7.5	6.7	6.6	-0.1
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
% saying most or all	18	1.7	1.4	0.9	1.9	1.2	1.5	1.5	2.6	1.5	0.9	1.2	1.3	+0.1
	19–22	0.1	0.3	0.1	0.3	0.7	0.7	0.4	*	0.1	0.3	0.3	0.3	0.0
	23–26	0.5	*	0.1	0.3	0.2	0.1	*	0.1	0.3	0.3	*	*	0.0
	27–30	*	*	*	*	*	0.3	*	*	0.1	*	*	*	0.0
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Approximate</i>	18	2,292	2,313	2,060	1,838	1,923	1,968	2,233	2,271	2,266	2,266	2,253	2,125	
<i>Weighted N =</i>	19–22	467	437	426	402	402	375	388	443	395	377	362	375	
	23–26	419	394	414	387	403	358	362	411	361	336	340	355	
	27–30	454	428	424	363	359	348	369	396	363	350	324	332	
	35	1,209	1,067	1,071	1,033	1,005	918	968	985	1,041	953	884	905	
	40		1,098	1,156	1,144	1,119	1,083	945	1,004	975	951	896	924	
	45							976	1,074	1,052	1,009	999	904	
	50												940	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

"—" indicates data not available.

"*" indicates a prevalence rate of less than 0.05%.

^aAnswer alternatives were: (1) None, (2) A few, (3) Some, (4) Most, (5) All. The "any" percentage combines categories (2)–(5).

The "most or all" percentage combines categories (4) and (5).

^bThese estimates were derived from responses to the questions listed above. For the young adult sample, "any illicit drug" includes all of the drugs listed except cigarettes and alcohol. For the 35-, 40-, 45-, and 50-year-olds, "any illicit drug" includes marijuana, tranquilizers, crack, cocaine powder, and "other illicit drugs."

^cIn 2001 the question text was changed from "other psychedelics" to "other hallucinogens," and "shrooms" was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.

^dIn 2004 the question text was changed from "barbiturates" to "sedatives/barbiturates" and the list of examples was changed from "downers, goofballs, rebs, yellows, etc." to just "downers." These changes likely explain the discontinuity in the 2004 results.

TABLE 7-3
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

		Percentage saying exposed to drug ^a																	
		Age Group																	
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	
Any illicit drug^b	% saying any	18	84.3	82.7	81.4	79.4	77.9	77.7	75.5	73.9	71.3	68.6	67.6	64.2	61.3	66.1	70.8	75.3	78.0
		19–22	80.6	81.0	81.5	76.5	76.3	77.4	74.6	72.7	69.5	61.5	60.8	58.9	58.6	58.4	60.7	66.4	67.2
		23–26	—	—	—	—	68.9	70.2	68.0	62.4	62.7	58.3	54.6	52.1	48.2	49.9	47.1	54.2	50.3
		27–30	—	—	—	—	—	—	—	—	52.4	50.2	47.0	39.6	41.7	38.9	45.6	42.4	44.9
		% saying often	18	36.3	36.1	31.4	29.8	28.3	27.2	26.3	23.3	20.8	22.0	20.7	18.2	18.0	24.0	29.3	32.3
		19–22	34.6	34.0	32.1	24.4	24.4	23.7	21.1	18.9	19.9	16.2	16.4	17.6	21.4	16.1	18.1	23.7	20.4
		23–26	—	—	—	—	20.7	23.3	18.5	17.4	18.2	13.8	13.7	13.3	12.2	11.1	11.1	12.5	12.8
		27–30	—	—	—	—	—	—	—	13.7	12.0	10.8	8.2	10.5	9.0	12.5	8.5	10.1	
Any illicit drug^b other than marijuana	% saying any	18	58.5	62.6	62.5	59.4	59.8	59.3	55.3	51.7	47.8	47.1	45.4	40.0	41.6	42.6	45.3	47.2	49.7
		19–22	56.9	58.4	61.6	54.9	57.1	53.3	53.4	48.5	46.4	36.5	39.4	33.8	37.1	29.4	33.9	36.8	36.5
		23–26	—	—	—	—	51.5	51.9	51.5	43.6	42.9	36.8	34.0	30.0	27.3	27.8	24.9	26.8	23.2
		27–30	—	—	—	—	—	—	—	—	35.8	33.7	31.5	25.8	26.6	24.2	25.8	21.1	21.8
		% saying often	18	14.1	17.1	16.6	14.2	14.6	12.9	12.1	10.2	9.6	10.7	9.2	7.9	7.5	9.6	9.4	11.1
		19–22	11.8	15.6	13.5	11.1	10.7	10.2	8.2	8.1	7.5	6.7	4.5	4.4	5.5	4.1	5.1	7.7	3.9
		23–26	—	—	—	—	9.0	10.4	9.3	8.5	6.7	5.0	5.1	3.5	2.6	3.0	2.2	3.5	3.4
		27–30	—	—	—	—	—	—	—	6.0	4.7	4.1	3.2	3.7	2.4	3.4	2.9	3.4	
Marijuana	% saying any	18	82.0	80.2	77.9	76.2	74.4	73.5	72.0	70.4	67.0	64.8	63.4	59.6	56.8	61.0	67.2	72.7	75.6
		19–22	79.8	79.8	78.7	72.7	74.1	75.5	72.4	70.5	66.3	59.3	57.5	55.0	56.4	55.4	56.8	64.0	64.8
		23–26	—	—	—	—	65.3	66.0	64.1	59.0	57.6	55.0	50.6	47.9	44.6	45.9	44.4	51.0	47.8
		27–30	—	—	—	—	—	—	—	—	49.1	47.4	42.1	36.0	38.2	35.3	41.9	38.3	41.8
		% saying often	18	33.8	33.1	28.0	26.1	24.8	24.2	24.0	20.6	17.9	19.5	17.8	16.0	15.6	20.9	27.6	30.7
		19–22	32.6	30.5	30.3	21.1	21.9	20.3	18.6	16.4	18.3	14.2	14.7	15.9	19.9	14.7	17.0	22.1	20.3
		23–26	—	—	—	—	17.5	20.6	14.6	14.8	15.6	11.6	11.2	11.6	10.9	10.4	10.4	11.1	11.5
		27–30	—	—	—	—	—	—	—	10.9	9.8	8.5	6.7	8.9	7.6	10.7	7.4	9.1	
LSD	% saying any	18	17.2	17.4	16.1	13.8	12.5	13.2	13.1	12.9	13.4	15.0	14.9	15.7	17.8	21.0	24.2	26.1	27.6
		19–22	17.4	15.8	16.0	13.5	12.8	12.7	10.8	10.9	12.0	12.0	12.1	13.1	19.3	13.4	16.5	18.6	20.7
		23–26	—	—	—	—	8.3	9.3	8.8	7.3	6.3	6.7	8.4	8.6	8.8	7.8	8.4	9.9	8.6
		27–30	—	—	—	—	—	—	—	—	3.6	3.2	3.3	3.6	3.9	4.9	5.3	5.5	4.3
		% saying often	18	1.4	2.0	1.9	1.4	1.5	1.3	1.6	1.8	1.6	2.2	2.6	2.9	3.0	3.9	4.2	6.1
		19–22	1.4	1.5	1.4	0.6	0.8	0.7	0.5	1.2	0.6	1.1	1.2	1.0	2.0	1.1	0.4	3.6	1.4
		23–26	—	—	—	—	0.3	0.4	0.4	0.7	0.6	0.3	0.5	0.2	0.8	0.3	0.5	0.5	0.4
		27–30	—	—	—	—	—	—	—	0.3	0.2	0.5	0.2	0.2	0.5	0.5	0.2	0.2	
Other hallucinogens^c	% saying any	18	20.4	17.6	16.8	13.1	12.7	12.5	11.8	10.0	9.0	8.8	9.4	9.4	9.7	12.1	14.0	15.8	16.6
		19–22	18.3	16.3	16.3	12.5	10.5	11.0	9.2	9.1	7.7	8.4	8.3	8.9	10.6	6.7	8.3	12.8	13.1
		23–26	—	—	—	—	8.4	8.9	9.1	6.0	5.1	4.8	5.7	5.5	5.1	5.7	5.2	5.5	6.9
		27–30	—	—	—	—	—	—	—	—	5.0	3.4	3.4	3.4	2.1	3.7	3.4	4.2	3.2
		% saying often	18	2.2	2.0	2.6	1.1	1.7	1.4	1.5	1.2	1.1	1.3	1.2	1.3	1.1	1.9	2.3	2.5
		19–22	1.1	0.9	0.9	0.7	0.8	0.8	0.2	0.8	0.3	0.4	0.4	0.5	0.7	0.4	0.2	1.6	0.7
		23–26	—	—	—	—	0.1	0.3	0.5	0.6	0.8	0.1	0.4	0.4	*	0.2	0.4	0.3	0.3
		27–30	—	—	—	—	—	—	—	0.2	0.4	0.5	0.3	0.1	0.5	0.2	0.3	0.2	

cont. →

(Table continued on next page.)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

	Age Group	Percentage saying exposed to drug ^a												2007–2008 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Any illicit drug^b														
% saying any	18	78.8	77.2	77.9	76.0	76.5	76.5	73.6	74.3	73.0	73.7	70.8	71.9	+1.1
	19–22	65.3	69.1	65.8	64.7	69.7	65.7	68.0	67.6	68.8	67.1	67.4	66.2	-1.2
	23–26	55.4	50.6	50.5	55.1	56.4	56.5	57.0	53.5	53.9	56.7	58.3	56.3	-2.0
	27–30	41.6	37.5	41.1	40.8	42.2	47.0	46.7	43.3	45.7	48.4	44.1	48.7	+4.6
% saying often	18	34.7	33.2	35.6	32.6	33.6	32.6	31.8	30.4	29.9	29.7	27.8	28.6	+0.8
	19–22	25.3	24.2	24.0	21.3	26.1	25.2	26.5	26.8	25.2	24.2	22.8	20.1	-2.7
	23–26	14.3	14.2	15.0	15.9	16.4	15.9	17.8	15.1	18.7	14.9	18.9	15.4	-3.5
	27–30	10.3	8.5	9.6	9.4	10.4	13.8	13.9	10.3	14.5	13.2	9.7	9.7	-0.1
Any illicit drug^b other than marijuana														
% saying any	18	47.9	47.3	46.5	47.2	49.9	49.3	46.3	48.3	45.9	45.4	45.4	43.8	-1.6
	19–22	39.4	40.0	36.4	38.1	39.2	38.0	40.2	40.9	41.1	38.5	42.7	38.2	-4.5
	23–26	25.6	27.1	28.0	31.0	31.4	31.5	32.2	32.6	32.3	34.5	33.1	31.3	-1.8
	27–30	21.4	15.4	19.5	17.2	22.2	23.1	26.1	23.2	27.1	27.4	24.8	27.7	+2.9
% saying often	18	11.7	9.9	11.7	10.5	11.9	12.6	10.8	11.4	10.6	11.4	10.8	8.2	-2.6 s
	19–22	7.6	7.0	4.8	6.4	7.8	8.6	5.2	7.9	8.0	6.7	6.9	6.6	-0.3
	23–26	3.1	3.1	4.3	3.5	3.4	5.0	5.4	5.4	4.0	5.4	6.7	5.4	-1.4
	27–30	3.2	1.0	2.5	1.6	3.7	4.7	4.9	2.4	5.6	4.0	3.4	2.3	-1.2
Marijuana														
% saying any	18	76.8	75.5	75.8	73.8	74.9	74.2	71.4	72.2	70.8	71.4	68.4	69.8	+1.4
	19–22	63.4	67.1	63.5	63.9	68.0	64.6	64.8	65.1	66.8	65.4	66.3	64.3	-1.9
	23–26	53.1	48.8	48.1	51.8	54.2	53.5	54.4	50.6	49.7	51.9	53.3	54.0	+0.7
	27–30	39.1	35.7	38.7	38.8	37.0	44.6	44.1	40.4	42.4	44.1	40.7	44.8	+4.1
% saying often	18	32.9	31.4	34.4	30.3	30.8	30.7	30.4	28.0	27.0	27.8	25.1	27.0	+1.9
	19–22	23.7	22.8	23.0	20.4	24.5	24.8	24.2	24.5	23.6	23.1	20.1	18.3	-1.8
	23–26	12.9	13.6	13.2	15.2	15.6	14.9	16.2	13.7	17.8	12.5	16.2	13.7	-2.5
	27–30	8.9	8.1	8.8	8.6	8.4	11.7	11.7	9.6	12.2	11.5	8.2	8.5	+0.3
LSD														
% saying any	18	25.9	23.1	23.6	22.0	21.6	17.2	14.2	12.4	10.8	11.6	12.4	12.1	-0.2
	19–22	22.3	21.0	20.1	15.9	15.2	13.6	10.0	8.5	7.2	10.4	6.3	9.2	+2.9
	23–26	7.6	9.8	9.4	9.8	11.1	9.3	5.5	4.4	4.7	5.6	4.5	4.8	+0.4
	27–30	3.9	3.2	3.7	3.2	4.3	4.8	3.0	4.7	4.0	3.4	3.9	1.7	-2.2
% saying often	18	5.1	3.2	4.1	3.3	2.8	2.6	1.8	1.6	1.5	1.9	1.7	0.8	-0.8 s
	19–22	1.8	2.0	1.7	1.4	2.4	0.9	0.2	0.1	0.7	0.7	0.3	0.7	+0.4
	23–26	0.2	0.1	0.3	0.2	*	0.3	0.3	*	0.3	*	0.5	0.6	+0.1
	27–30	*	*	0.1	*	*	*	0.3	0.3	0.6	*	0.1	*	-0.1
Other hallucinogens^c														
% saying any	18	17.8	15.9	17.7	16.3	28.1	26.4	25.8	24.8	24.3	23.8	23.5	23.6	+0.2
	19–22	15.0	15.0	12.4	11.8	22.8	23.4	18.9	18.7	19.5	17.8	20.2	17.5	-2.8
	23–26	5.6	8.7	5.8	8.9	14.8	14.7	11.9	10.1	11.3	10.3	9.8	9.8	+0.1
	27–30	2.9	2.6	3.0	3.0	6.4	7.7	6.3	7.9	8.8	7.8	6.8	5.2	-1.6
% saying often	18	2.8	1.7	2.7	2.1	3.6	4.5	3.2	3.2	2.6	4.1	3.0	1.9	-1.1
	19–22	0.7	0.5	0.6	0.8	2.6	2.4	0.4	0.7	1.2	0.7	0.7	0.9	+0.2
	23–26	0.2	*	*	0.4	0.2	0.4	*	*	0.5	*	0.6	0.7	+0.1
	27–30	0.5	*	0.1	*	0.4	*	*	0.3	0.6	*	0.4	0.3	-0.1

↓
(List of drugs continued.)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

	Age Group	Percentage saying exposed to drug ^a																
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Cocaine																		
% saying any	18	37.7	36.3	34.9	33.3	35.6	38.3	37.4	34.9	30.2	30.2	27.7	21.3	19.8	19.2	18.8	21.6	25.0
	19–22	37.6	42.3	43.6	36.6	38.9	39.4	41.5	37.0	36.2	26.6	24.0	18.5	19.8	13.5	14.7	14.1	19.3
	23–26	—	—	—	—	38.5	40.6	42.0	34.5	35.9	28.0	24.0	19.9	16.7	14.6	14.3	14.1	12.5
	27–30	—	—	—	—	—	—	—	—	28.9	28.3	24.2	18.6	19.4	16.6	14.3	11.4	12.1
% saying often	18	5.9	6.6	6.6	5.2	6.7	7.1	7.8	5.9	5.1	5.4	4.7	3.4	2.7	2.9	2.5	3.2	4.0
	19–22	5.8	7.6	6.5	4.3	6.5	7.0	5.4	5.2	4.8	4.3	2.2	1.6	1.7	1.7	1.8	1.7	1.2
	23–26	—	—	—	—	5.3	8.5	7.0	6.0	5.4	3.5	2.5	1.7	1.4	1.7	1.0	1.7	1.3
	27–30	—	—	—	—	—	—	—	—	4.4	3.9	2.9	2.2	2.0	1.2	1.5	1.4	1.9
Heroin																		
% saying any	18	7.4	6.6	7.1	5.1	6.0	5.5	6.0	5.8	5.7	6.5	5.4	5.1	5.4	5.7	7.3	7.9	8.6
	19–22	4.4	3.3	4.1	2.9	3.1	4.8	2.9	2.9	2.9	2.9	2.5	3.0	2.7	2.0	3.7	3.8	3.6
	23–26	—	—	—	—	2.3	3.3	3.2	2.9	1.7	2.3	2.3	1.8	1.7	1.5	1.9	2.8	2.9
	27–30	—	—	—	—	—	—	—	—	2.1	1.4	1.5	0.9	1.0	2.0	2.0	1.7	1.5
% saying often	18	0.4	0.6	1.0	0.7	1.1	0.5	1.0	0.9	0.8	1.0	0.5	0.9	0.7	1.1	0.7	1.2	1.6
	19–22	0.2	0.3	0.3	0.1	0.2	0.5	0.2	0.1	0.2	0.1	0.2	0.4	0.6	0.4	0.6	1.2	0.2
	23–26	—	—	—	—	*	0.7	0.3	0.6	0.4	0.3	0.6	0.3	*	*	*	0.2	0.2
	27–30	—	—	—	—	—	—	—	—	0.3	0.3	0.5	0.2	0.2	0.9	0.3	0.6	0.6
Other narcotics																		
% saying any	18	19.6	17.5	18.5	17.3	18.0	18.4	15.6	14.4	14.8	13.8	14.2	11.3	11.1	12.4	14.9	15.5	18.5
	19–22	14.4	14.4	15.2	10.9	12.4	13.7	9.8	12.2	11.2	9.0	9.4	9.2	8.5	6.8	10.1	12.1	11.5
	23–26	—	—	—	—	9.0	12.3	9.2	9.7	7.4	8.0	5.9	8.3	7.0	4.6	6.9	7.8	7.4
	27–30	—	—	—	—	—	—	—	—	6.5	6.5	5.8	5.5	3.7	5.6	5.9	5.7	4.7
% saying often	18	1.7	1.7	2.4	2.2	2.0	1.8	2.1	1.7	1.7	1.7	1.6	1.4	1.3	1.7	1.7	2.1	3.4
	19–22	0.7	0.5	0.5	0.9	0.7	1.0	0.5	0.4	0.9	0.3	0.2	1.0	0.9	0.6	0.8	1.4	0.7
	23–26	—	—	—	—	0.4	0.5	1.3	0.8	0.8	0.5	1.6	0.7	0.1	0.3	0.1	0.1	0.3
	27–30	—	—	—	—	—	—	—	—	0.7	0.5	1.0	0.3	0.8	1.2	0.8	0.8	0.7
Amphetamines																		
% saying any	18	40.8	49.5	50.2	46.1	45.0	41.0	36.5	31.7	27.9	27.4	28.3	23.6	24.5	24.7	28.2	28.1	31.5
	19–22	42.3	48.6	48.4	39.7	41.3	35.9	31.3	26.7	21.2	18.5	19.5	17.4	21.3	15.1	20.3	21.0	22.3
	23–26	—	—	—	—	32.3	30.5	29.1	20.9	18.8	14.0	16.8	14.6	11.8	13.2	11.2	13.0	11.1
	27–30	—	—	—	—	—	—	—	—	15.6	14.3	13.5	10.7	11.4	11.3	11.0	10.6	7.6
% saying often	18	8.3	12.1	12.3	10.1	9.0	6.5	5.8	4.5	4.1	4.7	4.1	3.1	3.0	3.9	4.1	4.5	5.6
	19–22	7.4	9.9	7.7	6.9	5.4	4.4	3.1	3.3	2.2	1.5	1.1	1.9	2.6	1.5	3.3	5.0	1.3
	23–26	—	—	—	—	3.9	3.2	2.2	3.3	1.9	0.7	2.0	1.3	0.2	0.8	0.9	1.6	1.3
	27–30	—	—	—	—	—	—	—	—	2.0	2.0	1.2	0.8	0.8	1.3	0.7	1.6	1.8

(Table continued on next page.)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

	Age Group	Percentage saying exposed to drug ^a												2007–2008 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Cocaine														
% saying any	18	25.6	26.6	25.8	24.2	24.5	24.9	24.8	24.4	25.7	28.2	25.2	24.1	-1.2
	19–22	18.8	21.6	18.5	19.1	20.6	22.5	18.4	23.6	22.7	22.9	22.5	22.7	+0.2
	23–26	14.0	16.0	18.2	16.4	16.9	18.3	17.4	18.7	19.2	19.3	19.0	18.2	-0.8
	27–30	11.4	8.6	11.6	10.2	11.6	12.2	12.6	13.0	15.8	16.0	14.1	14.8	+0.7
% saying often	18	4.2	3.7	4.6	4.6	4.5	5.3	5.0	4.7	4.2	5.3	4.6	3.6	-1.0
	19–22	2.4	3.2	1.4	3.8	3.0	4.1	1.6	2.6	4.0	2.6	1.8	2.6	+0.8
	23–26	1.8	1.5	2.2	1.8	1.0	2.5	1.9	2.9	1.8	2.0	3.2	1.0	-2.2 s
	27–30	1.6	0.8	1.5	0.3	1.6	2.4	1.7	0.7	2.4	0.8	0.7	1.4	+0.7
Heroin														
% saying any	18	9.1	8.7	8.1	9.1	8.7	8.3	7.3	6.6	7.3	9.0	8.6	6.8	-1.8
	19–22	3.7	6.4	3.2	5.2	3.2	5.3	3.4	3.0	3.2	3.7	3.3	4.8	+1.5
	23–26	2.7	3.1	2.9	2.6	2.4	3.8	2.0	3.1	2.6	3.5	3.6	1.8	-1.8
	27–30	1.3	1.4	1.9	1.9	2.3	2.7	1.3	3.2	2.9	2.3	3.0	2.2	-0.8
% saying often	18	1.2	0.9	1.3	1.5	0.7	1.3	1.2	1.2	0.8	1.7	1.1	0.8	-0.3
	19–22	0.4	0.7	0.8	0.7	0.8	0.6	0.2	*	0.8	0.1	*	0.6	+0.6
	23–26	0.3	0.5	1.0	*	*	0.8	0.5	0.5	0.3	0.6	0.3	*	-0.3
	27–30	*	*	0.2	*	*	0.7	0.3	*	0.4	0.3	0.4	0.3	-0.1
Other narcotics														
% saying any	18	20.4	20.7	21.9	21.1	21.6	22.5	21.8	20.3	19.0	18.9	18.9	16.3	-2.6
	19–22	14.5	15.3	13.9	17.0	18.3	18.7	13.6	14.5	16.8	15.3	12.5	13.2	+0.7
	23–26	6.5	8.1	9.4	10.9	12.2	12.0	12.6	12.6	12.4	13.0	14.4	11.2	-3.2
	27–30	4.9	3.6	5.2	6.5	9.0	7.9	9.5	8.8	11.6	10.6	9.2	9.1	-0.2
% saying often	18	2.5	2.8	3.9	2.9	3.0	3.8	3.0	3.3	2.6	3.4	3.4	2.1	-1.3 s
	19–22	1.5	1.7	1.1	2.4	1.6	3.0	1.2	0.8	2.4	1.9	1.7	1.9	+0.2
	23–26	0.7	0.5	1.1	0.7	1.0	0.9	1.6	1.4	1.3	1.1	1.8	1.0	-0.9
	27–30	0.5	*	0.2	1.1	1.0	0.7	1.2	0.1	1.7	0.7	0.8	0.4	-0.3
Amphetamines														
% saying any	18	31.0	29.9	30.1	29.5	31.5	30.6	27.4	27.2	26.4	26.6	23.8	23.3	-0.5
	19–22	24.6	24.8	21.2	24.8	23.3	25.5	21.6	23.7	22.2	22.7	22.8	17.6	-5.2
	23–26	11.7	14.6	12.3	18.5	18.2	17.9	15.4	18.8	15.6	18.7	16.6	13.7	-2.9
	27–30	9.1	6.6	10.4	7.4	11.1	11.5	12.2	11.4	12.2	14.1	10.0	10.3	+0.3
% saying often	18	5.2	4.7	6.3	4.4	6.0	6.4	4.9	5.3	4.1	5.6	4.3	3.0	-1.3
	19–22	4.1	2.9	2.2	2.4	2.6	5.6	1.7	4.1	3.1	2.9	2.3	2.1	-0.2
	23–26	1.4	2.2	1.7	1.4	2.2	0.7	1.3	1.7	1.6	2.6	1.6	1.8	+0.1
	27–30	1.0	0.2	1.1	0.4	0.6	1.5	1.0	1.2	1.0	0.8	1.1	0.3	-0.8

↓
(List of drugs continued.)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

		Percentage saying exposed to drug ^a																
		Age Group																
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Sedatives/barbiturates^d																		
% saying any	18	25.2	25.9	25.7	22.5	21.2	18.9	15.8	13.1	12.4	11.8	13.3	10.0	10.2	11.9	13.0	14.5	15.5
	19–22	25.6	23.1	21.8	18.3	15.7	14.7	12.8	12.0	8.2	8.3	6.5	7.9	7.3	7.2	7.4	10.1	8.8
	23–26	—	—	—	—	16.1	13.1	11.0	7.1	7.1	6.6	6.9	5.9	6.5	3.8	4.2	5.7	6.6
	27–30	—	—	—	—	—	—	—	—	8.0	6.8	5.9	5.4	5.2	5.7	4.5	5.2	3.5
% saying often	18	3.4	4.0	4.3	3.0	2.7	1.7	2.1	1.5	1.4	1.7	1.7	1.2	1.1	1.6	1.7	2.0	2.9
	19–22	2.5	2.8	1.1	1.4	0.7	1.3	0.5	0.7	0.7	0.3	0.7	0.4	0.7	0.7	1.3	1.3	0.4
	23–26	—	—	—	—	0.7	0.9	1.7	0.8	0.6	0.3	1.1	0.3	0.3	*	*	0.2	0.3
	27–30	—	—	—	—	—	—	—	—	0.7	0.4	0.6	0.2	0.4	1.2	0.2	0.6	0.5
Tranquilizers^e																		
% saying any	18	29.1	29.0	26.6	23.5	23.1	23.4	19.6	18.4	18.2	15.1	16.3	14.2	12.7	13.8	16.5	15.7	17.9
	19–22	29.6	26.9	28.5	19.5	21.2	19.5	16.4	18.5	13.8	12.0	12.7	12.6	11.0	10.0	12.0	11.8	10.7
	23–26	—	—	—	—	23.1	21.0	16.9	15.9	13.4	12.9	12.0	10.4	9.7	10.9	9.8	10.3	10.1
	27–30	—	—	—	—	—	—	—	—	15.0	11.6	11.1	9.7	10.3	10.4	9.0	11.2	9.6
% saying often	18	3.2	4.2	3.5	2.9	2.9	2.2	2.5	2.6	2.2	2.1	1.9	1.4	1.9	1.7	1.8	2.3	3.5
	19–22	3.2	2.6	1.8	2.1	1.5	1.7	0.9	1.1	1.8	1.0	1.1	1.1	1.5	1.1	1.3	1.5	0.5
	23–26	—	—	—	—	2.0	1.6	2.6	1.8	1.2	0.8	0.5	1.0	0.6	0.7	0.1	1.1	1.5
	27–30	—	—	—	—	—	—	—	—	1.4	0.3	1.7	0.8	1.3	1.3	1.0	1.1	0.8
Alcoholic beverages																		
% saying any	18	94.7	94.0	94.0	94.0	94.0	94.0	94.1	93.9	93.1	92.3	93.6	91.7	90.6	91.8	90.0	91.2	91.5
	19–22	94.3	93.8	94.5	93.4	94.2	92.7	93.6	94.4	92.5	91.8	92.4	94.0	93.3	92.9	93.7	93.1	93.7
	23–26	—	—	—	—	90.3	92.7	91.4	90.6	91.1	92.9	91.3	91.0	91.4	90.3	89.5	91.9	89.6
	27–30	—	—	—	—	—	—	—	—	87.1	88.4	86.2	87.7	87.3	86.6	86.2	89.3	89.2
% saying often	18	60.2	61.0	59.3	60.2	58.7	59.5	58.0	58.7	56.4	55.5	56.1	54.5	53.1	51.9	54.0	54.0	54.5
	19–22	59.6	61.2	62.5	56.6	59.3	61.8	59.9	61.4	55.4	53.8	56.0	53.9	56.1	56.8	57.0	56.3	52.3
	23–26	—	—	—	—	52.1	54.8	51.4	53.0	48.1	50.9	49.7	48.4	45.4	45.4	43.3	47.5	44.8
	27–30	—	—	—	—	—	—	—	—	39.9	39.5	38.7	38.0	39.9	38.1	39.3	38.0	34.7
Approximate	18	3,259	3,608	3,645	3,334	3,238	3,252	3,078	3,296	3,300	2,795	2,556	2,525	2,630	2,730	2,581	2,608	2,407
Weighted N =	19–22	582	574	601	569	578	549	591	582	556	567	567	532	528	489	460	464	485
	23–26					533	532	557	529	531	514	523	494	532	513	471	467	447
	27–30									522	507	506	478	502	457	425	452	432

(Table continued on next page.)

TABLE 7-3 (cont.)
Trends in Exposure to Drug Use
among Respondents in Modal Age Groups of 18, 19–22, 23–26, and 27–30

Q. During the LAST 12 MONTHS how often have you been around people who were taking each of the following to get high or for "kicks"?

		Percentage saying exposed to drug ^a												2007– 2008
														change
	Age Group	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Sedatives/barbiturates^d														
% saying any	18	16.1	16.1	17.1	16.3	17.1	17.7	14.8	21.5	20.4	21.3	18.8	16.7	-2.2
	19–22	11.7	13.4	11.6	13.1	13.1	16.0	11.9	17.2	17.8	16.0	16.1	15.2	-0.9
	23–26	4.9	8.5	7.1	9.3	9.0	9.8	7.9	15.9	12.5	14.8	13.1	12.4	-0.8
	27–30	3.8	2.7	4.1	2.9	5.3	6.0	6.1	9.2	12.4	11.9	10.3	10.1	-0.2
% saying often	18	2.5	2.7	3.8	2.7	2.7	4.6	2.8	4.1	3.7	3.9	3.9	2.1	-1.7 ss
	19–22	0.9	1.4	0.9	1.6	1.2	1.8	0.8	1.7	2.1	2.5	1.4	2.2	+0.8
	23–26	0.8	0.5	0.9	0.7	0.2	0.3	0.4	0.7	1.1	1.1	1.6	1.7	+0.1
	27–30	0.2	*	0.6	0.2	0.9	0.4	0.6	0.4	1.7	0.7	1.3	0.4	-0.9
Tranquilizers^e														
% saying any	18	18.9	17.3	18.2	17.7	23.8	22.7	21.0	22.1	20.9	21.8	19.3	19.9	+0.6
	19–22	15.6	16.9	14.3	18.5	21.3	23.6	20.0	21.9	20.6	23.1	21.4	20.0	-1.5
	23–26	9.4	10.9	10.8	12.3	16.4	20.1	18.7	19.9	20.1	19.9	18.8	18.4	-0.4
	27–30	9.6	6.1	8.8	7.6	12.6	13.6	15.3	14.6	18.1	19.2	16.7	16.8	+0.1
% saying often	18	3.2	2.8	3.7	3.5	4.9	5.8	4.2	4.1	4.5	5.4	4.9	3.7	-1.3
	19–22	1.3	1.6	1.5	1.7	3.1	3.6	2.3	2.7	2.7	3.2	3.0	3.2	+0.2
	23–26	0.7	1.1	1.5	1.7	1.3	2.1	1.6	2.0	1.3	2.6	2.4	3.6	+1.1
	27–30	1.2	0.2	0.9	0.4	1.6	1.6	1.9	0.8	3.5	2.9	2.6	1.0	-1.6
Alcoholic beverages														
% saying any	18	91.4	92.2	91.8	90.7	90.8	89.5	88.3	87.6	87.4	87.6	86.5	85.7	-0.8
	19–22	93.1	91.8	91.0	93.3	94.3	93.7	93.6	92.5	92.7	92.0	91.8	90.5	-1.4
	23–26	93.1	89.1	91.5	92.1	90.1	91.9	91.8	92.2	90.0	94.0	94.5	92.0	-2.5
	27–30	86.4	88.4	88.7	89.8	91.2	89.0	90.0	85.3	92.2	91.8	89.6	94.4	+4.8 s
% saying often	18	53.9	54.5	53.5	50.2	52.7	50.8	49.0	48.2	49.1	47.8	46.4	45.4	-1.0
	19–22	54.2	57.9	54.7	54.3	53.4	54.9	55.7	54.3	58.9	55.0	60.7	53.9	-6.8
	23–26	49.8	44.6	45.7	49.6	48.8	46.3	50.5	48.3	46.4	57.1	54.2	49.6	-4.7
	27–30	37.1	36.6	38.3	34.4	40.0	39.6	40.6	36.8	43.6	47.3	44.3	47.8	+3.5
<i>Approximate</i>	18	2,595	2,541	2,312	2,153	2,147	2,162	2,454	2,456	2,469	2,469	2,448	2,332	
<i>Weighted N =</i>	19–22	471	445	450	415	412	403	396	432	377	378	333	365	
	23–26	424	400	398	389	406	345	385	404	374	363	327	333	
	27–30	455	449	430	395	369	359	347	370	370	330	356	339	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

** indicates a prevalence rate of less than 0.05%.

^aAnswer alternatives were: (1) Not at all, (2) Once or twice, (3) Occasionally, (4) Often. The "any" percentage combines categories (2)–(4).

^bThese estimates were derived from responses to the question for the following drugs: marijuana, LSD, other hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), and tranquilizers.

^cIn 2001 the question text was changed from "other psychedelics" to "other hallucinogens," and "shrooms" was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.

^dIn 2004 the question text was changed from "barbiturates" to "sedatives/barbiturates" and the list of examples was changed from "downers, goofballs, reids, yellows, etc." to just "downers." These changes likely explain the discontinuity in the 2004 results.

^eIn 2001 Xanax was added to the list of examples. This change likely explains the discontinuity in the 2001 results.

TABLE 7-4
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

		Percentage saying "fairly easy" or "very easy" to get ^a																	
<i>Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?</i>		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Marijuana	18	89.0	89.2	88.5	86.2	84.6	85.5	85.2	84.8	85.0	84.3	84.4	83.3	82.7	83.0	85.5	88.5	88.7	
	19–22	95.6	91.1	92.4	89.7	88.3	89.5	87.2	85.9	87.1	87.1	86.2	86.0	87.8	85.6	87.2	87.9	89.3	
	23–26	—	—	—	—	92.5	88.8	88.8	90.3	86.9	88.7	83.3	82.5	83.8	84.6	87.1	86.2	85.3	
	27–30	—	—	—	—	—	—	—	—	89.3	86.0	83.1	83.8	80.7	82.8	80.3	83.3	82.6	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	75.7	75.6	73.0	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Amyl & butyl nitrites	18	—	—	—	—	—	—	—	23.9	25.9	26.8	24.4	22.7	25.9	25.9	26.7	26.0	23.9	
	19–22	—	—	—	—	—	—	—	22.8	26.0	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	23.1	28.0	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	26.7	—	—	—	—	—	—	—	—	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
LSD	18	35.3	35.0	34.2	30.9	30.6	30.5	28.5	31.4	33.3	38.3	40.7	39.5	44.5	49.2	50.8	53.8	51.3	
	19–22	39.6	38.4	35.1	31.8	32.7	29.6	30.5	29.9	33.9	36.4	36.6	37.8	42.5	44.9	43.7	50.5	50.8	
	23–26	—	—	—	—	32.7	29.1	30.0	27.5	32.7	32.6	30.2	32.8	33.5	33.4	40.1	41.0	43.6	
	27–30	—	—	—	—	—	—	—	—	29.4	29.9	32.3	27.0	30.9	30.5	27.2	35.6	33.6	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	33.8	32.4	28.4	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Other hallucinogens^b	18	35.0	32.7	30.6	26.6	26.6	26.1	24.9	25.0	26.2	28.2	28.3	28.0	29.9	33.5	33.8	35.8	33.9	
	19–22	42.1	37.7	33.5	31.0	28.9	28.7	26.3	27.5	28.7	28.1	28.9	26.6	28.3	29.5	28.6	31.5	31.5	
	23–26	—	—	—	—	31.8	29.6	26.4	25.6	29.6	28.7	27.0	25.7	27.7	25.3	28.3	29.2	32.6	
	27–30	—	—	—	—	—	—	—	—	28.6	29.6	30.8	24.9	24.8	25.4	24.7	29.3	25.9	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
PCP	18	—	—	—	—	—	—	—	22.8	24.9	28.9	27.7	27.6	31.7	31.7	31.4	31.0	30.5	
	19–22	—	—	—	—	—	—	—	21.7	24.6	—	—	—	—	—	—	—	—	
	23–26	—	—	—	—	—	—	—	21.2	27.6	—	—	—	—	—	—	—	—	
	27–30	—	—	—	—	—	—	—	—	24.3	—	—	—	—	—	—	—	—	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Ecstasy (MDMA)	18	—	—	—	—	—	—	—	—	—	21.7	22.0	22.1	24.2	28.1	31.2	34.2	36.9	
	19–22	—	—	—	—	—	—	—	—	—	—	26.6	24.9	27.1	23.9	27.0	29.3	33.4	
	23–26	—	—	—	—	—	—	—	—	—	—	21.4	23.1	26.4	24.0	26.0	27.8	28.7	
	27–30	—	—	—	—	—	—	—	—	—	—	27.1	20.8	22.2	22.8	21.9	27.1	29.3	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	

(Table continued on next page.)

TABLE 7-4 (cont.)
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?	Age Group	Percentage saying "fairly easy" or "very easy" to get ^a												2007–2008 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Marijuana	18	89.6	90.4	88.9	88.5	88.5	87.2	87.1	85.8	85.6	84.9	83.9	83.9	-0.1
	19–22	90.6	89.9	87.4	89.6	91.7	88.1	87.7	87.3	88.0	86.8	88.4	87.5	-0.9
	23–26	84.4	87.5	85.9	88.4	87.0	89.1	87.2	88.8	87.0	86.8	87.6	85.3	-2.3
	27–30	84.5	82.1	83.0	81.5	84.8	83.6	81.8	86.0	84.6	87.6	87.8	86.4	-1.4
	35	77.1	76.0	74.9	77.1	75.3	76.5	75.1	75.6	73.8	75.1	75.5	76.4	+0.8
	40	—	73.4	71.7	73.1	70.4	72.1	72.3	68.9	73.6	69.7	71.2	72.5	+1.3
	45	—	—	—	—	—	—	68.5	69.9	70.1	67.9	70.1	68.1	-2.0
	50	—	—	—	—	—	—	—	—	—	—	—	64.4	—
Amyl & butyl nitrites	18	23.8	25.1	21.4	23.3	22.5	22.3	19.7	20.0	19.7	18.4	18.1	16.9	-1.1
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
LSD	18	50.7	48.8	44.7	46.9	44.7	39.6	33.6	33.1	28.6	29.0	28.7	28.5	-0.2
	19–22	47.7	51.1	43.8	47.1	42.5	37.9	34.1	30.3	27.7	29.0	23.0	19.7	-3.3
	23–26	39.2	40.4	41.2	40.4	38.3	37.2	34.1	38.5	26.5	30.3	25.2	24.1	-1.1
	27–30	35.2	32.9	35.7	35.6	38.3	32.3	33.5	30.0	29.3	29.7	26.8	28.1	+1.3
	35	32.9	31.2	27.7	32.2	28.7	29.1	29.8	25.6	24.0	28.7	26.6	26.4	-0.3
	40	—	31.1	31.0	28.5	25.7	27.4	25.0	24.4	24.3	23.9	21.5	25.1	+3.5
	45	—	—	—	—	—	—	24.2	27.0	25.4	23.7	23.6	21.1	-2.5
	50	—	—	—	—	—	—	—	—	—	—	—	19.0	—
Other hallucinogens^b	18	33.9	35.1	29.5	34.5	48.5	47.7	47.2	49.4	45.0	43.9	43.7	42.8	-0.8
	19–22	33.4	34.1	31.1	33.4	45.9	48.8	45.1	46.9	48.5	41.9	39.3	34.7	-4.6
	23–26	31.0	32.4	31.5	28.5	38.3	39.7	39.2	44.4	39.2	41.5	36.8	39.3	+2.5
	27–30	28.0	25.2	30.3	25.0	38.6	33.3	35.6	31.2	30.8	32.1	30.0	36.2	+6.3
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
PCP	18	30.0	30.7	26.7	28.8	27.2	25.8	21.9	24.2	23.2	23.1	21.0	20.6	-0.4
	19–22	—	—	—	—	—	—	—	—	—	—	—	—	—
	23–26	—	—	—	—	—	—	—	—	—	—	—	—	—
	27–30	—	—	—	—	—	—	—	—	—	—	—	—	—
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Ecstasy (MDMA)	18	38.8	38.2	40.1	51.4	61.5	59.1	57.5	47.9	40.3	40.3	40.9	41.9	+1.0
	19–22	35.6	39.4	43.2	49.9	55.5	59.7	52.1	45.8	43.5	41.2	38.4	34.7	-3.6
	23–26	31.1	30.1	34.9	41.8	51.5	52.9	49.3	51.3	46.4	44.6	42.2	41.5	-0.7
	27–30	24.3	26.4	30.0	35.5	40.6	41.2	41.0	41.1	38.0	40.5	40.7	42.2	+1.5
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—

↓
(List of drugs continued.)

TABLE 7-4 (cont.)
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

		Percentage saying "fairly easy" or "very easy" to get ^a																		
<i>Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?</i>		Age Group	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	cont.
Cocaine	18	47.9	47.5	47.4	43.1	45.0	48.9	51.5	54.2	55.0	58.7	54.5	51.0	52.7	48.5	46.6	47.7	48.1		
	19–22	55.7	56.2	57.1	55.2	56.2	56.9	60.4	65.0	64.9	66.8	61.7	54.3	54.5	49.2	49.9	49.4	44.4		
	23–26	—	—	—	—	63.7	67.2	65.8	69.0	71.7	70.0	65.6	58.0	61.1	53.8	54.4	54.7	50.2		
	27–30	—	—	—	—	—	—	—	—	68.6	68.2	64.0	60.0	63.1	56.8	53.1	57.0	53.0		
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Crack	18	—	—	—	—	—	—	—	41.1	42.1	47.0	42.4	39.9	43.5	43.6	40.5	41.9	40.7		
	19–22	—	—	—	—	—	—	—	41.9	47.3	47.2	46.9	42.1	42.1	38.4	41.6	40.7	32.9		
	23–26	—	—	—	—	—	—	—	44.5	53.0	49.9	46.9	42.0	42.6	42.5	42.4	42.3	37.9		
	27–30	—	—	—	—	—	—	—	—	46.5	46.8	46.8	43.1	45.2	45.8	41.1	44.7	39.9		
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	49.6	48.2	43.1		
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Cocaine powder	18	—	—	—	—	—	—	—	52.9	50.3	53.7	49.0	46.0	48.0	45.4	43.7	43.8	44.4		
	19–22	—	—	—	—	—	—	—	58.7	60.2	61.7	56.5	52.5	48.9	45.7	47.8	45.5	41.3		
	23–26	—	—	—	—	—	—	—	64.9	69.1	60.1	58.6	53.2	56.4	50.5	49.7	49.6	45.9		
	27–30	—	—	—	—	—	—	—	—	63.5	62.8	57.9	55.8	56.8	55.0	48.9	52.9	48.4		
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	53.9	52.1	46.7		
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Heroin	18	21.2	19.2	20.8	19.3	19.9	21.0	22.0	23.7	28.0	31.4	31.9	30.6	34.9	33.7	34.1	35.1	32.2		
	19–22	18.9	19.4	19.3	16.4	17.2	20.8	21.2	24.4	28.5	31.6	30.7	25.3	30.2	30.0	33.2	35.2	29.1		
	23–26	—	—	—	—	18.6	18.1	21.0	22.3	28.4	31.2	28.1	25.6	25.7	25.7	29.2	29.3	32.3		
	27–30	—	—	—	—	—	—	—	—	23.6	27.4	29.5	22.1	25.6	28.5	24.4	30.7	29.5		
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Some other narcotic	18	29.4	29.6	30.4	30.0	32.1	33.1	32.2	33.0	35.8	38.3	38.1	34.6	37.1	37.5	38.0	39.8	40.0		
	19–22	32.7	32.4	30.8	31.0	28.7	34.3	32.6	33.8	37.9	37.9	35.6	35.4	35.2	33.5	35.1	38.7	37.3		
	23–26	—	—	—	—	32.8	32.1	33.6	32.2	35.9	36.4	34.7	33.2	33.9	33.1	35.8	32.6	36.7		
	27–30	—	—	—	—	—	—	—	—	31.6	36.2	36.1	29.0	31.8	33.0	34.8	36.9	37.2		
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
Amphetamines	18	61.3	69.5	70.8	68.5	68.2	66.4	64.3	64.5	63.9	64.3	59.7	57.3	58.8	61.5	62.0	62.8	59.4		
	19–22	71.7	72.6	73.5	69.7	69.1	69.1	63.1	61.8	61.3	62.2	57.7	58.3	56.3	56.0	56.6	60.3	56.9		
	23–26	—	—	—	—	65.8	66.0	64.5	65.3	62.2	60.1	55.8	54.8	54.5	52.6	52.9	56.0	52.8		
	27–30	—	—	—	—	—	—	—	—	54.3	58.6	55.3	54.4	50.4	52.9	48.3	53.7	51.7		
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	45.6	43.5	39.1		
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—		

(Table continued on next page.)

TABLE 7-4 (cont.)
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?	Age Group	Percentage saying "fairly easy" or "very easy" to get ^a												2007–2008 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Cocaine	18	48.5	51.3	47.6	47.8	46.2	44.6	43.3	47.8	44.7	46.5	47.1	42.4	-4.7 ss
	19–22	49.7	47.7	52.6	52.1	49.6	47.6	46.7	47.0	50.0	47.4	47.3	44.0	-3.2
	23–26	46.9	51.8	45.7	45.0	44.6	47.8	40.8	50.7	48.4	51.2	47.4	45.5	-1.9
	27–30	50.4	46.9	50.0	44.6	45.5	46.3	42.9	38.0	43.1	43.2	45.8	50.6	+4.8
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Crack	18	40.6	43.8	41.1	42.6	40.2	38.5	35.3	39.2	39.3	38.8	37.5	35.2	-2.3
	19–22	39.9	40.0	40.8	40.2	37.3	35.7	37.5	33.7	34.0	35.2	35.7	31.4	-4.3
	23–26	37.2	38.4	35.0	31.9	37.1	33.9	32.8	36.5	35.1	34.0	31.4	33.1	+1.7
	27–30	36.5	33.3	38.8	35.9	36.9	33.4	33.7	28.0	34.4	29.6	36.4	36.1	-0.3
	35	44.3	45.0	41.6	45.0	41.2	38.9	40.5	36.1	34.2	37.1	35.1	33.2	-1.9
	40	—	43.3	44.3	42.0	38.7	39.5	39.0	35.8	38.6	37.1	32.7	35.2	+2.6
	45	—	—	—	—	—	—	37.0	40.0	40.6	36.2	37.0	34.2	-2.9
	50	—	—	—	—	—	—	—	—	—	—	—	32.8	—
Cocaine powder	18	43.3	45.7	43.7	44.6	40.7	40.2	37.4	41.7	41.6	42.5	41.2	38.9	-2.3
	19–22	46.0	47.1	45.2	45.2	43.3	43.9	45.5	43.2	44.3	44.2	44.5	39.0	-5.6
	23–26	43.6	44.4	44.3	41.8	44.4	40.7	43.4	48.5	45.1	46.4	45.0	41.4	-3.6
	27–30	45.1	43.9	46.5	43.9	42.7	42.4	39.7	37.9	40.2	42.7	43.0	47.5	+4.5
	35	48.3	47.0	43.4	47.9	43.1	41.7	42.0	39.6	35.8	39.5	37.4	38.6	+1.3
	40	—	46.0	46.7	44.7	41.5	41.5	40.7	38.5	40.3	37.8	35.2	36.5	+1.4
	45	—	—	—	—	—	—	39.0	40.2	40.6	37.3	38.2	34.1	-4.1
	50	—	—	—	—	—	—	—	—	—	—	—	32.6	—
Heroin	18	33.8	35.6	32.1	33.5	32.3	29.0	27.9	29.6	27.3	27.4	29.7	25.4	-4.2 ss
	19–22	31.4	32.1	32.7	29.4	30.2	26.4	26.9	22.6	25.4	25.3	26.5	24.2	-2.3
	23–26	30.5	35.1	31.9	25.7	26.6	27.2	25.5	30.9	22.5	28.1	22.2	23.4	+1.2
	27–30	30.0	28.3	33.0	29.3	29.9	27.0	27.5	22.0	27.8	25.4	27.5	26.3	-1.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Some other narcotic	18	38.9	42.8	40.8	43.9	40.5	44.0	39.3	40.2	39.2	39.6	37.3	34.9	-2.4
	19–22	38.3	38.9	39.5	41.1	44.1	40.4	40.6	39.4	41.4	38.5	38.3	38.0	-0.2
	23–26	35.7	39.9	38.2	38.1	35.8	40.0	40.3	47.7	44.7	45.5	41.7	41.2	-0.5
	27–30	35.2	32.2	36.9	32.4	39.4	38.5	38.9	35.8	37.7	39.8	41.3	39.4	-1.9
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Amphetamines	18	59.8	60.8	58.1	57.1	57.1	57.4	55.0	55.4	51.2	52.9	49.6	47.9	-1.7
	19–22	55.5	56.3	57.6	60.2	56.5	53.7	55.1	53.9	56.9	52.3	55.8	49.5	-6.3
	23–26	51.2	53.2	49.1	51.1	49.4	48.2	50.3	51.8	51.9	58.0	53.7	46.9	-6.8
	27–30	48.1	41.4	48.2	47.6	49.3	45.6	48.7	43.9	45.3	49.2	48.1	45.0	-3.1
	35	40.9	39.4	38.5	42.2	39.6	39.2	39.2	35.4	35.4	40.3	40.4	40.6	+0.2
	40	—	41.0	41.9	39.4	37.5	39.4	38.7	37.9	41.1	38.4	37.6	39.2	+1.6
	45	—	—	—	—	—	—	35.8	39.8	39.3	37.1	38.3	36.8	-1.5
	50	—	—	—	—	—	—	—	—	—	—	—	32.8	—

(List of drugs continued.)

TABLE 7-4 (cont.)
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?	Age Group	Percentage saying "fairly easy" or "very easy" to get ^a																cont. →
		1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	
Crystal meth. (ice)	18	—	—	—	—	—	—	—	—	—	24.0	24.3	26.0	26.6	25.6	27.0	26.9	
	19–22	—	—	—	—	—	—	—	—	—	24.0	21.8	22.5	20.9	24.7	25.5	25.4	
	23–26	—	—	—	—	—	—	—	—	—	22.3	20.0	21.3	22.9	24.5	24.7	24.7	
	27–30	—	—	—	—	—	—	—	—	—	27.3	19.7	22.0	21.2	21.7	25.8	26.1	
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Sedatives/ barbiturates^c	18	49.1	54.9	55.2	52.5	51.9	51.3	48.3	48.2	47.8	48.4	45.9	42.4	44.0	44.5	43.3	42.3	41.4
	19–22	59.5	61.1	56.8	54.2	48.1	52.7	46.8	44.6	45.5	47.7	44.2	41.7	43.4	41.9	40.6	42.9	41.1
	23–26	—	—	—	—	52.7	47.7	46.4	45.9	47.4	44.8	41.6	39.6	42.0	38.8	40.3	42.1	40.6
	27–30	—	—	—	—	—	—	—	—	43.2	44.5	44.2	38.5	37.8	39.7	37.4	39.9	41.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers	18	59.1	60.8	58.9	55.3	54.5	54.7	51.2	48.6	49.1	45.3	44.7	40.8	40.9	41.1	39.2	37.8	36.0
	19–22	67.4	62.8	62.0	62.3	52.5	55.6	52.9	50.3	50.0	49.4	45.4	44.8	40.7	40.9	41.0	40.2	37.6
	23–26	—	—	—	—	60.2	54.3	54.1	56.3	52.8	51.4	47.8	45.1	48.1	43.2	45.9	44.3	42.3
	27–30	—	—	—	—	—	—	—	—	55.3	54.4	54.9	47.5	47.8	47.4	44.4	44.8	46.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids	18	—	—	—	—	—	—	—	—	—	—	46.7	46.8	44.8	42.9	45.5	40.3	
	19–22	—	—	—	—	—	—	—	—	—	—	44.1	44.8	46.3	41.7	40.9	41.8	40.8
	23–26	—	—	—	—	—	—	—	—	—	—	37.6	35.8	39.3	35.8	37.0	37.4	33.9
	27–30	—	—	—	—	—	—	—	—	—	—	36.4	30.6	35.0	31.6	30.5	33.1	35.6
	35	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Approximate Weighted N =</i>	18	3,240	3,578	3,602	3,385	3,269	3,274	3,077	3,271	3,231	2,806	2,549	2,476	2,586	2,670	2,526	2,552	2,340
	19–22	582	601	582	588	559	571	592	581	568	572	571	534	512	480	459	470	467
	23–26					540	541	548	539	526	514	532	511	523	500	463	449	418
	27–30									519	513	510	487	475	473	437	446	468
	35															1,142	1,141	1,146
	40																	
	45																	
	50																	

TABLE 7-4 (cont.)
Trends in Availability of Drugs as Perceived by
Respondents in Modal Age Groups of 18, 19–22, 23–26, 27–30, 35, 40, 45, and 50

Q. How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?	Age Group	Percentage saying "fairly easy" or "very easy" to get ^a												2007–2008 change
		1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Crystal meth. (ice)	18	27.6	29.8	27.6	27.8	28.3	28.3	26.1	26.7	27.2	26.7	25.1	23.3	-1.8
	19–22	29.3	31.0	31.8	27.4	28.4	31.2	26.5	27.1	28.9	29.1	27.7	24.1	-3.6
	23–26	25.8	30.2	28.5	25.8	26.4	25.1	26.4	32.3	27.8	32.3	27.8	27.7	-0.1
	27–30	25.1	22.6	29.1	25.3	27.6	29.5	30.9	25.5	27.4	31.8	29.7	31.4	+1.7
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Sedatives/ barbiturates^c	18	40.0	40.7	37.9	37.4	35.7	36.6	35.3	46.3	44.4	43.8	41.7	38.8	-2.9
	19–22	39.8	39.2	42.3	40.6	39.3	40.8	38.4	43.8	47.8	42.6	47.5	43.2	-4.3
	23–26	39.1	42.6	39.7	37.6	36.1	36.4	37.8	49.4	48.4	51.4	46.5	43.3	-3.2
	27–30	39.1	33.9	38.4	36.1	38.1	34.8	35.6	40.5	42.9	43.3	46.4	44.7	-1.7
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers	18	35.4	36.2	32.7	33.8	33.1	32.9	29.8	30.1	25.7	24.4	23.6	22.4	-1.2
	19–22	37.8	36.8	37.1	36.5	34.9	34.6	34.2	29.7	30.1	22.8	28.5	23.3	-5.2
	23–26	36.4	39.4	38.3	37.6	38.7	33.7	32.5	36.6	32.9	33.0	31.7	30.3	-1.4
	27–30	41.9	39.9	41.5	36.7	42.9	38.1	35.9	30.6	33.5	32.1	32.4	33.1	+0.7
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids	18	41.7	44.5	44.6	44.8	44.4	45.5	40.7	42.6	39.7	41.1	40.1	35.2	-4.9 ss
	19–22	39.2	39.2	40.5	40.3	38.1	41.4	39.4	37.8	37.6	37.1	37.9	33.5	-4.4
	23–26	35.5	34.9	37.1	34.0	34.7	33.1	31.1	34.7	31.2	34.2	33.3	30.2	-3.1
	27–30	32.5	30.5	34.5	36.2	34.6	33.0	32.6	30.6	32.4	29.7	30.9	31.0	+0.2
	35	—	—	—	—	—	—	—	—	—	—	—	—	—
	40	—	—	—	—	—	—	—	—	—	—	—	—	—
	45	—	—	—	—	—	—	—	—	—	—	—	—	—
	50	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Approximate</i>	18	2,517	2,520	2,215	2,095	1,850	2,138	2,391	2,169	2,161	2,161	2,420	2,276	
<i>Weighted N =</i>	19–22	463	433	425	400	398	375	386	441	392	376	362	380	
	23–26	419	395	415	388	401	362	356	411	359	335	338	355	
	27–30	459	425	424	365	357	349	368	393	359	347	324	334	
	35	1,150	1,032	1,022	981	977	890	934	963	1,009	925	863	898	
	40		1,029	1,093	1,096	1,065	1,037	898	967	928	919	868	881	
	45							911	1,026	1,005	972	954	851	
	50												902	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

"—" indicates data not available.

^aAnswer alternatives were: (1) Probably impossible, (2) Very difficult, (3) Fairly difficult, (4) Fairly easy, and (5) Very easy.

^bIn 2001 the question text was changed from "other psychedelics" to "other hallucinogens," and "shrooms" was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.

^cIn 2004 the question text was changed from "barbiturates" to "sedatives/barbiturates" and the list of examples was changed from "downers, goofballs, reds, yellows, etc." to just "downers." These changes likely explain the discontinuity in the 2004 results.

Chapter 8

PREVALENCE OF DRUG USE AMONG COLLEGE STUDENTS

We believe that one of the more important functions of the Monitoring the Future (MTF) study has been to track various forms of substance use among the nation's college students, in part because they have often been the harbingers of social and political changes that spread eventually to other segments of the population up and down the age spectrum. This was certainly the case for the epidemic of illicit drug use that emerged in the American population in the late 1960s.

MTF has been able to generate an unparalleled national sample of college students every year since 1980 by following representative samples of high school classes after they graduate. The graduating class of 1976 was the first such class followed after graduation, and by 1980 the survey included college students one to four years post-high school. The 2008 survey is thus the 29th such survey covering this important segment of the general population.

The absence of dropouts in the original high school senior samples has practically no effect on the representativeness of these college samples because very few dropouts go on to college. One notable limitation of the present design for the purpose of characterizing college students is that it limits the age range of the college sample. For trend estimation purposes, we decided to limit the age band to the most typical one for college attendance, that is, one to four years past high school, which corresponds to modal ages 19 to 22. According to the latest statistics available from the United States Census Bureau,⁵⁴ this age band should encompass about 67% of all undergraduate college students enrolled full-time in 2007, down some from the 79% covered in 1989. Although expanding the age band to include an additional two years would cover 76% of all enrolled college students, it would also reduce by two years the interval over which we could report trend data. Some special analyses conducted in 1985 indicated that the differences in *prevalence-of-use* estimates under the two definitions were extremely small. The annual prevalence of all drugs except cocaine shifted only about one or two tenths of a percent. Cocaine, which had the greatest amount of age-related change, would have had an annual prevalence rate only 0.8 percentage points higher if the six-year age span were included rather than the four-year age span. A replication of these analyses in 1997 yielded virtually the same results. Thus, for purposes of estimating all prevalence rates except lifetime prevalence, the four- and six-year intervals are nearly interchangeable, suggesting that this limitation is negligible for our purposes.

On the positive side, maintaining a consistent age band allows for trend estimation by controlling for changes in the age composition of college students over the years. Otherwise, college students characterized in one year might represent a noncomparable segment of the larger population when compared to college students surveyed in another year.

⁵⁴U.S. Census Bureau, October 2007. Available at <http://www.census.gov>.

Definition of college students. *College students are defined here as those follow-up respondents one to four years past high school who say they were registered as full-time students in a two- or four-year undergraduate college at the beginning of March in the year in question.* Note that students at two-year colleges, such as community colleges, are included. The definition excludes those who are currently enrolled in college part-time and those who previously may have been college students or may have completed college.

Prevalence-of-use rates for college students, as well as their same-age peers who are also high school graduates, are provided in Tables 8-1 to 8-4. Having statistics for both groups, a unique feature of the MTF panels, makes it possible to see whether college students' substance use rates are higher or lower than those of their age peers (one to four years past high school, i.e., of modal ages 19 to 22). The college-enrolled sample now constitutes well over half (62%) of the entire follow-up sample one to four years past high school. The differences reported here pertain to those who are full-time college students versus those who are not, *among high school graduates*. If data from the missing high school dropout segment were available for inclusion as part of the noncollege segment, any difference between the two groups in terms of their substance use would likely be enlarged; therefore, any differences observed here are only an indication of the direction and relative size of differences between the college and the *entire* noncollege population, not an absolute estimate of them.

PREVALENCE OF DRUG USE AMONG COLLEGE STUDENTS VERSUS THOSE NOT ENROLLED IN COLLEGE

In 2008, lifetime prevalence of use among college students is lower for all illicit drugs compared with use among their age peers, but the degree of difference varies considerably by drug, as Table 8-1 shows. Lifetime use of alcohol is higher among college students. Annual rates of use of *LSD*, *Ritalin*, *Rohypnol*, and *alcohol* are higher among college students than among those not enrolled in college (see Tables 8-2 and 8-3).

- In 2008, the annual prevalence of use for *any illicit drug* is similar among college students (35%) and high school graduates not in college (37%), but a considerably higher proportion of noncollege respondents have used *any illicit drug other than marijuana* in the past year (23%) than college students (15%).
- Annual *marijuana* use is very close among college students (32%) and high school graduates of the same age who are not in college (33%). The rate of current *daily marijuana* use, however, is considerably lower among college students (3.9% versus 7.0%) (see Table 8-4 for the prevalence of current daily use).
- A number of drugs show annual use rates for noncollege respondents that are two or more times those for college students, including *inhalants*, *cocaine*, *crack cocaine*, *other cocaine*, *heroin*, *heroin with a needle*, *heroin without a needle*, *methamphetamine*, *crystal methamphetamine (ice)*, *sedatives (barbiturates)*, and *steroids*.

- It is clear that use of a number of the *illicit drugs other than marijuana* tends to be distinctly higher among those not in college. As previously noted, such differences would likely be still larger if the noncollege group included high school dropouts.
- **Ritalin**, a drug in the amphetamine class that was added to the MTF questionnaires in 2002, has typically shown a different pattern, with use higher among college students than among those not in college, quite possibly explained by college students using Ritalin to stay awake late at night to finish assignments or study for tests. This pattern continued in 2008, with college students' annual use at 3.2% and the noncollege group at 2.1%. Use was particularly low among noncollege females—0.8% compared to 3.8% among female college students.
- In 2008, college students were modestly higher in lifetime, annual, and 30-day use of **alcohol** than the noncollege group, with the difference being largest in the 30-day rate (69% vs. 55%).
- College students had a significantly higher prevalence of **occasions of heavy drinking** (five or more drinks in a row in the past two weeks)—40% versus 30% among their age peers. *Indeed, it is noteworthy that nearly half of all college students (45%) report having been drunk in the prior 30 days, compared to 31% of the noncollege respondents.* Rates of **daily drinking** were much closer (4.0% vs. 3.7%; see Table 8-4). In high school, college-bound students, especially in earlier grades, were far less likely to drink alcohol at any level compared to their non-college-bound peers; thus, the relative and absolute increases in alcohol use in the first few years following high school are quite striking for college students.
- Because of increasing attention being paid to the problem of **extreme binge drinking**, in 2005 we introduced a set of questions on the subject into one of the six questionnaire forms used with young adults, including college students. The questions ask the respondent about the frequency in the past two weeks of their having (a) four or more drinks in a row, (b) five or more drinks in a row, (c) 10 or more drinks in a row, and (d) 15 or more drinks in a row. The low *Ns* resulting from a single questionnaire form necessitate combining four years of data (2005–2008), and even then only 877 weighted cases are available from the college student population and 511 for their noncollege peers. However, they give us an idea of the prevalence and frequency of these binge drinking measures.

About one in nine college students (11%) reported having **10 or more drinks in a row** at least once in the prior two weeks, and 5% (or 1 in 20) reported **15 or more drinks in a row**. Their noncollege peers had about the same rates (13% and 6%, respectively). While we lack earlier data on these measures to determine whether this type of extreme drinking behavior has changed over time, it is clear that it is quite high today among both college students and their noncollege age peers. As will be discussed below, there is a major difference between males and females in the prevalence of these behaviors.

- In 2008, nearly two thirds (65%) of college students reported using *flavored alcoholic beverages* in the prior year, compared to a lower rate for the noncollege group (51%).
- Among all substances studied, except for alcohol, the largest absolute differences in 30-day and daily prevalence rates between the two groups occur for *cigarette smoking*. For example, the prevalence of daily smoking for college students is 9% versus 25% for their age-mates not enrolled full-time in college. Smoking at the rate of a half pack per day stands at 4% versus 15% for these two groups, respectively. Recall that the 12th-grade data show the college-bound have much lower smoking rates in high school than the non-college-bound; thus, in contrast to what was true for alcohol use, these substantial differences observed at college age actually preceded college attendance.⁵⁵ The smoking differences would likely be even greater if dropouts were included in the noncollege group, because dropouts have consistently shown an exceptionally high rate of smoking.⁵⁶
- The prevalence of using *steroids* in the past year is higher in the noncollege segment (1.3% compared to 0.1% among college students).

In sum, the noncollege segment is generally more drug-involved than the college student segment. This pattern is a continuation of the high school scenario in which those without college plans are more likely to use drugs. The only substance for which college students are significantly and substantially more likely to be users is *alcohol* (particularly *getting drunk* and *binge drinking*), and this difference emerges primarily after high school.

GENDER DIFFERENCES IN PREVALENCE OF USE AMONG COLLEGE STUDENTS

Tabular data are provided separately in Tables 8-1 to 8-4 for male and female college students and their same-age peers.

- Most of the gender differences *among* college students replicate those discussed earlier for all young adults 1 to 12 years past high school, and they in turn replicate gender differences among secondary school students for the most part. That means that among college students, males have higher annual prevalence rates for most of the illicit drugs. The annual prevalence rates for use of *any illicit drug* are 39% for males versus 33% for females; for *any illicit drug other than marijuana*, 18% versus 14%; for *marijuana*, 37% versus 29%; and for *narcotics other than heroin*, 8.2% versus 5.5%. Some drugs show a larger difference in the same direction: *hallucinogens*, 8.2% versus 3.0%; *LSD* specifically, 3.7% versus 1.8%; and *hallucinogens other than LSD*, 7.4% versus 2.4%.

⁵⁵See also Bachman, J. G., Wadsworth, K. N., O'Malley, P. M., Johnston, L. D., & Schulenberg, J. E. (1997). *Smoking, drinking, and drug use in young adulthood: The impacts of new freedoms and new responsibilities*. Mahwah, NJ: Lawrence Erlbaum Associates.

⁵⁶For a recent analysis showing much higher smoking rates among 8th graders who later dropped out before completing high school, see Bachman, J. G., O'Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education-drug use connection: How successes and failures in school relate to adolescent smoking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

- Last year the gender difference was particularly large for **Vicodin**—10.0% for males versus 4.8% for females—but in 2008 the genders showed virtually the same usage rate (6.7% vs. 6.6%).
- Annual prevalence of **marijuana** use does not differ significantly between males and females among college students, nor among the noncollege respondents.
- **Daily marijuana** use is much higher among male college students compared to female college students (7.3% versus 1.7%). There is considerably less difference among the noncollege group, with 8.3% of males reporting daily use versus 6.1% of females.
- **Amphetamine** use is a little higher among college males (7.2%) than college females (4.8%), but somewhat lower among males in the noncollege segment (6.3%) than among females (7.7%).
- College males and females report a similar prevalence of **getting drunk** in the prior 30 days (48% and 44%); noncollege respondents report lower rates, 33% for males and 29% for females. Both male and female college students have higher rates of **occasional binge drinking** than their counterparts not in college (34% for college females versus 28% for noncollege females and 49% versus 32% for males, respectively) (see Table 8-4). More **extreme levels of binge drinking**, however, show a large gender difference in both groups. Among college students, the prevalence of having **10 or more drinks in a row** in the prior two weeks was 6% for college females versus 25% for college males for 2005–2008 combined; similar rates were reported by noncollege respondents, 7% for females and 23% for males. The prevalence of having 15 or more drinks in a row was 1.0% for college females versus 11% for college males; again, similar rates were reported by noncollege respondents, 2% for females and 12% for males.
- **Flavored alcoholic beverages** are consumed by more females than males, regardless of whether they attend college (71% of females versus 56% of males in college reporting past-year use, and 59% versus 43%, respectively, for the noncollege group).
- Among college students, 30-day prevalence of **cigarette smoking** is slightly higher for males (20%) than for females (17%), and the same holds among the noncollege segment (33% and 30%, respectively). **Daily smoking** is reported by a slightly higher proportion of males than females in the college segment (9.8% versus 8.8%); whereas among the noncollege segment both genders have the same prevalence (25%). Rates of **smoking a half pack or more per day** are 4.3% and 4.2% for males and females among college students, and 16% and 15% for the noncollege segment.

TABLE 8-1
Lifetime Prevalence of Use for Various Types of Drugs, 2008:
Full-Time College Students vs. Others
among Respondents 1 to 4 Years beyond High School
(Entries are percentages.)

	Total		Males		Females	
	Full-Time		Full-Time		Full-Time	
	<u>College</u>	<u>Others</u>	<u>College</u>	<u>Others</u>	<u>College</u>	<u>Others</u>
Any Illicit Drug ^a	49.5	58.5	50.7	55.6	48.8	60.7
Any Illicit Drug ^a other than Marijuana	22.6	35.9	25.2	32.7	21.0	38.3
Marijuana	46.8	55.2	49.2	53.6	45.3	56.3
Inhalants ^b	4.9	9.5	5.4	9.7	4.7	9.3
Hallucinogens	8.5	12.1	13.4	13.4	5.3	11.1
LSD	4.3	5.3	6.6	5.9	2.8	4.9
Hallucinogens other than LSD	8.2	11.1	12.8	12.5	5.1	10.0
Ecstasy (MDMA) ^b	6.2	13.3	7.0	12.9	5.6	13.6
Cocaine	7.2	16.6	9.5	16.2	5.7	16.9
Crack ^c	1.4	6.4	1.7	6.9	1.2	6.0
Other Cocaine ^d	7.1	16.1	9.1	15.7	5.8	16.5
Heroin	0.7	3.0	1.1	3.6	0.3	2.6
With a Needle ^e	*	0.8	*	0.8	*	0.8
Without a Needle ^e	0.7	2.5	1.2	3.8	0.3	1.4
Narcotics other than Heroin ^f	12.4	20.0	14.5	18.3	11.1	21.3
Amphetamines, Adjusted ^{f,g}	9.1	16.6	10.4	15.8	8.3	17.2
Methamphetamine ^e	1.9	4.0	1.5	4.0	2.2	4.1
Crystal Methamphetamine (Ice) ^e	1.1	4.7	1.6	3.1	0.8	5.9
Sedatives (Barbiturates) ^f	6.4	13.7	7.4	12.9	5.7	14.2
Tranquilizers ^f	8.6	16.8	10.2	15.2	7.6	18.0
Alcohol	85.3	81.7	85.6	78.4	85.1	84.3
Been Drunk ^b	72.5	69.6	73.7	64.8	71.7	73.4
Flavored Alcoholic Bvg. ^h	78.6	77.0	75.9	63.9	80.5	89.8
Cigarettes	—	—	—	—	—	—
Steroids ^e	1.6	1.3	3.2	2.2	0.6	0.6
<i>Approximate Weighted N =</i>	<i>1,270</i>	<i>780</i>	<i>510</i>	<i>330</i>	<i>760</i>	<i>450</i>

Source. The Monitoring the Future study, the University of Michigan.

“*” indicates a prevalence rate of less than 0.05%.

“—” indicates data not available.

See footnotes on next page.

Footnotes for Tables 8-1 through 8-4

^aUse of “any illicit drug” includes any use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor’s orders.

^bThis drug was asked about in three of the six questionnaire forms. Total *N* in 2008 for college students is approximately 630.

^cThis drug was asked about in five of the six questionnaire forms. Total *N* in 2008 for college students is approximately 1,060.

^dThis drug was asked about in four of the six questionnaire forms. Total *N* in 2008 for college students is approximately 850.

^eThis drug was asked about in two of the six questionnaire forms. Total *N* in 2008 for college students is approximately 420.

^fOnly drug use that was not under a doctor’s orders is included here.

^gBased on the data from the revised question, which attempts to exclude inappropriate reporting of nonprescription amphetamines.

^hThis drug was asked about in one of the six questionnaire forms. Total *N* in 2008 for college students is approximately 210.

ⁱDaily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes, for which actual daily use is measured, and for 5+ drinks, for which the prevalence of having five or more drinks in a row in the last two weeks is measured.

TABLE 8-2
Annual Prevalence of Use for Various Types of Drugs, 2008:
Full-Time College Students vs. Others
among Respondents 1 to 4 Years beyond High School
(Entries are percentages.)

	Total		Males		Females	
	Full-Time College	Others	Full-Time College	Others	Full-Time College	Others
Any Illicit Drug ^a	35.2	37.1	38.7	36.8	32.9	37.4
Any Illicit Drug ^a other than Marijuana	15.3	22.5	17.8	22.0	13.7	22.8
Marijuana	32.3	33.1	37.0	33.0	29.1	33.2
Inhalants ^b	1.1	2.5	1.5	2.3	0.9	2.7
Hallucinogens	5.1	5.7	8.2	6.5	3.0	5.1
LSD	2.6	1.9	3.7	2.4	1.8	1.5
Hallucinogens other than LSD	4.4	5.1	7.4	6.0	2.4	4.5
Ecstasy (MDMA) ^b	3.7	6.3	4.4	4.8	3.3	7.6
Cocaine	4.4	9.0	6.4	9.3	3.0	8.7
Crack ^c	0.5	2.0	0.5	1.9	0.5	2.1
Other Cocaine ^d	4.2	8.7	6.0	9.0	3.0	8.6
Heroin	0.3	1.5	0.7	1.3	0.1	1.6
With a Needle ^e	*	0.3	*	*	*	0.6
Without a Needle ^e	0.3	0.8	0.9	1.1	*	0.5
Narcotics other than Heroin ^f	6.5	11.7	8.2	11.8	5.5	11.7
OxyContin ^{e,f}	3.6	6.5	4.9	6.9	2.7	6.2
Vicodin ^{e,f}	6.7	13.0	6.7	13.5	6.6	12.5
Amphetamines, Adjusted ^{f,g}	5.7	7.1	7.2	6.3	4.8	7.7
Ritalin ^{e,f}	3.2	2.1	2.4	3.6	3.8	0.8
Methamphetamine ^e	0.5	1.2	0.4	0.8	0.5	1.5
Crystal Methamphetamine (Ice) ^e	0.1	1.1	*	1.5	0.1	0.7
Sedatives (Barbiturates) ^f	3.7	7.4	4.0	6.3	3.5	8.2
Tranquilizers ^f	5.0	9.4	5.8	8.7	4.5	9.9
Rohypnol ^e	0.3	0.1	*	*	0.4	0.3
GHB ^e	0.2	0.2	0.3	*	0.2	0.3
Ketamine ^e	0.4	0.3	0.9	0.8	*	*
Alcohol	82.1	77.0	82.8	74.8	81.7	78.6
Been Drunk ^b	66.8	55.5	68.9	52.1	65.3	58.2
Flavored Alcoholic Beverage. ^h	65.0	50.9	56.3	43.2	70.8	58.5
Cigarettes	30.0	41.9	30.5	44.2	29.7	40.1
Steroids ^e	0.1	1.3	0.2	2.2	*	0.6
<i>Approximate Weighted N =</i>	<i>1,270</i>	<i>780</i>	<i>510</i>	<i>330</i>	<i>760</i>	<i>450</i>

Source. The Monitoring the Future study, the University of Michigan.

“*” indicates a prevalence rate of less than 0.05%.

See relevant footnotes at the end of Table 8-1.

TABLE 8-3
Thirty-Day Prevalence of Use for Various Types of Drugs, 2008:
Full-Time College Students vs. Others
among Respondents 1 to 4 Years beyond High School
(Entries are percentages.)

	Total		Males		Females	
	Full-Time College	Others	Full-Time College	Others	Full-Time College	Others
Any Illicit Drug ^a	18.9	22.8	23.1	22.1	16.2	23.3
Any Illicit Drug ^a other than Marijuana	7.3	12.0	9.6	10.4	5.8	13.3
Marijuana	17.0	19.2	22.1	19.2	13.6	19.2
Inhalants ^b	0.4	0.6	0.5	0.8	0.3	0.4
Hallucinogens	1.7	1.1	3.0	1.7	0.8	0.6
LSD	0.8	0.4	1.2	0.6	0.5	0.2
Hallucinogens other than LSD	1.3	0.9	2.6	1.6	0.4	0.4
Ecstasy (MDMA) ^b	0.6	1.8	0.6	1.6	0.6	1.9
Cocaine	1.2	3.0	2.0	2.5	0.6	3.3
Crack ^c	0.1	0.8	0.3	0.4	*	1.1
Other Cocaine ^d	1.1	3.1	1.9	2.8	0.6	3.3
Heroin	*	0.3	0.1	0.3	*	0.3
Narcotics other than Heroin ^f	2.3	5.5	3.0	4.8	1.9	6.0
Amphetamines, Adjusted ^{f,g}	2.8	2.7	4.0	2.0	1.9	3.3
Crystal Methamphetamine (Ice) ^e	*	0.1	*	0.3	*	*
Sedatives (Barbiturates) ^f	1.4	2.8	1.6	2.4	1.3	3.0
Tranquilizers ^f	1.6	4.6	1.8	4.3	1.4	4.9
Alcohol	69.0	55.0	71.2	54.4	67.5	55.5
Been Drunk ^b	45.3	30.8	47.9	32.7	43.5	29.4
Flavored Alcoholic Beverages ^h	35.8	28.1	28.9	20.1	40.4	36.0
Cigarettes	17.9	31.4	20.0	32.8	16.6	30.3
<i>Approximate Weighted N =</i>	<i>1,270</i>	<i>780</i>	<i>510</i>	<i>330</i>	<i>760</i>	<i>450</i>

Source. The Monitoring the Future study, the University of Michigan.

“*” indicates a prevalence rate of less than 0.05%.

See relevant footnotes at the end of Table 8-1.

TABLE 8-4
Thirty-Day Prevalence of Daily Use for Various Types of Drugs, 2008:
Full-Time College Students vs. Others
among Respondents 1 to 4 Years beyond High School
(Entries are percentages.)

	Total		Males		Females	
	Full-Time College	Others	Full-Time College	Others	Full-Time College	Others
Marijuana ⁱ	3.9	7.0	7.3	8.3	1.7	6.1
Cocaine ⁱ	*	*	0.1	0.1	*	*
Amphetamines, Adjusted ^{i,f,g}	0.2	0.3	0.1	0.2	0.2	0.3
Alcohol						
Daily ⁱ	4.0	3.7	5.1	5.0	3.3	2.7
5+ Drinks in a Row in Last 2 Weeks	40.0	29.9	48.6	32.3	34.4	28.0
Cigarettes						
Daily	9.2	24.7	9.8	24.5	8.8	24.9
1/2 Pack+/Day	4.3	15.2	4.3	15.6	4.2	15.0
<i>Approximate Weighted N =</i>	<i>1,270</i>	<i>780</i>	<i>510</i>	<i>330</i>	<i>760</i>	<i>450</i>

Source. The Monitoring the Future study, the University of Michigan.

"*" indicates a prevalence rate of less than 0.05%.

See relevant footnotes at the end of Table 8-1.

Chapter 9

TRENDS IN DRUG USE AMONG COLLEGE STUDENTS

College students often initiate social change, and such was the case with the illicit drug epidemic. Illicit drug use increased dramatically among American college students in the mid-1960s, then spread quickly to their noncollege age peers and eventually down the age spectrum to high school and even middle school students. As illustrated in this chapter, the diffusion process seems to have reversed during the more recent “relapse” of the epidemic in the 1990s, when use increased first among those in early adolescence and then radiated *up* the age spectrum as those cohorts grew older. Use has subsequently declined among adolescents; this decline, like the recent increase, is radiating up the age spectrum.

Again, we define *college students* as high school graduates one to four years past high school who are enrolled full-time in a two- or four-year college at the beginning of March in the year in question. For comparison purposes, trend data are provided on the remaining follow-up respondents in this age band, who are also one to four years past high school (see Figures 9-1 through 9-15c). Because the proportion of an age group in college declines steadily with the number of years beyond high school, this comparison group is slightly older on average than the college-enrolled group. Also, the proportion of young adult high school graduates one to four years beyond high school who are enrolled full-time in college has increased considerably since the MTF follow-ups began. In 2008, about 62% of the weighted number of follow-up respondents one to four years past high school met our definition of college students, compared with only 38% in the 1980 survey.

The reader is reminded that the difference between the college group and the other group provides an estimate of the degree to which college students are above or below average for other high school *graduates* in this age band. Were we able to include the high school dropout segment in the calculation for the noncollege group, many of the differences with the college-enrolled would be accentuated.⁵⁷

For each year, there are approximately 1,100–1,500 weighted respondents constituting the college student sample (see Table 9-5 for *Ns* per year) and roughly 700–1,700 respondents constituting the noncollege group among those one to four years beyond high school. Trend comparisons for these two groups are provided in this chapter. Because it was not until 1980 that enough follow-up years had accrued to characterize young people one to four years past high school, the comparisons begin with that year.

In some of the discussion that follows, as well as a number of the figures, trends for the 12th-grade samples are included for comparison purposes. It is important to keep in mind that the total 12th-grade samples are shown, and that there are substantial differences in rates of substance use

⁵⁷Panel analyses of samples from the high school classes of 1995–1997, followed for an eight-year period beginning when they were in 8th grade, show clearly that those who dropped out of high school had distinctly higher rates of substance use both before and after they left school. See Bachman, J. G., O’Malley, P. M., Schulenberg, J. E., Johnston, L. D., Freedman-Doan, P., & Messersmith, E. E. (2008). *The education–drug use connection: How successes and failures in school relate to adolescent smoking, drinking, drug use, and delinquency*. New York: Lawrence Erlbaum Associates/Taylor & Francis.

within those samples between the college-bound and those who do not plan to complete college. As shown extensively in *Volume I*, 12th-grade students expecting to complete college are far less likely to smoke cigarettes, and also less likely to use most other substances. So when considering figures that show higher rates of use among 12th graders than among college students, it should not be concluded that usage declined after college entrance; the more likely interpretation is that the college-bound were already lower in usage rates than other 12th graders.

TRENDS IN PREVALENCE, 1980–2008: COLLEGE STUDENTS VERSUS THOSE NOT ENROLLED IN COLLEGE

- The proportion of college students using *any illicit drug* in the 12 months prior to the survey dropped fairly steadily between 1980 and 1991—from 56% to 29%, a decrease of nearly half (see Table 9-2). After 1991, annual (and also 30-day) prevalence held fairly steady for a couple of years before beginning to rise, reaching 38% in 1998 and again in 2001—still well below the peak of 56% in 1980. There has not been a great deal of change since then, although some decline has occurred recently (the rate was 35% in 2008). The noncollege group moved similarly. Twelfth graders also showed a similar trajectory in the decline through 1991 and the leveling after 1998, but their rise in use in 1992–1997 was distinctly sharper, as Figure 9-1 illustrates. In 2000, the noncollege group exhibited a four-percentage-point increase that was due largely to their sharper increases in marijuana, amphetamine, and tranquilizer use; their level remained above the college student sector for several years, until they showed a similar-sized decline in 2007, just about eliminating the gap. Use among high school seniors has declined some since 1999 (by about six percentage points), whereas among college students there was little decline until 2006, when the rate fell nearly three percentage points. As a result, all three groups have quite similar prevalence rates in 2008. We believe the divergence among the three groups and subsequent convergence of college students and high school seniors, at least, reflect cohort effects.
- Use of *any illicit drug other than marijuana* declined fairly steadily and appreciably among college students between 1980 and 1994, with annual prevalence dropping by nearly two thirds from 32% to 12% (Table 9-2). This generally paralleled the trends for the noncollege group and the 12th graders. All three groups showed some increase in use during the early 1990s; however, the rise in use of illicit drugs other than marijuana was not as sharp among college students (Figure 9-2). After 1999, as use among 12th graders leveled off, the college students and noncollege segment showed some further increase. In fact, the college students and noncollege respondents continued to show an increase in their annual prevalence rate from 1998 through 2004, before declining from 2005 through 2007 among the noncollege group and from 2005 through 2008 among the college students. College students and 12th graders now have comparable prevalence rates, following more than a decade in which the 12th graders had higher rates. Again, this divergence and then convergence most likely reflect some cohort effects working their way up the age spectrum. Of the three groups, the noncollege group has shown the highest rates of using illicit drugs other than marijuana since 2000.

- Trends during the 1980s for most individual classes of illicit drugs tended to be similar among the three groups. During the 1990s, however, there was more divergence in the trends, with college students usually showing less increase than the 12th graders, and, for some drugs, less increase than their age-mates not in college.
- The annual prevalence of *marijuana* use among college students decreased steadily from 1981 through 1991, dropping by nearly half from 51% to 27% (Figure 9-3a). Their noncollege peers showed a comparable decline over the same time interval, as did the 12th graders. Use among 12th graders rose sharply after 1992, while use among college students and noncollege respondents rose more gradually. From 1991 through 1998, annual prevalence rose by nearly 10 percentage points among college students, by 7 percentage points among the noncollege group, but by 14 percentage points among 12th graders. As a result, the 12th graders came to exhibit the highest rate of marijuana use in the last half of the 1990s. The 12th graders were the first to show a leveling off in marijuana use (in 1998), followed by the college students in 1999 and the noncollege group in 2002. All three groups had very similar rates of use in 2005. The college students and high school seniors both showed some decline in 2006, followed by some increase in their marijuana use from 2006 through 2008. This may indicate that the gradual improvement in marijuana use we found earlier in the decade is over.
- *Daily marijuana* use among college students (Figure 9-3b) fell appreciably between 1980 and 1986, from 7.2% to 2.1%, as it did for noncollege respondents and 12th graders. (The latter two groups were able to show sharper declines because they started higher than college students in 1980. This sharper decline nearly eliminated the differences by 1989.) After 1986, the decline decelerated, and by 1991 the rate stood at 1.8% among college students. In sum, the proportion of American college students who actively smoked marijuana on a daily basis dropped by about three fourths between 1980 and 1991. Daily use then leveled until 1994 and began increasing thereafter, reaching 4.6% in 2000. By 2008, daily use among college students was slightly lower, at 3.9%. The other two groups showed considerably larger increases after 1993, with 12th graders' daily use rates leveling after 2000 and declining after 2004. The noncollege segment showed further increase in 2001, reaching 9.4%; their use was down to 7.0% by 2008, still well above college students. Of the three groups, the college students have had the lowest rate of daily marijuana use throughout the MTF study, and the noncollege segment the highest. As is often the case, these subgroup differences have narrowed considerably during overall declines in use and widened during periods of increasing use.
- A dramatic decline occurred for *amphetamine* use between 1981 and 1991 (Figure 9-11). Annual prevalence among college students dropped by more than four fifths, from 22% in 1981 to 4% in 1991. Proportionately, this was a larger drop than among 12th graders, but fairly parallel to the overall change among the noncollege group. Amphetamine use among college students and their noncollege age peers leveled for a year before beginning to increase in both groups after 1992 and 1993, respectively, through 2001, with a leveling in 2002. During the 1990s and early 2000s, the prevalence rates for amphetamines in all three groups have remained well below the rates observed in the early 1980s. Since 2002 there have been some small nonparallel changes among the three

groups, with amphetamine use among college students (who have consistently had the lowest rate of use since the mid-1980s) holding steady through 2007, while use among 12th graders and the noncollege group declined. In 2008, use among the college students also showed some decline, and the prevalence rates are now very similar across the three groups.

- Use of *inhalants* has been very low among college and noncollege respondents since 1980, when rates were first measured (Figure 9-4). Twelfth graders have consistently had higher rates of inhalant use than either of these segments of the young adult population. All three groups have trended in parallel, though, with an increase in use from around 1981 through 1995, followed by a long decline thereafter. The increase and decline were more pronounced among 12th graders.
- During the early 1980s, one of the largest proportional declines observed among college students was for *LSD* (see Figure 9-6). Annual prevalence fell from 6.3% in 1982 to 2.2% in 1985. After 1985, their use began to increase, reaching 6.9% by 1995. After 1995, use fell gradually among college students, their age-mates, and 12th graders. In 2002 there was a particularly sharp decrease in all groups, resulting in a considerable convergence in usage rates. All three groups now have annual prevalence rates between 1.3% and 2.1%. College students maintained lower levels of use than the other two groups from the mid-1990s through 2007. Use rose slightly in all three groups in 2008.
- The use of *ecstasy (MDMA)* by American college students and their noncollege age peers began to rise after 1994 (Figure 9-8). After 1997 there was a sharp increase in the college group. Their annual prevalence rose three- to fourfold in just three years, from 2.4% in 1997 to 9.2% in 2001; it then fell to 2.2% by 2004, and is at 3.7% in 2008. The trends among the noncollege segment have run fairly parallel to those for college students and 12th graders through 2008, except that use did not continue to rise among 12th graders in 2008 as in the other two groups. The noncollege young adults had a later start in the sharp increase in ecstasy use (after 1999), and they eventually reached a higher level of use. From 2000 through 2006, the noncollege segment exhibited the highest rate of ecstasy use—reaching 14% in 2001, when use among college students and 12th graders was at 9%. All three groups showed sharp declines in the following three years.
- When the college data were first available in 1980, *sedative (barbiturate)* use was already quite low among college students (at 2.9% annual prevalence) (see Figure 9-12), but it still fell by more than half to 1.3% by 1985. This proportional decline was sharper than among 12th graders and less sharp than among the noncollege respondents, both of whom started at a considerably higher level of use. Annual prevalence remained essentially unchanged between 1985 and 1993 for all three groups. All groups then showed a gradual increase in use between 1993 (or 1994 in the case of the college students) and 2001, with 12th graders showing a significant increase in 2002 and use in the other two groups leveling off. The college students showed a fairly steady increase over the 10-year period 1994–2004, with pauses in 1998 and 2002 and then a leveling in 2005; the other two groups remained at higher levels than the college students throughout this period. After

2005, modest declines in use appeared in all three groups, but the college students and their noncollege peers showed some upturn in 2008.

- Figure 9-13 shows that the annual prevalence of *tranquilizer* use among college students dropped by nearly three fourths in 1980–1994, from 6.9% to 1.8%. After this long period of decline, tranquilizer use then began to increase gradually, returning to 6.9% by 2003. Use by the noncollege segment and by 12th graders dropped more sharply, eliminating the differences among the three groups by 1992. Use rose after 1992 for all but the college students, again creating some differences, and then rose in all three groups after 1994; in 2002, tranquilizer use was once again at or near its recent high in all three groups. From 1999 through 2002 the increase in use was particularly sharp among the noncollege segment, making them the highest using group. In 2003, however, the noncollege group and the 12th graders showed their first declines in recent years, thus narrowing the differences among the three groups. From 2004 to 2007, all three groups showed modest declines in use; but in 2008 the decline continued only among the college students.
- The overall trends in use of *narcotics other than heroin*⁵⁸ have been quite parallel to those for sedatives (barbiturates) and tranquilizers. By 1994 the use of narcotics other than heroin (Figure 9-10a) by college students was about half what it was in 1980 (2.4% in 1994 versus 5.1% in 1980) as a result of a fairly gradual decline over that 14-year interval. This trend closely parallels use among their noncollege counterparts and 12th graders. As with a number of other drugs, use among 12th graders began to rise after 1992, but use among college students did not begin to increase until after 1994, likely due to a cohort effect. In 2003, annual prevalence among college students reached an historic high point of 8.7% before leveling for three years. It then declined some from 8.8% in 2006 to 6.5% by 2008. Use among 12th graders leveled after reaching an historic high in 2004 of 9.5% (9.1% in 2008). The noncollege group emerged after 2000 as the heaviest using group for the first time, as their use kept increasing through 2005, reaching an all-time high of 13%. In 2006 and 2007 their use declined, but the decline did not continue in 2008; their use remains higher than the college group.
- Although data were not collected until 2002, it is clear that *OxyContin* and *Vicodin* (Figures 9-10b and 9-10c and Table 8-2) help to explain the difference between the college and noncollege segments in their use of narcotics other than heroin. The noncollege group has had annual prevalence rates up to twice that for the college students in the use of both drugs. Annual prevalence of *OxyContin* use among college students rose fairly steadily, from 1.5% in 2002 to 3.6% in 2008. Use in the noncollege segment also rose from 2002 (3.3%) to 2005 (6.2%), dipped in 2006–2007, but is back to 6.5% in 2008. Among 12th graders, it also rose from 2002 (4.0%) to 2005 (5.5%), then falling to 4.7% in 2008. The college segment remains lower than the other two groups. *Vicodin* use showed a somewhat different pattern, with annual prevalence among all three groups remaining fairly level—and substantially higher—from 2002 through 2008. As with *OxyContin*, the noncollege segment has had consistently higher *Vicodin* use than the

⁵⁸As discussed in chapters 4 and 5, because the questions about narcotics other than heroin were changed in 2002, the prevalence figures are adjusted estimates. See the earlier discussion for details.

college students. The 12th graders have been between the two young adult groups in their rate of Vicodin use. Because of the limited numbers of cases, the trends in use of these two drugs have generally been uneven in the young adult groups.

- Like the 12th graders, college students showed a relatively stable pattern of *cocaine* use between 1980 and 1986, when their usage levels (and those of their noncollege age peers) were considerably higher than those observed among 12th graders (see Figure 9-9). This level period was followed by a dramatic drop of nearly nine tenths in annual prevalence among college students, from 17.1% in 1986 to 2.0% in 1994. Their noncollege counterparts also showed a large but somewhat less dramatic decline, from 18.9% in 1986 to 5.1% in 1994. Because use among college students also dropped more sharply than among 12th graders, there was little or no difference between those two groups in annual prevalence rates for cocaine use between 1990 and 1995. After 1995, cocaine use rose the least among the college students, creating a reversal of the previous gap, with the 12th graders having higher usage levels than the college students; this lasted for a few years before the college students caught up. Between 1994 and 1998, annual cocaine prevalence for college students increased significantly, from a 14-year low of 2.0% in 1994 to 4.5% by 1998, roughly where it stayed through 2002. Their use then showed a gradual rise after 2002, with annual prevalence increasing from 4.8% to 6.6% in 2004; it then dropped off some to 4.4% by 2008. Twelfth graders and the noncollege segment also exhibited an increase in annual prevalence of cocaine use after 1992 and 1993, respectively. Use was level among 12th graders from 2000 to 2007, but continued to increase among those not in college between 1999 and 2006, considerably widening the gap between the noncollege segment and the other two groups. In 2008, use among college students finally showed some decline in parallel with the decline among 12th graders.
- College students have shown some unique shifts in *alcohol* use. As can be seen in Figure 9-14d, both the noncollege segment and 12th graders showed fairly substantial declines from 1981 through 1990 in the prevalence of having *five or more drinks in a row* at least once during the prior two weeks. (The 12th graders then showed further decline for three more years, bringing them below the noncollege segment.) In contrast, college students showed no decline in binge drinking from 1981 to 1986, and then only a modest decline of five percentage points from 1986 through 1993. Between 1981 (when all three populations were very close in use) and 1992, this measure of heavy drinking dropped by 14 percentage points among 12th graders, by 11 percentage points among noncollege 19- to 22-year-olds, but by only 2 percentage points among full-time college students of the same age. After 1992, binge drinking began to rise among 12th graders, while still declining some among college students—likely reflecting a cohort effect emerging during this period, similar to that observed for a number of illicit drugs—narrowing the gap somewhat. Binge drinking subsequently began to increase among the noncollege segment after 1995, and by less among college students after 1996—increases that continued into 2001. Since 2001, college students have held fairly steady in their rates of binge drinking, while the noncollege segment has shown some decline. Meanwhile, among 12th graders, binge drinking started a gradual decline after 1998 that continued through 2006, enlarging the difference between them and the other two groups. Once again there is

evidence of cohort effects at work here since the beginning of the 1990s, with the inflection points being later for the older strata. Despite the different patterns of trends, perhaps the most noteworthy facts are that college students have exhibited the highest level and greatest constancy in binge drinking throughout the entire 29-year interval that college student samples have been included in MTF. Over that interval, from 1980 to 2008, college students' rates of binge drinking declined only 4% (from 44% to 40%), while noncollege respondents' rates declined 11% (41% to 30%) and high school seniors' rates declined by 16% (41% to 25%).

- It is interesting to conjecture why college students did not show much decline in heavy drinking for a decade (1981–1991) while their noncollege peers and 12th graders did. One possibility is that campuses provided some insulation from the effects of changes in the drinking age laws that took place during that interval. Similarly, entrenched in many college campuses is a culture of binge drinking which has proven impervious to many societal trends (and intervention attempts).⁵⁹ Also, individuals who are under the legal drinking age in college are mixed in with peers who are of legal age to purchase alcohol; this is no longer true in high schools and less true, perhaps, for many of those ages 19 to 22 who are not in college. Finally, much alcohol advertising and promotion was and is directed specifically at the college student population.
- College students generally have had slightly lower rates of *daily drinking* than the noncollege respondents, though that was not true in 2008 (Figure 9-14c). Daily drinking among noncollege young adults (one to four years past high school) declined from 8.7% in 1981 to 6.5% in 1984, remained essentially unchanged through 1988, declined further (to 3.2%) by 1994, then increased to 5.8% by 2004, about where it was in 2007. College students' daily drinking estimates—which appear a little less stable, perhaps due to smaller sample sizes in the 1980s—showed little or no decline between 1980 (6.5%) and 1984 (6.6%), but a considerable decline from 1984 through 1995 (to 3.0%), followed by a period of some increase, reaching 5.0% in 2002. After 2002 their daily drinking dropped to 3.7% in 2004, but increased slightly to 4.3% in 2007, about where it remained in 2008. Twelfth graders showed a somewhat similar pattern of daily drinking with a long period of decline, followed by a somewhat earlier reversal, beginning in 1994. After 1998 their daily drinking rate actually declined a little, and then remained fairly level through 2008. Of the three groups, the noncollege 19- to 22-year-olds have had the highest rate of current daily drinking over most of the past 29 years.
- *Cigarette smoking* among American college students (Figure 9-15a) declined modestly in the first half of the 1980s. Thirty-day prevalence fell from 26% to 22% between 1980 and 1984, remained fairly stable through 1990 (22%), then increased gradually but substantially, reaching 31% by 1999. It was not until 2000 that the first evidence of a decline in smoking among college students began to appear, two years after smoking had begun to decline among 12th graders. This lag no doubt reflects a cohort effect. The noncollege group showed little consistent change after the mid-1990s until evidence of gradual decline began to emerge sometime after 1999. Twelfth graders have shown a

⁵⁹Schulenberg, J. E., & Maggs, J. L. (2002). A developmental perspective on alcohol use and heavy drinking during adolescence and the transition to young adulthood. *Journal of Studies on Alcohol, Supplement 14*, 54–70.

fairly steady decline since 1997, while use among college students has declined since 1999. Because the noncollege segment has shown only a moderate decline so far, their smoking rate is now much higher than the college students.

- While smoking rates have consistently been lower among college students than the noncollege segment, the trends for these two groups converged some after 1984, as smoking rates more or less stabilized among college students but continued to decline among young adults not in college (see Figure 9-15a). In fact, between 1989 and 1991, use began to rise among college students while continuing to decline among their peers. Both groups showed fairly parallel increases in smoking between about 1991 and 1999, after which use continued to increase among the noncollege segment, but began to decline among college students. (Twelfth graders exhibited an increase from 1992 to 1997, and their use has declined significantly since.) The popularity of Camel cigarettes among the college-bound, which we have reported elsewhere, may help to explain some of the narrowing of the gap between college students and their age peers.⁶⁰ The Joe Camel advertising and promotion campaign, commenced in the late 1980s and ended in the late 1990s, may have succeeded in initiating more college students (particularly males) to smoking than had been the case previously or since.
- For many prescription-type drugs—*amphetamines*, *sedatives (barbiturates)*, and *tranquilizers*—differences between college students and their noncollege age peers narrowed over the years, particularly through the early 1990s. Much of this is due to general overall declines in usage rates during the 1980s, but may also reflect the increasing proportion of the age group going to college. Since then, the differences between these two groups have increased for sedatives (barbiturates) and tranquilizers as use in general has grown.
- The overall drug use trends among college students are also parallel, for the most part, to the trends among 12th graders; still, declines in many drugs from 1980 to 1990 were proportionately larger among 19- to 22-year-olds (college and noncollege) than among 12th graders. Despite parallel trends in the early 1990s, 12th graders have shown larger, and often earlier, increases in the use of a number of drugs in the years since; as indicated in *Volume I*, 8th and 10th graders showed increases a year earlier than 12th graders. Clearly this most recent upsurge, or what we have called a “relapse phase” in the illicit drug epidemic, did not originate on the nation’s college campuses, as did the earlier epidemic. It originated among secondary school students—and the younger ones at that—and has been carried up the age spectrum through generational replacement, at least in part.

⁶⁰Johnston, L. D., O’Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (1999). *Cigarette brand preferences among adolescents* (Monitoring the Future Occasional Paper No. 45). Ann Arbor, MI: Institute for Social Research.

GENDER DIFFERENCES IN TRENDS AMONG COLLEGE STUDENTS

Since MTF began, there has been a slow rise in the proportion of college students who are female. Females constituted 50% of our 1980 sample of college students compared to 60% of our 2008 sample. Given that substantial gender differences exist in the use of some drugs, we have been concerned that apparent long-term trends in the levels of drug use among college students (and/or among the noncollege group) might actually be attributable to changes in the gender composition of that population. For this reason, in particular, we have consistently presented separate trend lines for males and females. Gender differences in trends are illustrated in the lower panels of Figures 9-1 through 9-15c, and are discussed next.

In general, trends in use of the various drugs and in the overall drug use indexes have been highly parallel for male and female college students, as an examination of the relevant figures will show. The most noteworthy exceptions are mentioned below.

- Certain drug use measures showed a convergence between the genders as use rates declined to low levels. This has been true for the use of *any illicit drug* and *any illicit drug other than marijuana*. *Marijuana* use was another example, with some convergence in the rates between 1980 and 1991 as overall use declined, and then some divergence between 1991 and 1999 as usage rates rose. After 2001, however, the two genders diverged somewhat, with use among males remaining essentially unchanged through 2008 and use among females decreasing (see Figure 9-3a). *Daily marijuana* use (Figure 9-3b) presents a clearer example, with the steep decline among males between 1980 and 1986 narrowing the gap between the genders substantially. Between 1986 and 1993 there was no further narrowing; but as use began to rise in the mid-1990s, a greater increase among college males widened the gap again through about 2000. After 2000 the gender gap remained fairly stable until 2008, when daily marijuana use rose among males and fell among females. Whether this unusually sharp divergence holds in the 2009 survey will be of interest.
- *LSD* use dropped more steeply among males from 1999 to 2005 than among females, bringing the genders close together at very low prevalence rates (Figure 9-6). Prior to that period of convergence, use had quite consistently been higher among males. The small increases in use that have occurred since 2005 have been greater among males.
- Use of *hallucinogens other than LSD* has dropped for both genders since 2003, but the decline has been greater among female college students, leading to an enlargement of the gender difference—just the opposite of what usually occurs during a period of declining use (Figure 9-7).
- Rates of *ecstasy (MDMA)* use have been quite similar for male and female college students since measures were first introduced in 1989 (Figure 9-8).
- After 1986, *cocaine* use, which had been substantially higher among males, dropped more steeply for males than for females in general and among male college students in particular, considerably narrowing the sizable gap between genders (see Figure 9-9).

Since 1991 both genders have moved in parallel, with males reporting somewhat higher usage rates.

- **Amphetamine** use (Figure 9-11) also showed some convergence in the early 1980s due to a greater decline among males. Since 1989 the trends have been quite parallel, with males generally having a slightly higher annual prevalence rate.
- Among college students, the annual prevalence of **alcohol** use has been virtually identical for the two genders since MTF began (Figure 9-14a). Prior to 2000, the 30-day prevalence rate showed a modest difference, with males slightly higher (Figure 9-14b); but that difference disappeared by 2000 as drinking rose some among females. College males have consistently had considerably higher rates of daily drinking and binge drinking than college females (Figures 9-14c and 9-14d). If anything, the gender difference in **daily drinking** has expanded after 2000, with males increasing and females showing some net decrease. Since about 2006, however, they have converged, as use among female college students has remained steady while use among males has declined considerably. In 2008, the rates were 5.1% for college males versus 3.3% for college females.
- From 1988 through 1994, **binge drinking** among college females decreased some (from 37% to 31%); but binge drinking among college males declined more, from a high point in 1986 of 58% to 47% in 1995 (see Figure 9-14d). Since 1998, there has been some closing of the gender gap in binge drinking, as the rate among college females has risen from 31% in 1998 to 34% in 2008, while it actually declined some from 52% to 49% among college males.
- Between 1980 and 1992, the 30-day prevalence of **cigarette smoking** was consistently higher among college females than males (Figure 9-15a). However, the gap in 30-day prevalence narrowed because use by female college students declined considerably between 1980 and 1989, while use by male college students did not decline. After 1989, the gap remained quite small and the genders reversed position, with college males catching up to, and passing, females in their rate of smoking by 1994 and then remaining higher through at least 2000. (A similar reversal had occurred among 12th graders a few years earlier, so this reversal probably reflected a cohort effect.) Both genders exhibited a considerable decrease in 30-day smoking between 1999 and 2003, leaving very little difference between them (22% for males, 23% for females), although the trend line for college males was irregular during this interval. Since 2006, males have been smoking at higher levels than females, but the differences are still very small.
- While the rise in smoking among college students was longer term and more gradual than in the other two groups, it was nevertheless substantial, rising by nearly half between 1989 (21%) and 1999 (31%). The increase in smoking after 1988 was sharper among college males than among college females, consistent with the notion that Camel cigarettes' promotion and advertising—which ended in the late 1990s as a part of the tobacco settlement—played a role in the overall increase. Camels proved considerably

more popular among males, especially among those college-bound and from more educated families.⁶¹

⁶¹Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (1999). *Cigarette brand preferences among adolescents* (Monitoring the Future Occasional Paper No. 45). Ann Arbor, MI: Institute for Social Research.

TABLE 9-1
Trends in Lifetime Prevalence of Various Types of Drugs
among College Students 1 to 4 Years beyond High School
 (Entries are percentages.)

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007- 2008 change	
<i>Approximate</i>	1,040	1,130	1,150	1,170	1,110	1,080	1,190	1,220	1,310	1,300	1,400	1,410	1,490	1,490	1,410	1,450	1,460	1,480	1,440	1,440	1,350	1,340	1,260	1,270	1,400	1,360	1,280	1,250	1,270		
<i>Weighted N =</i>	69.4	66.8	64.6	66.9	62.7	65.2	61.8	60.0	58.4	55.6	54.0	50.4	48.8	45.9	45.5	47.4	49.0	52.9	53.2	53.2	53.7	53.6	51.8	53.9	52.2	52.3	50.6	50.5	49.5	-0.9	
Any illicit Drug ^a	42.2	41.3	39.6	41.7	38.6	40.0	37.5	35.7	33.4	30.5	28.4	25.8	26.1	24.3	22.0	24.5	22.7	24.4	24.8	25.5	25.8	26.3	26.9	27.6	28.0	26.5	26.3	25.3	22.6	-2.7	
Any illicit Drug ^a other than Marijuana	65.0	63.3	60.5	63.1	59.0	60.6	57.9	55.8	54.3	51.3	49.1	46.3	44.1	42.0	42.2	41.7	45.1	46.1	49.9	50.8	51.2	51.0	49.5	50.7	49.1	49.1	46.9	47.5	46.8	-0.7	
Marijuana	10.2	8.8	10.6	11.0	10.4	10.6	11.0	13.2	12.6	15.0	13.9	14.4	14.2	14.8	12.0	13.8	11.4	12.4	12.8	12.4	12.9	9.6	7.7	9.7	8.5	7.1	7.4	6.3	4.9	-1.4	
Inhalants ^b	15.0	12.0	15.0	12.2	12.9	11.4	11.2	10.9	10.2	10.7	11.2	11.3	12.0	11.8	10.0	13.0	12.6	13.8	15.2	14.8	14.4	14.8	13.6	14.5	12.0	11.0	10.6	9.1	8.5	-0.6	
Hallucinogens ^c	10.3	8.5	11.5	8.8	9.4	7.4	7.7	8.0	7.5	7.8	9.1	9.6	10.6	10.6	9.2	11.5	10.8	11.7	13.1	12.7	11.8	12.2	8.6	8.7	5.6	3.7	3.5	3.3	4.3	+1.0	
LSD	11.6	9.0	10.6	8.3	9.2	8.1	7.8	6.8	6.2	6.2	6.0	6.0	5.7	5.4	4.4	6.5	6.5	7.5	8.7	8.8	8.2	10.7	11.0	12.8	10.1	10.6	10.1	8.5	8.2	-0.4	
Hallucinogens other than LSD ^d	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Ecstasy (MDMA) ^e	22.0	21.5	22.4	23.1	21.7	22.9	23.3	20.6	15.8	14.6	11.4	9.4	7.9	6.3	5.0	5.5	5.0	5.6	8.1	8.4	9.1	8.6	8.2	9.2	9.5	8.8	7.7	8.5	7.2	-1.3	
Cocaine	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Crack ^f	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other Cocaine ^g	0.9	0.6	0.5	0.3	0.5	0.4	0.4	0.6	0.3	0.7	0.3	0.5	0.5	0.6	0.1	0.6	0.7	0.9	1.7	0.9	1.7	1.2	1.0	1.0	0.9	0.5	0.7	0.5	0.7	+0.2	
Heroin	8.9	8.3	8.1	8.4	8.9	6.3	8.8	7.6	6.3	7.6	6.8	7.3	7.3	6.2	5.1	7.2	5.7	8.2	8.7	8.7	8.9	11.0	12.2	14.2	13.8	14.4	14.6	14.1	12.4	-1.7	
Narcotics other than Heroin ^h	29.5	29.4	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Amphetamines ⁱ	—	—	30.1	27.8	27.8	25.4	22.3	19.8	17.7	14.6	13.2	13.0	10.5	10.1	9.2	10.7	9.5	10.6	10.6	11.9	12.3	12.4	11.9	12.3	12.7	12.3	10.7	11.2	9.1	-2.1	
Amphetamines, Adj. ^{g,i}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Methamphetamine ^j	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Crystal Methamphetamine	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Methamphetamine (ice)	—	—	—	—	—	—	—	—	—	—	1.0	1.3	0.6	1.6	1.3	1.0	0.8	1.6	2.2	2.8	1.3	2.3	2.0	2.9	2.2	2.4	1.7	1.3	1.1	-0.2	
Sedatives	8.1	7.8	8.2	6.6	6.4	4.9	5.4	3.5	3.6	3.2	3.8	3.5	3.8	3.5	3.2	4.0	4.6	5.2	5.7	6.7	6.9	6.0	5.9	5.7	7.2	8.5	6.3	5.9	6.4	+0.5	
(Barbiturates) ^g	13.7	14.2	14.1	12.2	10.8	9.3	8.0	6.1	4.7	4.1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Sedatives, Adj. ^{g,k}	10.3	10.4	11.1	9.2	9.0	7.2	5.8	4.1	2.2	2.4	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Methaqualone ^l	15.2	11.4	11.7	10.8	10.8	9.8	10.7	8.7	8.0	8.0	7.1	6.8	6.9	6.3	4.4	5.4	5.4	6.9	7.7	8.2	8.8	9.7	10.7	11.0	10.6	11.9	10.0	9.1	8.6	-0.4	
Tranquilizers ^l	94.3	95.2	95.2	95.0	94.2	95.3	94.9	94.1	94.9	93.7	93.1	93.6	91.8	89.3	88.2	88.5	88.4	87.3	88.5	88.0	86.6	86.1	86.0	86.2	84.6	86.6	84.7	83.1	85.3	+2.1	
Alcohol ^m	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Been Drunk ^k	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Flavored Alcoholic Beverage ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cigarettes	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids ⁿ	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

Source: The Monitoring the Future study, the University of Michigan.

Notes: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

— indicates data not available.

See footnotes on next page.

Footnotes for Tables 9-1 through 9-5

^a“Any illicit drug” includes use of marijuana, hallucinogens, cocaine, heroin or other narcotics, amphetamines, sedatives (barbiturates), methaqualone (until 1990), or tranquilizers not under a doctor’s orders.

^bThis drug was asked about in four of the five questionnaire forms in 1980–1989, in five of the six forms in 1990–1998, and in three of the six forms in 1999–2008. Total *N* in 2008 is approximately 630.

^cIn 2001 the question text was changed on three of the six questionnaire forms. “Other psychedelics” was changed to “other hallucinogens,” and “shrooms” was added to the list of examples. Beginning in 2002 the remaining forms were changed to the new wording.

^dThis drug was asked about in two of the five questionnaire forms in 1989, in two of the six questionnaire forms in 1990–2001, and in three of the six questionnaire forms in 2002–2008. Total *N* in 2008 is approximately 630.

^eThis drug was asked about in one of the five questionnaires for annual use only in 1986, two of the five questionnaire forms in 1987–1989, in all six questionnaire forms in 1990–2001, and in five of the six questionnaire forms in 2002–2008. Total *N* in 2008 is approximately 1,060.

^fThis drug was asked about in one of the five questionnaire forms in 1987–1989 and in four of six questionnaire forms in 1990–2008. Total *N* in 2008 is approximately 850.

^gOnly drug use that was not under a doctor’s orders is included here.

^hIn 2002 the question text was changed on three of the six questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only; *N* is three sixths of *N* indicated. In 2003 the remaining forms were changed to the new wording. The data are based on all forms in 2003 and beyond.

ⁱBased on the data from the revised question, which attempts to exclude inappropriate reporting of nonprescription amphetamines.

^jThis drug was asked about in two of the six questionnaire forms. Total *N* in 2008 is approximately 420.

^k“Sedatives, adjusted” data are a combination of barbiturate and methaqualone data.

^lIn 2001 the question text was changed on three of the six questionnaire forms. “Miltown” was replaced with “Xanax” in the list of examples. Beginning in 2002 the remaining forms were changed to the new wording.

^mIn 1993 and 1994, the question text was changed slightly in three of the six questionnaire forms to indicate that a “drink” meant “more than just a few sips.” Because this revision resulted in rather little change in reported prevalence in the surveys of high school graduates, the data for all forms combined are used in order to provide the most reliable estimate of change. After 1994 the new question text was used in all six of the questionnaire forms.

ⁿThis drug was asked about in three of the six questionnaire forms. Total *N* in 2008 is approximately 630.

^oThis drug was asked about in one of the six questionnaire forms. Total *N* in 2008 is approximately 210.

^pThis drug was asked about in one of the five questionnaire forms in 1989 and in two of the six questionnaire forms in 1990–2008. Total *N* in 2008 is approximately 420.

^qDaily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes, for which actual daily use is measured, and for 5+ drinks, for which the prevalence of having five or more drinks in a row in the last two weeks is measured.

^rRevised questions about amphetamine use were introduced in 1982 to more completely exclude inappropriate reporting of nonprescription amphetamines. The data in italics are therefore not strictly comparable to the other data.

TABLE 9-3
Trends in 30-Day Prevalence of Various Types of Drugs
among College Students 1 to 4 Years beyond High School
 (Entries are percentages.)

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007-2008 change	
<i>Approximate</i>																															
<i>Weighted N =</i>	1,040	1,130	1,150	1,170	1,110	1,080	1,190	1,220	1,310	1,300	1,400	1,410	1,490	1,490	1,410	1,450	1,450	1,480	1,440	1,440	1,350	1,340	1,260	1,270	1,400	1,360	1,280	1,250	1,270		
Any Illicit Drug ^a	38.4	37.6	31.3	29.3	27.0	26.1	25.9	22.4	18.5	18.2	15.2	15.2	16.1	15.1	16.0	19.1	17.6	19.2	19.7	21.6	21.5	21.9	21.5	21.4	21.2	19.5	19.2	19.3	18.9	-0.3	
Any Illicit Drug ^a other than Marijuana	20.7	18.6	17.1	13.9	13.8	11.8	11.6	8.8	8.5	6.9	4.4	4.3	4.6	5.4	4.6	6.3	4.5	6.8	6.1	6.4	6.9	7.5	7.8	8.2	9.1	8.2	8.2	8.1	7.3	-0.8	
Marijuana	34.0	33.2	26.8	26.2	23.0	23.6	22.3	20.3	16.8	16.3	14.0	14.1	14.6	14.2	15.1	18.6	17.5	17.7	18.6	20.7	20.0	20.2	19.7	19.3	18.9	17.1	16.7	16.8	17.0	+0.2	
Inhalants ^b	1.5	0.9	0.8	0.7	0.7	1.0	1.1	0.9	1.3	0.8	1.0	0.9	1.1	1.3	0.6	1.6	0.8	0.7	0.6	1.5	0.9	0.4	0.7	0.4	0.4	0.3	0.4	0.1	0.4	+0.2	
Hallucinogens ^c	2.7	2.3	2.6	1.8	1.8	1.3	2.2	2.0	1.7	2.3	1.4	1.2	2.3	2.1	2.1	3.3	1.9	2.1	2.1	2.0	1.4	1.8	1.2	1.8	1.3	1.2	0.9	1.3	1.7	+0.4	
LSD	1.4	1.4	1.7	0.9	0.8	0.7	1.4	1.4	1.1	1.4	1.1	0.8	1.8	1.6	1.8	2.5	0.9	1.1	1.5	1.2	0.9	1.0	0.2	0.2	0.2	0.1	0.3	0.3	0.8	+0.5	
Hallucinogens other than LSD ^d	1.9	1.2	1.4	1.0	1.2	0.7	1.2	0.8	0.8	1.1	0.8	0.6	0.7	1.1	0.8	1.6	1.2	1.2	0.7	1.2	0.8	0.8	1.1	1.7	1.2	1.1	0.7	1.1	1.3	+0.2	
Ecstasy (MDMA) ^f	—	—	—	—	—	—	—	—	—	0.3	0.6	0.2	0.4	0.3	0.2	0.7	0.7	0.8	0.8	2.1	2.5	1.5	0.7	1.0	0.7	0.8	0.6	0.4	0.6	+0.2	
Cocaine	6.9	7.3	7.9	6.5	7.6	6.9	7.0	4.6	4.2	2.8	1.2	1.0	1.0	0.7	0.6	0.7	0.8	1.6	1.6	1.2	1.4	1.9	1.6	1.9	2.4	1.8	1.8	1.7	1.2	-0.6	
Crack ^g	—	—	—	—	—	—	1.3	0.4	0.5	0.2	0.1	0.3	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.1	0.3	0.4	0.4	0.1	*	0.1	0.1	0.0	
Other Cocaine ^h	—	—	—	—	—	—	—	3.5	3.2	3.2	1.0	1.0	0.9	0.6	0.3	0.8	0.6	1.3	1.5	1.0	0.9	1.5	1.4	1.9	2.2	1.8	1.3	1.6	1.1	-0.5	
Heroin	0.3	*	*	*	*	*	*	0.1	0.1	0.1	*	0.1	*	*	*	0.1	*	0.2	0.1	0.1	0.2	0.1	*	*	0.1	0.1	0.2	0.1	*	-0.1	
Narcotics other than Heroin ^h	1.8	1.1	0.9	1.1	1.4	0.7	0.6	0.8	0.8	0.7	0.5	0.6	1.0	0.7	0.4	1.2	0.7	1.3	1.1	1.0	1.7	1.7	3.2	2.3	3.0	3.1	3.1	2.2	2.3	+0.1	
Amphetamines ^b	13.4	12.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Amphetamines, Adj. ^{g,i}	—	—	9.9	7.0	5.5	4.2	3.7	2.3	1.8	1.3	1.4	1.0	1.1	1.5	1.5	2.2	0.9	2.1	1.7	2.3	2.9	3.3	3.0	3.1	3.2	2.9	2.5	3.1	2.8	-0.3	
Methamphetamine ^d	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.2	0.2	0.5	0.2	0.6	0.2	0.1	0.2	0.1	*	-0.1
Crystal Methamphetamine (Ice)	—	—	—	—	—	—	—	—	—	—	*	*	*	*	0.3	0.5	0.3	0.1	0.2	0.3	*	0.1	*	0.3	0.1	0.2	*	0.1	*	-0.1	
Sedatives (Barbiturates) ^g	0.9	0.8	1.0	0.5	0.7	0.4	0.6	0.5	0.5	0.2	0.2	0.3	0.7	0.4	0.4	0.5	0.8	1.2	1.1	1.1	1.1	1.5	1.7	1.7	1.5	1.3	1.3	1.4	1.4	+0.1	
Sedatives, Adj. ^{g,k}	3.8	3.4	2.5	1.1	1.0	0.7	0.6	0.6	0.6	0.2	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Methaqualone ^g	3.1	3.0	1.9	0.7	0.5	0.3	0.1	0.2	0.1	0.0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
Tranquilizers ^{g,l}	2.0	1.4	1.4	1.2	1.1	1.4	1.9	1.0	1.1	0.8	0.5	0.6	0.6	0.4	0.4	0.5	0.7	1.2	1.3	1.1	2.0	1.5	3.0	2.8	2.7	2.2	2.1	1.8	1.6	-0.3	
Alcohol ^h	81.8	81.9	82.8	80.3	79.1	80.3	79.7	78.4	77.0	76.2	74.5	74.7	71.4	70.1	67.8	67.5	67.0	65.8	68.1	69.6	67.4	67.0	68.9	66.2	67.7	67.9	65.4	66.6	69.0	+2.4	
Been Drunk ^h Flavored Alcoholic Beverage ^g	—	—	—	—	—	—	—	—	—	—	—	45.0	45.0	43.8	42.8	37.9	40.3	46.4	44.3	44.6	43.9	44.7	44.4	40.4	47.4	43.1	47.6	46.8	45.3	-1.5	
Cigarettes	25.8	25.9	24.4	24.7	21.5	22.4	22.4	24.0	22.6	21.1	21.5	23.2	23.5	24.5	23.5	26.8	27.9	28.3	30.0	30.6	28.2	25.7	26.7	22.5	24.3	23.8	19.2	19.9	17.9	-2.0	
Steroids ^g	—	—	—	—	—	—	—	—	—	*	0.2	0.3	0.2	0.2	0.2	0.1	*	0.2	0.2	0.4	*	0.3	*	0.1	*	*	*	0.1	*	-0.1	

Source: The Monitoring the Future study, the University of Michigan.
 Notes: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.
 *** indicates a prevalence rate of less than 0.05%.
 "—" indicates data not available.
 See relevant footnotes at the end of Table 9-1.

TABLE 9-4
Trends in 30-Day Prevalence of Daily Use of Various Types of Drugs
among College Students 1 to 4 Years beyond High School
 (Entries are percentages.)

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007- 2008 change	
<i>Approximate</i>																															
<i>Weighted N =</i>	1,040	1,130	1,150	1,170	1,110	1,080	1,190	1,220	1,310	1,300	1,400	1,410	1,490	1,490	1,410	1,450	1,450	1,480	1,440	1,440	1,350	1,340	1,260	1,270	1,400	1,360	1,280	1,250	1,270		
Marijuana ^a	7.2	5.6	4.2	3.8	3.6	3.1	2.1	2.3	1.8	2.6	1.7	1.8	1.6	1.9	1.8	3.7	2.8	3.7	4.0	4.0	4.6	4.5	4.1	4.7	4.5	4.0	4.3	3.5	3.9	+0.4	
Cocaine ^b	0.2	*	0.3	0.1	0.4	0.1	0.1	0.1	0.1	*	*	*	*	*	0.1	*	*	*	*	*	*	*	*	*	*	*	0.1	0.1	*	*	0.0
Amphetamines ^{b,c}	0.5	0.4	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Amphetamines, Adj. ^{d,e}	—	—	0.3	0.2	0.2	*	0.1	0.1	*	*	*	0.1	*	0.1	0.1	0.1	*	0.2	0.1	0.1	0.1	0.2	0.1	0.3	0.2	0.2	0.4	0.1	0.2	0.0	0.0
Alcohol ^f	6.5	5.5	6.1	6.1	6.6	5.0	4.6	6.0	4.9	4.0	3.8	4.1	3.7	3.9	3.7	3.0	3.2	4.5	3.9	4.5	3.6	4.7	5.0	4.3	3.7	4.6	4.8	4.3	4.0	-0.3	
Been Drunk ^{g,h}	—	—	—	—	—	—	—	—	—	—	—	—	0.5	0.2	0.3	0.8	0.5	0.1	1.3	0.8	1.0	0.7	0.5	0.8	1.1	0.8	0.5	0.6	0.7	0.5	-0.2
5+ Drinks in a Row	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
in Last 2 Weeks	43.9	43.6	44.0	43.1	45.4	44.6	45.0	42.8	43.2	41.7	41.0	42.8	41.4	40.2	40.2	38.6	38.3	40.7	38.9	40.0	39.3	40.9	40.1	38.5	41.7	40.1	40.2	41.1	40.0	-1.1	
Cigarettes	18.3	17.1	16.2	15.3	14.7	14.2	12.7	13.9	12.4	12.2	12.1	13.8	14.1	15.2	13.2	15.8	15.9	15.2	18.0	19.3	17.8	15.0	15.9	13.8	13.8	12.4	9.2	9.3	9.2	-0.1	
Daily	12.7	11.9	10.5	9.6	10.2	9.4	8.3	8.2	7.3	6.7	8.2	8.0	8.9	8.9	8.0	10.2	8.5	9.1	11.3	11.0	10.1	7.8	7.9	7.6	6.8	6.7	4.9	4.3	4.3	0.0	
1/2Pack+/Day																															

Source: The Monitoring the Future study, the University of Michigan.
 Notes: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

*** indicates a prevalence rate of less than 0.05%.

"—" indicates data not available.


See relevant footnotes at the end of Table 9-1.

**TABLE 9-5
Trends in Lifetime, Annual, and 30-Day Prevalence of an Illicit Drug Use Index^a
among College Students 1 to 4 Years beyond High School, by Gender**


	1980 ^b	1981 ^b	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2007- 2008 change
	Percentage who used in lifetime																													
Any Illicit Drug	69.4	66.8	64.6	66.9	62.7	65.2	61.8	60.0	58.4	55.6	54.0	50.4	48.8	45.9	45.5	47.4	49.0	52.9	53.2	53.7	53.6	51.8	53.9	52.2	52.3	50.6	50.5	49.5	-0.9	
Males	71.0	67.5	68.1	71.3	66.4	69.8	64.7	63.5	56.0	56.5	52.5	51.3	50.8	45.7	49.5	47.3	50.3	52.1	54.4	58.4	54.4	53.9	54.3	54.1	54.9	54.2	55.0	52.3	50.7	-1.7
Females	67.5	66.3	61.5	63.0	59.2	61.6	59.4	57.4	60.2	54.9	55.1	49.7	47.1	46.0	42.6	44.3	45.6	46.7	52.0	49.6	53.2	53.5	50.2	53.7	50.6	51.3	47.8	49.4	48.8	-0.6
Any Illicit Drug other than Marijuana	42.2	41.3	39.6	41.7	38.6	40.0	37.5	35.7	33.4	30.5	28.4	25.8	26.1	24.3	22.0	24.5	22.7	24.4	24.8	25.5	25.8	26.3	26.9	27.6	28.0	26.5	26.3	25.3	22.6	-2.7
Males	42.8	39.8	45.1	44.6	40.9	42.1	38.2	37.2	31.8	30.6	26.2	27.6	26.3	24.3	24.6	26.6	25.0	27.3	27.3	29.4	28.9	27.0	30.4	27.6	31.1	29.0	29.2	26.5	25.2	-1.4
Females	41.6	42.6	34.7	39.2	36.4	38.3	37.0	34.6	34.6	30.4	30.1	24.3	26.1	24.3	20.1	22.9	21.2	22.2	23.3	22.8	23.5	25.9	24.6	27.5	26.2	25.1	24.4	24.6	21.0	-3.6
	Percentage who used in last 12 months																													
Any Illicit Drug	56.2	55.0	49.5	49.8	45.1	46.3	45.0	40.1	37.4	36.7	33.3	29.2	30.6	30.6	31.4	33.5	34.2	34.1	37.8	36.9	36.1	37.9	37.0	36.5	36.2	36.6	33.9	35.0	35.2	+0.2
Males	58.9	56.2	54.6	53.4	48.4	50.9	49.8	43.3	37.0	38.2	34.2	30.2	32.8	32.6	33.9	36.1	36.6	38.3	40.1	42.5	38.0	38.8	39.5	39.2	40.9	40.7	39.2	38.0	38.7	+0.6
Females	53.3	54.0	44.9	46.7	41.9	42.7	41.1	37.7	37.6	35.4	32.5	28.4	28.7	29.1	29.5	31.7	32.7	31.1	36.4	33.2	34.7	37.3	35.4	34.8	33.4	34.2	30.6	33.1	32.9	-0.2
Any Illicit Drug other than Marijuana	32.3	31.7	29.9	29.9	27.2	26.7	25.0	21.3	19.2	16.4	15.2	13.2	13.1	12.5	12.2	15.9	12.8	15.8	14.0	15.4	15.6	16.4	16.6	17.9	18.6	18.5	18.1	17.3	15.3	-2.0
Males	33.7	32.8	33.4	33.5	29.2	29.7	28.6	23.5	19.4	18.7	15.7	14.4	13.8	15.0	14.9	19.5	15.1	18.1	17.0	19.0	18.6	17.2	19.2	19.3	22.1	21.1	22.6	19.0	17.8	-1.3
Females	31.1	30.8	26.9	26.8	25.2	24.4	22.1	19.6	19.0	14.6	14.8	12.1	12.6	10.5	10.2	13.3	11.3	14.1	12.1	12.8	13.5	15.8	15.0	17.1	16.5	16.9	15.2	16.3	13.7	-2.6
	Percentage who used in last 30 days																													
Any Illicit Drug	38.4	37.6	31.3	29.3	27.0	26.1	25.9	22.4	18.5	18.2	15.2	15.2	16.1	15.1	16.0	19.1	17.6	19.2	19.7	21.6	21.5	21.9	21.5	21.4	21.2	19.5	19.2	19.3	18.9	-0.3
Males	42.9	40.6	37.7	33.8	30.4	29.9	31.0	24.0	18.8	20.0	18.2	16.0	18.0	16.0	20.5	23.7	20.6	23.4	23.1	26.7	24.0	25.0	25.1	22.8	26.1	22.9	23.4	22.7	23.1	+0.4
Females	34.0	34.8	25.6	25.5	23.7	23.2	21.7	21.1	18.3	16.7	12.7	14.6	14.5	14.5	12.7	15.7	15.8	16.2	17.6	18.1	19.6	19.8	19.3	20.5	18.4	17.5	16.6	17.1	16.2	-1.0
Any Illicit Drug other than Marijuana	20.7	18.6	17.1	13.9	13.8	11.8	11.6	8.8	8.5	6.9	4.4	4.3	4.6	5.4	4.6	6.3	4.5	6.8	6.1	6.4	6.9	7.5	7.8	8.2	9.1	8.2	8.2	8.1	7.3	-0.8
Males	22.8	18.6	20.2	16.0	16.1	12.6	14.4	9.0	8.2	8.0	4.9	4.8	5.1	7.3	6.2	8.8	6.1	7.8	8.6	7.5	8.2	9.0	8.4	8.1	11.3	10.3	9.5	9.6	+0.1	
Females	18.7	18.5	14.2	12.1	11.5	11.2	9.3	8.5	8.8	6.0	4.0	3.9	4.2	3.8	3.4	4.5	3.4	6.1	4.6	5.6	6.0	6.4	7.4	8.3	7.8	7.0	6.9	7.2	5.8	-1.5
	Approximate Weighted N																													
All Respondents	1,040	1,130	1,150	1,170	1,110	1,080	1,190	1,220	1,310	1,300	1,400	1,410	1,490	1,490	1,410	1,450	1,450	1,480	1,440	1,440	1,350	1,340	1,260	1,270	1,400	1,360	1,280	1,250	1,270	
Males	520	530	550	550	540	490	540	520	560	580	620	640	680	660	590	610	560	630	570	590	560	540	490	480	520	500	500	470	510	
Females	520	600	610	620	570	600	650	700	750	720	780	770	810	830	820	840	890	860	880	850	790	800	770	790	880	860	780	770	760	

Source: The Monitoring the Future study, the University of Michigan.
Notes: Level of significance of difference between the two most recent years: s = .05, ss = .01, sss = .001. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. See relevant footnotes at the end of Table 9-1.

FIGURE 9-1
Any Illicit Drug: Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Any Illicit Drug: Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-2
Any Illicit Drug other than Marijuana: Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Any Illicit Drug other than Marijuana: Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-3a
Marijuana: Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Marijuana: Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-3b
Marijuana: Trends in 30-Day Prevalence of Daily Use
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Marijuana: Trends in 30-Day Prevalence of Daily Use
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-4
Inhalants:^a Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



Inhalants:^a Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.


Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

^aUnadjusted for the possible underreporting of amyl and butyl nitrites.

FIGURE 9-5
Hallucinogens:^a Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



Hallucinogens:^a Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.


Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

^aUnadjusted for the possible underreporting of PCP.

FIGURE 9-6
LSD: Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




LSD: Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-7
Hallucinogens other than LSD:^a Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



Hallucinogens other than LSD:^a Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.


Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

^aUnadjusted for the possible underreporting of PCP.

FIGURE 9-8
Ecstasy (MDMA): Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Ecstasy (MDMA): Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-9
Cocaine: Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Cocaine: Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-10a
Narcotics other than Heroin:^a Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



Narcotics other than Heroin:^a Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

^aIn 2002 the question text was changed on half of the questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only. In 2003 the remaining forms were changed to the new wording.


FIGURE 9-10b
Vicodin: Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.


FIGURE 9-10c
OxyContin: Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-11
Amphetamines: Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Amphetamines: Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-12
Sedatives (Barbiturates): Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Sedatives (Barbiturates): Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-13
Tranquilizers: Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Tranquilizers: Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-14a
Alcohol: Trends in Annual Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Alcohol: Trends in Annual Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-14b
Alcohol: Trends in 30-Day Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Alcohol: Trends in 30-Day Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-14c
Alcohol: Trends in 30-Day Prevalence of Daily Use
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Alcohol: Trends in 30-Day Prevalence of Daily Use
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-14d
Alcohol: Trends in 2-Week Prevalence of 5 or More Drinks in a Row
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.^a)




Alcohol: Trends in 2-Week Prevalence of 5 or More Drinks in a Row
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.
 Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.
^aDue to a coding error for 18-year-olds, previous versions of this figure contained a value that was slightly off for the measure of five or more drinks in a row for 2005. This has been corrected here.

FIGURE 9-15a
Cigarettes: Trends in 30-Day Prevalence
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Cigarettes: Trends in 30-Day Prevalence
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-15b
Cigarettes: Trends in 30-Day Prevalence of Daily Use
among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)




Cigarettes: Trends in 30-Day Prevalence of Daily Use
among Male vs. Female College Students




Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

FIGURE 9-15c
Cigarettes: Trends in 30-Day Prevalence of Smoking a Half Pack
or More per Day among College Students vs. Others
1 to 4 Years beyond High School
 (Twelfth graders included for comparison.)



Cigarettes: Trends in 30-Day Prevalence of Smoking a Half Pack
or More per Day among Male vs. Female College Students



Source. The Monitoring the Future study, the University of Michigan.

Note. "Others" refers to high school graduates one to four years beyond high school not currently enrolled full-time in college.

INDEX⁶²

A

alcohol

daily use

college enrollment, 23, 249, 264,
table 8-4, table 9-4, figure 9-14c

disapproval, 190, table 6-2

friends' disapproval, perceived, 204,
table 7-1

gender, 23, 84, 137, 267, table 4-1,
table 4-5, table 8-4, figure 9-14c

harmfulness, perceived, 187,
table 6-1

population density (urbanicity), 86,
146, table 4-5

prevalence, 31, 82, table 4-1,
table 4-5, figure 4-20b

region, 141, table 4-5

trends, 22, 131, table 2-4, table 5-4,
figure 5-18c, figure 9-14c

disapproval, 189–190, table 6-2

drunk, getting

friends' use, 208, 210, table 7-2

gender, 251

prevalence, 29, table 4-1
college enrollment, 249

race/ethnicity, 29

exposure to use, table 7-3 (*see also*
friends' use)

flavored alcoholic beverages

college enrollment, 250,
tables 8-1–8-3, tables 9-1–9-3

gender, 251, tables 4-1–4-4,
tables 8-1–8-3

population density (urbanicity),
tables 4-2–4-4

prevalence, tables 4-1–4-4,
tables 8-1–8-3

region, tables 4-2–4-4

trends, tables 2-1–2-3,
tables 5-1–5-3

friends' use, 208, 210, table 7-2

harmfulness, perceived, table 6-1

heavy drinking

college enrollment, 22, 249,
263–264, table 8-4, table 9-4,
figure 9-14d

disapproval, 190, table 6-2

friends' disapproval, perceived, 204,
table 7-1

gender, 23–24, 84, 137, 251, 267,
table 4-1, table 4-5, table 8-4,
figure 9-14d

harmfulness, perceived, 187,
table 6-1

population density (urbanicity), 86,
146, table 4-5

prevalence, 22, 23, 31–32, 34, 78,
82, table 4-1, table 4-5,
figure 4-20b

race/ethnicity, 29

region, 86, 141, table 4-5

trends, 22, 130, table 2-4, table 5-4,
table 9-4, figure 5-18d

prevalence, 22–24, 29, 30, 31–32, 78,
82, figure 4-20a

college enrollment, 22, 248, 249,
250, tables 8-1–8-3

gender, tables 4-1–4-4,
tables 8-1–8-3

population density (urbanicity), 86,
tables 4-2–4-4

race/ethnicity, 29

region, 85–86, tables 4-2–4-4

trends, 22–24, 130, tables 2-1–2-3,
tables 5-1–5-3, figures
5-18a–5-18d

college enrollment, 263–264,
tables 9-1–9-3, figures
9-14a–9-14b

gender, 136–137, 267,
figures 9-14a–9-14b

population density (urbanicity), 146
region, 141

⁶²This index comprises drugs and their variables mentioned in the text, figures, and selected tables.

amphetamines

availability, perceived, 212, 214–215, table 7-4
daily use, table 5-4, table 8-4, table 9-4
disapproval, 12, 188, 189, table 6-2
exposure to use, table 7-3 (*see also* friends' use)
friends' disapproval, perceived, 204, 205, table 7-1
friends' use, 207, 210, table 7-2
harmfulness, perceived, 12, 184, 191, table 6-1
prevalence, 19, 30, 31, 79, 80, figure 4-4, tables 8-1–8-3
college enrollment, 21, tables 8-1–8-4
gender, 22, 84, 251, tables 4-1–4-4, tables 8-1–8-4
population density (urbanicity), 86, tables 4-2–4-4
race/ethnicity, 28, 29
region, tables 4-2–4-4
trends, 12, 124, 126, 127–128, tables 2-1–2-3, tables 5-1–5-4, figure 5-13
college enrollment, 260–261, 265, tables 9-1–9-4, figure 9-11
gender, 135, 267, figure 9-11
population density (urbanicity), 145
region, 138, 139

amyl nitrites (*See* nitrites)

anabolic steroids (*See* steroids)

androstenedione

prevalence, 20
gender, 20

any illicit drug

exposure to use, 209, table 7-3 (*see also* friends' use)
friends' use, 207, 208, table 7-2
prevalence, 6, 7, 34, 78, 79, tables 4-1–4-4, figure 4-1
college enrollment, 21, 248, tables 8-1–8-3
gender, 84, 250, tables 4-1–4-4, tables 8-1–8-3

population density (urbanicity), 86, tables 4-2–4-4

race, 29

region, 85, tables 4-2–4-4

trends, 6, 7, 124–125, 126, tables 2-1–2-3, tables 5-1–5-3, figure 2-1, figure 5-1

college enrollment, 258, tables 9-1–9-3, table 9-5, figure 9-1

gender, 133, 266, table 5-5, table 9-5, figure 9-1

population density (urbanicity), 142
region, 138

any illicit drug other than marijuana

disapproval, 188

exposure to use, 209, 210, table 7-3 (*see also* friends' use)

friends' disapproval, perceived, 204

friends' use, 207, table 7-2

prevalence, 6, 30, 31, 34, 78, 79, tables 4-1–4-4, figure 4-2

college enrollment, 248, 249, tables 8-1–8-3

gender, 22, 250, tables 4-1–4-4, tables 8-1–8-3

population density (urbanicity), tables 4-2–4-4

region, 85, tables 4-2–4-4

trends, 7, 9, 32, 124, 125, 126, tables 2-1–2-3, tables 5-1–5-3, figure 5-2

college enrollment, 259, tables 9-1–9-3, table 9-5, figure 9-2

gender, 133, 134, 266, table 5-5, table 9-5, figure 9-2

population density (urbanicity), 142
region, 138

B

barbiturates (*See* sedatives)

bidis, table 2-2

butyl nitrites (*See* nitrites)

C

cigarettes

- availability, perceived, 26
- daily use
 - college enrollment, 27, 250, table 8-4, table 9-4, figures 9-15b–9-15c
 - disapproval, 190–191, table 6-2
 - friends' disapproval, perceived, 204, table 7-1
 - gender, 137, 251, table 4-1, table 4-5, table 8-4, figures 9-15b–9-15c
 - harmfulness, perceived, 25, 187, table 6-1
 - population density (urbanicity), 87, 146–147, table 4-5
 - prevalence, 32, 82–83, table 4-1, table 4-5, figure 4-21
 - region, 141–142, table 4-5
 - trends, 7–8, 24, 25, table 2-4, table 5-4, table 9-4, figures 5-19b–5-19c
- disapproval, 25, 190–191, table 6-2
- friends' disapproval, perceived, 204, 206, table 7-1
- friends' use, 209, table 7-2
- harmfulness, perceived, 25, 187, table 6-1
- prevalence, 8, 30, 32, 34, 82–83, tables 4-1–4-5, figure 4-21
 - college enrollment, 27, 250, tables 8-1–8-3
 - gender, 84, 251, tables 4-1–4-5, tables 8-1–8-3
 - population density (urbanicity), tables 4-2–4-5
 - race/ethnicity, 28, 29
 - region, 86, tables 4-2–4-5
- trends, 7–8, 24–28, 32, 131–132, tables 2-1–2-4, tables 5-1–5-4, figure 5-19a
 - college enrollment, 27, 264–265, tables 2-1–2-4, tables 9-1–9-4, figure 9-15a

- gender, 27–28, 137–138, 267, figure 9-15a
 - population density (urbanicity), 146–147
 - race/ethnicity, 28
 - region, 141–142
- club drugs (*See* GHB; ketamine; Rohypnol)
- cocaine
- availability, perceived, 15, 212, 213, table 7-4
 - daily use, table 5-4, table 8-4, table 9-4
 - disapproval, 15, 188, 189, table 6-2
 - exposure to use, 209, table 7-3 (*see also* friends' use)
 - friends' disapproval, perceived, 204, 205, table 7-1
 - friends' use, 207, 208, 210, table 7-2
 - harmfulness, perceived, 15, 185, 191, table 6-1
 - prevalence, 14, 19, 31, 34, 78, 79, 81, figure 4-7
 - college enrollment, 248, tables 8-1–8-3
 - gender, 22, 84, tables 4-1–4-4, tables 8-1–8-3
 - population density (urbanicity), tables 4-2–4-4
 - race/ethnicity, 29
 - region, tables 4-2–4-4
 - trends, 14, 32, 126, tables 2-1–2-3, tables 5-1–5-3, figure 5-9
 - college enrollment, 14, 21, 263, tables 9-1–9-3, figure 9-9
 - gender, 134, 266–267, figure 9-9
 - population density (urbanicity), 144–145
 - region, 138, 139
- See also* cocaine powder; crack cocaine
- cocaine, other (*See* cocaine powder)
- cocaine powder
- availability, perceived, 212, table 7-4
 - disapproval, 33
 - friends' use, table 7-2
 - harmfulness, perceived, 184, table 6-1
 - prevalence, 30, 31, 79, figure 4-9

- college enrollment, 248, tables 8-1-8-3
 - gender, 22, 84, tables 4-1-4-4, tables 8-1-8-3
 - population density (urbanicity), tables 4-2-4-4
 - race/ethnicity, 28
 - region, tables 4-2-4-4
 - trends, 33, 124, tables 2-1-2-3, tables 5-1-5-3
 - college enrollment, tables 9-1-9-3
 - codeine, 17
 - cough/cold medicines, table 2-2
 - crack cocaine
 - availability, perceived, 212, 213, table 7-4
 - disapproval, 14, 32-33
 - friends' use, 207, 208, 210, table 7-2
 - harmfulness, perceived, 13, 14, 32-33, 184, 185, 191, table 6-1
 - prevalence, 13-14, 30, 31, 34, 79, 81, figure 4-8
 - college enrollment, 248, tables 8-1-8-3
 - gender, 22, 84, tables 4-1-4-4, tables 8-1-8-3
 - population density (urbanicity), tables 4-2-4-4
 - race/ethnicity, 29
 - region, tables 4-2-4-4
 - trends, 13-14, 33, 124, 126, tables 2-1-2-3, tables 5-1-5-3, figure 5-10
 - college enrollment, 13, tables 9-1-9-3
 - gender, 134-135
 - population density (urbanicity), 145
 - region, 138, 139
 - creatine
 - gender, 20
 - crystal methamphetamine (ice)
 - availability, perceived, 212, 215, table 7-4
 - harmfulness, perceived, 185-186, 191, table 6-1
 - prevalence, 79, 80, figure 4-6
 - college enrollment, 248, tables 8-1-8-3
 - gender, tables 4-1-4-4, tables 8-1-8-3
 - population density (urbanicity), 86, tables 4-2-4-4
 - race/ethnicity, 29
 - region, 85, tables 4-2-4-4
 - trends, 12-13, 126, 128, tables 2-1-2-3, tables 5-1-5-3, figure 5-14
 - college enrollment, tables 9-1-9-3
 - gender, 136
 - population density (urbanicity), 145
 - region, 140
- D**
- dextromethorphan, 20
 - diet pills, 20
- E**
- ecstasy
 - availability, perceived, 11, 212, 214, table 7-4
 - disapproval, 189, table 6-2
 - friends' use, 207, 210, table 7-2
 - harmfulness, perceived, 127, 144, 184, 186, 191, table 6-1
 - prevalence, 11, 30, 79, 81-82, tables 4-1-4-4, figure 4-17
 - college enrollment, tables 8-1-8-3
 - gender, 84, tables 4-1-4-4, tables 8-1-8-3
 - population density (urbanicity), tables 4-2-4-4
 - race/ethnicity, 28, 29
 - region, 85, tables 4-2-4-4
 - trends, 10-11, 19, 127, tables 2-1-2-3, tables 5-1-5-3, figure 5-8
 - college enrollment, 10-11, 19, 261, tables 2-1-2-3, tables 9-1-9-3, figure 9-8
 - gender, 134, 265, figure 9-8
 - population density (urbanicity), 143-144
 - region, 139

exposure to drug use, table 7-3 (*See also*
friends' use)

F

flavored alcoholic beverages (*See* alcohol)

G

GHB

prevalence, 19, table 4-3
college enrollment, table 8-2
gender, 84, table 4-3, table 8-2
population density (urbanicity),
table 4-3
region, 85, table 4-3
trends, 19, table 2-2, table 5-2
college enrollment, table 9-2
gender, 136
region, 141

H

hallucinogens

prevalence, 31, 79, figure 4-10
college enrollment, 21, 251,
tables 8-1-8-3
gender, 251, tables 4-1-4-4,
tables 8-1-8-3
population density (urbanicity),
tables 4-2-4-4
race/ethnicity, 29
region, 85, tables 4-2-4-4
trends, 124, 126, tables 2-1-2-3,
tables 5-1-5-3, figure 5-5
college enrollment, tables 9-1-9-3,
figure 9-5
gender, figure 9-5

hallucinogens other than LSD

availability, perceived, 212, 214,
table 7-4
exposure to use, 207, table 7-3 (*see also*
friends' use)
friends' use, table 7-2
prevalence, 30, figure 4-12
college enrollment, 251,
tables 8-1-8-3
gender, 84, 251, tables 4-1-4-4,
tables 8-1-8-3

population density (urbanicity),
tables 4-2-4-4

race/ethnicity, 28, 29

region, tables 4-2-4-4

trends, 32, tables 2-1-2-3,
tables 5-1-5-3, figure 5-7

college enrollment, 266,

tables 9-1-9-3, figure 9-7

gender, 266, figure 9-7

population density (urbanicity), 143

hashish (*See* marijuana)

heroin (including with and without a needle)

availability, perceived, 212, 214,
table 7-4

disapproval, 188, 189, table 6-2

exposure to use, table 7-3 (*see also*
friends' use)

friends' use, 208, table 7-2

harmfulness, perceived, 16-17, 184,
186-187, 191, table 6-1

prevalence, 30, 31, 79, figure 4-19

college enrollment, 248,

tables 8-1-8-3

gender, 22, 84, tables 4-1-4-4,

tables 8-1-8-3

population density (urbanicity),

tables 4-2-4-4

race/ethnicity, 29

region, tables 4-2-4-4

trends, 15-16, 128, tables 2-1-2-3,

tables 5-1-5-3, figure 5-11

college enrollment, tables 9-1-9-3

gender, 135

population density (urbanicity),
145-146

region, 140

I

ice (*See* crystal methamphetamine)

inhalants

friends' use, 208, table 7-2

harmfulness, perceived, 13

prevalence, 19, 30, 79-80,
tables 4-1-4-4, figure 4-13

college enrollment, 248,

tables 8-1-8-3

gender, 22, 84, tables 4-1-4-4,
tables 8-1-8-3
population density (urbanicity),
tables 4-2-4-4
race/ethnicity, 28
region, tables 4-2-4-4
trends, 13, 32, 127, tables 2-1-2-3,
tables 5-1-5-3, figure 5-4
college enrollment, 261,
tables 9-1-9-3, figure 9-4
gender, 136, figure 9-4
population density (urbanicity), 146
region, 139

K

ketamine (Special K)

prevalence, 19
college enrollment, table 8-2
gender, 84, table 4-3, table 8-2
population density (urbanicity),
table 4-3
region, 85, table 4-3
trends, table 2-2, table 5-2
college enrollment, table 9-2
gender, 136
region, 141

kreteks, table 2-2

L

look-alike drugs, 20

LSD

availability, perceived, 10, 212, 213,
table 7-4
disapproval, 10, 188-189, table 6-2
exposure to use, table 7-3 (*see also*
friends' use)
friends' disapproval, perceived, 204,
205, table 7-1
friends' use, 207, table 7-2
harmfulness, perceived, 10, 184-185,
191, table 6-1
prevalence, 19, 30, 79, tables 4-1-4-4,
figure 4-11
college enrollment, 248, 250,
tables 8-1-8-3

gender, 84, 250, tables 4-1-4-4,
tables 8-1-8-3
population density (urbanicity),
tables 4-2-4-4
race/ethnicity, 28, 29
region, tables 4-2-4-4
trends, 9-10, 32, 125-126,
tables 2-1-2-3, tables 5-1-5-3,
figure 5-6
college enrollment, 261, 266,
tables 9-1-9-3, figure 9-6
gender, 134, 265, figure 9-6
population density (urbanicity), 143
region, 138

M

marijuana (hashish)

availability, perceived, 211, 212-213,
table 7-4
daily use
college enrollment, 248, 251, 260,
266, table 2-4, table 8-4,
table 9-4, figure 9-3b
gender, 22, 84, 133-134, 251, 266,
table 4-5, table 8-4, figure 9-3b
population density (urbanicity), 143,
table 4-5
prevalence, 8, 31, 34, 79, table 4-5
region, table 4-5
trends, 8-9, 125, 132-133, table 2-4,
table 5-4, figure 5-3c
disapproval, 9, 188, table 6-2
exposure to use, 209, table 7-3 (*see also*
friends' use)
friends' disapproval, perceived, 204,
205, table 7-1
friends' use, 207, 209-210, table 7-2
harmfulness, perceived, 9, 183-184,
191, table 6-1
prevalence, 8, 19, 30, 31, 78, 79,
tables 4-1-4-4, figure 4-3
college enrollment, 21, 248, 250,
251, tables 8-1-8-3
gender, 22, 250, 251, tables 4-1-4-4,
tables 8-1-8-3

- population density (urbanicity), 86, tables 4-2-4-4
 - race/ethnicity, 29
 - region, 85, tables 4-2-4-4
 - trends, 7, 8, 32, 124, 125, 126, tables 2-1-2-3, tables 5-1-5-3, figure 5-3a-5-3b
 - college enrollment, 21, 260, 266, tables 2-1-2-3, tables 9-1-9-3, figure 9-3a
 - gender, 133, 266, figure 9-3a
 - population density (urbanicity), 142-143
 - region, 138
- MDMA (*See* ecstasy)
- methamphetamine
- prevalence, 30, 80, tables 4-1-4-4, figure 4-5
 - college enrollment, 248, tables 8-1-8-2
 - gender, 22, tables 4-1-4-4, tables 8-1-8-2
 - population density (urbanicity), 86, tables 4-2-4-4
 - race/ethnicity, 28
 - region, 85, tables 4-2-4-4
 - trends, 12, 128, tables 2-1-2-3, tables 5-1-5-3
 - college enrollment, tables 2-1-2-3, tables 9-1-9-3
 - population density (urbanicity), 145
 - region, 140
- methaqualone
- prevalence, 18
 - trends, 18, 126, tables 2-1-2-3, tables 5-1-5-3
 - college enrollment, tables 9-1-9-3
 - population density (urbanicity), 145
- N**
- narcotics other than heroin
- availability, perceived, 212, 214, table 7-4
 - exposure to use, table 7-3 (*see also* friends' use)
 - friends' use, 207, 208, 210, table 7-2
- prevalence, 17, 19, 31, 79, 81, tables 4-1-4-4, figure 4-15
- college enrollment, 250, tables 8-1-8-3
- gender, 84, 250, tables 4-1-4-4, tables 8-1-8-3
- population density (urbanicity), tables 4-2-4-4
- race/ethnicity, 29
- region, tables 4-2-4-4
- trends, 17, 19, 35, 124, 126, 128-129, tables 2-1-2-3, tables 5-1-5-3, figure 5-12
- college enrollment, 21, 262, tables 2-1-2-3, tables 9-1-9-3, figure 9-10a
- gender, 135, figure 9-10a
- population density (urbanicity), 146
- region, 140-141
- nitrites
- availability, perceived, table 7-4
 - friends' use, table 7-2
 - prevalence, 13
 - trends, 13, tables 2-1-2-3, tables 5-1-5-3
- See also* inhalants
- O**
- OxyContin
- prevalence, 17, 19, table 4-3
 - college enrollment, table 8-2
 - gender, 84, table 4-3, table 8-2
 - population density (urbanicity), table 4-3
 - race/ethnicity, 28, 29
 - region, 85, table 4-3
 - trends, 17, 129, table 2-2, table 5-2
 - college enrollment, 262, table 2-2, table 9-2, figure 9-10c
 - gender, 135
 - population density (urbanicity), 146
 - region, 141
- P**
- PCP
- availability, perceived, table 7-4
 - friends' use, table 7-2

harmfulness, perceived, table 6-1
prevalence
 gender, 84, tables 4-1-4-4
 population density (urbanicity),
 tables 4-2-4-4
 region, tables 4-2-4-4
 trends, 15, tables 2-1-2-3, tables 5-1-5-3
perceived harmfulness (*See* harmfulness,
 perceived)
perceived risk (*See* harmfulness, perceived)
Percocet, 17
powder cocaine (*See* cocaine powder)
psychedelics (*See* hallucinogens)

Q

quaaludes, table 7-2

R

risk, perceived (*See* harmfulness, perceived)

Ritalin

 prevalence, 80, table 4-3
 college enrollment, 248, 249,
 table 8-2
 gender, 22, table 4-3, table 8-2
 population density (urbanicity),
 table 4-3
 race/ethnicity, 29
 region, table 4-3
 trends, 12, table 2-2, table 5-2
 college enrollment, table 9-2
 gender, 135-136

Rohypnol

 prevalence, 19, 30, table 4-3
 college enrollment, 248, table 8-2
 gender, table 4-3, table 8-2
 population density, table 4-3
 region, 85, table 4-3
 trends, tables 2-1-2-3, table 5-2
 college enrollment, table 9-2
 gender, 136
 region, 141

S

sedatives (barbiturates)

 availability, perceived, 212, 215,
 table 7-4

disapproval, 188, 189, table 6-2
exposure to use, table 7-3 (*see also*
 friends' use)

friends' use, 207, 210, table 7-2
harmfulness, perceived, 184, 191,
 table 6-1

prevalence, 18-19, 31, 79, 80-81,
 figure 4-14

 college enrollment, 248,
 tables 8-1-8-3

 gender, tables 4-1-4-4,
 tables 8-1-8-3

 population density (urbanicity),
 tables 4-2-4-4

 race/ethnicity, 29

 region, tables 4-2-4-4

 trends, 18, 35, 124, 126, 129,
 tables 2-1-2-3, tables 5-1-5-3,
 figure 5-15

 college enrollment, 21, 261-262,
 265, tables 9-1-9-3, figure 9-12

 gender, 135, figure 9-12

 population density (urbanicity), 145
 region, 140

smokeless tobacco

 harmfulness, perceived, 187-188,
 table 6-1

 prevalence

 gender, 30

 race/ethnicity, 29

 trends, table 2-1, tables 2-3-2-4

stay-awake pills, 20

steroids

 availability, perceived, 212, 215,
 table 7-4

 friends' use, 207, table 7-2

 prevalence, 30, 83, figure 4-18

 college enrollment, 248, 250,
 tables 8-1-8-2

 gender, 84, tables 4-1-4-4,
 tables 8-1-8-2

 population density (urbanicity),
 tables 4-2-4-4

 region, 85, tables 4-2-4-4

 trends, 129-130, tables 2-1-2-3,
 tables 5-1-5-3, figure 5-17

college enrollment, tables 9-1–9-3
stimulants, nonprescription, 20

T

tobacco (*See* cigarettes; smokeless tobacco)

tranquilizers

availability, perceived, 212, 215,
table 7-4

exposure to use, table 7-3 (*see also*
friends' use)

friends' use, 207, 210, table 7-2

prevalence, 19, 30, 31, 79, 81,

tables 4-1–4-4, figure 4-16

college enrollment, tables 8-1–8-3

gender, 22, tables 4-1–4-4,
tables 8-1–8-3

population density (urbanicity),
tables 4-2–4-4

race/ethnicity, 28, 29

region, tables 4-2–4-4

trends, 17–18, 35, 124, 129,

tables 2-1–2-3, tables 5-1–5-3,
figure 5-16

college enrollment, 21, 262, 265,
tables 9-1–9-3, figure 9-13

gender, 136, figure 9-13

population density (urbanicity), 145

region, 140

V

Valium, 18

Vicodin

prevalence, 17, 19

college enrollment, 262, table 8-2

gender, 84, 251, table 4-3, table 8-2

population density (urbanicity),
table 4-3

race/ethnicity, 28, 29

region, table 4-3

trends, 17, 129, table 2-2, table 5-2

college enrollment, 262–263,
table 9-2, figure 9-10b

gender, 135

population density (urbanicity), 146

region, 141

X

Xanax, 17–18

Monitoring the Future Web site:
<http://www.monitoringthefuture.org>



a continuing study of American youth

NIDA NATIONAL INSTITUTE
ON DRUG ABUSE

NIH Publication No. 09-7403

Printed November 2009