

Accepted Article

SCHOLARONE™
Manuscripts

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version record](#). Please cite this article as [doi:10.1002/lio2.123](https://doi.org/10.1002/lio2.123).

Perception and respiratory responses of the upper airway mechanism to added resistance with aging

Siiri Murtolahti, BSc; Ulla K. Crouse, PhD, DMD, MSc; Riitta Pahkala, PhD, DMD; Donald W. Warren, PhD, DDS, MSc; Maija T. Laine-Alava, PhD, DMD

From Institute of Clinical Medicine, University of Eastern Finland (SM, RP, MTLA) and Kuopio University Hospital (RP), Kuopio, Finland, Department of Orthodontics, University of Michigan, Ann Arbor, MI, USA (UKC), UNC Craniofacial Center, University of North Carolina, Chapel Hill, NC, USA (DWW)

Short running title: Respiratory responses to added resistance

The authors have no funding, financial relationship or conflicts of interest to disclose

Abstract

Objectives: To assess breathing behaviors and perception of added respiratory loads in young compared to old individuals, and to determine whether aging affects the perception and response to changes in nasal airway resistance.

Study design: In a clinical study, 40 young (11-20 years) and 40 older (59-82 years) subjects were evaluated during rest breathing and during the application of added airway resistance loads.

Methods: The pressure-flow technique (Microtronics Co., Chapel Hill, NC) was used to measure airflow rate (mL/s) and oral-nasal pressures (cmH₂O) to calculate nasal resistance (cmH₂O/L/s). To create calibrated resistance loads for the test conditions, we used a device modified from a precision iris diaphragm.

Results: During rest breathing airflow rate was significantly lower for the younger group compared to older group. Using the loading device, 11-20-year-olds detected increased resistance at the level of 2.26 cmH₂O/L/s compared to 4.55 cmH₂O/L/s in 59-82-year-olds. In contrast to the younger group, mean airflow rate was higher during expiration than during inspiration among 59-82-year-olds except at rest breathing.

Conclusions: The data revealed that the perception and respiratory response to increased airway resistance changed with aging. Younger subjects were more sensitive to changes within the airway. In both groups, subjects responded to increased airway resistance by decreasing airflow rate. However, expiratory phase became more active than inspiratory phase only in the older group.

Key words: Nasal resistance, nasal airflow rate, threshold load, pressure-flow technique, weber fraction.

Level of evidence: N/A

INTRODUCTION

Although most healthy individuals are primarily nasal breathers, airway impairment caused by allergies or infection results in a change to both nasal and oral breathing. Combined nasal and oral breathing also occurs during exercise,¹ when breathing colder air,^{2,3} or in supine position.⁴⁻⁶ Also nasal airway impairment in connection with mucosal swelling due to allergies can trigger mouth breathing.^{7,8} Switching from nasal to nasal and oral breathing occurs when nasal resistance reaches a threshold level which differs slightly among individuals and age.⁹⁻¹¹ A variety of techniques have been used to determine thresholds including body plethysmography and several rhinomanometric approaches.¹²⁻¹⁶ An increase in nasal airway size with age in children has been reported in several studies in different population groups.¹⁷⁻²⁴ Once adulthood is reached, there is little change in nasal airway size²⁵ although there are inconsistencies between nasal airway resistance and age.^{26,27} Other parameters of respiratory function have been less studied using aging as a factor.

Ventilatory responses due to external mechanical loading have been studied experimentally,^{13,15,28,29} in patients with increased airway resistance^{7,10,30,31} or with neurological abnormalities.³² In young adults Bennett et al.³³ showed that mean non-elastic resistance was 1.5-3.4 cmH₂O/L/s and the 50 % detection represents 25 % change in non-elastic resistance. Wiley and Zechman⁹ found that the 50 % detection represents about 25-30 % change in non-elastic load. All these studies indicate that breathing pattern changed although there is some disagreement on which variable is monitored and regulated during breathing.

The purpose of the present study was to assess breathing behaviors and the perception of added respiratory loads and, in particular to determine whether sensitivity and compensatory mechanisms change with aging.

MATERIALS AND METHODS

Two different age groups were compared with each other: 40 adolescents and 40 older adults, representing a homogenous population of Caucasian origin. The younger group included 40 healthy volunteers, 21 girls and 19 boys, with a mean age of 17.6 (SD 2.1) years and age range from 11.2 to 20.3 years. Forty older adult subjects, 29 women and 11 men, on an average 69.9 (SD 5.9) years of age, ranging from 59 to 82 years, participated in this study. Height (cm) was measured. A structured questionnaire was used to get information about history (0=no, 1=yes) of allergies, nasal symptoms, smoking habit and medical conditions including asthma, heart disease, rheumatism, diabetes, as well as lung, thyroid gland and biliary diseases. In the younger group, 11 subjects had seasonal allergies and 14 individuals smoking habit, in the older group the corresponding numbers were 9 and 4, respectively. None in the younger group while 13 subjects in the older group had chronic medical conditions, controlled by medication. Two older adults but none of the adolescents reported asthma. To include the volunteers in the study, they had to be free from nasal symptoms and seasonal allergies at the time of the measurements.

Airflow rate and oral and nasal pressures were recorded using the pressure-flow technique originally described by Warren et al.¹⁵ using the PERCI-PC and PERCI-SARS software (Microtronics Co., Chapel Hill, NC). The ICC value of 0.80 (95 % CI 0.58-0.94) has proved the reproducibility of the method to be good.² The pressure-drop across the nasal airway was measured by differential transducers connected to two catheters, by placing one catheter midway in the mouth and another catheter within a well-fitted nasal mask as in posterior rhinomanometry. Nasal airflow was measured with a heated pneumotachograph connected to the nasal mask which formed an airtight seal. Nasal airway resistance was determined using the following hydrokinetic equation:

$$R = \Delta P / V$$

where R = resistance $\text{cmH}_2\text{O}/\text{L}/\text{s}$, P = oral-nasal pressure (cmH_2O), and V = total airflow rate (mL/s).

The device used to create resistance loads was modified from a precision iris diaphragm (Model no. N36.624, o.d. 60 lever bridge) with a maximum opening of 8.0 mm diameter, corresponding an area of 0.50 cm^2 , that could be opened and closed in 0.2 mm increments in the diameter. The diaphragm was mounted halfway between the nasal mask and the pneumotachograph. The catheters of the pressure-flow instrumentation to measure pressures were connected to the tubing before and after the diaphragm. Table I presents data on the relationship between aperture area and resistance load, calibrated at an airflow rate of 500 mL/s . All measurements were recorded in an upright position in periods of 10 seconds for each subject. After recording rest breathing, the device to create added resistances was added, and the aperture size of the diaphragm was manually adjusted in a random sequence of higher and lower loads. In each instance the loaded condition was compared to a control, “unloaded”, condition with the diaphragm wide open. The subjects were asked to indicate when they detected a change in resistance. The same value had to be detected three times consecutively to be accepted as a threshold value.

The increment threshold for detecting a difference in nasal resistance was calculated for each individual as a Weber Fraction:

$$\text{WF} = (R_i - R_o) / R_o$$

where R_i = the resistance of the system corresponding to the just noticeable resistance during added load plus nasal resistance during rest breathing, and R_o = the resistance of the system corresponding to the diaphragm setting maximally open plus nasal resistance during each individual’s rest breathing.

Ethical approval for this study was granted by the Research Ethics Committee of the University of Kuopio and the Kuopio University Hospital, Kuopio, Finland, and the informed consent forms were signed by the participant or the parent.

Statistical methods

Differences between inspiratory and expiratory values of resistance and airflow rate during rest breathing and at three test conditions were assessed by paired t-test. Linear regression models were used to estimate associations between respiratory variables [resistance, airflow rate as well as the differential (inspiration minus expiration) values of resistance and airflow rate] during rest breathing and at the three test conditions according to age group (1 = 11-20-year-olds, 2 = 59-82-year-olds) and gender (0=females, 1=males), with occurrences (0=no, 1=yes) of smoking habit and any medical condition (see MATERIAL AND METHODS) as confounding factors, with height (cm) as covariate. Because none of the adolescents but 13 individuals among older adults had chronic diseases, controlled by medication, linear regression models were performed also separately for the older group. None of the adolescents and two older adults had asthma, the number of individuals being too small to be included in the analyses. For all analyses, p-values \leq 0.05 were considered statistically significant.

RESULTS

Table II shows mean values of nasal resistance and airflow rate during inspiration and expiration at rest breathing and all test conditions, namely at (1) the unloaded condition with the diaphragm the device to create resistance loads wide open, (2) at the time when added resistance was detected and (3) just before detecting the change, for both groups separately.

When comparing the inspiratory and expiratory variables (Table II), among 11-20-year-olds inspiratory resistance values were higher compared to expiration. The difference was statistically significant at all conditions except when the individuals detected the added respiratory load. Inspiratory airflow rate was higher than expiratory airflow rate but differed significantly only during rest breathing. For 59-82-year-olds, inspiratory resistance was significantly higher at rest breathing and at the unloaded condition but significantly lower at detection of the load and just before detection compared to expiration. For the older group, inspiratory airflow rate was lower compared to expiratory values except during rest breathing, and differed significantly only at the unloaded condition.

When comparing differential (inspiration minus expiration) values of the respiratory variables between the groups, linear regression models showed that the older group had significantly higher differential resistance at detection of the added load and just before detection and differential airflow rate at unloaded condition (Table III).

During inspiration at rest breathing and all test conditions (Table II), resistance values for the younger group were lower, 2.10, 1.54, 2.10 and 2.26 cmH₂O/L/s, respectively, compared to the corresponding values of older adults, 2.34, 1.75, 4.13 and 4.55 cmH₂O/L/s. Linear regression models showed that the difference between the groups was statistically significant at all conditions except during rest breathing (Table IV). Inspiratory airflow rate was lower for the adolescents compared to the older adults (Table II) but linear regression models (Table IV) showed that it was statistically significantly different only at rest breathing (431 and 490 mL/s, respectively), and decreased to about the same values of 300 and 308 mL/s just prior to detection, and to 302 and 303 mL/s, respectively, at detection of the added load.

The only statistically significant effect of gender was on differential resistance just prior to detecting the added load. Height, smoking habit, upper airway allergies, or other nasal symptoms did not have a statistically significant effect on any of the respiratory variables. Among older

adults, history of chronic diseases was related to nasal airflow only at the unloaded condition ($p = 0.035$).

Weber fraction, the just noticeable difference in added upper airway resistance, varied from 0.02 to 1.16 in adolescents and from 0.01 to 2.48 in older adults. The mean values were 0.23 (SD 0.26) for adolescents and 0.84 (SD 0.79) for older adults ($p < 0.001$). Scattering of Weber fraction according to baseline resistance values illustrated the constant nature of Weber fraction (Fig. 1) among adolescents, while there was a wider inter-individual variation among older adults (Fig. 2), indicating that higher added loads were needed for detection of change.

DISCUSSION

We determined that a sample size of 40 in each group was sufficient for this clinical study which, while non-invasive, was very time consuming for each subject. Power analyses confirmed this assumption. Earlier studies^{7,10,13,31,32} involved groups from 8 to 51 subjects. In the present study, all participants were clinically healthy and free of nasal symptoms at the time of the measurements. Previous studies have shown that when individuals with acute nasal congestion are excluded, medical history was not associated with measurements of nasal patency in children.^{22,34}

The questionnaire on medical history did not include psychological factors, including anxiety disorders, which can affect perception of breathing and alter respiratory mode³⁵. Also, use of anxiolytics is unknown because they were not separated from other medicines. When recording rest breathing followed by test measurements, performed by the same experienced examiners, special attention was paid to that the study subjects were relaxed.

The finding that in the present sample among older adults history of chronic diseases was related to nasal airflow only at the unloaded condition could bias the comparisons slightly. Linear

regression models showed that other variables of the medical history as well as smoking habit did not have significant effects on respiratory variables.

In studies on respiratory function, body size has been measured as height, weight or body mass index (BMI). Somewhat surprisingly, stature, estimated in the present study as height, was not related to respiratory function which is parallel with findings of Zapletal and Chapulova³⁶ in 2-19-year old study group. In 108 20-45-year old adults it has been reported that nasal resistance decreased when the height and weight increased except in individuals with weight of 85-95 kg.³⁷ Our finding is parallel with studies using BMI in adult population^{38,39} and also with results on subjects with sleepapnea and BMI ≥ 30 kg/m^{2,40} but contradictory to a study on sample of wide age range, from 16 to 82 years.⁴¹ When diagnosing problems in the upper airway, body size may only have significance when subjects are obese.

In our study groups, gender distribution was even among adolescents, while majority of the older subjects were women which is typical in older western populations. Only one variable associated with gender was statistically significantly different, nasal resistance just prior to detection of an added load.

This study revealed that older individuals detected added load at much higher resistance, 4.55 compared to 2.26 cmH₂O/L/s in adolescents, indicating less sensitivity in the perception of increased nasal resistance in older adults. Both groups responded in similar fashion to increased load resistance by decreasing airflow rate to about 300 mL/s prior to detection indicating that there is a physiologic response to the change even before one is aware of a change in the airway environment. Thus, despite the higher airflow rate in older adults during rest breathing and despite their weaker perception of changes in upper airway resistance, the physiologic response to changes was surprisingly similar in both age groups not only at detection of the added load but also with added load prior to detection of change.

Attempts to correlate subjective sensation of nasal obstruction with objective measurements, whether defined as nasal resistance dependent on minimum nasal cross-sectional area when nasal mucosa is included in the measurements as in our method, or as nasal airflow volume or rate, are undermined by wide interindividual variation in respiratory variables and by the complex nature of sensory neural functions⁴². Still, threshold values could provide helpful guidelines. The threshold for increased nasal resistance to provoke an individual to switch from nasal to oral breathing with its' harmful effects has been suggested to be around 3.5-4.5 cmH₂O/L/s in adults⁴³, and has been reported to be 4.7 cmH₂O/L/s in adolescents⁴⁴. Thus, among the present study groups the 59-82-year-olds but not the 11-20-year-olds would probably change their breathing mode at the time of perception of increased resistance.

In the present study, resistance values were higher during inspiration compared to expiration. However, in the older group, resistance values during expiration were higher just prior to detection and at detection. Airflow rates were higher during inspiration than expiration in adolescents at all test conditions except at detection of the added load when the values were almost equal. On the contrary, among older adults airflow values were higher during expiration than inspiration except at rest breathing. Only a few studies have investigated individual's ventilatory responses to added loads during inspiration and expiration. In agreement with our findings in adolescents, Ferris et al.⁴⁵ reported that nasal resistance was higher during inspiration compared to expiration, and Muza et al.⁴⁶ found that peak inspiratory airflow was consistently higher than peak expiratory flow. Contrary to our findings, healthy adults showed expiratory resistance values to be significantly higher than inspiratory values⁹ and to be more sensitive for expiratory than inspiratory loads.⁴⁷ In agreement with our findings, Tack et al.⁴⁸ compared younger and older adults and reported a difference for resistance, sensation intensity and peak mouth pressure loads.

In the present study, a Weber fraction was utilized to describe perception of the just noticeable increase in respiratory resistance. Weber's law expressing that the ratio of the increment threshold to the background intensity has been reported to be also valid in studies of perception of added loads to inspiratory resistance.⁹⁻¹¹ In individuals with asthma, the threshold values of detection have been reported to vary widely and to be much higher compared to controls, possibly partly due to chronic adaptation to the increased airway resistance.¹⁰ In their study, Weber's law seemed to be applicable when bronchodilatation was not used. Hallani et al.⁷ reported contradictory findings, namely enhanced detection of added nasal resistance in asthmatics compared to healthy subjects. In our study, Weber's law applied reasonably well among adolescents but not among the older group. The Weber fraction was clearly higher for older adults than adolescents, indicating that sensitivity to changes in airway resistance declines with aging.

Altogether, the clearly higher threshold for perception of increased upper airway resistance and increased effort for expiration in older people may pose a health hazard in individuals with cardiovascular or respiratory diseases during physical activities or stressful conditions.

CONCLUSIONS

This study revealed that aging significantly decreases the sensitivity to the recognition of changes in upper airway resistance. Before perception of an increased applied load, both groups attempted to compensate for the increase in resistance by lowering airflow rate. This respiratory response occurred at a lower load in the younger group but the differences in respiratory rate change were similar. That is, there was a lower sensitivity in the older group but the response was similar.

REFERENCES

1. Cole P, Forsyth R, Haight JS. Effects of cold air and exercise on nasal patency. *Ann Otol Rhinol Laryngol* 1983;92:196–198.
2. Laine MT, Huggare JA, Ruoppi P. A modification of the pressure-flow technique for measuring breathing of cold air and its effect on nasal cross-sectional area. *Am J Orthod Dentofacial Orthop* 1994;105:265–269.
3. Sue-Chu M. Winter sports athletes: long-term effects of cold air exposure. *Br J Sports Med* 2012;46:397–401.
4. Cole P, Haight JS. Posture and the nasal cycle. *Ann Otol Rhinol Laryngol*. 1986;95:33–37.
5. Duggan CJ, Watson RA, Pride NB. Postural changes in nasal and pulmonary resistance in subjects with asthma. *J Asthma* 2004;41:701–707.
6. Van Holsbeke CS, Verhulst SL, Vos WG, De Backer JW, Vinchurkar SC, Verdonck PR, van Doorn JW, Nadjmi N, De Backer WA. Change in upper airway geometry between upright and supine position during tidal nasal breathing. *J Aerosol Med Pulm Drug Deliv* 2014;7:51–57.
7. Hallani M, Wheatley JR, Amis TC. Initiating oral breathing in response to nasal loading: asthmatics versus healthy subjects. *Eur Respir J* 2008a;31:800–806.
8. Wandalsen GF, Mendes AI, Solé D. Correlation between nasal resistance and different acoustic rhinometry parameters in children and adolescents with and without allergic rhinitis. *Braz J Otorhinolaryngol* 2012;78:81–86
9. Wiley RL, Zechman FW Jr. Perception of added airflow resistance in humans. *Respir Physiol* 1966/67;2:73–87.
10. Burki NK, Mitchell K, Chaudhary BA, Zechman FW. The ability of asthmatics to detect added resistive loads. *Am Rev Respir Dis* 1978;117:71–75.

11. Gottfried SB, Altose MD, Kelsen SG, Cherniack NS. Perception of changes in airflow resistance in obstructive pulmonary disorders. *Am Rev Respir Dis* 1981;124:566–570.
12. Butler J. The work of breathing through the nose. *Clin Sci* 1960;19:55-62.
13. Watson RM, Warren DW, Fischer ND. Nasal resistance, skeletal classification and mouthbreathing in orthodontic patients. *Am J Orthod* 1968;54:367–379.
14. Warren DW, Duany LF, Fischer ND. Nasal pathway resistance in normal and cleft lip and palate subjects. *Cleft Palate J* 1969;6:134-140.
15. Warren DW, Lehman MD, Hinton VA. Analysis of simulated upper airway breathing. *Am J Orthod* 1984;86:197–206.
16. Haavisto LE, Vahlberg TJ, Sipila JI. Reference values for acoustic rhinometry in children at baseline and after decongestion. *Rhinology* 2011;49:243–247.
17. Warren DW, Hairfield WM, Dalston ET. Effect of age on nasal cross-sectional area and respiratory mode in children. *Laryngoscope* 1990;100:89-93.
18. Laine T, Warren DW. Effects of age, gender, and body size on nasal cross-sectional area in children. *Eur J Orthod* 1991;13:311-316.
19. Vig PS, Zajac DJ. Age and gender effects on nasal respiratory function in normal subjects. *Cleft Palate Craniofac J* 1993;30:279-284.
20. Laine-Alava MT, Minkkinen UK. Variation of nasal respiratory pattern with age during growth and development. *Laryngoscope* 1997;107:386-390.
21. Ho WK, Wei WI, Yuen AP, Chan KL, Hui Y. Measurement of nasal geometry by acoustic rhinometry in normal-breathing Asian children. *J Otolaryngol* 1999;28:232–237.
22. Straszék SP, Schlünssen V, Sigsgaard T, Pedersen O. Reference values for acoustic rhinometry in decongested school children and adults: the most sensitive measurement for change in nasal patency. *Rhinology* 2007;45:36-39.

23. Miyamoto Y, Takeuchi K, Majima Y. Measurement of nasal patency by acoustic rhinometry in Japanese schoolchildren. *Auris Nasus Larynx* 2009;36:406-410.
24. Haavisto LE, Sipilä JI. Acoustic rhinometry in children: some practical aspects and influence of age and body surface area on results. *Am J Rhinol* 2008;22:416-419.
25. Abramson Z, Susarla S, Troulis M, Kaban L. Age-related changes of the upper airway assessed by 3dimensional computed tomography. *J Craniofac Surg* 2009;20 Suppl 1:657–663.
26. Edelstein DR. Aging of the normal nose in adults. *Laryngoscope* 1996;106:1–25.
27. Kim SW, Mo JH, Kim JW, Kim DY, Rhee CS, Lee CH, Min YG. Change of nasal function with aging in Korean. *Acta Otolaryngol Suppl* 2007;Suppl 558:90–94.
28. Cole P, Forsyth R, Haight JS. Respiratory resistance of the oral airway. *Am Rev Respir Dis* 1982;125:363–365.
29. LaFramboise WA, Standaert TA, Guthrie RD, Woodrum DE. Developmental changes in the ventilatory response of the newborn to added airway resistance. *Am Rev Respir Dis* 1987;136:1075–1083.
30. O'Donnell DE, Sani R, Younes M. External mechanical loading in conscious humans: role of upper airway mechanoreceptors. *J Appl Physiol* 1988;65:541–548.
31. Hallani M, Wheatley J, Amis T. Enforced mouth breathing decreases lung function in mild asthmatics. *Respirology* 2008b;13:553–558.
32. Newsom Davis J. Contribution of somatic receptors in the chest wall to detection of added inspiratory airway resistance. *Clin Sci* 1967;33:259–260.
33. Bennett ED, Jayson MI, Rubenstein D, Campbell EJ. The ability of man to detect added non-elastic loads to breathing. *Clin Sci* 1962;23:155–162.
34. Laine-Alava MT, Minkkinen UK. Should history of nasal symptoms be considered when estimating nasal patency? *Angle Orthod* 1999;69:126–132.

35. Paulus MP. The breathing conundrum-interoceptive sensitivity and anxiety. *Depress Anxiety*. 2013;30(4):315-320.
36. Zapletal A, Chalupová J. Nasal airflow and resistance measured by active anterior rhinomanometry in healthy children and adolescents. *Pediatr Pulmonol* 2002;33:174-180.
37. Janosević L, Dotlić J, Janosević S, Dudvarski Z, Milovanović A, Pendjer I. Computerized rhinomanometry: a study of total nasal resistance normal values. *Acta Chir Jugosl* 2009;56:51-54.
38. Demir MG, Ylmaz HB. The Relation Between Body Mass Index and Nasal Airflow. *J Craniofac Surg* 2015;26:e295-297.
39. Raza MT, Wang DY. Is nasal cavity geometry associated with body mass index, height and weight? *Indian J Otolaryngol Head Neck Surg* 2012;64:266-269.
40. Tagaya M, Nakata S, Yasuma F, Noda A, Morinaga M, Yagi H, Sugiura M, Teranishi M, Nakashima T. Pathogenetic role of increased nasal resistance in obese patients with obstructive sleep apnea syndrome. *Am J Rhinol Allergy* 2010;24:51-54.
41. Crouse U, Laine-Alava MT. Effects of age, body mass index, and gender on nasal airflow rate and pressures. *Laryngoscope* 1999;109:1503-1508.
42. Baraniuk JN. Subjective nasal fullness and objective congestion. *Proc Am Thorac Soc*. 2011 Mar;8(1):62-69.
43. Laine MT, Warren DW. Perceptual and respiratory responses to added nasal airway resistance loads in older adults. *Laryngoscope*. 1995; 105:425-428.
44. Watson RM, Warren DW, Fischer ND. Nasal resistance, skeletal classification and mouthbreathing in orthodontic patients. *Am J Orthod*. 1968; 54:367-379.
45. Ferris BG Jr, Mead J, Opie LH. Partitioning of respiratory flow resistance in man. *J Appl Physiol* 1964;19:653-658.

46. Muza SR, McDonald S, Zechman FW. Comparison of subjects' perception of inspiratory and expiratory resistance. *J Appl Physiol Respir Environ Exerc Physiol* 1984;56:211–216.

47. Bonnel AM, Mathiot MJ, Grimaud C. Inspiratory and expiratory resistive load detection in normal and asthmatic subjects. A sensory decision theory analysis. *Respiration* 1985;48:12–23.

48. Tack M, Altose MD, Cherniack NS. Effect of aging on the perception of resistive ventilatory loads. *Am Rev Respir Dis* 1982;126:463–467.

Accepted Article

LEGENDS

Figure 1. Scattering of Weber fraction according to baseline resistance (cmH₂O/L/s) in 40 adolescents and young adults.

Figure 2. Scattering of Weber fraction according to baseline resistance (cmH₂O/L/s) in 40 older adults.

Accepted Article

Table I. Relation between cross-sectional area and resistance with the device used to create added resistance with different iris setting, calibrated at 500 mL/s.

Cross-Sectional Area of the Diaphragm (cm ²)	Resistance (cmH ₂ O/L/s)
0.50	2.2
0.48	2.5
0.45	2.9
0.43	3.4
0.41	3.6
0.38	4.2
0.36	4.6
0.34	5.1
0.32	6.3
0.31	7.3
0.28	8.9
0.26	9.5
0.25	10.2
0.23	12.9
0.21	16.4
0.20	19.1
0.18	22.1
0.16	25.1
0.15	30.6
0.14	36.9
0.13	46.3

Table II. Differences between inspiratory and expiratory airflow rate (mL/s) and nasal resistance (cmH₂O/mL/s) values among the younger (11-20 yrs) and older (59-82 yrs) study groups at different load conditions.

	Adolescents (n=40)		p*	Older adults (n=40)		p*
	Inspiration mean (SD)	Expiration mean (SD)		Inspiration mean (SD)	Expiration mean (SD)	
Rest breathing						
Resistance	2.10 (1.64)	1.77 (1.46)	0.000	2.34 (2.61)	2.00 (2.46)	<0.001
Airflow rate	431 (103)	387 (125)	0.000	490 (149)	486 (206)	0.827
Unloaded condition						
Resistance	1.54 (0.27)	1.40 (0.33)	0.000	1.75 (0.35)	1.55 (0.57)	0.001
Airflow rate	318 (58)	312 (73)	0.504	351 (82)	385 (108)	0.002
Loaded just before detection						
Resistance	2.10 (0.65)	1.95 (0.77)	0.005	4.13 (1.98)	4.70 (3.24)	0.038
Airflow rate	300 (62)	295 (64)	0.340	308 (82)	318 (110)	0.290
Loaded at detection						
Resistance	2.26 (0.73)	2.17 (0.85)	0.103	4.55 (2.29)	5.45 (4.13)	0.017
Airflow rate	302 (61)	303 (65)	0.953	303 (84)	315 (117)	0.212

*by paired t-test

Table III. Associations between differential (inspiration – expiration) resistance and airflow values (cmH₂O/L/s) according to age groups (1 = 11-20 yrs, 2 = 59-82 yrs) and gender (0=female, 1=male), considering the effects of smoking habit (0=no, 1=yes) medical history (0=no, 1=yes, see Subjects and Methods), with height (cm) as a covariate by linear regression analysis. Only statistically significant associations are given.

Dependent / Independent variable	Standardized coefficients Beta	p-value
RESISTANCE (cmH ₂ O/L/s)		
Loaded just prior to detection		
Group	0.277	0.020
Gender	0.339	0.026
Loaded at detection		
Group	0.301	0.013
AIRFLOW RATE (mL/s)		
Unloaded condition		
Group	0.283	0.020

Table IV. Associations between respiratory responses to added loads during inspiration according to age groups (1 = 11-20 yrs, 2 = 59-82 yrs) and gender (0=female, 1=male), considering the effects of smoking habit (0=no, 1=yes) and medical history (0=no, 1=yes, see Subjects and Methods), with height (cm) as a covariate by linear regression analyses. Only statistically significant associations are given.

Dependent / Independent variable	Standardized coefficients	p-value
RESISTANCE (cmH ₂ O/L/s)		
Unloaded condition		
Group	0.318	0.008
Loaded just prior to detection		
Group	0.592	<0.001
Loaded at detection		
Group	0.594	<0.001
AIRFLOW RATE (mL/s)		
Rest breathing		
Group	0.266	0.032

Comments to Reviewers:

Iscope-17-0574 Perception and respiratory responses of the upper airway mechanism to added resistance with aging

Reviewer: 1

Confidential Comments to the Author

This study appears to be well designed and is written in a clear voice.

The conclusions made are supported by the evidence presented.

The figures and tables are appropriate.

While the current study may serve to support a future line of questioning, the current manuscript lacks a clear clinical impact or implication.

Reviewer: 2

Confidential Comments to the Author

Murtolahti et al have investigated nasal airway breathing between a younger and older population by measuring airway resistance at rest and then how both populations react to a device to artificially create resistance. They found that airflow rate was lower in the younger population and the younger population was more sensitive to changes in airway pressure. This manuscript represents a significant research effort and the authors should be commended on their work. My comments are as follows:

-The age range of the younger population is quite broad (11-20 year). It seems the general consensus is to consider patients 18 or older as adults and those younger as children. I was curious why the authors included such young subjects.

-I think the discussion section would be strengthened if the authors discussed how their findings are clinically useful. In addition, a thoughtful discussion of the weaknesses and future directions would be helpful.

RESPONSE TO THE COMMENTS

Reviewer #1 and #2: Clinical implications have been added to the end of Discussion as follows:

Altogether, the clearly higher threshold for perception of increased upper airway resistance and increased effort for expiration in older people may pose a health hazard in individuals with cardiovascular or respiratory diseases during physical activities or stressful conditions.

Reviewer #2: The age range of the younger population (11-20 year): The age group between 10 and 20 years of age, defined as adolescents (WHO), is of special interest to our study group working mainly with craniofacial orthopedics. Some craniofacial structures and functions are mature as early as at 11 years of age, some continue to change until the end of puberty, and in case of mandibular growth, beyond 18 years of age in males. In our research on craniofacial growth and functions, adolescents provide an essentially important study group.

Regarding implications of the study, this paper is part of our cross-sectional, longitudinal and experimental studies on craniofacial growth, changes with age and aging, associations between craniofacial structures and dysfunctions, partly to support treatment of patients with congenital craniofacial anomalies and syndromes.

Perception and respiratory responses of the nasal airway mechanism to added resistance with aging

Abstract

Objectives: To assess breathing behaviors and perception of added respiratory loads in young compared to old individuals, and to determine whether aging affects the perception and response to changes in nasal airway resistance.

Study design: In a clinical study, 40 young (11-20 years) and 40 older (59-82 years) subjects were evaluated during rest breathing and during the application of added airway resistance loads.

Methods: The pressure-flow technique (Microtronics Co., Chapel Hill, NC) was used to measure airflow rate (mL/s) and oral-nasal pressures (cmH₂O) to calculate nasal resistance (cmH₂O/L/s). To create calibrated resistance loads for the test conditions, we used a device modified from a precision iris diaphragm.

Results: During rest breathing airflow rate was significantly lower for the younger group compared to older group. Using the loading device, 11-20-year-olds detected increased resistance at the level of 2.26 cmH₂O/L/s compared to 4.55 cmH₂O/L/s in 59-82-year-olds. In contrast to the younger group, mean airflow rate was higher during expiration than during inspiration among 59-82-year-olds except at rest breathing.

Conclusions: The data revealed that the perception and respiratory response to increased airway resistance changed with aging. Younger subjects were more sensitive to changes within the airway. In both groups, subjects responded to increased airway resistance by decreasing airflow rate. However, expiratory phase became more active than inspiratory phase only in the older group.

Key words: Nasal resistance, nasal airflow rate, threshold load, pressure-flow technique, weber fraction.

Level of evidence: N/A

Accepted Article

INTRODUCTION

Although most healthy individuals are primarily nasal breathers, airway impairment caused by allergies or infection results in a change to both nasal and oral breathing. Combined nasal and oral breathing also occurs during exercise,¹ when breathing colder air,^{2,3} or in supine position.⁴⁻⁶ Also nasal airway impairment in connection with mucosal swelling due to allergies can trigger mouth breathing.^{7,8} Switching from nasal to nasal and oral breathing occurs when nasal resistance reaches a threshold level which differs slightly among individuals and age.⁹⁻¹¹ A variety of techniques have been used to determine thresholds including body plethysmography and several rhinomanometric approaches.¹²⁻¹⁶ An increase in nasal airway size with age in children has been reported in several studies in different population groups.¹⁷⁻²⁴ Once adulthood is reached, there is little change in nasal airway size²⁵ although there are inconsistencies between nasal airway resistance and age.^{26,27} Other parameters of respiratory function have been less studied using aging as a factor.

Ventilatory responses due to external mechanical loading have been studied experimentally,^{13,15,28,29} in patients with increased airway resistance^{7,10,30,31} or with neurological abnormalities.³² In young adults Bennett et al.³³ showed that mean non-elastic resistance was 1.5-3.4 cmH₂O/L/s and the 50 % detection represents 25 % change in non-elastic resistance. Wiley and Zechman⁹ found that the 50 % detection represents about 25-30 % change in non-elastic load. All these studies indicate that breathing pattern changed although there is some disagreement on which variable is monitored and regulated during breathing.

The purpose of the present study was to assess breathing behaviors and the perception of added respiratory loads and, in particular to determine whether sensitivity and compensatory mechanisms change with aging.

MATERIALS AND METHODS

Two different age groups were compared with each other: 40 adolescents and 40 older adults, representing a homogenous population of Caucasian origin. The younger group included 40 healthy volunteers, 21 girls and 19 boys, with a mean age of 17.6 (SD 2.1) years and age range from 11.2 to 20.3 years. Forty older adult subjects, 29 women and 11 men, on an average 69.9 (SD 5.9) years of age, ranging from 59 to 82 years, participated in this study. Height (cm) was measured. A structured questionnaire was used to get information about history (0=no, 1=yes) of allergies, nasal symptoms, smoking habit and medical conditions including **asthma**, heart disease, rheumatism, diabetes, as well as lung, thyroid gland and biliary diseases. In the younger group, 11 subjects had seasonal allergies and 14 individuals smoking habit, in the older group the corresponding numbers were 9 and 4, respectively. None in the younger group while 13 subjects in the older group had chronic medical conditions, controlled by medication. **Two older adults but none of the adolescents reported asthma**. To include the volunteers in the study, they had to be free from nasal symptoms and seasonal allergies at the time of the measurements.

Airflow rate and oral and nasal pressures were recorded using the pressure-flow technique originally described by Warren et al.¹⁵ using the PERCI-PC and PERCI-SARS software (Microtronics Co., Chapel Hill, NC). The ICC value of 0.80 (95 % CI 0.58-0.94) has proved the reproducibility of the method to be good.² The pressure-drop across the nasal airway was measured by differential transducers connected to two catheters, by placing one catheter midway in the mouth and another catheter within a well-fitted nasal mask as in posterior rhinomanometry. Nasal airflow was measured with a heated pneumotachograph connected to the nasal mask which formed an airtight seal. Nasal airway resistance was determined using the following hydrokinetic equation:

$$R = \Delta P / V$$

where R = resistance $\text{cmH}_2\text{O}/\text{L}/\text{s}$, P = oral-nasal pressure (cmH_2O), and V = total airflow rate (mL/s).

The device used to create resistance loads was modified from a precision iris diaphragm (Model no. N36.624, o.d. 60 lever bridge) with a maximum opening of 8.0 mm diameter, corresponding an area of 0.50 cm^2 , that could be opened and closed in 0.2 mm increments in the diameter. The diaphragm was mounted halfway between the nasal mask and the pneumotachograph. The catheters of the pressure-flow instrumentation to measure pressures were connected to the tubing before and after the diaphragm. Table I presents data on the relationship between aperture area and resistance load, calibrated at an airflow rate of 500 mL/s . All measurements were recorded in an upright position in periods of 10 seconds for each subject. After recording rest breathing, the device to create added resistances was added, and the aperture size of the diaphragm was manually adjusted in a random sequence of higher and lower loads. In each instance the loaded condition was compared to a control, “unloaded”, condition with the diaphragm wide open. The subjects were asked to indicate when they detected a change in resistance. The same value had to be detected three times consecutively to be accepted as a threshold value.

The increment threshold for detecting a difference in nasal resistance was calculated for each individual as a Weber Fraction:

$$\text{WF} = (R_i - R_o) / R_o$$

where R_i = the resistance of the system corresponding to the just noticeable resistance during added load plus nasal resistance during rest breathing, and R_o = the resistance of the system corresponding to the diaphragm setting maximally open plus nasal resistance during each individual’s rest breathing.

Ethical approval for this study was granted by the Research Ethics Committee of the University of Kuopio and the Kuopio University Hospital, Kuopio, Finland, and the informed consent forms were signed by the participant or the parent.

Statistical methods

Differences between inspiratory and expiratory values of resistance and airflow rate during rest breathing and at three test conditions were assessed by paired t-test. Linear regression models were used to estimate associations between respiratory variables [resistance, airflow rate as well as the differential (inspiration minus expiration) values of resistance and airflow rate] during rest breathing and at the three test conditions according to age group (1 = 11-20-year-olds, 2 = 59-82-year-olds) and gender (0=females, 1=males), with occurrences (0=no, 1=yes) of smoking habit and any medical condition (see MATERIAL AND METHODS) as confounding factors, with height (cm) as covariate. Because none of the adolescents but 13 individuals among older adults had chronic diseases, controlled by medication, linear regression models were performed also separately for the older group. **None of the adolescents and two older adults had asthma, the number of individuals being too small to be included in the analyses.** For all analyses, p-values \leq 0.05 were considered statistically significant.

RESULTS

Table II shows mean values of nasal resistance and airflow rate during inspiration and expiration at rest breathing and all test conditions, namely at (1) the unloaded condition with the diaphragm the device to create resistance loads wide open, (2) at the time when added resistance was detected and (3) just before detecting the change, for both groups separately.

When comparing the inspiratory and expiratory variables (Table II), among 11-20-year-olds inspiratory resistance values were higher compared to expiration. The difference was statistically significant at all conditions except when the individuals detected the added respiratory load. Inspiratory airflow rate was higher than expiratory airflow rate but differed significantly only during rest breathing. For 59-82-year-olds, inspiratory resistance was significantly higher at rest breathing and at the unloaded condition but significantly lower at detection of the load and just before detection compared to expiration. For the older group, inspiratory airflow rate was lower compared to expiratory values except during rest breathing, and differed significantly only at the unloaded condition.

When comparing differential (inspiration minus expiration) values of the respiratory variables between the groups, linear regression models showed that the older group had significantly higher differential resistance at detection of the added load and just before detection and differential airflow rate at unloaded condition (Table III).

During inspiration at rest breathing and all test conditions (Table II), resistance values for the younger group were lower, 2.10, 1.54, 2.10 and 2.26 cmH₂O/L/s, respectively, compared to the corresponding values of older adults, 2.34, 1.75, 4.13 and 4.55 cmH₂O/L/s. Linear regression models showed that the difference between the groups was statistically significant at all conditions except during rest breathing (Table IV). Inspiratory airflow rate was lower for the adolescents compared to the older adults (Table II) but linear regression models (Table IV) showed that it was statistically significantly different only at rest breathing (431 and 490 mL/s, respectively), and decreased to about the same values of 300 and 308 mL/s just prior to detection, and to 302 and 303 mL/s, respectively, at detection of the added load.

The only statistically significant effect of gender was on differential resistance just prior to detecting the added load. Height, smoking habit, upper airway allergies, or other nasal symptoms did not have a statistically significant effect on any of the respiratory variables. Among older

adults, history of chronic diseases was related to nasal airflow only at the unloaded condition ($p = 0.035$).

Weber fraction, the just noticeable difference in added upper airway resistance, varied from 0.02 to 1.16 in adolescents and from 0.01 to 2.48 in older adults. The mean values were 0.23 (SD 0.26) for adolescents and 0.84 (SD 0.79) for older adults ($p < 0.001$). Scattering of Weber fraction according to baseline resistance values illustrated the constant nature of Weber fraction (Fig. 1) among adolescents, while there was a wider inter-individual variation among older adults (Fig. 2), indicating that higher added loads were needed for detection of change.

DISCUSSION

We determined that a sample size of 40 in each group was sufficient for this clinical study which, while non-invasive, was very time consuming for each subject. Power analyses confirmed this assumption. Earlier studies^{7,10,13,31,32} involved groups from 8 to 51 subjects. In the present study, all participants were clinically healthy and free of nasal symptoms at the time of the measurements. Previous studies have shown that when individuals with acute nasal congestion are excluded, medical history was not associated with measurements of nasal patency in children.^{22,34}

The questionnaire on medical history did not include psychological factors, including anxiety disorders, which can affect perception of breathing and alter respiratory mode³⁵. Also, use of anxiolytics is unknown because they were not separated from other medicines. When recording rest breathing followed by test measurements, performed by the same experienced examiners, special attention was paid to that the study subjects were relaxed.

The finding that in the present sample among older adults history of chronic diseases was related to nasal airflow only at the unloaded condition could bias the comparisons slightly. Linear

regression models showed that other variables of the medical history as well as smoking habit did not have significant effects on respiratory variables.

In studies on respiratory function, body size has been measured as height, weight or body mass index (BMI). Somewhat surprisingly, stature, estimated in the present study as height, was not related to respiratory function which is parallel with findings of Zapletal and Chapulova³⁶ in 2-19-year old study group. In 108 20-45-year old adults it has been reported that nasal resistance decreased when the height and weight increased except in individuals with weight of 85-95 kg.³⁷ Our finding is parallel with studies using BMI in adult population^{38,39} and also with results on subjects with sleepapnea and BMI ≥ 30 kg/m^{2,40} but contradictory to a study on sample of wide age range, from 16 to 82 years.⁴¹ When diagnosing problems in the upper airway, body size may only have significance when subjects are obese.

In our study groups, gender distribution was even among adolescents, while majority of the older subjects were women which is typical in older western populations. Only one variable associated with gender was statistically significantly different, nasal resistance just prior to detection of an added load.

This study revealed that older individuals detected added load at much higher resistance, 4.55 compared to 2.26 cmH₂O/L/s in adolescents, indicating less sensitivity in the perception of increased nasal resistance in older adults. Both groups responded in similar fashion to increased load resistance by decreasing airflow rate to about 300 mL/s prior to detection indicating that there is a physiologic response to the change even before one is aware of a change in the airway environment. Thus, despite the higher airflow rate in older adults during rest breathing and despite their weaker perception of changes in upper airway resistance, the physiologic response to changes was surprisingly similar in both age groups not only at detection of the added load but also with added load prior to detection of change.

Attempts to correlate subjective sensation of nasal obstruction with objective measurements, whether defined as nasal resistance dependent on minimum nasal cross-sectional area when nasal mucosa is included in the measurements as in our method, or as nasal airflow volume or rate, are undermined by wide interindividual variation in respiratory variables and by the complex nature of sensory neural functions⁴². Still, threshold values could provide helpful guidelines. The threshold for increased nasal resistance to provoke an individual to switch from nasal to oral breathing with its' harmful effects has been suggested to be around 3.5-4.5 cmH₂O/L/s in adults⁴³, and has been reported to be 4.7 cmH₂O/L/s in adolescents⁴⁴. Thus, among the present study groups the 59-82-year-olds but not the 11-20-year-olds would probably change their breathing mode at the time of perception of increased resistance.

In the present study, resistance values were higher during inspiration compared to expiration. However, in the older group, resistance values during expiration were higher just prior to detection and at detection. Airflow rates were higher during inspiration than expiration in adolescents at all test conditions except at detection of the added load when the values were almost equal. On the contrary, among older adults airflow values were higher during expiration than inspiration except at rest breathing. Only a few studies have investigated individual's ventilatory responses to added loads during inspiration and expiration. In agreement with our findings in adolescents, Ferris et al.⁴⁵ reported that nasal resistance was higher during inspiration compared to expiration, and Muza et al.⁴⁶ found that peak inspiratory airflow was consistently higher than peak expiratory flow. Contrary to our findings, healthy adults showed expiratory resistance values to be significantly higher than inspiratory values⁹ and to be more sensitive for expiratory than inspiratory loads.⁴⁷ In agreement with our findings, Tack et al.⁴⁸ compared younger and older adults and reported a difference for resistance, sensation intensity and peak mouth pressure loads.

In the present study, a Weber fraction was utilized to describe perception of the just noticeable increase in respiratory resistance. Weber's law expressing that the ratio of the increment threshold to the background intensity has been reported to be also valid in studies of perception of added loads to inspiratory resistance.⁹⁻¹¹ In individuals with asthma, the threshold values of detection have been reported to vary widely and to be much higher compared to controls, possibly partly due to chronic adaptation to the increased airway resistance.¹⁰ In their study, Weber's law seemed to be applicable when bronchodilatation was not used. Hallani et al.⁷ reported contradictory findings, namely enhanced detection of added nasal resistance in asthmatics compared to healthy subjects. In our study, Weber's law applied reasonably well among adolescents but not among the older group. The Weber fraction was clearly higher for older adults than adolescents, indicating that sensitivity to changes in airway resistance declines with aging.

Altogether, the clearly higher threshold for perception of increased upper airway resistance and increased effort for expiration in older people may pose a health hazard in individuals with cardiovascular or respiratory diseases during physical activities or stressful conditions.

CONCLUSIONS

This study revealed that aging significantly decreases the sensitivity to the recognition of changes in upper airway resistance. Before perception of an increased applied load, both groups attempted to compensate for the increase in resistance by lowering airflow rate. This respiratory response occurred at a lower load in the younger group but the differences in respiratory rate change were similar. That is, there was a lower sensitivity in the older group but the response was similar.

REFERENCES

1. Cole P, Forsyth R, Haight JS. Effects of cold air and exercise on nasal patency. *Ann Otol Rhinol Laryngol* 1983;92:196–198.
2. Laine MT, Huggare JA, Ruoppi P. A modification of the pressure-flow technique for measuring breathing of cold air and its effect on nasal cross-sectional area. *Am J Orthod Dentofacial Orthop* 1994;105:265–269.
3. Sue-Chu M. Winter sports athletes: long-term effects of cold air exposure. *Br J Sports Med* 2012;46:397–401.
4. Cole P, Haight JS. Posture and the nasal cycle. *Ann Otol Rhinol Laryngol*. 1986;95:33–37.
5. Duggan CJ, Watson RA, Pride NB. Postural changes in nasal and pulmonary resistance in subjects with asthma. *J Asthma* 2004;41:701–707.
6. Van Holsbeke CS, Verhulst SL, Vos WG, De Backer JW, Vinchurkar SC, Verdonck PR, van Doorn JW, Nadjmi N, De Backer WA. Change in upper airway geometry between upright and supine position during tidal nasal breathing. *J Aerosol Med Pulm Drug Deliv* 2014;7:51–57.
7. Hallani M, Wheatley JR, Amis TC. Initiating oral breathing in response to nasal loading: asthmatics versus healthy subjects. *Eur Respir J* 2008a;31:800–806.
8. Wandalsen GF, Mendes AI, Solé D. Correlation between nasal resistance and different acoustic rhinometry parameters in children and adolescents with and without allergic rhinitis. *Braz J Otorhinolaryngol* 2012;78:81–86
9. Wiley RL, Zechman FW Jr. Perception of added airflow resistance in humans. *Respir Physiol* 1966/67;2:73–87.
10. Burki NK, Mitchell K, Chaudhary BA, Zechman FW. The ability of asthmatics to detect added resistive loads. *Am Rev Respir Dis* 1978;117:71–75.

11. Gottfried SB, Altose MD, Kelsen SG, Cherniack NS. Perception of changes in airflow resistance in obstructive pulmonary disorders. *Am Rev Respir Dis* 1981;124:566–570.
12. Butler J. The work of breathing through the nose. *Clin Sci* 1960;19:55-62.
13. Watson RM, Warren DW, Fischer ND. Nasal resistance, skeletal classification and mouthbreathing in orthodontic patients. *Am J Orthod* 1968;54:367–379.
14. Warren DW, Duany LF, Fischer ND. Nasal pathway resistance in normal and cleft lip and palate subjects. *Cleft Palate J* 1969;6:134-140.
15. Warren DW, Lehman MD, Hinton VA. Analysis of simulated upper airway breathing. *Am J Orthod* 1984;86:197–206.
16. Haavisto LE, Vahlberg TJ, Sipila JI. Reference values for acoustic rhinometry in children at baseline and after decongestion. *Rhinology* 2011;49:243–247.
17. Warren DW, Hairfield WM, Dalston ET. Effect of age on nasal cross-sectional area and respiratory mode in children. *Laryngoscope* 1990;100:89-93.
18. Laine T, Warren DW. Effects of age, gender, and body size on nasal cross-sectional area in children. *Eur J Orthod* 1991;13:311-316.
19. Vig PS, Zajac DJ. Age and gender effects on nasal respiratory function in normal subjects. *Cleft Palate Craniofac J* 1993;30:279-284.
20. Laine-Alava MT, Minkkinen UK. Variation of nasal respiratory pattern with age during growth and development. *Laryngoscope* 1997;107:386-390.
21. Ho WK, Wei WI, Yuen AP, Chan KL, Hui Y. Measurement of nasal geometry by acoustic rhinometry in normal-breathing Asian children. *J Otolaryngol* 1999;28:232–237.
22. Straszék SP, Schlünssen V, Sigsgaard T, Pedersen O. Reference values for acoustic rhinometry in decongested school children and adults: the most sensitive measurement for change in nasal patency. *Rhinology* 2007;45:36-39.

23. Miyamoto Y, Takeuchi K, Majima Y. Measurement of nasal patency by acoustic rhinometry in Japanese schoolchildren. *Auris Nasus Larynx* 2009;36:406-410.
24. Haavisto LE, Sipilä JI. Acoustic rhinometry in children: some practical aspects and influence of age and body surface area on results. *Am J Rhinol* 2008;22:416-419.
25. Abramson Z, Susarla S, Troulis M, Kaban L. Age-related changes of the upper airway assessed by 3dimensional computed tomography. *J Craniofac Surg* 2009;20 Suppl 1:657–663.
26. Edelstein DR. Aging of the normal nose in adults. *Laryngoscope* 1996;106:1–25.
27. Kim SW, Mo JH, Kim JW, Kim DY, Rhee CS, Lee CH, Min YG. Change of nasal function with aging in Korean. *Acta Otolaryngol Suppl* 2007;Suppl 558:90–94.
28. Cole P, Forsyth R, Haight JS. Respiratory resistance of the oral airway. *Am Rev Respir Dis* 1982;125:363–365.
29. LaFramboise WA, Standaert TA, Guthrie RD, Woodrum DE. Developmental changes in the ventilatory response of the newborn to added airway resistance. *Am Rev Respir Dis* 1987;136:1075–1083.
30. O'Donnell DE, Sani R, Younes M. External mechanical loading in conscious humans: role of upper airway mechanoreceptors. *J Appl Physiol* 1988;65:541–548.
31. Hallani M, Wheatley J, Amis T. Enforced mouth breathing decreases lung function in mild asthmatics. *Respirology* 2008b;13:553–558.
32. Newsom Davis J. Contribution of somatic receptors in the chest wall to detection of added inspiratory airway resistance. *Clin Sci* 1967;33:259–260.
33. Bennett ED, Jayson MI, Rubenstein D, Campbell EJ. The ability of man to detect added non-elastic loads to breathing. *Clin Sci* 1962;23:155–162.
34. Laine-Alava MT, Minkkinen UK. Should history of nasal symptoms be considered when estimating nasal patency? *Angle Orthod* 1999;69:126–132.

35. Paulus MP. The breathing conundrum-interoceptive sensitivity and anxiety. *Depress Anxiety*. 2013;30(4):315-320.
36. Zapletal A, Chalupová J. Nasal airflow and resistance measured by active anterior rhinomanometry in healthy children and adolescents. *Pediatr Pulmonol* 2002;33:174-180.
37. Janosević L, Dotlić J, Janosević S, Dudvarski Z, Milovanović A, Pendjer I. Computerized rhinomanometry: a study of total nasal resistance normal values. *Acta Chir Iugosl* 2009;56:51-54.
38. Demir MG, Ylmaz HB. The Relation Between Body Mass Index and Nasal Airflow. *J Craniofac Surg* 2015;26:e295-297.
39. Raza MT, Wang DY. Is nasal cavity geometry associated with body mass index, height and weight? *Indian J Otolaryngol Head Neck Surg* 2012;64:266-269.
40. Tagaya M, Nakata S, Yasuma F, Noda A, Morinaga M, Yagi H, Sugiura M, Teranishi M, Nakashima T. Pathogenetic role of increased nasal resistance in obese patients with obstructive sleep apnea syndrome. *Am J Rhinol Allergy* 2010;24:51-54.
41. Crouse U, Laine-Alava MT. Effects of age, body mass index, and gender on nasal airflow rate and pressures. *Laryngoscope* 1999;109:1503-1508.
42. Baraniuk JN. Subjective nasal fullness and objective congestion. *Proc Am Thorac Soc*. 2011 Mar;8(1):62-69.
43. Laine MT, Warren DW. Perceptual and respiratory responses to added nasal airway resistance loads in older adults. *Laryngoscope*. 1995; 105:425-428.
44. Watson RM, Warren DW, Fischer ND. Nasal resistance, skeletal classification and mouthbreathing in orthodontic patients. *Am J Orthod*. 1968; 54:367-379.
45. Ferris BG Jr, Mead J, Opie LH. Partitioning of respiratory flow resistance in man. *J Appl Physiol* 1964;19:653-658.

46. Muza SR, McDonald S, Zechman FW. Comparison of subjects' perception of inspiratory and expiratory resistance. *J Appl Physiol Respir Environ Exerc Physiol* 1984;56:211–216.

47. Bonnel AM, Mathiot MJ, Grimaud C. Inspiratory and expiratory resistive load detection in normal and asthmatic subjects. A sensory decision theory analysis. *Respiration* 1985;48:12–23.

48. Tack M, Altose MD, Cherniack NS. Effect of aging on the perception of resistive ventilatory loads. *Am Rev Respir Dis* 1982;126:463–467.

Accepted Article

LEGENDS

Figure 1. Scattering of Weber fraction according to baseline resistance (cmH₂O/L/s) in 40 adolescents and young adults.

Figure 2. Scattering of Weber fraction according to baseline resistance (cmH₂O/L/s) in 40 older adults.

Accepted Article

Table I. Relation between and cross-sectional area and resistance with the device used to create added resistance with different iris setting, calibrated at 500 mL/s.

Cross-Sectional Area of the Diaphragm (cm ²)	Resistance (cmH ₂ O/L/s)
0.50	2.2
0.48	2.5
0.45	2.9
0.43	3.4
0.41	3.6
0.38	4.2
0.36	4.6
0.34	5.1
0.32	6.3
0.31	7.3
0.28	8.9
0.26	9.5
0.25	10.2
0.23	12.9
0.21	16.4
0.20	19.1
0.18	22.1
0.16	25.1
0.15	30.6
0.14	36.9
0.13	46.3

Table II. Differences between inspiratory and expiratory airflow rate (mL/s) and nasal resistance (cmH₂O/mL/s) values among the younger (11-20 yrs) and older (59-82 yrs) study groups at different load conditions.

	Adolescents (n=40)			Older adults (n=40)		
	Inspiration mean (SD)	Expiration mean (SD)	p*	Inspiration mean (SD)	Expiration mean (SD)	p*
Rest breathing						
Resistance	2.10 (1.64)	1.77 (1.46)	0.000	2.34 (2.61)	2.00 (2.46)	<0.001
Airflow rate	431 (103)	387 (125)	0.000	490 (149)	486 (206)	0.827
Unloaded condition						
Resistance	1.54 (0.27)	1.40 (0.33)	0.000	1.75 (0.35)	1.55 (0.57)	0.001
Airflow rate	318 (58)	312 (73)	0.504	351 (82)	385 (108)	0.002
Loaded just before detection						
Resistance	2.10 (0.65)	1.95 (0.77)	0.005	4.13 (1.98)	4.70 (3.24)	0.038
Airflow rate	300 (62)	295 (64)	0.340	308 (82)	318 (110)	0.290
Loaded at detection						
Resistance	2.26 (0.73)	2.17 (0.85)	0.103	4.55 (2.29)	5.45 (4.13)	0.017
Airflow rate	302 (61)	303 (65)	0.953	303 (84)	315 (117)	0.212


*by paired t-test

Table III. Associations between differential (inspiration – expiration) resistance and airflow values ($\text{cmH}_2\text{O/L/s}$) according to age groups (1 = 11-20 yrs, 2 = 59-82 yrs) and gender (0=female, 1=male), considering the effects of smoking habit (0=no, 1=yes) medical history (0=no, 1=yes, see Subjects and Methods), with height (cm) as a covariate by linear regression analysis. Only statistically significant associations are given.

Dependent / Independent variable	Standardized coefficients <u>Beta</u>	p-value
RESISTANCE ($\text{cmH}_2\text{O/L/s}$)		
Loaded just prior to detection		
Group	0.277	0.020
<u>Gender</u>	0.339	0.026
<u>Loaded at detection</u>		
Group	0.301	0.013
AIRFLOW RATE (mL/s)		
<u>Unloaded condition</u>		
Group	0.283	0.020


Table IV. Associations between respiratory responses to added loads during inspiration according to age groups (1 = 11-20 yrs, 2 = 59-82 yrs) and gender (0=female, 1=male), considering the effects of smoking habit (0=no, 1=yes) and medical history (0=no, 1=yes, see Subjects and Methods), with height (cm) as a covariate by linear regression analyses. Only statistically significant associations are given.

Dependent / Independent variable	Standardized coefficients	p-value
RESISTANCE (cmH ₂ O/L/s)		
<u>Unloaded condition</u>		
Group	0.318	0.008
Loaded just prior to detection		
Group	0.592	<0.001
<u>Loaded at detection</u>		
Group	0.594	<0.001
AIRFLOW RATE (mL/s)		
<u>Rest breathing</u>		
Group	0.266	0.032



165x132mm (96 x 96 DPI)

Accep



165x132mm (96 x 96 DPI)

Accep