

Simultaneous Alcohol and Marijuana Use Among Young Adult Drinkers: Age-Specific Changes in Prevalence from 1977 to 2016

Yvonne M. Terry-McElrath  and Megan E. Patrick 

Background: The overall prevalence of U.S. young adult alcohol use has decreased, but little is known about historical change in related behaviors such as simultaneous alcohol and marijuana (SAM) use that may increase alcohol-related risks and societal costs. The purpose of this paper was to examine historical change in SAM use prevalence among U.S. young adult alcohol users from 1977 to 2016, and consider the extent to which observed historical change in SAM use among alcohol users reflects co-occurring change in marijuana use during these years.

Methods: Data on past 12-month alcohol, marijuana, and SAM use at up to 6 modal ages (19/20, 21/22, 23/24, 25/26, 27/28, and 29/30) were collected from 11,789 individuals (45.0% men) participating in the Monitoring the Future panel study. Annual prevalence estimates within modal age group were obtained; historical SAM use trends among alcohol and marijuana users were estimated.

Results: From 2014 to 2016, SAM use was reported by approximately 30% of alcohol users aged 19/20 and 21/22, and 20 to 25% of alcohol users aged 23/24 through 29/30. Since the mid-1990s, age-specific historical trends in SAM use prevalence among alcohol users followed 1 of 4 patterns: significant increase followed by oscillating increases/decreases (at modal age 19/20), consistent and significant increases (at modal ages 21/22, 23/24, and 25/26), stability followed by increase (at modal ages 27/28), or stability (at modal ages 29/30). In contrast, SAM use trends among marijuana users primarily reflected stability, with some evidence of a decrease across time at modal ages 19/20 and 23/24. Historical change in SAM prevalence among alcohol users was strongly and positively correlated with changes in overall marijuana use prevalence.

Conclusions: A growing proportion of early and mid-young adult alcohol users reported SAM use, with the highest risk among those in the early years of young adulthood. Young adult SAM use may continue to increase in proportion to the degree that young adult marijuana use continues to increase.

Key Words: Young Adult, Alcohol, Marijuana, Simultaneous Use.

THE WORLDWIDE ECONOMIC, health, and societal costs of alcohol-related harms are staggering (Anderson and Baumberg, 2006; Centers for Disease Control and Prevention [CDC], 2016; Institute of Alcohol Studies [IAS], 2016, 2017a,b; Manning et al., 2013; Sacks et al., 2015; World Health Organization [WHO], 2014). Average high-volume consumption, as well as drinking patterns characterized by heavy episodic drinking and high-intensity drinking, is particularly likely to be associated with alcohol-related harms (Hingson et al., 2017; Linden-Carmichael et al., 2017;

Miller et al., 2007; Patrick et al., 2016a; Rossow et al., 2013; Sacks et al., 2015; WHO, 2014).

The normative developmental trajectories for alcohol behaviors commonly associated with increased risk, such as heavy episodic and high-intensity drinking, increase from age 18 through 21/22 and decrease thereafter (Miech et al., 2017; Patrick et al., 2016b, 2017a). Thus—due to a range of both developmental and cultural reasons—the age group most likely to engage in these risky behaviors is young adults or those aged 19 to 30 (Schulenberg and Maggs, 2002). Monitoring change in both volume and prevalence of alcohol use among this subgroup is important; such information is critical for effective planning and provision of future health service and treatment needs (Dawson et al., 2015). Recent research indicates that, among young adults overall, prevalence of heavy episodic drinking and high-intensity drinking has been decreasing (Center for Behavioral Health Statistics and Quality, 2015). Data that examine differential rates of change across young adulthood provide a more nuanced picture, indicating that heavy episodic drinking and high-intensity drinking have been decreasing historically for individuals in their early 20s (Patrick and Terry-

From the Institute for Social Research (YMT-M, MEP), University of Michigan, Ann Arbor, Michigan.

Received for publication May 16, 2018; accepted August 21, 2018.

Reprint requests: Yvonne M. Terry-McElrath, MSA, Institute for Social Research, University of Michigan, Room 2312, PO Box 1248, Ann Arbor, MI 48106-1248; Tel.: 734-647-9142; Fax: 734-936-0043; E-mail: yterry@umich.edu.

Current address: Megan E. Patrick, Institute for Translational Research in Children's Mental Health & Institute of Child Development, University of Minnesota, Minneapolis, MN

© 2018 by the Research Society on Alcoholism.

DOI: 10.1111/acer.13879

McElrath, 2017; Patrick et al., 2017a), but holding steady for those in their mid-20s (Patrick et al., 2017a; Terry-McElrath and Patrick, 2016), and increasing actually for individuals at the end of young adulthood (ages 29/30; Patrick et al., 2017a).

Alcohol consumption risk also differs by the presence or absence of simultaneous use with other substances. The most common form of such simultaneous use involves using alcohol together with marijuana (Collins et al., 1998; Earleywine and Newcomb, 1997; Martin et al., 1996; Midanik et al., 2007; Substance Abuse and Mental Health Services Administration [SAMHSA], 2009). Simultaneous alcohol and marijuana (SAM) use involves the use of both substances at the same time, such that effects of the 2 substances overlap. SAM use has significant additive (and possibly synergistic) effects on cognitive, perceptual, and motor functions, over and above that of either alcohol or marijuana use alone (Belgrave et al., 1979; Chesher et al., 1976, 1977; Hartman and Huestis, 2013; Kelly et al., 2004; Lamers and Ramaekers, 2001; Ramaekers et al., 2000; Robbe, 1998). SAM use is associated with a range of serious public health concerns, including unsafe driving (Subbaraman and Kerr, 2015; Terry-McElrath et al., 2014); social consequences, depression, and physical health problems (Earleywine and Newcomb, 1997; Midanik et al., 2007; Subbaraman and Kerr, 2015); high rates of alcohol consumption (Patrick et al., 2017b; Subbaraman and Kerr, 2015); and substance dependence (Agosti et al., 2002; Martin et al., 1996; Midanik et al., 2007).

The literature on the extent of SAM use among the general adult population indicates that young adulthood has consistently been a period of elevated prevalence. The 1982 National Household Survey on Drug Abuse found that, among those aged 12 and older, the prevalence of SAM use was higher among those aged 18 to 25 and 26 to 34 than younger or older individuals (Norton and Collier, 1988). The 2000 National Alcohol Survey found while 7% of all U.S. adults reported past 12-month SAM use, rates were highest (14.8%) among those aged 18 to 29 (Midanik et al., 2007). Research combining the 2005 and 2010 National Alcohol Surveys found those aged 18 to 29 continued to report the highest prevalence of SAM use among U.S. adults at 15.3% (Subbaraman and Kerr, 2015). While research to date indicates that young adulthood is a key age for the risks and associated harms of SAM use, no data have been available that examine the extent to which this behavior is changing over time, and thus potentially affecting overall levels of harm associated with alcohol use.

While SAM use prevalence has been found to be significantly higher among those who consume high quantities of both alcohol and marijuana (Patrick et al., 2017b), SAM use is not confined to individuals reporting high consumption levels (Patrick et al., 2018; Terry-McElrath et al., 2013). SAM use is, to some degree, likely incidental to the general use of both substances (i.e., both alcohol and marijuana use prevalence may be high enough that simultaneous use is not uncommon; Hoffman et al., 2000). To the extent that SAM

use is incidental to both alcohol use and marijuana use, historical changes in marijuana use may be expected to be associated with changes in SAM use among alcohol users (both high- and low-frequency users). Published historical trend data indicate that, since the late 1970s, prevalence of past 12-month alcohol use has generally decreased for those aged 19/20, generally remained stable for those aged 21/22 through 27/28, and slightly increased for those aged 29/30 (Schulenberg et al., 2017). In contrast, historical trends for past 12-month marijuana use over the same time period show strong declines in marijuana use prevalence from the late 1970s through the early 1990s across ages. Thereafter, historical trends for marijuana use increase at varying levels and for different lengths of time across ages 19/20 through 29/30, with virtually all age groups reporting increasing prevalence from 2010 onward (Schulenberg et al., 2017). These historical increases in marijuana use prevalence bring with them the possibility that SAM use prevalence also may be increasing, thus raising the potential risk of adverse consequences among alcohol users even if overall alcohol use has been decreasing or holding steady.

The current paper aims to contribute to the literature on young adult alcohol use risk by examining historical change from 1977 through 2016 in the proportion of young adult alcohol drinkers in a general U.S. young adult sample who engaged in SAM use. Three research aims guided the analysis (i) estimate recent SAM prevalence levels among past 12-month alcohol users, by 6 modal age groups; (ii) model historical SAM use trends by modal age group among past 12-month alcohol users; and (iii) examine the degree to which observed historical SAM use trends appear to reflect historical changes in marijuana use prevalence.

MATERIALS AND METHODS

Data, Setting, and Study Population

The study utilizes data from Monitoring the Future (MTF), a national cohort-sequential study (for detailed methodology, see Bachman et al., 2015; Schulenberg et al., 2017). Briefly, a nationally representative sample of approximately 15,000 12th graders (modal age 18) from about 130 schools in the contiguous 48 states has been surveyed annually since 1976 yielding sequential cohorts. Students complete self-administered surveys, typically during a normal class period. A subsample of about 2,400 12th graders is selected from each annual sample for longitudinal follow-up by mail; substance users are oversampled (analyses include weights accounting for sampling procedures). Respondents are randomly divided with half surveyed 1 year after graduation (modal age 19) and then every 2 years after that to age 29, and half surveyed 2 years after graduation (modal age 20) and then every 2 years following to age 30, resulting in 6 follow-up surveys: 1 at each of modal ages 19/20, 21/22, 23/24, 25/26, 27/28, and 29/30. Follow-up questionnaires are mailed in the spring with a modest monetary incentive. A University of Michigan Institutional Review Board approved the study.

SAM use was included in only 1 of the 6 different questionnaire forms used in the study (randomly distributed within classroom at age 18; respondents were sent the same survey questionnaire form they responded to at age 18 for all follow-up surveys through age 30). The analytic sample was limited to 12th-grade cohorts from

1976 to 2015 who received the relevant form and responded to at least 1 young adult survey from 1977 to 2016. The average age 18 response rate for these cohorts was 82.7%. A total of 16,797 individuals from the 1976 to 2015 cohorts responded to the relevant form at age 18 and were selected for longitudinal participation. Of these individuals, 12,763 (76.0%) responded to at least 1 of the 6 follow-up surveys. Data on the outcomes of interest on at least 1 follow-up survey were available for 11,798 responding individuals (70.2% of those selected for follow-up). Men made up 45.0% of the analytic sample. Attrition adjustments are discussed below.

Measures

At each follow-up, respondents self-reported past 12-month alcohol use and marijuana use frequency; any/none use dichotomies were coded for analysis. Respondents who reported any past 12-month marijuana use were asked, "How many of the times when you used marijuana or hashish during the last year did you use it along with alcohol—that is, so that their effects overlapped?" Responses were coded as any or no SAM use (respondents reporting no past 12-month marijuana use were coded as non-SAM users).

Analysis

All analyses were conducted using SAS 13.2 (SAS Institute Inc., Cary, NC) or Joinpoint 4.6.0.0 (Statistical Methodology and Applications Branch, Surveillance Research Program, National Cancer Institute, Bethesda, MD) (National Cancer Institute [NCI], 2018). SAS survey procedures enabled appropriate modeling of the MTF study's complex sampling design. For research aim (RA) 1 (estimation of recent SAM prevalence levels by modal age group), proportions of alcohol users within each modal age group reporting SAM use for the most recent 3 years combined (2014 to 2016) were estimated using the SURVEYMEANS procedure in SAS.

For RA2 (historical trends), proportions of past 12-month alcohol users reporting any SAM use were estimated by calendar year within modal age group using SURVEYMEANS. Using the obtained estimates, trends within modal age group were modeled using Joinpoint, wherein trend lines are connected together at "joinpoints," where a significant change in slope occurs (Kim et al., 2000). In these models (as well as those described below for RA3), the focus was on the prevalence of SAM use at the specified age across sequential cohorts: age 19/20 trends included 1976 to 2015 12th-grade cohorts; age 21/22 trends included 1976 to 2013 cohorts; age 23/24 trends included 1976 to 2011 cohorts; age 25/26 trends included 1976 to 2009 cohorts; age 27/28 trends included 1976 to 2007 cohorts; and 29/30 trends included 1976 to 2005 cohorts. All Joinpoint models specified grid search (i.e., joinpoints were required to occur exactly at observations vs. between adjacent observed values) and required a minimum of 3 observations from a joinpoint to either end of the data as well as between 2 joinpoints (standard errors of slope parameters and associated p -values cannot always be calculated with <3 observations per line segment [NCI, n.d.a]). Model selection was based on permutation testing (NCI, n.d.c), with the goal of selecting the most parsimonious model. The method uses sequential permutation tests (based on Monte Carlo draws, here set at the default of 4,499 permutations) to select the final model while ensuring that the probability of overall type I error is less than the specified significance level (here set at 0.05). For the final selected model, joinpoints are identified and slope estimates and standard errors are provided. In addition, p -values for each slope change are provided from t -tests based on asymptotic normality. There are times when the p -value for an identified slope is >0.05, but the permutation test procedure identifies a slope that is significantly different from the preceding slope. This is due to the permutation test procedure not requiring asymptotic normality while

maintaining correct type I error probability; the number of joinpoints determined by the software is more reliable in such situations (NCI, n.d.b).

For RA3 (comparison of extent to which SAM use trends reflect marijuana use trends), proportions of past 12-month marijuana users reporting any SAM use were estimated by calendar year within modal age group using SURVEYMEANS, and trends within modal age group then were modeled using Joinpoint as described above. In addition, proportions of past 12-month marijuana users and past 12-month alcohol users reporting SAM use were estimated by calendar year within modal age group using SURVEYMEANS, and correlations between annual outcome prevalence estimates were estimated using the PROC CORR procedure in SAS.

All analyses were weighted using previously calculated follow-up specific attrition weights, calculated as the inverse of the probability of responding at each modal age based on covariates measured at age 18 (sex, race/ethnicity, college plans, high school grades, number of parents in the home, religiosity, parental education, alcohol use, cigarette use, marijuana use, region of country, cohort, and sampling weight correcting for oversampling of age 18 substance users).

RESULTS

Young Adult SAM Use Estimates Across Age Among Alcohol Users (RA1)

Among young adult alcohol users from 2014 to 2016 combined, SAM use prevalence was highest among early young adults, averaging 30% at ages 19/20 to 21/22 and then decreased throughout the remainder of young adulthood, reaching 19% at age 29/30 (see Table 1).

Young Adult SAM Use Across Historical Time Among Alcohol Users (RA2)

Slope estimates for historical trends from 1977 to 2016 for SAM use among past 12-month alcohol users by young adult age group are provided in Table 2; trends are presented graphically in Fig. 1.

Among alcohol users aged 19/20, SAM prevalence decreased significantly from 1977 through 1991 (modal age 19/20: slope 1 = -0.0174 , $p < 0.001$; joinpoint 1 = 1991) and then increased significantly from 1991 through 2003 (slope 2 = 0.0114 , $p < 0.001$; joinpoint 2 = 2003). SAM use among those aged 19/20 then entered a period of oscillating

Table 1. Prevalence of Simultaneous Alcohol and Marijuana (SAM) Use Among Past 12-Month Alcohol Users: U.S. Young Adults, 2014 to 2016

Modal age group	SAM prevalence among alcohol users	
	%	95% CI
19/20	30.0	24.2, 35.7
21/22	30.5	25.2, 35.9
23/24	24.2	19.7, 28.8
25/26	25.8	21.0, 30.5
27/28	24.5	19.3, 29.7
29/30	19.4	15.0, 23.8

Range of unweighted total n per modal age group: 300 to 435.

Table 2. Slope Estimates for Historical Trends in Simultaneous Alcohol and Marijuana (SAM) Use Among Past 12-Month Alcohol Users: U.S. Young Adults, 1977 to 2016

Modal age group	Slope 1 (SE)	JP 1 ^a	Slope 2 (SE)	JP 2	Slope 3 (SE)	JP 3	Slope 4 (SE)	JP 4	Slope 5 (SE)
19/20	-0.0174 (0.0016)***	1991	0.0114 (0.0026)***	2003	-0.0202 (0.0221)	2007	0.0190 (0.0159)	2012	-0.0274 (0.0163)
21/22	-0.0230 (0.0021)***	1990	0.0037 (0.0007)***						
23/24	-0.0172 (0.0023)***	1994	0.0047 (0.0012)***						
25/26	-0.0185 (0.0028)***	1993	0.0037 (0.0009)***						
27/28	-0.0237 (0.0075)**	1991	-0.0007 (0.0013)	2011	0.0240 (0.0147)				
29/30	-0.0007 (0.0008)								

^aJP = joinpoint or year in which significant change in slope occurred. For example, for the modal age group 19/20, SAM use among past 12-month alcohol users decreased significantly (slope 1 = -0.0174***) from 1977 through 1991 (JP 1) and then increased significantly (slope 2 = 0.0114***) through 2003 (JP 2). Thereafter, 3 slopes and 2 additional joinpoints were identified; *t*-test *p*-values of slope change assuming asymptotic normality were not significant, but permutation tests indicated meaningful change in trends between joinpoints. If no joinpoint is noted for a specific age group, no statistically significant change in slope estimate was observed from 1977 to 2016.

p* < 0.01, *p* < 0.001.

A total of 11,763 respondents who reported past 12-month alcohol use provided data on past 12-month SAM use.

increasing and decreasing trends (decreasing SAM use from 2003 through 2007 [slope 3 = -0.0202]; increasing SAM use from 2007 through 2012 [slope 4 = 0.0190]; decreasing SAM use from 2012 through 2016 [slope 5 = -0.0274]). During these oscillating years, *t*-test *p*-values of slope change assuming asymptotic normality were not significant, but permutation tests indicated meaningful change in trends between joinpoints.

Among alcohol users aged 21/22, 23/24, and 25/26, values for slope 1 in Table 2 show that SAM prevalence decreased significantly through the early to mid-1990s (joinpoint 1 = 1990, 1994, and 1993, for age groups 21/22, 23/24, and 25/26, respectively). For each of these 3 age groups, SAM prevalence then significantly increased from the first joinpoint through 2016 (see values for slope 2).

Among alcohol users aged 27/28, SAM prevalence decreased significantly through 1991 (slope 1 = -0.0237, *p* < 0.001). From 1991 through 2011, SAM prevalence remained statistically flat for those in this age group (slope 2 = -0.0007), and from 2011 through 2016, the slope indicated increasing prevalence (slope 3 = 0.0240; while the *p*-value of slope change was not significant, permutation tests indicated a meaningful difference in trend starting in 2011).

In contrast to age groups 19/20 through 27/28, SAM prevalence among alcohol users aged 29/30 remained statistically unchanged from 1977 through 2016 (slope 1 = -0.0007), with no joinpoints identified.

Comparison of Extent to Which SAM Use Trends Reflected Marijuana Use Trends (RA3)

Slope estimates for historical trends from 1977 to 2016 for SAM use among past 12-month marijuana users by young adult age group are provided in Table 3; trends are presented graphically in Fig. 2. Any SAM use was reported by approximately three-quarters of marijuana users across all young adult age groups. Significant but small decreases in SAM use prevalence among marijuana users were observed for 2 age groups: 19/20 (slope 1 = -0.0015, *p* < 0.05) and 23/24 (slope

1 = -0.0025, *p* < 0.001), with no joinpoints identified. For all other age groups (21/22, 25/26, 27/28, and 29/30), SAM use remained statistically unchanged, with no joinpoints identified.

Thus, while prior research showed significant changes in overall marijuana use prevalence for these age groups (significant declines from the late 1970s through the early 1990s, followed by varying levels and durations of significant increase; Schulenberg et al., 2017), among marijuana users, the prevalence of any SAM use within age groups appeared either to remain relatively consistent across time or to reflect significant but relatively small decreases across time. The Pearson correlations between year-level aggregated proportions of past 12-month marijuana users and past 12-month alcohol users reporting SAM use were significant (*p* < 0.001) and positive for all age groups (ranging from *r* = 0.845 to 0.954). Such high correlations clearly indicated that the prevalence of any SAM use among alcohol users was strongly linked with changes in the prevalence of past 12-month marijuana use.

DISCUSSION

On average from 2014 to 2016, any SAM use was reported by approximately 30% of alcohol users aged 19/20 and 21/22, and between approximately 20 and 25% of alcohol users aged 23/24 through 29/30. Historical trends in annual SAM prevalence among young adult alcohol users followed 1 of 4 patterns from the mid-1990s onward: (i) significant prevalence increase followed by oscillating change (ages 19/20); (ii) significant and consistent prevalence increase (ages 21/22 through 25/26); (iii) stability followed by increasing prevalence (ages 27/28); or (iv) prevalence stability (ages 29/30). In contrast, SAM use prevalence among marijuana users was generally stable across time within age (excluding ages 19/20 and 23/24, which showed small but significant decreases across time). Among young adult alcohol users, changes in SAM prevalence were strongly and positively correlated with changes in overall marijuana use trends. Trend changes in




Fig. 1. Trends by modal age group in prevalence of simultaneous alcohol and marijuana (SAM) use among past 12-month alcohol users: U.S. young adults, 1977 to 2016. Note: Slope estimates for all trends are provided in Table 2.

the use of a secondary substance—including marijuana—may meaningfully affect risks associated with alcohol use.

Based on the current study, between one-fifth and almost one-third of recent U.S. young adult alcohol users reported using alcohol and marijuana so that their effects overlapped, thus placing themselves and others at risk of associated consequences. SAM use is particularly associated with situations that involve heavy episodic drinking or high-intensity drinking (Patrick et al., 2017b). Drinking at these levels is associated with severe and even life-threatening impairment, resulting in possible alcohol poisoning and suppression of vital life functions (National Institute on Alcohol Abuse and Alcoholism, 2015). The additive effects of SAM use on cognitive, perceptual, and motor functions (Belgrave et al., 1979; Chesher et al., 1976, 1977; Kelly et al., 2004; Lamers and

Ramaekers, 2001; Ramaekers et al., 2000; Robbe, 1998) result in risk not only to individual users, but the public at large. SAM use is frequently associated with situations that involve the public, such as parks/beaches, parties, and—most alarmingly—cars (Pakula et al., 2009; Terry-McElrath et al., 2013). SAM use is strongly linked to increased drunk driving (Subbaraman and Kerr, 2015), tickets/warnings, and accidents (Terry-McElrath et al., 2014). For example, data from the state of Washington (which voted to legalize recreational marijuana use in 2012) indicated that 9% of drivers who tested positive for alcohol and drugs in fatal crashes from 2010 to 2014 tested positive for cannabinoids only, compared with 13% who tested positive for cannabinoids and alcohol, and an additional 4% who tested positive for cannabinoids, alcohol, and other drugs (Washington Traffic Safety

Table 3. Slope Estimates for Historical Trends in Simultaneous Alcohol and Marijuana (SAM) Use Among Past 12-Month Marijuana Users: U.S. Young Adults, 1977 to 2016

Modal age group	Slope 1 (SE)
19/20	-0.0015 (0.0007)*
21/22	-0.0005 (0.0006)
23/24	-0.0025 (0.0007)***
25/26	-0.0005 (0.0008)
27/28	-0.0002 (0.0013)
29/30	-0.0001 (0.0017)

* $p < 0.05$, *** $p < 0.001$.

A total of 6,578 respondents who reported past 12-month marijuana use provided data on past 12-month SAM use.

No joinpoints are noted for the above trends as no statistically significant changes in slope estimates were observed over time.

Commission, 2016). In both Washington and Colorado (which also voted to legalize recreational marijuana use in 2012), there have been indications that impaired driving involving marijuana—particularly marijuana combined with alcohol—is increasing (Rocky Mountain High Intensity Drug Trafficking Area, 2015; Washington Traffic Safety Commission, 2016).

The current study found that following the mid-1990s, the percentage of alcohol users also reporting SAM use significantly increased for multiple years among all but young adults in their very late 20s. Among these early to mid-young adults, the potential level of risk associated with alcohol use increased due to increasing SAM use prevalence. These findings, together with results of age-specific research on trends in heavy episodic drinking or high-intensity drinking (Patrick et al., 2017a; Terry-McElrath and Patrick, 2016), may offer a potential explanation for at least part of the observed but unexplained recent increases in demand for alcohol-related emergency department (ED) and inpatient services at U.S. hospitals (Mullins et al., 2016; National Institutes of Health, 2013; White et al., 2018). National ED visits for U.S. adults aged 18 and older due to alcohol-related causes increased at a significantly greater rate than overall ED visits from 2001 to 2011 (Mullins et al., 2016), and overall ED visits involving both acute and chronic alcohol consumption rose dramatically between 2006 and 2014 (White et al., 2018). These increases have been particularly puzzling since (as noted previously) there have not been co-occurring increases observed in alcohol-related behaviors likely to result in ED use such as overall heavy episodic drinking or high-intensity drinking (e.g., Center for Behavioral Health Statistics and Quality, 2015; Hingson and White, 2013; Kann et al., 2016; Patrick and Terry-McElrath, 2017; Patrick et al., 2013). White and colleagues (2018) reported that approximately 1 in 6 alcohol-related ED visits involved alcohol use combined with other drugs. It is possible that changes in the prevalence or frequency of combining alcohol with other drugs could lead to changes in acute and chronic alcohol-related problems, whether or not alcohol intensity increases (Patrick and Terry-

McElrath, 2018). While marijuana is only one of many substances that can be used simultaneously with alcohol, combined results of research on age-specific trends of heavy episodic drinking and high-intensity drinking (Patrick et al., 2017a; Terry-McElrath and Patrick, 2016), together with the current study, indicate that overall alcohol risk indeed may have increased across young adulthood. Further research that could quantify the changes in overall alcohol-related harms—such as health services utilization—associated with trend changes in heavy episodic drinking or high-intensity drinking compared with trend changes in SAM use is needed. Related to this issue is the need for studies examining how simultaneous use of alcohol and other substances (such as nonmedical use of prescription drugs) may be impacting health services utilization.

The results of the current study indicate that the percentage of alcohol users engaging in SAM use will likely continue to increase in proportion to the degree that young adult marijuana use continues to increase. The ages of peak past 12-month marijuana use prevalence among the U.S. adult population (i.e., ages 19 to 21 as reported in SAMHSA, 2016) were strongly reflected in the current study's observed ages of peak SAM use among alcohol users. In addition, the strong positive correlations between annual prevalence of marijuana use and SAM use among young adult alcohol users indicates that changes in marijuana use prevalence have had a meaningful impact on alcohol use risk. Across young adults, the prevalence of past 12-month marijuana use has increased notably since 2010: from 31 to 41% among those aged 19/20, 34 to 41% among those aged 21/22, 31 to 36% for those aged 23/24, 26 to 29% for those aged 25/26, 22 to 30% for those aged 27/28, and 22 to 26% for those aged 29/30 (Schulenberg et al., 2017). These increases, together with the strong correlations between marijuana use and SAM use among alcohol users, indicate growing overall risk from alcohol use among young adults at all ages.

Contrary to these overall trends, there is some indication that SAM use prevalence decreased somewhat from 1976 to 2016 among marijuana users aged 19/20 and 23/24, and from 2012 to 2016 among alcohol users aged 19/20. The observed decreases in SAM prevalence trends for individuals in these specific subgroups may indicate increasing substitution of marijuana for alcohol use (Baggio et al., 2018; Kendall, 2016) and thus an observed decrease in SAM use. Yet, recent changes in the legal status of marijuana in various states (via medicalization, decriminalization, and legalization) have been forecasted to result in increased intentions to use and actual marijuana use (Hasin et al., 2015; Keyes et al., 2016; Miech et al., 2015; Palamar et al., 2014). Whether or not such forecasts are correct, to the extent that marijuana use increases, alcohol-related harms associated with SAM use will likely increase apace.

While SAM use prevalence was found among all young adult age groups, these data showed that rates




Fig. 2. Trends by modal age group in prevalence of simultaneous alcohol and marijuana (SAM) use among past 12-month marijuana users: U.S. young adults, 1977 to 2016. Note: Slope estimates for all trends are provided in Table 3.

were highest among those ages 19/20 and 21/22. Efforts to prevent and/or reduce SAM use among all young adult alcohol users are clearly warranted that effectively communicate and seek to mitigate the risks associated with SAM use over and above those associated with use of either substance separately. The developmental processes underlying lower SAM use prevalence among alcohol users at the end of young adulthood may be similar to previously observed developmental changes in marijuana and alcohol use. Research has proposed both circumstantial (passive settings, environmental constraints, influences) and functional (active decision making regarding substances, temporal order of use) explanations of how and why substances are used simultaneously (Pakula et al., 2009). Some research has found adolescent SAM use is incidental to general alcohol or marijuana use

(Hoffman et al., 2000); other studies indicate the behavior is both incidental to general use and associated with specific simultaneous use reasons or expectancies (Terry-McElrath et al., 2013). Adolescent and adult SAM use is strongly associated with social contexts, negative emotional states, and perceived dependence (Pakula et al., 2009; Terry-McElrath et al., 2013). Reasons for alcohol and marijuana use (separately) change significantly with age. Moving from early to later young adulthood, individuals are (i) more likely to report using both substances to relax, using alcohol to sleep and because it tastes good, and using marijuana to get high; and (ii) less likely to report use of both substances for dealing with negative emotional states, perceived dependence, and to increase the effect of other drugs (Patrick et al., 2011). While such changes in reasons for general alcohol

and marijuana use could help explain the observed significant decrease in any SAM use across young adulthood in the current study, research is needed that investigates reasons for combining these substances, in particular.

Limitations

The current study's findings should be considered within their limitations. Findings may not generalize to individuals who drop out of high school prior to 12th grade; lower educational attainment is associated with higher marijuana and other substance use (Tice et al., 2017). Attrition weighting was based on variables measured at 12th grade (including substance use) and was not able to take into account that risk of dropout also likely correlates with substance use at later ages. Across the key outcomes of interest (alcohol, marijuana, and SAM use), the largest loss of data to missingness occurred between the 12th-grade survey and first follow-up at age 19/20. Thereafter, the percentage of data lost to missingness on these key measures increased approximately 5 percentage points per follow-up. The SAM measure available did not allow for assessing SAM use frequency, so results are limited to examining any versus no SAM use. Further, all data are based on self-reports, which have been found to be reasonably reliable and valid under conditions which the MTF study strives to provide (Brener et al., 2003; Miech et al., 2017; O'Malley et al., 1983). These limitations notwithstanding, the current study contributes significantly to available knowledge on SAM use among young adults in the United States.

CONCLUSIONS

A growing proportion of early and mid-young adult drinkers report SAM use, with the highest risk belonging to those in the early years of this developmental period. Efforts to understand the costs and consequences of alcohol use may be significantly enhanced by actively seeking to model how changes in the use of a secondary substance (such as marijuana) may affect the nature of how alcohol is used (at least for some individuals) and thus the level of risk associated with use.

FUNDING

Development of this manuscript was supported by research grants R01AA023504 and R01AA025037 from the National Institute on Alcohol Abuse and Alcoholism and R01DA001411 and R01DA016575 from the National Institute on Drug Abuse. The study sponsors had no role in the study design, collection, analysis, or interpretation of the data, writing of the manuscript, or the decision to submit the paper for publication. The content is solely the responsibility of the authors and does not necessarily represent the official views of the study sponsors.

AUTHOR CONTRIBUTIONS

YMT-M and MEP contributed equally to this manuscript with respect to concept, interpretation of data, and the write-up of results. YMT-M conducted the analyses.

CONFLICT OF INTEREST

None.

REFERENCES

- Agosti V, Nunes E, Levin F (2002) Rates of psychiatric comorbidity among US residents with lifetime cannabis dependence. *Am J Drug Alcohol Abuse* 28:643–652.
- Anderson P, Baumberg B (2006) Alcohol in Europe: A Public Health Perspective. A Report for the European Commission. Institute of Alcohol Studies, London.
- Bachman JG, Johnston LD, O'Malley PM, Schulenberg JE, Miech RA (2015) The monitoring the future project after four decades: design and procedures (Monitoring the future occasional paper no. 82). Institute for Social Research, University of Michigan, Ann Arbor, MI. Available at: <http://monitoringthefuture.org/pubs/occpapers/mtf-occ82.pdf>. Accessed March 27, 2018.
- Baggio M, Chong A, Kwon S (2018) Marijuana and alcohol evidence using border analysis and retail sales data. Available at: <https://ssrn.com/abstract=3063288> or <https://doi.org/10.2139/ssrn.3063288>. Accessed September 5, 2018.
- Belgrave BE, Bird KD, Chesher GB, Jackson DM, Lubbe KE, Starmer GA, Teo RK (1979) The effect of (–) trans-delta9-tetrahydrocannabinol, alone and in combination with ethanol, on human performance. *Psychopharmacology* 62:53–60.
- Brener ND, Billy JOG, Grady WR (2003) Assessment of factors affecting the validity of self-reported health-risk behavior among adolescents: evidence from the scientific literature. *J Adolesc Health* 33:436–457.
- Center for Behavioral Health Statistics and Quality (2015) Behavioral health trends in the United States: results from the 2014 National Survey on Drug Use and Health (HHS Publication No. SMA 15-4927, NSDUH Series H-50). Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Rockville, MD.
- Centers for Disease Control and Prevention (2016) Excessive drinking is draining the US economy [CDC Web site]. January 12. Available at: <https://onlinelibrary.wiley.com/page/journal/15300277/homepage/forauthors.html>. Accessed March 27, 2018.
- Chesher GB, Franks HM, Hensley VR, Hensley WJ, Jackson DM, Starmer GA, Teo RK (1976) The interaction of ethanol and delta9-tetrahydrocannabinol in man: effects on perceptual, cognitive and motor functions. *Med J Aust* 2:159–163.
- Chesher GB, Franks HM, Jackson DM, Starmer GA, Teo RK (1977) Ethanol and delta9-tetrahydrocannabinol: interactive effects on human perceptual, cognitive and motor functions. *Med J Aust* 1:478–481.
- Collins RL, Ellickson PL, Bell RM (1998) Simultaneous polydrug use among teens: prevalence and predictors. *J Subst Abuse* 10:233–253.
- Dawson DA, Goldstein RB, Saha TD, Grant BF (2015) Changes in alcohol consumption: United States, 2001–2002 to 2012–2013. *Drug Alcohol Depend* 148:56–61.
- Earleywine M, Newcomb MD (1997) Concurrent versus simultaneous polydrug use: prevalence, correlates, discriminant validity, and prospective effects on health outcomes. *Exp Clin Psychopharmacol* 5:353–364.
- Hartman RL, Huestis MA (2013) Cannabis effects on driving skills. *Clin Chem* 59:478–492.
- Hasin DS, Wall M, Keyes KM, Cerdá M, Schulenberg J, O'Malley PM, Galea S, Pacula R, Feng T (2015) Medical marijuana laws and adolescent marijuana use in the USA from 1991 to 2014: results from annual, repeated cross-sectional surveys. *Lancet Psychiatry* 2:601–608.

- Hingson R, Zha W, Smyth D (2017) Magnitude and trends in heavy episodic drinking, alcohol-impaired driving, and alcohol-related mortality and overdose hospitalizations among emerging adults of college ages 18-24 in the United States, 1998-2014. *J Stud Alcohol Drugs* 78:540-548.
- Hingson RW, White A (2013) Trends in extreme binge drinking among US high school seniors. *JAMA Pediatr* 167(11):996-998.
- Hoffman JH, Barnes GM, Welte JW, Dintcheff BA (2000) Trends in combinational use of alcohol and illicit drugs among minority adolescents, 1983-1994. *Am J Drug Alcohol Abuse* 26:311-324.
- Institute of Alcohol Studies (2016) *The Economic Impacts of Alcohol*. Factsheet Series. Institute of Alcohol Studies, London.
- Institute of Alcohol Studies (2017a) *Crime and Social Impacts of Alcohol*. Factsheet Series. Institute of Alcohol Studies, London.
- Institute of Alcohol Studies (2017b) *The Health Impacts of Alcohol*. Factsheet Series. Institute of Alcohol Studies, London.
- Kann L, McManus T, Harris WA, Shanklin SL, Flint KH, Hawkins J, Queen B, Lowry R, Olsen EO, Chyen D, Whittle L, Thornton J, Lim C, Yamakawa Y, Brener N, Zaza S (2016) Youth risk surveillance—United States, 2015. *MMWR Morb Mortal Wkly Rep* 65:1-174.
- Kelly E, Drake S, Ross J (2004) A review of drug use and driving: epidemiology, impairment and risk perceptions. *Drug Alcohol Rev* 23:319-344.
- Kendall J (2016) Report: beer volumes declining in markets where recreational cannabis is legal. Available at: <https://www.brewbound.com/news/report-beer-volumes-declining-markets-recreational-cannabis-legal/#Report:%20Beer%20Volumes%20Declining%20in%20Markets%20Where%20Recreational%20Cannabis%20is%20Legal>. Accessed September 5, 2018.
- Keyes KM, Wall M, Cerdá M, Schulenberg J, O'Malley PM, Galea S, Feng T, Hasin DS (2016) How does state marijuana policy affect US youth? Medical marijuana laws, marijuana use and perceived harmfulness: 1991-2014. *Addiction* 111:2187-2195.
- Kim HJ, Fay MP, Feuer EJ, Midthune DN (2000) Permutation tests for jointpoint regression with applications to cancer rates. *Stat Med* 19:335-351 (correction: 2001;20:655).
- Lamers C, Ramaekers J (2001) Visual search and urban city driving under the influence of marijuana and alcohol. *Hum Psychopharmacol* 16:393-401.
- Linden-Carmichael AN, Vasilenko SA, Lanza ST, Maggs JL (2017) High-intensity drinking versus heavy episodic drinking: prevalence rates and relative odds of alcohol use disorder across adulthood. *Alcohol Clin Exp Res* 41:1754-1759.
- Manning M, Smith C, Mazerolle P (2013) The societal costs of alcohol misuse in Australia. *Trends and Issues in Crime and Criminal Justice* April (No. 454):1-6.
- Martin CS, Kaczynski NA, Maisto SA, Tarter RE (1996) Polydrug use in adolescent drinkers with and without DSM-IV alcohol abuse and dependence. *Alcohol Clin Exp Res* 20:1099-1108.
- Midanik LT, Tam TW, Weisner C (2007) Concurrent and simultaneous drug and alcohol use: results of the 2000 National Alcohol Survey. *Drug Alcohol Depend* 90:72-80.
- Miech RA, Johnston L, O'Malley PM, Bachman JG, Schulenberg J, Patrick ME (2015) Trends in use of marijuana and attitudes toward marijuana among youth before and after decriminalization: the case of California 2007-2013. *Int J Drug Policy* 26:336-344.
- Miech RA, Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE (2017) *Monitoring the Future National Survey Results on Drug Use, 1975-2014: Volume I, Secondary School Students*. Institute for Social Research, The University of Michigan, Ann Arbor, MI. Available at: http://monitoringthefuture.org/pubs/monographs/mtf-vol1_2016.pdf. Accessed March 27, 2018.
- Miller JW, Naimi TS, Brewer RD, Jones SE (2007) Binge drinking and associated health risk behaviors among high school students. *Pediatrics* 119:76-85.
- Mullins PM, Mazer-Amirshahi M, Pines JM (2016) Alcohol-related visits to US emergency departments, 2001-2011. *Alcohol Alcohol* 52:119-125.
- National Cancer Institute (2018) *Joinpoint Regression Program*. Version 4.6.0.0. Statistical Research and Applications Branch, National Cancer Institute, Bethesda, MD.
- National Cancer Institute (n.d.a) Grid search regression method – details. Available at: <https://surveillance.cancer.gov/help/joinpoint/setting-parameters/method-and-parameters-tab/method/method-grid-search-or-hudsons>. Accessed September 5, 2018.
- National Cancer Institute (n.d.c) Permutation test details. Available at: <https://surveillance.cancer.gov/help/joinpoint/setting-parameters/method-and-parameters-tab/model-selection-method/permutation-tests/permutation-test-details>. Accessed September 5, 2018.
- National Institute on Alcohol Abuse and Alcoholism (2015) *Alcohol overdose: the dangers of drinking too much* (NIAAA Web site) October. Available at: <https://pubs.niaaa.nih.gov/publications/alcoholoverdosefactsheet/overdosefact.htm>. Accessed March 27, 2018.
- National Cancer Institute (n.d.b) Non-significant changes in slopes. Available at: <https://surveillance.cancer.gov/help/joinpoint/tech-help/frequently-asked-questions/non-significant-change-in-slopes>. Accessed September 5, 2018.
- National Institutes of Health (2013) *Alcohol-related emergency department visits and hospitalizations and their co-occurring drug-related, mental health, and injury conditions in the United States: findings from the 2006-2010 Nationwide Emergency Department Sample and Nationwide Inpatient Sample*. US Alcohol Epidemiologic Data Reference Manual, Vol. 9. National Institutes of Health, National Institute on Alcohol Abuse and Alcoholism, Bethesda, MD.
- Norton R, Colliver J (1988) Prevalence and patterns of combined alcohol and marijuana use. *J Stud Alcohol* 49:378-380.
- O'Malley PM, Bachman JG, Johnston LD (1983) Reliability and consistency of self-reports of drug use. *Int J Addict* 18:805-824.
- Pakula B, MacDonald S, Stockwell T (2009) Settings and functions related to simultaneous use of alcohol with marijuana or cocaine among clients in treatment for substance abuse. *Subst Use Misuse* 44:212-226.
- Palamar JJ, Ompad DC, Petkova E (2014) Correlates of intentions to use cannabis among US high school seniors in the case of cannabis legalization. *Int J Drug Policy* 25:424-435.
- Patrick ME, Counce JM, Fairlie AM, Atkins DC, Lee CM (2016a) Day-to-day variation in high-intensity drinking, expectancies, and positive and negative alcohol-related consequences. *Addict Behav* 58:110-116.
- Patrick ME, Kloska DD, Terry-McElrath YM, Lee CM, O'Malley PM, Johnston LD (2018) Patterns of simultaneous and concurrent alcohol and marijuana use among adolescents. *Am J Drug Alcohol Abuse* 44:441-451.
- Patrick ME, Schulenberg JE, Martz ME, Maggs JL, O'Malley PM, Johnston LD (2013) Extreme binge drinking among 12th-grade students in the United States: prevalence and predictors. *JAMA Pediatr* 167:1019-1025.
- Patrick ME, Schulenberg JE, O'Malley PM, Maggs JL, Kloska DD, Johnston LD, Bachman JG (2011) Age-related changes in reasons for using alcohol and marijuana from ages 18 to 30 in a national sample. *Psychol Addict Behav* 25:330-339.
- Patrick ME, Terry-McElrath YM (2017) High-intensity drinking by underage young adults in the United States. *Addiction* 112:82-93.
- Patrick ME, Terry-McElrath YM (2018) Commentary on White and colleagues: trends in alcohol-related emergency department visits in the United States: results from the nationwide emergency department sample, 2006-2014. *Alcohol Clin Exp Res* 42:489-491.
- Patrick ME, Terry-McElrath YM, Kloska DD, Schulenberg JE (2016b) High-intensity drinking among young adults in the United States: prevalence, frequency, and developmental change. *Alcohol Clin Exp Res* 40:1905-1912.
- Patrick ME, Terry-McElrath YM, Miech RA, Schulenberg JE, O'Malley PM, Johnston LD (2017a) Age-specific prevalence of binge and high-intensity drinking among U.S. young adults: changes from 2005 to 2015. *Alcohol Clin Exp Res* 41:1319-1328.
- Patrick ME, Veliz PT, Terry-McElrath YM (2017b) High-intensity and simultaneous alcohol and marijuana use among high school seniors in the United States. *Subst Abuse* 38:498-503.

- Ramaekers J, Robbe H, O'Hanlon J (2000) Marijuana, alcohol and actual driving performance. *Hum Psychopharmacol* 15:551–558.
- Robbe H (1998) Marijuana's impairing effects on driving are moderate when taken alone but severe when combined with alcohol. *Hum Psychopharmacol* 13:S70–S78.
- Rocky Mountain High Intensity Drug Trafficking Area (2015) *The Legalization of Marijuana in Colorado: The Impact. Vol 3 Preview 2015*. Rocky Mountain High Intensity Drug Trafficking Area, Investigative Support Center, Denver, CO.
- Rosow I, Bogstrand ST, Ekeberg Ø, Normann PT (2013) Associations between heavy episodic drinking and alcohol related injuries: a case control study. *BMC Public Health* 13:1076.
- Sacks JJ, Gonzales KR, Bouchery EE, Tomedi LE, Brewer RD (2015) 2010 National and state costs of excessive alcohol consumption. *Am J Prev Med* 49:e78–e79.
- Schulenberg JE, Johnston LD, O'Malley PM, Bachman JG, Miech RA, Patrick ME (2017) *Monitoring the Future National Survey Results on Drug Use, 1975-2016: Volume II, College Students and Adults Ages 19-55*. Institute for Social Research, The University of Michigan, Ann Arbor, MI. Available at: http://monitoringthefuture.org/pubs/monographs/mtf-vol2_2016.pdf. Accessed March 28, 2018.
- Schulenberg ME, Maggs JL (2002) A developmental perspective on alcohol use and heavy drinking during adolescence and the transition to young adulthood. *J Stud Alcohol* (14):54–70.
- Subbaraman MS, Kerr WC (2015) Simultaneous versus concurrent use of alcohol and cannabis in the National Alcohol Survey. *Alcohol Clin Exp Res* 39:872–879.
- Substance Abuse and Mental Health Services Administration (2009) *The NSDUH report: concurrent illicit drug and alcohol use*. Substance Abuse and Mental Health Services Administration, Office of Applied Studies, Rockville, MD.
- Substance Abuse and Mental Health Services Administration (2016) *Results from the 2015 National Survey on Drug Use and Health: detailed tables*. Substance Abuse and Mental Health Services Administration, Office of Applied Studies, Rockville, MD. Available at: [https://www.samhsa.gov/data/sites/default/files/NSDUH-DetTabs-2015/NSDUH-DetTabs-2015.pdf](https://www.samhsa.gov/data/sites/default/files/NSDUH-DetTabs-2015/NSDUH-DetTabs-2015/NSDUH-DetTabs-2015.pdf). Accessed July 9, 2018.
- Terry-McElrath YM, O'Malley PM, Johnston LD (2013) Simultaneous alcohol and marijuana use among US high school seniors from 1976-2011: trends, reasons, and situations. *Drug Alcohol Depend* 133:71–79.
- Terry-McElrath YM, O'Malley PM, Johnston LD (2014) Alcohol and marijuana use patterns associated with unsafe driving among US high school seniors: high use frequency, concurrent use, and simultaneous use. *J Stud Alcohol Drugs* 75:378–389.
- Terry-McElrath YM, Patrick ME (2016) Intoxication and binge and high-intensity drinking among US young adults in their mid-20s. *Subst Abuse* 37:597–605.
- Tice P, Lipari RN, Van Horn SL (2017) *Substance use among 12th grade aged youths, by dropout status. The CBHSQ report*. US Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Rockville, MD. Available at: https://www.samhsa.gov/data/sites/default/files/report_3196/ShortReport-3196.html. Accessed March 27, 2018.
- Washington Traffic Safety Commission (2016) *Driver Toxicology Testing and the Involvement of Marijuana in Fatal Crashes, 2010-2014*. Washington Traffic Safety Commission, Olympia, WA.
- White AM, Slater ME, Ng G, Hingson R, Breslow R (2018) Trends in alcohol-related emergency department visits in the United States: results from the nationwide emergency department sample, 2006 to 2014. *Alcohol Clin Exp Res* 42:352–359.
- World Health Organization (2014) *Global Status Report on Alcohol and Health 2014. Management of Substance Abuse*, Department of Mental Health and Substance Abuse, World Health Organization, Geneva.