

REVIEW ARTICLE

PI3K signalling in leptin receptor cells: Role in growth and reproduction

David Garcia-Galiano¹  | Beatriz C. Borges^{1,2} | Susan J. Allen¹  | Carol F. Elias^{1,3} 

¹Department of Molecular & Integrative Physiology, University of Michigan, Ann Arbor, Michigan

²Kresge Hearing Research Institute and Department of Otolaryngology - Head and Neck Surgery, University of Michigan, Ann Arbor, Michigan

³Department of Obstetrics and Gynecology, University of Michigan, Ann Arbor, Michigan

Correspondence

Carol F. Elias, Department of Molecular and Integrative Physiology, University of Michigan, Ann Arbor, MI.
Email: cfelias@umich.edu

Funding information

Grants from NIH (R01 HD069702; R21 HD090567; R03 HD092855; P30 DK020572) and pilot grants from the Reproductive Sciences Program (RSP) of the University of Michigan, USA. Postdoctoral fellowships from The Lalor Foundation, USA (DGG) and the São Paulo Research Foundation, Brazil (BCB, 2016/10398-0).

Nutrition and growth are important signals for pubertal development, although how they are perceived and integrated in brain circuits has not been well defined. Growth hormones and metabolic cues both recruit phosphatidylinositol 3-kinase (PI3K) signalling in hypothalamic sites, although whether they converge into the same neuronal population(s) is also not known. In this review, we discuss recent findings from our laboratory showing the role of PI3K subunits in cells directly responsive to the adipocyte-derived hormone leptin in the coordination of growth, pubertal development and fertility. Mice with deletion of PI3K p110 α and p110 β catalytic subunits in leptin receptor cells (LR $\Delta\alpha+\beta$) have a lean phenotype associated with increased energy expenditure, locomotor activity and thermogenesis. The LR $\Delta\alpha+\beta$ mice also show deficient growth and delayed puberty. Deletion of a single subunit (ie, p110 α) in LR cells (LR $\Delta\alpha$) causes a similar phenotype of increased energy expenditure, deficient growth and delayed pubertal development, indicating that these functions are preferably controlled by p110 α . The LR $\Delta\alpha$ mice show enhanced leptin sensitivity in metabolic regulation but, remarkably, these mice are unresponsive to the effects of leptin on growth and puberty. PI3K is also recruited by insulin and a subpopulation of LR neurons is responsive to i.c.v. insulin administration. Deletion of insulin receptor in LR cells causes no changes in body weight or linear growth and induces only a mild delay in pubertal completion. Our findings demonstrate that PI3K in LR cells plays an essential role in growth and reproduction. We will also discuss the potential neural pathways underlying these effects.

KEYWORDS

bone, hypothalamus, insulin, metabolism, reproduction

1 | INTRODUCTION

The role of metabolic cues in pubertal progression has been well described in different mammalian species.¹⁻⁴ Among these cues, the adipocyte-derived hormone leptin plays a critical role. The circulating levels of leptin are positively correlated with the amount of body fat.⁵⁻⁷ Disruption of leptin signalling causes obesity, diabetes and a number of neuroendocrine dysfunctions, including changes in bone accrual and growth, pubertal arrest and infertility.^{5,8-17} Genetic mutations of leptin or leptin receptor in humans are rare.^{11,18,19} Obese

individuals exhibit hyperleptinaemia with a low or absent response to leptin administration.^{20,21} The underlying mechanisms are diverse, although impairment of leptin receptor signalling appears to play a major role. Therefore, leptin “resistance” as a result of the disruption of leptin receptor (LR) function or signalling capacity may cause most of the metabolic and neuroendocrine deficits observed in obese subjects.^{22,23} Excess adiposity and high leptin levels also impact reproductive function and growth by disrupting steroidogenesis in males and females, aggravating ovulatory disorders, and altering bone formation.^{17,23-27} The increasing prevalence of childhood

obesity has been associated with early puberty in girls.²⁸⁻³⁰ Thus, it has become clear that metabolic cues are key for the coordinated timing of growth and puberty and maintenance of a healthy reproductive function in adult life.

The LR is a member of the class I cytokine receptor family comprised of several isoforms.^{10,31-35} The LR long form (LRb) contains three intracellular tyrosine residues that are phosphorylated by JAK2. Selective blockade of each one of these residues has produced compelling findings regarding the effects of specific signalling pathways in leptin function.³⁶⁻³⁸ For example, disruption of Tyr1138 or leptin-induced signal transducer and activator of transcription (STAT) 3 signalling generates severely obese mice, with disruption of thyroid and adrenal axes. However, these mice show improved glucose homeostasis, growth and fertility compared to loss-of-function mutation of leptin or leptin receptor genes (ie, *ob/ob* or *Lep^{ob}* and *db/db* or *Lep^{db}* mice). Lack of either leptin-induced SHP2-ERK (Tyr985) or STAT5 (Tyr1077) caused only mild metabolic or reproductive phenotypes.³⁷⁻³⁹

Leptin also recruits phosphatidylinositol 3-kinase (PI3K) signalling,⁴⁰⁻⁴³ a major molecular pathway associated with metabolic regulation, insulin signalling, glucose homeostasis and neuroendocrine function.^{40,42,44-49} However, the requirement of PI3K in leptin responsive cells for growth and pubertal maturation had not been described until very recently.⁵⁰ In this review, we discuss findings showing that PI3K signalling functions as a key molecular pathway linking leptin and the neuroendocrine axes. Because PI3K is also recruited by insulin, we discuss its potential role in the integration of both metabolic cues.^{43,51,52}

2 | PI3K IN LEPTIN ACTION: MOLECULAR COMPONENTS

Studies from different groups support the hypothesis that leptin recruits the PI3K signalling pathway to exert some of its effects.^{40,42} For example, leptin hyperpolarises a subset of arcuate nucleus neurones via activation of an ATP sensitive potassium channel, resulting in reduced firing rates. Inhibitors of PI3K blocked this effect.^{44,53,54} Similarly, leptin-mediated depolarising effects on pro-opiomelanocortin (POMC) neurones are dependent on PI3K signalling.⁵⁵ Pharmacological inhibition of PI3K blocked the ability of i.c.v. leptin to reduce food intake^{40,44} and precluded the effects of hypothalamic leptin administration to inhibit white adipose tissue lipogenesis.⁵⁶

Multiple classes of PI3Ks exist. In particular, class IA is primarily associated with metabolic regulation.⁵⁷⁻⁵⁹ PI3K class IA consists of heterodimers of one regulatory and one catalytic subunit. There are typically five regulatory subunits, often called the p85s, and the catalytic subunits comprise three variants referred to as p110s.^{57,59} Activation of the heterodimers occurs when the p85 binds to the insulin receptor substrate (IRS) and position the catalytic subunit in the cellular membrane, where PI3K catalyses the phosphorylation of the phosphatidylinositol bisphosphate to PIP3, which, in turn, recruits and activates downstream pathways.

The p110 α and p110 β subunits are ubiquitously expressed, whereas the p110 γ is almost absent in the brain.^{60,61} Because of the distribution pattern and the lack of p110 γ in hypothalamic sites, most of the studies in metabolic control have focused on the roles of p110 α and p110 β subunits. Mice lacking p110 α or p110 β die early during embryonic development.^{62,63} However, mice carrying a *knockin* mutation causing a 50% loss-of-function of p110 α activity (kinase-dead/D993A) are viable, although they display suppressed IRS signalling, decreased responsiveness to insulin and leptin, reduced linear growth, hyperphagia and increased adiposity.⁴⁷ Deletion of p110 α in cells expressing POMC or steroidogenesis factor 1 (SF1) decreased energy expenditure and increased susceptibility to high-fat diet.^{64,65}

These initial findings suggested that the p110 α was the main isoform downstream of leptin or insulin action. However, subsequent studies showed that the isoform interactions are more complex than previously anticipated. Insulin-induced phosphorylation of Akt (pAkt) is only partially blocked by inhibitors of either p110 α or p110 β , whereas the administration of both inhibitors completely prevented insulin signalling. Likewise, the combined administration of p110 α and p110 β inhibitors blocked the acute anorexigenic action of leptin and insulin.⁶⁶ Intact p110 β or both p110 α and p110 β subunits in neurones expressing agouti-related protein (AgRP) appear to be required for metabolic control.^{67,68} In SF1 neurones, distinct catalytic subunits are necessary for leptin or insulin cellular effects.⁶⁹ Thus, it soon became clear that blockade of PI3K downstream of leptin would only be attained by deletion of both subunits. This concept is further strengthened by data showing that LR neurones coexpress both p110 α and p110 β subunits.⁷⁰

3 | EFFECTS OF DELETION OF P110 α AND P110 β SUBUNITS IN LR NEURONES

To assess the direct role of PI3K downstream of leptin, we generated mice with deletion of both p110 α and p110 β catalytic subunits in LR cells.^{50,70,71} The experimental group was comprised of mice homozygous for three alleles: LR-Cre, p110 α -floxed and p110 β -floxed,^{48,72,73} named LR $\Delta\alpha+\beta$. We used LR-Cre homozygous mice because, in preliminary studies, we found only partial deletion of PI3K in LR cells of heterozygous LR^{Cre/+} mice harvested by fluorescence activated cell sorting.⁵⁰ This finding is also in agreement with previous studies using a different LR-Cre mouse line.⁷⁴⁻⁷⁶ To assess whether Cre in homozygosity alters LR function, we performed a systematic evaluation of LR^{Cre/Cre} mouse phenotype. No ectopic expression of Cre (using reporter genes) and no metabolic, growth or reproductive deficits were observed.⁵⁰

Male and female mice were evaluated separately due to the sexually dimorphic responses in metabolic and neuroendocrine functions.⁷⁷ Precautions were taken to avoid inconsistencies and exogenous interference in the physiological measurements. For example, on postnatal day 1 (P1), litter size was kept consistent (equal in number) among groups and cohorts to avoid early-life nutritional

effects on body weight and metabolic programming.^{1,4,78} After weaning (P21), males and females were fed phyto-oestrogen-reduced diet to prevent the effects of exogenous oestrogens on pubertal development and fertility. Body weights were measured weekly, and the metabolic phenotype was monitored by CLAMS (Comprehensive Laboratory Animal Monitoring System; Columbus Instruments, Columbus, OH, USA) using average values obtained from 3–4 days of metabolic recordings to avoid individual variations or random errors.

Briefly, male and female LR^{Δα+β} mice showed a decreased body weight, starting at 4–5 weeks of age, associated with decreased lean and fat mass and decreased linear growth. They also showed an increased food intake (when normalised by body weight), energy expenditure, locomotor activity and brown adipose tissue *Ucp1* expression, suggesting increased thermogenesis. Females (but not males) had delayed puberty determined by late vaginal opening (a sign of puberty onset) and delayed first oestrus (puberty completion).⁵⁰ Whether disruption of the actions of oestradiol as a result of PI3K imbalance in LR cells is the cause of the sexually dimorphic phenotype in reproductive function requires further evaluation.

These findings raised several questions that require additional studies. We initially focused on answering the following: (i) are individual PI3K catalytic subunits associated with specific physiological control and (ii) is the delayed puberty in females caused by the decrease in body weight, fat mass or leptin levels?

4 | EFFECTS OF DELETION OF P110α SUBUNITS IN LR NEURONES

To address the initial question, we generated a mouse line with deletion of a single PI3K subunit, p110α, in LR cells. We chose p110α because of data from D993A mice showing that a half dosage of p110α induces changes in metabolic and growth phenotypes.⁴⁷ To avoid potential confounders originating from the use of different genetic backgrounds, the mouse colony with deletion of p110α in LR cells (LR^{Δα}) was derived from the LR^{Δα+β} mice.

The LR^{Δα} male and female mice showed decreased body weight, lean and fat mass, and reduced linear growth. Changes in growth were observed only in adults (around and after P60); no differences were observed before or during pubertal maturation (P40). Low bone mineral density was detected in the femoral trabecular and cortical layers associated with a decrease in hepatic growth hormone receptor (*Ghr*) and insulin-like growth factor 1 (*Igf1*) expression in adult females. Similar to LR^{Δα+β} mice, food intake was increased when normalised by body weight. Increased energy expenditure and delayed pubertal maturation were observed. Female LR^{Δα} mice also showed prolonged oestrous cycles and progressive subfertility. No changes in locomotor activity and brown adipose tissue uncoupling protein 1 (*Ucp1*) gene expression were detected. Taken together, these findings indicate that, apart from energy balance, the p110α subunit in LR cells is also necessary for typical growth and reproduction.

Because the LR^{Δα} (and the LR^{Δα+β}) mice had a low body weight, fat mass and leptin levels, we performed several metabolic manipulations to assess whether the delayed puberty and reproductive phenotypes were secondary to the metabolic disruption. We started by applying the early postnatal overnutrition approach via manipulation of the litter size.^{1,4,78} A cohort of mice was maintained in small litters in the attempt to increase or normalise the body weight of LR^{Δα} mice prior to puberty onset. This manipulation was successful until weaning day, when LR^{Δα} mice start to consistently decrease the weight gain. Timing of pubertal development and leptin levels were marginally improved, although not corrected. Again, because we still found a lean phenotype in the postnatal overnutrition paradigm after weaning, our question remained unanswered. We decided to use the leptin challenge instead.

Leptin treatment in small doses that do not alter metabolic responses was shown to advance puberty in rodents.^{79,80} We used a very similar approach and found that, although control mice had early puberty onset, LR^{Δα} female mice were unresponsive to the effects of leptin on the timing of pubertal development. Another interesting finding from this experiment was the increased leptin sensitivity in metabolic regulation of the mutant mice potentially as a result of the increased basal levels of pAkt and leptin-induced phosphorylated STAT-3 and the decreased levels of the PIP3 phosphatase PTEN.⁵⁰

Of further interest was the increased expression of AgRP mRNA and peptide in fed female LR^{Δα} mice. The AgRP neurones are located in the arcuate nucleus and coexpress LR.^{45,81–83} Ablation of AgRP neurones (AgRP^{DTR} mice) ameliorates the metabolic and reproductive phenotypes of leptin-deficient *ob/ob* mice. Improved fertility was also observed in LR-deficient *db/db* mice with global deletion of *AgRP* gene, and deletion of LR in AgRP neurones alters fertility.^{84–88} Thus, changes in AgRP levels and/or signalling in LR^{Δα} mice may explain the increased food intake, and the disruption of growth and pubertal development observed in LR^{Δα} mice.^{87,89} It is also in agreement with findings showing that PI3K is required for leptin actions on *AgRP* gene expression.⁴⁵ Alternatively, ablation of PI3K may have affected the acute actions of leptin in the ventral premammillary nucleus, a hypothalamic site associated with leptin action in female reproductive function.^{70,74,90,91}

5 | PROS AND CONS OF USING THE LR-CRE LINE AS A METABOLICALLY RELEVANT TARGET

LR expression is observed in many peripheral tissues and in the brain, with a high density in hypothalamic sites.^{10,92,93} Thus, whether the effects observed using the LR-Cre mice were a result of the deletion of PI3K in neurones or in other peripheral organs may appear somewhat unclear. In this regard, it is important to emphasise that studies using conditional deletion or re-expression of LR have shown that the effects of leptin in the neuroendocrine axes are mediated by the brain.^{94–96} The deletion of LR from gonadotrophs caused no changes in body weight, timing of pubertal maturation or oestrous

cycle duration.⁹⁷ We have also shown that LR expression only in gonadotrophs is not sufficient to improve the metabolic or reproductive phenotypes of the LR null mice.⁹⁸ Similarly, initial studies have suggested that the reproductive deficits caused by a lack of leptin signalling are not mediated by the gonads,⁹⁹ indicating that the brain is the main target of leptin in neuroendocrine regulation. However, the LR^{Δα} mouse phenotype may not be entirely related to the disruption of leptin signalling. We could hypothesise that the lack of PI3K downstream of other hormones and/or growth factors in ovaries, for example, has contributed to the LR^{Δα} mouse phenotype. LR-Cre reporter gene is expressed in theca cells⁹⁸ and studies have suggested that insulin signalling in theca cells is associated with an obesity-induced increase in oestrous cycle length.¹⁰⁰ Deletion of insulin receptor in theca cells or gonadotrophs blocked this response, allowing females to maintain normal oestrous cycles under obese conditions.^{100,101} Taken together, these findings suggest that the reproductive phenotype of the LR^{Δα} mice is not associated with the deletion of PI3K in gonadotrophs or theca cells. As discussed earlier in this review, blunted PI3K signalling in LR cells of the ventral premammillary nucleus or the arcuate nucleus (ie, those coexpressing AgRP) are prime candidates.

Leptin receptor is also expressed in the adipocytes, liver and osteoblasts.^{10,102,103} Whether deletion of PI3K subunits in LR cells of peripheral tissues may have impacted the growth and reproductive phenotypes observed in our studies needs further evaluation. However, it should be noted that deletion of p110α only in adipocytes resulted in a very distinct phenotype compared to that observed in our studies, such that delayed puberty and infertility were observed only in male (not female) mice.¹⁰⁴ The mechanism(s) associated with this phenotype remain unknown. Moreover, no reproductive or linear growth deficits have been described in mice with deletion of LR or disruption of PI3K class I in the liver.^{105,106}

6 | COMPLEXITY OF PI3K SIGNALLING AND POTENTIAL CONFOUNDERS

The PI3K signalling pathway is comprised of intricate interactions of independent subunits and molecular targets. Thus, genetic modifications of specific subunits may cause an imbalance of the entire complex, generating unexpected phenotypes. For example, mice with deletion of p85α or p85β regulatory subunits have improved insulin sensitivity and hypoglycaemia, despite PI3K being a key pathway for the effects of insulin in glucose homeostasis.^{58,107,108} The selective deletion of the p110 catalytic subunits in LR cells may have caused a similar response, such as an improvement of leptin sensitivity in energy homeostasis because of increases in energy expenditure and basal pAkt, a decrease in body weight and the potentiation of leptin-induced phosphorylated STAT3. Notably, although the effects of leptin in metabolism were amplified in LR^{Δα} mice, linear growth and reproductive function were compromised. These findings suggest that PI3K is a crucial downstream signal of metabolic cues to growth and the reproductive neuroendocrine axes.

Another potential confounder is the expression of LR in neurones with distinct or opposite functions, such as the AgRP and POMC neurones in the arcuate nucleus. Previous studies have assessed the role of PI3K subunits in AgRP, POMC and other neuronal populations (ie, SF1).^{64,65,67,68} Because only subsets of these neurones express LR, the results are ambiguous. It is not possible to determine whether the observed effects are associated with a lack of leptin, insulin or growth factors.^{69,109} Using the LR-Cre mouse model, we expect the data to be more specific, generating insights into the direct effects of leptin-induced PI3K in physiology. Further studies will be necessary to dissociate the role of specific LR neurones with respect to each of the phenotypic changes observed.

LR cells likely coexpress a number of receptors that also recruit PI3K signalling (eg, receptor tyrosine kinases or G protein-coupled receptors). Therefore, it is possible that the deletion of PI3K subunits has altered the signalling of other hormones and/or growth factors, producing a phenotype unrelated to the action of leptin. As already noted, one key hormone is insulin. By deleting PI3K from LR cells, we may have blocked the actions of insulin in subsets of LR neurones. To test this hypothesis, we used the Cre-loxP system to delete InsR from LR cells.

7 | EFFECTS OF DELETION OF INSR IN LR CELLS

Previous studies using electrophysiological recordings have suggested that leptin or insulin target distinct POMC and SF1 neuronal populations.^{69,109} However, a lack of changes in membrane potential does not preclude other cellular responses, such as gene expression and/or post-translational modifications. To assess whether downstream targets of insulin-induced PI3K with genomic actions may be detected in LR neurones, we performed a colocalisation study in the LR reporter mouse. Mice were treated with i.c.v. insulin to avoid potential confounders with respect to peripheral insulin administration, and FoxO1 translocation or pAkt were identified in LR-Cre reporter neurones.⁵⁰ Partial colocalisation was detected, reinforcing the hypothesis that the effects of PI3K deletion in LR cells were a result of a partial blockade of insulin signalling. To test this, we generated mice with deletion of InsR in LR cells, using the same LR-Cre mouse line bred with previously validated InsR-floxed mice.¹¹⁰ No deficits in growth and only minor changes in reproductive physiology were observed in the mutant mice. Females showed a mild delay in puberty completion, suggesting that a lack of insulin signalling in LR cells (LR^{ΔInsR}) may add to the effect observed in LR^{Δα} and LR^{Δα+β} mice showing a more severe disruption of pubertal timing. Interestingly, female LR^{ΔInsR} mice showed almost no metabolic dysfunction, whereas males had an increased fat mass and glucose oxidation but no reproductive deficits.^{50,111} We concluded that the changes in metabolic, growth and reproductive phenotypes of the LR^{Δα} and LR^{Δα+β} mice are not a result of the blockade of insulin signalling. Because the conditional deletion of InsR in the brain causes metabolic and reproductive deficits,¹¹⁰ our findings also indicate

that the actions of insulin in these physiological systems are attained by targeting cells distinct from those expressing LR. Whether other PI3K recruiting factors account for the effects observed in LR^{Δα} and LR^{Δαβ} mice requires further evaluation.

Genomic screening in humans has identified the PI3K as a key signalling pathway associated with pubertal development.¹¹² In agreement, our findings obtained in mice show that PI3K signalling selectively in LR cells plays a major role. The decreased growth of LR^{Δα} mice and the higher sensitivity to leptin in metabolic regulation make PI3K a potential target under conditions of delayed growth and puberty in humans, as well as in hypothalamic amenorrhoea associated with sustained negative energy balance.

ACKNOWLEDGEMENTS

We thank all the current and former members of the Elias Lab who have contributed to the findings discussed in this review. Particularly, we thank the former postdoctoral fellows: José Donato Jr (University of São Paulo, Brazil) and Yun-Hee Choi (Anti-Aging Research Institute of BIO-FD&C Co. Ltd, South Korea) for the generation of the mouse colony and the production of preliminary data. We also thank Dr Jennifer Hill (University of Toledo) for insightful discussions.

CONFLICT OF INTERESTS

The authors declare that they have no conflicts of interest.

ORCID

David Garcia-Galiano  <https://orcid.org/0000-0001-9506-8785>

Susan J. Allen  <https://orcid.org/0000-0002-8271-0115>

Carol F. Elias  <https://orcid.org/0000-0001-9878-9203>

REFERENCES

- Kennedy GC, Mitra J. Body weight and food intake as initiating factors for puberty in the rat. *J Physiol* 1963;166:408-418.
- Frisch RE. The right weight: body fat, menarche and fertility. *Proc Nutr Soc*. 1994;53:113-129.
- Terasawa E, Kurian JR, Keen KL, Shiel NA, Colman RJ, Capuano SV. Body weight impact on puberty: effects of high-calorie diet on puberty onset in female rhesus monkeys. *Endocrinology*. 2012;153:1696-1705.
- Castellano JM, Bentsen AH, Sánchez-Garrido MA, et al. Early metabolic programming of puberty onset: impact of changes in postnatal feeding and rearing conditions on the timing of puberty and development of the hypothalamic kisspeptin system. *Endocrinology*. 2011;152:3396-3408.
- Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, Friedman JM. Positional cloning of the mouse obese gene and its human homologue [published erratum appears in *Nature* 1995 Mar 30;374(6521):479]. *Nature*. 1994;372:425-432.
- Maffei M, Halaas J, Ravussin E, et al. Leptin levels in human and rodent: measurement of plasma leptin and ob RNA in obese and weight-reduced subjects. *Nat Med*. 1995;1:1155-1161.
- Considine RV, Sinha MK, Heiman ML, et al. Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *N Engl J Med*. 1996;334:292-295.
- Coleman DL. Obese and diabetes: two mutant genes causing diabetes-obesity syndromes in mice. *Diabetologia*. 1978;14:141-148.
- Batt RAL, Everard DM, Gillies G, Wilkinson M, Wilson CA, Yeo TA. Investigation into the hypogonadism of the obese mouse (genotype ob/ob). *J Reprod Fertil*. 1982;64:363-371.
- Tartaglia LA, Dembski M, Weng X, et al. Identification and expression cloning of a leptin receptor, OB-R. *Cell*. 1995;83:1263-1271.
- Montague CT, Farooqi IS, Whitehead JP, et al. Congenital leptin deficiency is associated with severe early-onset obesity in humans. *Nature*. 1997;387:903-908.
- Gill MS, Hall CM, Tillmann V, Clayton PE. Constitutional delay in growth and puberty (CDGP) is associated with hypoleptinaemia. *Clin Endocrinol*. 1999;50:721-726.
- Barash IA, Cheung CC, Weigle DS, et al. Leptin is a metabolic signal to the reproductive system. *Endocrinology*. 1996;137:3144-3147.
- Chehab FF, Lim ME, Lu R. Correction of the sterility defect in homozygous obese female mice by treatment with the human recombinant leptin. *Nat Genet*. 1996;12:318-320.
- Cornish J, Callon KE, Bava U, et al. Leptin directly regulates bone cell function in vitro and reduces bone fragility in vivo. *J Endocrinol*. 2002;175:405-415.
- Reseland JE, Gordeladze JO. Role of leptin in bone growth: central player or peripheral supporter? *FEBS Lett*. 2002;528:40-42.
- Thomas T, Burguera B. Is leptin the link between fat and bone mass? *J Bone Miner Res*. 2002;17:1563-1569.
- Ozata M, Ozdemir IC, Licinio J. Human leptin deficiency caused by a missense mutation: multiple endocrine defects, decreased sympathetic tone, and immune system dysfunction indicate new targets for leptin action, greater central than peripheral resistance to the effects of leptin, and spontaneous correction of leptin-mediated defects. *J Clin Endocrinol Metab*. 1999;84:3686-3695.
- Farooqi IS, Wangenstein T, Collins S, et al. Clinical and molecular genetic spectrum of congenital deficiency of the leptin receptor. *N Engl J Med*. 2007;356:237-247.
- Myers MG, Cowley MA, Munzberg H. Mechanisms of leptin action and leptin resistance. *Annu Rev Physiol* 2008;70:537-556.
- Moon HS, Dalamaga M, Kim SY, et al. Leptin's role in lipodystrophic and nonlipodystrophic insulin-resistant and diabetic individuals. *Endocr Rev*. 2013;34:377-412.
- Chan JL, Mantzoros CS. Leptin and the hypothalamic-pituitary regulation of the gonadotropin-gonadal axis. *Pituitary*. 2001;4:87-92.
- Blüher S, Mantzoros CS. Leptin in reproduction. *Curr Opin Endocrinol Diabetes Obes* 2007;14:458-464.
- Rittmaster RS, Deshwal N, Lehman L. The role of adrenal hyperandrogenism, insulin resistance, and obesity in the pathogenesis of polycystic ovarian syndrome. *J Clin Endocrinol Metab*. 1993;76:1295-1300.
- Kasturi SS, Tannir J, Brannigan RE. The metabolic syndrome and male infertility. *J Androl*. 2008; 29:251-259.
- Loret de Mola JR. Obesity and its relationship to infertility in men and women. *Obstet Gynecol Clin North Am*. 2009;36:333-346.
- Borges BC, Garcia-Galiano D, da Silveira Cruz-Machado S, et al. Obesity-induced infertility in male mice is associated with disruption of crisp4 expression and sperm fertilization capacity. *Endocrinology*. 2017;158:2930-2943.
- Herman-Giddens ME, Slora EJ, Wasserman RC, et al. Secondary sexual characteristics and menses in young girls seen in office practice: a study from the pediatric research in office settings network. *Pediatrics*. 1997;99:505-512.
- Biro FM, Khoury P, Morrison JA. Influence of obesity on timing of puberty. *Int J Androl*. 2006;29:272-277.

30. Biro FM, Galvez MP, Greenspan LC, et al. Pubertal assessment method and baseline characteristics in a mixed longitudinal study of girls. *Pediatrics*. 2010;126:e583-e590.
31. Ihle JN. Cytokine receptor signalling. *Nature*. 1995;377:591-594.
32. Chua SC Jr, Chung WK, Wu-Peng XS, et al. Phenotypes of mouse diabetes and rat fatty due to mutations in the OB (leptin) receptor. *Science*. 1996;271:994-996.
33. Lee GH, Proenca R, Montez JM, et al. Abnormal splicing of the leptin receptor in diabetic mice. *Nature*. 1996;379:632-635.
34. Vaisse C, Halaas JL, Horvath CM, Darnell JE Jr, Stoffel M, Friedman JM. Leptin activation of Stat3 in the hypothalamus of wild-type and ob/ob mice but not db/db mice. *Nat Genet*. 1996;14:95-97.
35. Bjorbaek C, Elmquist JK, Frantz JD, Shoelson SE, Flier JS. Identification of SOCS-3 as a potential mediator of central leptin resistance. *Mol Cell*. 1998;1:619-625.
36. Bates SH, Stearns WH, Dundon TA, et al. STAT3 signalling is required for leptin regulation of energy balance but not reproduction. *Nature*. 2003;421:856-859.
37. Patterson CM, Villanueva EC, Greenwald-Yarnell M, et al. Leptin action via LepR-b Tyr(1077) contributes to the control of energy balance and female reproduction. *Molecular Metabolism*. 2012;1:61-69.
38. Bjornholm M, Munzberg H, Leshan RL, et al. Mice lacking inhibitory leptin receptor signals are lean with normal endocrine function. *J Clin Invest*. 2007;117:1354-1360.
39. Singireddy AV, Inglis MA, Zuure WA, Kim JS, Anderson GM. Neither signal transducer and activator of transcription 3 (STAT3) or STAT5 signaling pathways are required for leptin's effects on fertility in mice. *Endocrinology*. 2013;154:2434-2445.
40. Zhao AZ, Huan JN, Gupta S, Pal R, Sahu A. A phosphatidylinositol 3-kinase phosphodiesterase 3B-cyclic AMP pathway in hypothalamic action of leptin on feeding. *Nat Neurosci* 2002;5:727-728.
41. Niswender KD, Baskin DG, Schwartz MW. Insulin and its evolving partnership with leptin in the hypothalamic control of energy homeostasis. *Trends Endocrinol Metab*. 2004;15:362-369.
42. Xu A, Kaelin CB, Takeda K, Akira S, Schwartz MW, Barsh GS. PI3K integrates the action of insulin and leptin on hypothalamic neurons. *J Clin Invest*. 2005;115:951-958.
43. Sahu A. Intracellular leptin-signaling pathways in hypothalamic neurons: the emerging role of phosphatidylinositol-3 kinase-phosphodiesterase-3B-cAMP pathway. *Neuroendocrinology*. 2011;93:201-210.
44. Niswender KD, Morton GJ, Stearns WH, Rhodes CJ, Myers MG Jr, Schwartz MW. Intracellular signalling. Key enzyme in leptin-induced anorexia. *Nature*. 2001;413:794-795.
45. Morrison CD, Morton GJ, Niswender KD, Gelling RW, Schwartz MW. Leptin inhibits hypothalamic Npy and AgRP gene expression via a mechanism that requires phosphatidylinositol 3-OH-kinase signaling. *Am J Physiol Endocrinol Metab*. 2005;289:E1051-E1057.
46. Morton GJ, Gelling RW, Niswender KD, Morrison CD, Rhodes CJ, Schwartz MW. Leptin regulates insulin sensitivity via phosphatidylinositol-3-OH kinase signaling in mediobasal hypothalamic neurons. *Cell Metab*. 2005;2:411-420.
47. Foukas LC, Claret M, Pearce W, et al. Critical role for the p110[alpha] phosphoinositide-3-OH kinase in growth and metabolic regulation. *Nature*. 2006;441:366-370.
48. Jia S, Liu Z, Zhang S, et al. Essential roles of PI(3)K-p110beta in cell growth, metabolism and tumorigenesis. *Nature*. 2008;454:776-779.
49. Pal A, Barber TM, Van de Bunt M, et al. PTEN mutations as a cause of constitutive insulin sensitivity and obesity. *N Engl J Med*. 2012;367:1002-1011.
50. Garcia-Galiano D, Borges BC, Donato J Jr., et al. PI3Kalpha inactivation in leptin receptor cells increases leptin sensitivity but disrupts growth and reproduction. *JCI Insight* 2017;2:Pii: 96728.
51. Plum L, Belgardt BF, Bruning JC. Central insulin action in energy and glucose homeostasis. *J Clin Invest*. 2006;116:1761-1766.
52. Acosta-Martinez M. PI3K: an attractive candidate for the central integration of metabolism and reproduction. *Front Endocrinol*. 2011;2:110.
53. Spanswick D, Smith MA, Groppi VE, Logan SD, Ashford ML. Leptin inhibits hypothalamic neurons by activation of ATP-sensitive potassium channels. *Nature*. 1997;390:521-525.
54. Spanswick D, Smith MA, Mirshamsi S, Routh VH, Ashford ML. Insulin activates ATP-sensitive K⁺ channels in hypothalamic neurons of lean, but not obese rats. *Nat Neurosci*. 2000;3:757-758.
55. Hill JW, Williams KW, Ye C, et al. Acute effects of leptin require PI3K signaling in hypothalamic proopiomelanocortin neurons in mice. *J Clin Invest* 2008;118:1796-1805.
56. Buettner C, Muse ED, Cheng A, et al. Leptin controls adipose tissue lipogenesis via central, STAT3-independent mechanisms. *Nat Med*. 2008;14:667-675.
57. Cantley LC. The phosphoinositide 3-kinase pathway. *Science* 2002;296:1655-1657.
58. Ueki K, Yballe C, Brachmann S, et al. Increased insulin sensitivity in mice lacking p85beta subunit of phosphoinositide 3-kinase. *Proc Natl Acad Sci U S A*. 2002;99:419-424.
59. Vanhaesebroeck B, Ali K, Bilancio A, Geering B, Foukas LC. Signalling by PI3K isoforms: insights from gene-targeted mice. *Trends Biochem Sci*. 2005;30:194-204.
60. Chantry D, Vojtek A, Kashishian A, et al. p110delta, a novel phosphatidylinositol 3-kinase catalytic subunit that associates with p85 and is expressed predominantly in leukocytes. *J Biol Chem*. 1997;272:19236-19241.
61. Horsch D, Kahn CR. Region-specific mRNA expression of phosphatidylinositol 3-kinase regulatory isoforms in the central nervous system of C57BL/6J mice. *J Comp Neurol*. 1999;415:105-120.
62. Bi L, Okabe I, Bernard DJ, Wynshaw-Boris A, Nussbaum RL. Proliferative defect and embryonic lethality in mice homozygous for a deletion in the p110alpha subunit of phosphoinositide 3-kinase. *J Biol Chem*. 1999;274:10963-10968.
63. Bi L, Okabe I, Bernard DJ, Nussbaum RL. Early embryonic lethality in mice deficient in the p110β catalytic subunit of PI 3-kinase. *Mamm Genome*. 2002;13:169-172.
64. Xu Y, Hill JW, Fukuda M, et al. PI3K signaling in the ventromedial hypothalamic nucleus is required for normal energy homeostasis. *Cell Metab*. 2010;12:88-95.
65. Hill JW, Xu Y, Preitner F, et al. Phosphatidylinositol 3-kinase signaling in hypothalamic proopiomelanocortin neurons contributes to the regulation of glucose homeostasis. *Endocrinology*. 2009;150:4874-4882.
66. Tups A, Anderson GM, Rizwan M, et al. Both p110alpha and p110beta isoforms of phosphatidylinositol 3-OH-kinase are required for insulin signalling in the hypothalamus. *J Neuroendocrinol*. 2010;22:534-542.
67. Al-Qassab H, Smith MA, Irvine EE, et al. Dominant role of the p110β isoform of PI3K over p110α in energy homeostasis regulation by POMC and AgRP neurons. *Cell Metab*. 2009;10:343-354.
68. Huang Y, He Z, Gao Y, et al. Phosphoinositide 3-kinase is integral for the acute activity of leptin and insulin in male arcuate NPY/AgRP neurons. *Journal of the Endocrine Society*. 2018;2(6):518-532.
69. Sohn J-W, Oh Y, Kim KW, Lee S, Williams KW, Elmquist JK. Leptin and insulin engage specific PI3K subunits in hypothalamic SF1 neurons. *Molecular Metabolism*. 2016;5:669-679.
70. Williams KW, Sohn JW, Donato J Jr, et al. The acute effects of leptin require PI3K signaling in the hypothalamic ventral premammillary nucleus. *J Neurosci*. 2011;31:13147-13156.
71. Borges BC, Garcia-Galiano D, Rorato R, Elias LL, Elias CF. PI3K p110beta subunit in leptin receptor expressing cells is required

- for the acute hypophagia induced by endotoxemia. *Mol Metab.* 2016;5:379-391.
72. Zhao J, Cheng H, Jia S, et al. The p110alpha isoform of PI3K is essential for proper growth factor signaling and oncogenic transformation. *Proc Natl Acad Sci U S A.* 2006;103:16296-16300.
 73. Scott MM, Lachey JL, Sternson SM, et al. Leptin targets in the mouse brain. *J Comp Neurol.* 2009;514:518-532.
 74. Leshan RL, Louis GW, Jo Y-H, Rhodes CJ, Munzberg H, Myers MG Jr. Direct innervation of GnRH neurons by metabolic- and sexual odorant-sensing leptin receptor neurons in the hypothalamic ventral premammillary nucleus. *J Neurosci.* 2009;29:3138-3147.
 75. Louis GW, Greenwald-Yarnell M, Phillips RJ, Coolen LM, Lehman MN, Myers MG. Molecular mapping of the neural pathways linking leptin to the neuroendocrine reproductive axis. *Endocrinology.* 2011;152:2302-2310.
 76. Allison MB, Patterson CM, Krashes MJ, Lowell BB, Myers MG Jr, Olson DP. TRAP-seq defines markers for novel populations of hypothalamic and brainstem LepRb neurons. *Mol Metab.* 2015;4:299-309.
 77. Clegg DJ, Brown LM, Woods SC, Benoit SC. Gonadal hormones determine sensitivity to central leptin and insulin. *Diabetes.* 2006;55:978-987.
 78. Caron E, Ciofi P, Prevot V, Bouret SG. Alteration in neonatal nutrition causes perturbations in hypothalamic neural circuits controlling reproductive function. *J Neurosci.* 2012;32:11486-11494.
 79. Ahima RS, Dushay J, Flier SN, Prabakaran D, Flier JS. Leptin accelerates the onset of puberty in normal female mice. *J Clin Invest.* 1997;99:391-395.
 80. Cheung CC, Thornton JE, Kuijper JL, Weigle DS, Clifton DK, Steiner RA. Leptin is a metabolic gate for the onset of puberty in the female rat. *Endocrinology.* 1997;138:855-858.
 81. Elias CF, Aschkenasi C, Lee C, et al. Leptin differentially regulates NPY and POMC neurons projecting to the lateral hypothalamic area. *Neuron.* 1999;23:775-786.
 82. Mizuno TM, Mobbs CV. Hypothalamic agouti-related protein messenger ribonucleic acid is inhibited by leptin and stimulated by fasting. *Endocrinology.* 1999;140:814-817.
 83. Cone RD, Cowley MA, Butler AA, Fan W, Marks DL, Low MJ. The arcuate nucleus as a conduit for diverse signals relevant to energy homeostasis. *Int J Obes Relat Metab Disord* 2001;25(Suppl 5):S63-S67.
 84. Wu Q, Whiddon BB, Palmiter RD. Ablation of neurons expressing agouti-related protein, but not melanin concentrating hormone, in leptin-deficient mice restores metabolic functions and fertility. *Proc Natl Acad Sci U S A.* 2012;109:3155-3160.
 85. Israel DD, Sheffer-Babila S, de Luca C, et al. Effects of leptin and melanocortin signaling interactions on pubertal development and reproduction. *Endocrinology.* 2012;153:2408-2419.
 86. Sheffer-Babila S, Sun Y, Israel DD, Liu S-M, Neal-Perry G, Chua SC. Agouti-related peptide plays a critical role in leptin's effects on female puberty and reproduction. *American Journal of Physiology - Endocrinology and Metabolism.* 2013;305:E1512-E1520.
 87. Padilla SL, Qiu J, Nestor CC, et al. AgRP to Kiss1 neuron signaling links nutritional state and fertility. *Proc Natl Acad Sci U S A.* 2017;114:2413-2418.
 88. Egan OK, Inglis MA, Anderson GM. Leptin signaling in AgRP neurons modulates puberty onset and adult fertility in mice. *The Journal of Neuroscience.* 2017;37:3875-3886.
 89. Sasanuma H, Nakata M, Parmila K, Nakae J, Yada T. PDK1-FoxO1 pathway in AgRP neurons of arcuate nucleus promotes bone formation via GHRH-GH-IGF1 axis. *Mol Metab.* 2017;6:428-439.
 90. Donato J Jr, Cravo RM, Frazao R, et al. Leptin's effect on puberty in mice is relayed by the ventral premammillary nucleus and does not require signaling in Kiss1 neurons. *J Clin Invest.* 2011;121:355-368.
 91. Ross RA, Leon S, Madara JC, et al. PACAP neurons in the ventral premammillary nucleus regulate reproductive function in the female mouse. *ELife* 2018;7:e35960.
 92. Mercer JG, Hoggard N, Williams LM, Lawrence CB, Hannah LT, Trayhurn P. Localization of leptin receptor mRNA and the long form splice variant (Ob-Rb) in mouse hypothalamus and adjacent brain regions by in situ hybridization. *FEBS Lett.* 1996;387:113-116.
 93. Zamorano PL, Mahesh VB, De Sevilla LM, Chorich LP, Bhat GK, Brann DW. Expression and localization of the leptin receptor in endocrine and neuroendocrine tissues of the rat. *Neuroendocrinology.* 1997;65:223-228.
 94. Cohen P, Zhao C, Cai X, et al. Selective deletion of leptin receptor in neurons leads to obesity. *J Clin Invest.* 2001;108:1113-1121.
 95. Kowalski TJ, Liu SM, Leibel RL, Chua SC Jr. Transgenic complementation of leptin-receptor deficiency. I. Rescue of the obesity/diabetes phenotype of LEPR-null mice expressing a LEPR-B transgene. *Diabetes* 2001;50:425-435.
 96. Quenell JH, Mulligan AC, Tups A, et al. Leptin indirectly regulates gonadotropin-releasing hormone neuronal function. *Endocrinology.* 2009;150:2805-2812.
 97. Akhter N, CarlLee T, Syed MM, et al. Selective deletion of leptin receptors in gonadotropes reveals activin and GnRH-binding sites as leptin targets in support of fertility. *Endocrinology.* 2014;155:4027-4042.
 98. Allen SJ, Garcia-Galiano D, Borges BC, Burger LL, Boehm U, Elias CF. Leptin receptor null mice with re-expression of LepR in GnRH-R expressing cells display elevated FSH levels but remain in a prepubertal state. *Am J Physiol Regul Integr Comp Physiol* 2016;310:R1258-R1266.
 99. Zhang Y, Hu M, Ma H, et al. The impairment of reproduction in db/db mice is not mediated by intraovarian defective leptin signaling. *Fertil Steril.* 2012;97:1183-1191.
 100. Wu S, Divall S, Nwaopara A, et al. Obesity-induced infertility and hyperandrogenism are corrected by deletion of the insulin receptor in the ovarian theca cell. *Diabetes.* 2014;63:1270-1282.
 101. Brothers KJ, Wu S, DiVall SA, et al. Rescue of obesity-induced infertility in female mice due to a pituitary-specific knockout of the insulin receptor. *Cell Metab.* 2010;12:295-305.
 102. Kielar D, Clark JS, Ciechanowicz A, Kurzawski G, Sulikowski T, Naruszewicz M. Leptin receptor isoforms expressed in human adipose tissue. *Metabolism.* 1998;47:844-847.
 103. Gordeladze JO, Reseland JE. A unified model for the action of leptin on bone turnover. *J Cell Biochem.* 2003;88:706-712.
 104. Nelson VL, Negron AL, Reid I, et al. Loss of PI3K p110alpha in the adipose tissue results in infertility and delayed puberty onset in male mice. *Biomed Res Int.* 2017;2017:13.
 105. Mittenbühler MJ, Sprenger H-G, Gruber S, et al. Hepatic leptin receptor expression can partially compensate for IL-6Rα deficiency in DEN-induced hepatocellular carcinoma. *Molecular Metabolism* 2018;17:122-133.
 106. Kenerson HL, Yeh MM, Kazami M, et al. Akt and mTORC1 have different roles during liver tumorigenesis in mice. *Gastroenterology.* 2013;144:1055-1065.
 107. Fruman DA, Mauvais-Jarvis F, Pollard DA, et al. Hypoglycaemia, liver necrosis and perinatal death in mice lacking all isoforms of phosphoinositide 3-kinase p85 alpha. *Nat Genet.* 2000;26:379-382.
 108. Ueki K, Fruman DA, Yballe CM, et al. Positive and negative roles of p85 alpha and p85 beta regulatory subunits of phosphoinositide 3-kinase in insulin signaling. *J Biol Chem.* 2003;278:48453-48466.
 109. Williams KW, Margatho LO, Lee CE, et al. Segregation of acute leptin and insulin effects in distinct populations of arcuate proopiomelanocortin neurons. *J Neurosci.* 2010;30:2472-2479.
 110. Bruning JC, Gautam D, Burks DJ, et al. Role of brain insulin receptor in control of body weight and reproduction. *Science.* 2000;289:2122-2125.

111. Borges BC, Han X, Allen SJ, Garcia-Galiano D, Elias CF. Insulin signaling in LepR cells modulates fat and glucose homeostasis independent of leptin. *Am J Physiol Endocrinol Metab.* 2018;316:E121-E134.
112. Day FR, Thompson DJ, Helgason H, et al., LifeLines Cohort Study; InterAct Consortium; kConFab/AOCS Investigators; Endometrial Cancer Association Consortium; Ovarian Cancer Association Consortium; PRACTICAL consortium,, Spurdle AB, Thorsteindottir U, Pollard KS, Easton DF, Tung JY, Chang-Claude J, Hinds D, Murray A, Murabito JM, Stefansson K, Ong KK, Perry JRB. Genomic analyses identify hundreds of variants associated

with age at menarche and support a role for puberty timing in cancer risk. *Nat Genet.* 2017;49:834.

How to cite this article: Garcia-Galiano D, Borges BC, Allen SJ, Elias CF. PI3K signalling in leptin receptor cells: Role in growth and reproduction. *J Neuroendocrinol.* 2019;31:e12685. <https://doi.org/10.1111/jne.12685>