counterparts $F_{(1, 258)}=5.74$, $p=0.017$. **Conclusions:** Although the specific symptomatology of each of these 3 dementia types differ, the collective emotional toll on caregivers does not fundamentally differ by disease. The caregiver experience is equally difficult for all types of dementia. The only caveat to this finding is that we used measures developed primarily to assess the experience for AD caregivers that may not have adequately captured unique facets of the DLB or the PDD caregiving experience.

P2-257

COGNITIVE TRAINING PROGRAMS FOR VERY OLD LONE ADULTS IN A KOREAN RURAL COMMUNITY

Kyung-Phil Kwak¹, Sumgmin Lee², Taewoo Kim¹, Nari Bae¹, ¹Dongguk University Gyeongju Hospital, Gyeongju, South Korea; ²Dongguk University Gyeongju Hospital, Gyeongju, South Korea.
Contact e-mail: dependence@gmail.com

Background: The Gyeongbukdo cognitive training program(GBD-CTP) of 80 sessions was provided for community dwelling very older adults and the effects on cognitive function, depression and quality of life were tested. **Methods:** The participants were 1459 community-based sample of 'lone' older adults from 75 to 92 years of age who have mild cognitive disorder(CDR=0.5). This program was applied two times a week for 40 weeks. The sample was randomized in two groups: 876 participants underwent full length period GBD-CTPA while 583 participants received only 20 weeks CTP and stop CTP next 20 weeks. Research tool was applied in order to verify the effects of the program on the subjects' ability toward ADL, cognitive function, and depression and quality of life. **Results:** The intervention effect(mean change from baseline to final test) was significant($P<.05$) on MMSE-DS(1.32), GDS(0.3), QOL-AD(3.5), S-IADL(2.5) and Greitric Dep Scale(4.2). Five participants(stop progressed to dementia). **Conclusions:** These results suggest that this long term CTP has an effect on the improvement of the ability toward cognition and ADL of mild cognitive impairment living alone in a community.

P2-258

MILD COGNITIVE IMPAIRMENT INFLUENCES ASSOCIATIONS BETWEEN BALANCE CONFIDENCE AND FUNCTIONAL PERFORMANCE IN COMMUNITY-DWELLING OLDER ADULTS

Jennifer Blackwood, University of Michigan-Flint, Flint, MI, USA.
Contact e-mail: jblackwo@umflint.edu

Background: Older adults with mild cognitive impairment (MCI) are at a greater risk for falling than cognitively intact older adults. High balance confidence has been reported to be associated with higher levels of physical function and decreased risk of falling. The purpose of this study was to examine how confidence in balance is influenced by the presence of MCI in community dwelling older adults. **Methods:** Forty-seven community dwelling older adults were assigned to the MCI or non-MCI group based on a cut-off score of 26 on the Montreal Cognitive Assessment tool. Balance confidence was assessed using the Activities Specific Balance Scale-6 (ABC-6) while physical function was assessed with measures of gait speed and the Timed Up and Go tests. Descriptive statistics, correlations, and comparisons of mean scores between groups on the measures were completed. **Results:** Mean scores on the physical function measures did not differ between groups. Balance confidence was moderately associated with gait speed ($r = 0.531$, $p<.05$) and TUG performance ($r = -0.674$, $p<.05$)

in all subjects, but was no longer significantly associated with gait speed in those who screened positive for MCI ($r = 0.153$, $p> .05$). **Conclusions:** In those screened positive for MCI, the association between balance confidence (ABC-6) and walking speed indicates that a self-assessment of balance confidence may not provide valid information to determine fall risk. Clinicians seeking to quantify physical function as it relates to fall risk, should consider other measures of fall risk besides the ABC-6.

P2-259

BRAINS FOR DEMENTIA RESEARCH: USE OF TELEPHONE INTERVIEW TO GATHER ASSESSMENT DATA

Gillian Hayes¹, Helen Costello², Paul Francis¹, ¹King's College London, London, England; ²King's College London, London, United Kingdom.
Contact e-mail: Gillian.Hayes@kcl.ac.uk

Background: Brains for Dementia Research (BDR) aims to provide high quality post-mortem brain tissue supported by detailed clinical data regarding cognition, behaviour and activities of daily living. With a cohort now in excess of 3,000 we sought to assess the feasibility of conducting assessments with participants by telephone rather than face to face. Specifically our aim was to compare satisfaction with telephone, video and face-to-face interviews. **Methods:** The study took place in 3 phases. Phase 1 focused on the development and feasibility of a telephone assessment package. Phase 2 was an RCT comparing satisfaction with telephone and face-to-face assessment and phase 3 examined the feasibility of and satisfaction with video based assessment. Feasibility data were collected during assessments and participants completed a self-report satisfaction questionnaire. **Results:** The majority of participants were happy to complete a telephone assessment (72.8%) and following the interview 100% were happy to be contacted for their next scheduled assessment. A total of 7.6% participants (94.8% male) declined a telephone assessment for various reasons.

- Participants were mostly female (64.2%), with an average age of 73.4 years (SD 6.7, range 64-95).
- Initial telephone screens lasted an average of 11 minutes (Range 6-35 minutes) with over two thirds taking place on the first or second attempt.
- Background questionnaires took an average of 15 minutes (Range 5-70 minutes) to complete, with over 90% returned before the assessment.
- Average length of participant telephone assessment was 33 minutes (Range 21-75 minutes).
- Study partner telephone assessments lasted 15 minutes (Range 7-45 minutes).
- High levels of participant satisfaction observed in relation to attitudes towards & practical aspects of telephone assessment and self-completion of the background questionnaire.
- 97% agreed/strongly agreed that the overall experience was positive.
- Major benefit was 'convenience'.

Conclusions: Preliminary results indicate that telephone assessment for control participants is a feasible mechanism for overcoming the challenges faced by BDR in relation to a growing, ageing cohort and uncertain resources. Overall, participants were highly satisfied with telephone assessment, although further analysis is required to compare satisfaction across face-to-face and telephone assessment.