was present in 264 out of 676 participants. The rate of both being frail or prefrail and impairment on CDT was 20.4%. During the follow up, 31 developed IADLs disability. Risk for incident IADLs disability was greater in the group with impairment on CDT (OR: 2.68, 95% CI 1.21-5.93), after adjustment for age, gender, education and number of chronic diseases. Compared with non-frail persons, those who were frail and pre-frail showed an increased risk of incident IADLs disability (adjusted OR: 2.56, 95% CI: 1.07-6.14). Frail and pre-frail persons with impairment on CDT were significantly more likely to develop IADLs disability (adjusted OR: 3.49, 95% CI: 1.54-7.89). **Conclusions:** Combination of CDT and frailty phenotype improves their individual predictive ability for disability in instrumental activities of daily living in this population.

P3-262

COMORBID SUBJECTIVE COGNITIVE DECLINE AND SLEEP APNEA SIGNIFICANTLY INCREASE THE RISK OF INCIDENT DEMENTIA: RESULTS FROM THE PREVENTION OF ALZHEIMER'S DISEASE WITH VITAMIN E AND SELENIUM STUDY

Erin L. Abner, Xiuhua Ding, Allison M. Caban-Holt, Frederick A. Schmitt, Richard J. Kryscio, *University of Kentucky, Lexington, KY, USA.*
Contact e-mail: Erin.abner@uky.edu

Background: Subjective memory complaints (SMC) reflect patient-identified deficits in memory and have been linked to increased risk of future dementia in nondemented (including cognitively intact) older adults, but the relationship between SMC and comorbid medical conditions is not well understood. **Methods:** Double-blind, placebo controlled 2x2 randomized controlled trial that transformed into an exposure cohort following discontinuation of supplementation in the SELECT parent trial. Setting: Prevention of Alzheimer's with Vitamin E and Selenium (PREADVISE) participants were assessed for incident dementia at 130 local clinical study sites in the United States, Canada, and Puerto Rico during the controlled trial phase and were later followed by telephone from a centralized location during the post exposure (observational) phase. PREADVISE enrolled a total of 7,547 nondemented men age 60 and older; 4,271 were able to participate in the observational study. Participants were interviewed at baseline for memory complaints ("Do you have more problems with your memory than most people?") and presence of comorbidities including sleep apnea, stroke, hypertension, diabetes, history of traumatic brain injury (TBI), coronary artery bypass graft (CABG), and congestive heart failure. **Results:** Men who reported memory problems (SMC) (n=123) at baseline were significantly more likely to also report history of sleep apnea and TBI and had significantly worse scores on the baseline memory screen (Table 1). Cox proportional hazards regression revealed that the effect of baseline SMC depended on the presence of sleep apnea (p for interaction=0.015). Men who reported sleep apnea and SMC at baseline had 14.9 times (95%CI: 6.0-37.0) the hazard of incident dementia compared to men with sleep apnea but no SMC. For men without sleep apnea, SMC more than tripled the hazard (HR=3.86). History of CABG and stroke also significantly increased the hazard of incident dementia (Table 2). **Conclusions:** Memory complaints and comorbidities in nondemented older men strongly predicted future dementia in the PREADVISE cohort. Special care should be paid to management of comorbidities in older adult men who report memory problems, as these may be modifiable risk factors for dementia onset.

Table 1

PREADVISE baseline participant characteristics (N=7,547)

Characteristic*	All Subjects (N=7,547)	No Memory Problem (n=7424)	Memory Problem (n=123)
Age, y	67.5±5.3	67.5±5.3	67.5±5.3
Education, y	15.0±2.7	15.0±2.7	14.6±2.9
Black race [†]	756 (10.0)	749 (10.1)	8 (6.5)
Baseline MIS [‡]	7.6±0.7	7.6±0.7	7.4±0.8
APOE-e4	2,029 (26.9)	1890 (25.5)	40 (32.5)
Hypertension	2,998 (39.7)	2951 (39.8)	48 (39.0)
Diabetes	858 (11.4)	842 (11.3)	16 (13.0)
Coronary Artery Bypass Graft	318 (4.2)	314 (4.2)	4(3.3)
Stroke	43 (0.6)	41 (0.6)	2 (1.6)
Congestive Heart Failure	62 (0.8)	61 (0.8)	1 (0.8)
Sleep apnea [§]	552 (7.3)	533 (7.2)	19 (15.5)
Traumatic brain injury [§]	997 (13.2)	971 (13.1)	26 (21.4)

*Results presented are mean±SD or frequencies with proportions. [†]Proportion of black participants is significantly higher in the No Memory Complaint group than either Memory Change or Problem (p<0.001). [‡]Mean baseline MIS scores were significantly lower in men who reported a Memory Problem than men who did not complain (p=0.004). [§]Sleep apnea and history of traumatic brain injury were more common among men who reported a Memory Problem than men who did not complain (p<0.01).

Table 2

Cox proportional hazards regression model results.

Comparison	Adjusted HR (95% CI)
Memory Problem vs. No Complaint (Sleep Apnea= Y)	14.90 (6.02, 37.04)
Memory Problem vs. No Complaint (Sleep Apnea= N)	3.86 (2.11, 7.09)
Baseline age, 1 year	1.11 (1.09, 1.13)
Black vs. not black race	1.85 (1.28, 2.68)
Education, 1 year	0.98 (0.94, 1.02)
APOE-e4 carrier vs. not	2.11 (1.68, 2.65)
CABG vs. none	1.61 (1.07, 2.44)
Stroke vs. none	3.45 (1.28, 9.28)
Head injury vs. none	1.49 (1.10, 2.01)

P3-263

FACTORS THAT INFLUENCE SCREENING OF COGNITION WITHIN PHYSICAL THERAPY PRACTICE IN THE UNITED STATES

Jennifer Blackwood, *University of Michigan-Flint, Flint, MI, USA.*
Contact e-mail: jblackwo@umflint.edu

Background: Having a cognitive impairment puts an older adult at an increased risk of falling. The prevalence of screening for cognitive deficits as a part of a fall risk assessment has been reported to be low in physical therapist practice. The purpose of this study was to examine the factors that influence the practice of screening cognition as a part of physical therapist practice in the United States. **Methods:** Five thousand physical therapists were sent an electronic survey regarding the use of cognitive and fall risk measures in physical therapist practice. Descriptive

statistics, correlations, Chi Square analyses, and logistic regression analyses were completed. **Results:** A total of 572 physical therapists with a mean age of 44.5 years and whose highest level of education was a doctoral degree (58.8%) completed the survey. The strongest predictor of whether a therapist was likely to screen for cognition was the time spent with older adults. Neither the therapists' education nor years in practice did not significantly influence their likelihood to screen cognition as a part of a fall risk assessment. **Conclusions:** Cognitive screening as a part of a fall risk assessment may not be routinely performed by physical therapists in the United States. In order to address this gap in practice, further education is needed to improve screening efforts as a part of practice.

P3-264

PERTH FAMILY STUDIES: IDENTIFICATION OF AN AT-RISK POPULATION FOR DEMENTIA PREVENTION STUDIES

Maggie Robertson¹, Lawrence Whalley¹, Peter Connelly², ¹University of the Highlands and Islands, Perth, United Kingdom; ²Tayside Health Board, Perth, United Kingdom. Contact e-mail: maggie.robertson@perth.uhi.ac.uk

Background: The biological mechanisms involved in late-onset dementia are identifiable in mid-life. The associated neurodegeneration is the result of a complex interplay between genetic and environmental factors. Identifying critical windows to target interventions offer the best hope for risk reduction and dementia prevention outcomes. The earliest clinical symptoms are more difficult to identify and often not apparent until substantial neurodegeneration has taken place. Research efforts will need to be more efficient and should consider focusing on populations most at risk of dementia for participation in longitudinal prevention studies rather than screening the general population. **Methods:** With ethics approval from NHS Tayside, we recruited families from Perth, Scotland into an observational study investigating their psycho-educational needs following a diagnosis of dementia. We asked people living with dementia to provide the medical history of their first-degree relatives. Family members corroborated the medical history information and we used death certificates to verify family histories. Data were collected over eighteen months from March 2013. We also collected supplementary information via a telephone interviews with adult offspring and siblings to ascertain perspectives on dementia risk prevention. **Results:** We have recruited 50 families. Interview data from 37 families stress the need for more information relating to the dementia continuum and estimations of dementia risk. We have also identified a cohort >70 unaffected at-risk adults aged between late 30's and mid 60's with a family history of late-onset dementia. **Conclusions:** We will discuss how the family history method has helped us to develop a strategy to identify at-risk offspring in families with a history of late-onset dementia. To our knowledge, we are the first group to have identified at-risk offspring and a verified family history in Scotland and possibly the UK. Most of these individuals (69) have said they want the opportunity to participate in longitudinal dementia prevention studies. We are seeking funding and collaboration to develop a prevention study using epigenetic biomarkers, neuroimaging and tests measuring cognitive function. Study design would stratify adult offspring into different risk levels so we can identify and respond with appropriate bespoke risk reduction interventions within the optimum therapeutic window.

P3-265

PSYCHOACTIVE MEDICATION AND MEMORY IMPAIRMENT AMONG MEMBERS OF AN ALZHEIMER PREVENTION REGISTRY

Lori A. Daiello^{1,2}, Sevdener Cizginer^{1,3}, Marissa A. Pelosi⁴, Brian R. Ott^{1,2}, ¹Alpert Medical School of Brown University, Providence, RI, USA; ²Alzheimers Disease and Memory Disorders Center at Rhode Island Hospital, Providence, RI, USA; ³Miriam Hospital, Providence, RI, USA; ⁴Rhode Island Hospital, Providence, RI, USA. Contact e-mail: lori_daiello@brown.edu

Background: Studies of psychoactive medication use among older adults with dementia have demonstrated that many take medications that may cause or contribute to memory impairment; however, patterns of psychoactive medication use in cognitively normal older adults volunteering for early intervention and AD prevention studies is unknown. **Objectives:** (1) Describe the prevalence of psychoactive medication use among members of the Rhode Island Hospital Alzheimer Prevention Registry; and (2) investigate relationships between sedative and anticholinergic medication burden and neuropsychological test performance at baseline and the first annual follow up visit. **Methods:** Subjects include 207 healthy, older adult (age 55-90 years) members of the Registry without previously diagnosed cognitive disorders, and interested in volunteering for AD prevention studies. Information on self-reported medication use, history of medical and psychiatric conditions, sociodemographics, APOE genotype, and neurocognitive test results, will be obtained from the Registry database. Prescription and over-the-counter medications are classified into two categories associated with cognitive toxicity in older adults: (1) sedative, and (2) anticholinergic drugs. For each participant, the degree of exposure to drugs in each category will be estimated using the Drug Burden Index (DBI), a dose-standardized measure that has been shown to predict cognitive and functional outcomes in previous studies of community-dwelling older adults. Relationships between the DBI and cognitive outcomes will be analyzed for each category of psychoactive medication, as well as the total psychoactive medication burden (DBI_{sedative} + DBI_{anticholinergic}) at baseline and one year. **Results:** Associations between the time-varying DBI scores and global cognitive functioning (Mattis Dementia Rating Scale-2), in addition to tests of cognitive domains (executive, attention, memory, psychomotor speed), will be analyzed with multivariable Poisson regression models, adjusted for potential confounding by indication with propensity score methods. Results will be presented in July 2015. **Conclusions:** The results of this study are of importance in developing selection criteria for AD prevention trials, because adverse cognitive effects of medications may exclude individuals who might otherwise qualify for participation. These findings also highlight the need for documenting medication use information in the databases of similar research registries that aim to recruit community-dwelling older adults concerned with brain health and aging.

P3-266

SELF-REPORTED INCREASED CONFUSION OR MEMORY LOSS AND CO-OCCURRING CONDITIONS AMONG ADULTS AGED 45 OR OLDER IN 21 STATES: 2012 BEHAVIORAL RISK FACTOR SURVEILLANCE SYSTEM

Angela J. Deokar, Lynda A. Anderson, Centers for Disease Control and Prevention, Atlanta, GA, USA. Contact e-mail: fjm4@cdc.gov

Background: Public health surveillance plays an important role in setting priorities and informing strategies for current and emerging