

Constraining Fossil Fuel CO₂ Emissions From Urban Area Using OCO-2 Observations of Total Column CO₂

Xinxin Ye^{1,2} , Thomas Lauvaux^{1,3} , Eric A. Kort⁴ , Tomohiro Oda^{5,6} , Sha Feng¹ , John C. Lin⁷ , Emily G. Yang⁴ , and Dien Wu⁷ 

Key Points:

- Inversion method is utilized to constrain whole-city fossil fuel emissions with measurement and transport model errors considered
- Potential of incorporating multiple tracks to obtain regular emission estimates is evaluated by pseudo data experiments
- Significant contribution of the biospheric fluxes variability to local X_{CO₂} variation is demonstrated

Supporting Information:

- Supporting Information S1

Correspondence to:

X. Ye,
xuy112@psu.edu;
xinxinye@ucla.edu

Citation:

Ye, X., Lauvaux, T., Kort, E. A., Oda, T., Feng, S., Lin, J. C., et al. (2020). Constraining fossil fuel CO₂ emissions from urban area using OCO-2 observations of total column CO₂. *Journal of Geophysical Research: Atmospheres*, 125, e2019JD030528. <https://doi.org/10.1029/2019JD030528>

Received 27 FEB 2019

Accepted 22 MAR 2020

Accepted article online 24 MAR 2020

¹Department of Meteorology and Atmospheric Science, Pennsylvania State University, University Park, PA, USA,

²Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, CA, USA, ³Laboratoire des Sciences du Climat et de l'Environnement, CEA, CNRS, UVSQ/IPSL, Université Paris-Saclay, Gif-sur-Yvette, France,

⁴Department of Climate and Space Sciences and Engineering, University of Michigan, Ann Arbor, MI, USA, ⁵Global Modeling and Assimilation Office, NASA Goddard Space Flight Center, Greenbelt, MD, USA, ⁶Goddard Earth Sciences Technology and Research, Universities Space Research Association, Columbia, MD, USA, ⁷Department of Atmospheric Sciences, University of Utah, Salt Lake City, UT, USA

Abstract Satellite observations of the total column dry-air CO₂ (X_{CO₂}) are expected to support the quantification and monitoring of fossil fuel CO₂ (ffCO₂) emissions from urban areas. We evaluate the utility of the Orbiting Carbon Observatory 2 (OCO-2) X_{CO₂} retrievals to optimize whole-city emissions, using a Bayesian inversion system and high-resolution transport modeling. The uncertainties of constrained emissions related to transport model, satellite measurements, and local biospheric fluxes are quantified. For the first two uncertainty sources, we examine cities of different landscapes: “plume city” located in relatively flat terrain, represented by Riyadh and Cairo; and “basin city” located in basin terrain, represented by Los Angeles (LA). The retrieved scaling factors of emissions and their uncertainties show prominent variabilities from track to track, due to the varying meteorological conditions and relative locations of the tracks transecting plumes. To explore the performance of multiple tracks in retrieving emissions, pseudo data experiments are carried out. The estimated least numbers of tracks required to constrain the total emissions for Riyadh (<10% uncertainty), Cairo (<10%), and LA (<5%) are 8, 5, and 7, respectively. Additionally, to evaluate the impact of biospheric fluxes on derivation of the ffX_{CO₂} enhancements, we conduct simulations for Pearl River Delta metropolitan area. Significant fractions of local X_{CO₂} enhancements associated with local biospheric X_{CO₂} variations are shown, which potentially lead to biased estimates of ffCO₂ emissions. We demonstrate that satellite measurements can be used to improve urban ffCO₂ emissions with a sufficient amount of measurements and appropriate representations of the uncertainty components.

1. Introduction

The global atmospheric CO₂ concentration has increased by more than 40% since the preindustrial era to more than 400 ppm in recent years and remains the main driver to current and future climate changes (Le Quéré et al., 2018). The increase in CO₂ concentration predominantly originates from anthropogenic CO₂ emissions by combustion of fossil fuels such as coal, petroleum, and natural gas (Andres et al., 2012; Ciais et al., 2013; Rotty, 1983). In order to foster the mitigation and management of anthropogenic CO₂ emissions, the international community has pursued treaties and agreements in the recent decades, such as the Kyoto Protocol (United Nations, 1998) and the Paris Agreement (UNFCCC, 2015).

A large percentage of the anthropogenic CO₂ is emitted from urban areas, about 40% as estimated by production-based figures (i.e., adding up emissions from entities located within cities) and as high as 60–70% with a consumption-based method (i.e., adding up emissions resulting from the production of all goods consumed by urban residents) (International Energy Agency, 2008; Satterthwaite, 2008; UN-Habitat, 2011). Given their significant contributions to fossil fuel CO₂ (ffCO₂), cities can perform as leading entities in implementing emission reduction plans. Comprehensive, accurate, and comparable emission quantifications are crucial for transparent monitoring of ffCO₂ emissions from urban areas and implementing effective mitigation schemes (Duren & Miller, 2012; Gurney et al., 2015; Pacala et al., 2010).

Emission inventory compilations with two methods referred to as “bottom-up” and “downscaling” are generally used to quantify ffCO₂ emissions. However, urban emissions usually bear large uncertainties, due to the missing socioeconomic information and inaccurate emission conversion factors, which are critical elements affecting the quality of emission inventories at the urban scale (Gately & Hutyra, 2017). For example, urban energy consumption and industrial activity data needed for “bottom-up” method are reported on a voluntary basis or under climate action activities in only a few cities, for example, those participating the Global Covenant of Mayors (<http://www.globalcovenantofmayors.org/>) and are usually spatially inexplicit, incomplete, and unverified (Hutyra et al., 2014). Similarly, for inventories developed by disaggregating (“downscaling”) national or regional emissions at fine scales, the uncertainty is large at high spatial and temporal resolutions caused by the disaggregation methods (Janssens-Maenhout et al., 2012; Kurokawa et al., 2013; Oda & Maksyutov, 2011), resulting in significant discrepancies among different emission inventories (Ackerman & Sundquist, 2008; van der Gon et al., 2011; Gurney et al., 2012, 2019; Hogue et al., 2016; Oda & Maksyutov, 2011; Oda et al., 2018; Turnbull, Karion et al., 2011). The two-sigma uncertainties of national annual ffCO₂ emissions are estimated to be 2–4% for countries with well-developed energy statistics and inventories (Rypdal & Winiwarter, 2001) and are at a possible order of 10% for countries with less well-developed energy data systems (IPCC, 2006). Nevertheless, urban-scale emission inventories can exhibit large differences (50–250%) compared to other downscaled data sets at local scales (Gately & Hutyra, 2017). The large uncertainties in ffCO₂ emissions not only impose difficulties on evaluating the effects of emission reduction strategies but also would lead to significant biases in the regional carbon budget estimations (Corbin et al., 2010). It is reported by European Space Agency (2015) that accuracies of inferred emissions in the order of 10% of the total would be needed for providing constraints that allows emission inventories to be evaluated at the time of an overpass. For cities that do not have inventories, an accuracy of 20% is already an important gain in information.

Atmospheric observations-based methods are becoming important ways to objectively obtain ffCO₂ emission estimates, allowing existing emission inventories to be improved. Some attempts have been made to derive local-scale emissions for the urban areas by utilizing the inverse modeling method (Bousquet, 2000; Ciais et al., 2010) with ground-based observations (Bréon et al., 2015; Lauvaux et al., 2016; McKain et al., 2012; Stauder et al., 2016; Wunch et al., 2009) or by the mass-balance approach with aircraft measurements (Cambaliza et al., 2014). However, one of the key limitations of these approaches is the unavailability of direct, continuous, and high-frequency atmospheric CO₂ measurements representing CO₂ enhancement in urban areas (Bréon et al., 2015), as only a handful of cities, mostly in Europe and North America, are instrumented with networks of CO₂ sensors (Bréon et al., 2015; Davis et al., 2017; McKain et al., 2012; Miles et al., 2017; Verhulst et al., 2017).

In recent decades, space-based satellite measurements of column-averaged CO₂ dry air mole fractions (X_{CO_2}) have been highly recommended for quantification and monitoring of urban ffCO₂ emissions, especially for cities where ground-based observations are sparse or unavailable (Duren & Miller, 2012; Kort et al., 2012; McKain et al., 2012; Schneising et al., 2013). Initial attempts have been made to relate satellite X_{CO_2} measurements to ffCO₂. Based on the retrievals from Scanning Imaging Absorption Spectrometer for Atmospheric Chartography (SCIAMACHY) on ENVISAT (2002–2012) (Bovensmann et al., 1999; Buchwitz et al., 2005), regional X_{CO_2} enhancements over the industrial areas in Germany were revealed to be correlated with yearly increase of anthropogenic CO₂ emissions (Schneising et al., 2008, 2013). In addition, with the launch of Greenhouse gases Observing SATellite (GOSAT) in 2009 (Kuze et al., 2009; Morino et al., 2011), discernible X_{CO_2} contrasts between the emission and background regions (Janardanan et al., 2016; Keppel-Aleks et al., 2013) and local X_{CO_2} enhancements over megacities (Kort et al., 2012) have been reported. However, due to coarse spatial resolution (~60 km × 30 km) and relatively low sensitivity (4–8 ppm) of the SCIAMACHY instrument, applications of its data are limited to large and intense emission regions. For the GOSAT instrument, the major limitation is its low sounding density, with a single 85-km² measurement per 250 km, resulting in fewer than 1,000 cloud-free soundings each day (Eldering, Wennberg, et al., 2017). Hence, these data sets are insufficient to enable accurate assessment of ffCO₂ urban emissions at high spatial and temporal resolutions.

National Aeronautics and Space Administration's Orbiting Carbon Observatory 2 (OCO-2) satellite mission (Crisp, 2008; Crisp et al., 2004) has been providing continuous and global retrievals of X_{CO_2} since September

2014 (Crisp, 2015). Although the OCO-2 mission is primarily developed for assessing regional carbon sources and sinks, its unique characteristics allow for detection of X_{CO_2} enhancements over cities at fine scales. The OCO-2 measurements have a higher spatial resolution than GOSAT and SCIAMACHY and collect more data per day. In particular, the small nadir footprint ($\sim 1.29 \text{ km} \times 2.25 \text{ km}$) helps to maximize the detectability of local emissions and increase the probability of cloud-free observations in the presence of patchy clouds. In addition, the high spectral resolution of OCO-2 enables high precision of X_{CO_2} with single sounding random errors of 0.5–1 ppm (Eldering, O'Dell, et al., 2017), which will greatly help to detect the small anthropogenic X_{CO_2} signals from the large background driven by biospheric sources and sinks and large-scale atmospheric transport (Keppel-Aleks et al., 2013; Turnbull et al., 2016). There have been some studies to examine X_{CO_2} imprints linked to anthropogenic emissions with the OCO-2 data. For example, spatial enhancements have been reported over the Northern Hemisphere regionally (Hakkarainen et al., 2016) and across the Los Angeles basin (Schwandner et al., 2017). For local-scale emissions and point sources, atmospheric transport modeling approach is applied to disentangle ffCO_2 emissions from background. Wu et al. (2018) developed a Lagrangian model to interpret the X_{CO_2} retrievals and constrain emissions from some cities in the Middle East. Nassar et al. (2017) presented the capability of quantifying ffCO_2 emissions from individual power plants by utilizing a Gaussian plume model.

Although the abovementioned studies have demonstrated the utility of OCO-2 X_{CO_2} on revealing ffCO_2 emissions, the emission estimation has not been carried out with high spatial resolution forward transport modeling so far, which has an advantage in capturing the fine-scale structure of ffCO_2 plume. Meanwhile, it is essential to evaluate the uncertainty in emission estimate related to atmospheric transport model error with the high-resolution plume simulations, which has been identified as a major source of uncertainty in inverse modeling (Gerbig et al., 2003; Houweling et al., 2010; Lauvaux & Davis, 2014; Lauvaux et al., 2012; Lin & Gerbig, 2005; S. M. Miller et al., 2015; Pacala et al., 2010). Additionally, the uncertainty in biosphere X_{CO_2} at local scale increases the difficulty of unambiguously disentangling fossil fuel emission signals in X_{CO_2} from the observations for cities in vegetated areas, which has not been evaluated in respect of space-based total column observations.

The previous studies were focused on cases using individual OCO-2 tracks. However, from the perspective of tracking ffCO_2 emissions for global cities on a regular basis, the potential of OCO-2 retrievals has yet to be investigated, which will be helpful to provide regular and policy-relevant references in support to improving the emission reduction strategies. Given current limitations in remote sensing that create a trade-off in sampling coverage and measurement precision, OCO-2 was designed as a sampling mission to provide measurements at high precision but only samples a small fraction of the globe each day (Eldering, Wennberg, et al., 2017), with a narrow swath ($\sim 10.3 \text{ km}$) and a long revisit cycle (~ 16 days). These features enable a small percentage of local emissions to be detected in each cycle (Pacala et al., 2010). Moreover, because the OCO-2 sampling locations vary among different observation modes (nadir, glint, and target), and the atmospheric transport condition changes, overpasses that happen to detect fossil fuel X_{CO_2} enhancements across a certain city are limited. For the cities examined in this work, about 5–15% of nearby OCO-2 tracks show detectable urban plumes, based on 15 months of OCO-2 data (September 2014 to November 2015). Hence, it would be still difficult to regularly monitor emissions by using individual overpasses. Despite these limitations, there is a potential to constrain fossil fuel emissions regularly by utilizing data collected along multiple tracks over one or more revisit cycle(s), which could enable tracking emission variations, although at a lower temporal resolution.

In this paper, we present the utility of OCO-2 X_{CO_2} data to constrain ffCO_2 emission estimations for urban areas. Several sources of uncertainty in emission estimate are evaluated, including the transport model errors, measurement errors, and variations in local biospheric carbon fluxes. High-resolution forward simulations are performed using the Weather Research and Forecast (WRF) model, which is capable of capturing fine-scale variability in X_{CO_2} distributions caused by transport and emission processes at urban scales. We first evaluate the emission estimate uncertainty related to transport model errors and measurement errors for three selected cities with different topographic influences and negligible impact of variations in local biospheric fluxes, that is, Riyadh and Cairo, classified as “plume cities” located in relatively flat terrain, and Los Angeles as a “basin city.” Based on these simulations, we carried out Observing System Simulation Experiments (OSSEs) to evaluate the potential of tracking urban emissions regularly by utilizing X_{CO_2}

data from multiple OCO-2 tracks. We also evaluated the uncertainty induced by local biospheric fluxes variability for the Pearl River Delta (PRD) metropolitan area. We discuss additional uncertainty sources in the inverse emission estimates, for example, the prior emission error correlations and the daytime-only sampling, and conclude with the implications for utilizing future satellite observations to monitor urban emissions.

2. Data and Method

2.1. OCO-2 X_{CO_2} Observations

The OCO-2 Lite files (Version 7r) from September 2014 to November 2015 are used in this study (obtained online at <https://co2.jpl.nasa.gov>). The OCO-2 satellite operates in a sun-synchronous polar orbit at the altitude of about 705 km and crosses the equator nominally at 13:36 LT (local time). It provides high-resolution spectroscopic measurements at eight adjacent 2.25-km-long footprints within a narrow swath every 0.333 s, with a cross-track resolution of 0.1–1.3 km (Crisp, 2008; Eldering, O'Dell, et al., 2017). The X_{CO_2} data are retrieved from the spectroscopic observations of reflected sunlight in near-infrared CO_2 and O_2 bands near midday, using the Atmospheric CO_2 Observations from Space algorithm (O'Dell et al., 2012). We used the bias-corrected X_{CO_2} data and filter them with “xco2_quality_flag” (QF) equals to 0, which is an indicator of data passing the internal quality check. The bias correction and data quality assessment have been detailed in the OCO-2 documentation (Mandrake et al., 2015). In addition, we analyzed each individual track over the selected cities for possible interferences by complex terrain, aerosols, and clouds. Some tracks were excluded due to contaminations by aerosols or clouds, as confirmed with the Cloud-Aerosol Lidar with Orthogonal Polarization data (Winker, 2016). The OCO-2 X_{CO_2} data are averaged over time windows of 1 s, consisting of 24 consecutive soundings (at most) representing ~ 10.32 km (cross track) \times 6.75 km (along track) in area. Note that we only derive 1-s averaged X_{CO_2} when there are at least 5 soundings (at most 24) in that time window passing the data selection criteria.

The measurement error of each X_{CO_2} sounding consists of two parts: a random error related to noise and a systematic error that is in principle bounded by the calculated interference error owing primarily to aerosol optical depth, surface albedo, and surface pressure (Boxe et al., 2010). For the random error, the OCO-2 data products include an estimate of the uncertainty on X_{CO_2} , which is generally smaller over water than the land surface and larger at the extreme latitudes (Eldering, O'Dell, et al., 2017). However, this estimation is a lower bound (Connor et al., 2016). Worden et al. (2017) evaluated the OCO-2 uncertainty by examining the X_{CO_2} variability within small neighborhoods of ~ 100 km \times 10.5 km, in which natural CO_2 variability is expected to be small. It is shown that the random error in the data product over land (~ 0.36 ppm) is smaller than the empirically derived X_{CO_2} random error (~ 0.75 ppm) by a factor of approximately 2. Conservatively, we consider a random error for each X_{CO_2} sounding with the standard deviation of 1 ppm, which leads to 0.20–0.45 ppm for the 1-s averaged data calculated with 5–24 soundings.

For the systematic error, as we used the bias-corrected data, some biases have been removed, for example, systematic footprint-to-footprint differences, mode-to-mode differences, and systematic differences that appear to be correlated to other retrieval variables (Mandrake et al., 2015). However, as validated using the Total Carbon Column Observing Network (TCCON) X_{CO_2} measurements by Wunch et al. (2017), residual biases remain in the OCO-2 retrievals after bias correction, with the absolute median differences < 0.4 ppm and RMS differences < 1.5 ppm. These biases appear to depend on latitude, surface properties, and scattering by aerosols. We note that for OCO-2 nadir and glint modes, these biases are evaluated using aggregated X_{CO_2} data within a box centered around the TCCON station that spans 5° in latitude (~ 555 km) and 10° in longitude ($\sim 1,100$ km) on the same day as a TCCON measurement. As we examine the variability in X_{CO_2} at a smaller spatial scale (~ 200 – 300 km, see Table 1 for the domain sizes), these biases in background can be removed when we extract local X_{CO_2} enhancements associated with fossil fuel emissions (ffX_{CO_2}) by subtracting the background X_{CO_2} . Due to the lack of an accurate representation of aerosol contamination in the current retrieval algorithm, we assume that the effects of urban aerosols on X_{CO_2} retrievals are negligible. Hence, the local ffX_{CO_2} derived from OCO-2 observations is assumed to be unbiased in this paper. Details of the derivation of background X_{CO_2} can be found in section 2.2.

Table 1
Summary of the Simulations Carried Out for the Selected Cities

City type	City/Metropolitan region	Land cover (% innermost domain) ^a	Transport error propagation method	Innermost domain size and resolution	Simulation time
Plume city	Riyadh	Barren or sparse (92.2%)	Perturbed plume	201 × 201 (1 km)	1–16 November 2014 17 December 2014 to 5 January 2015 27–30 January 2015
	Cairo	Barron or sparse (57.4%) Croplands (32.8%)	Perturbed plume	201 × 201 (1 km)	4–7 October 2014 16–19 March 2015 17–20 May 2015 13–16 July 2015 14–17 August 2015
Basin city	Los Angeles (LA) metropolitan region	Water (45.2%) Open shrublands (34.9%) Barren or sparse (6.1%) Woody savannas (3.6%) Croplands (2.9%)	Ensemble simulation	207 × 150 ^b (4 km)	3 July to 20 August 2015 6–19 October 2015
Multi-city	Pearl River Delta (PRD) metropolitan region	Water (26.7%) Croplands (26.2%) Evergreen broadleaf (20.0%) Woody savannas (10.1%) Urban (8.6%)	Perturbed plume	240 × 240 (1.333 km)	12–15 January 2015 1–4 August 2015

^aThe domain of interest for LA is set to 119.0°W to 116.3°W, 32.2–35.7°N, which is smaller than the innermost domain of simulations. ^bLand cover is based on MODIS IGBP 21-category data. The land cover types accounting for more than 90% of the innermost domain in aggregate are listed.

2.2. Background X_{CO2}

We extract ffX_{CO_2} , that is, X_{CO_2} enhancements caused by ffCO_2 emissions, by subtracting background X_{CO_2} from the OCO-2 X_{CO_2} retrievals. A typical method to derive ffCO_2 from in situ CO_2 measurements is to calculate the difference between CO_2 at an upwind site and a downwind site (e.g., Bréon et al., 2015; Lauvaux et al., 2016; Super et al., 2017). The best application condition of this background method is when the wind vector is aligned with the stations. Similarly, when the wind is aligned with the orbit, we could get background X_{CO_2} by averaging the measurements in the upwind region of a city. However, the alignment is very rare for OCO-2 (Nassar et al., 2017). In previous studies, a constant background is often used for X_{CO_2} retrievals collected over a time period, which is calculated as the median X_{CO_2} over a latitudinal band (Hakkarainen et al., 2016) or the average X_{CO_2} in a “background area,” for example, the desert located close to Los Angeles (Kort et al., 2012). However, for a single overpass, a constant background would not represent the spatial variability of the background concentrations. Thus, for each single overpass across a city of interest, we derived a “background line,” as shown with the black lines in Figures 4, 5, and 7. The “background line” is derived by a two-step linear regression. We firstly decompose the X_{CO_2} data into two parts, that is, $\text{X}_{\text{CO}_2} = \text{X}_{\text{CO}_2\text{-trend}} + \text{X}_{\text{CO}_2\text{-local}}$, where the $\text{X}_{\text{CO}_2\text{-trend}}$ is the nonlocal trend represented using a linear function: $\text{X}_{\text{CO}_2\text{-trend}} = a \cdot x + b$. Here x is latitude, and a and b are the slope and interception derived by linear regression. With the standard deviation of $\text{X}_{\text{CO}_2\text{-local}}$ (σ_{local}) representing the local-scale variability, we filtered the X_{CO_2} samples with $\text{X}_{\text{CO}_2} < \text{X}_{\text{CO}_2\text{-trend}} + 0.5\sigma_{\text{local}}$. These filtered data are chosen as “background samples” (the black triangles in Figures 4, 5, and 7), as they have lower spatial variability at local scales compared to the samples affected by urban ffCO_2 emissions. Then we recalculate the linear regression line, that is, the “background line,” based on the “background samples.” This “background line” method allows considering the spatial trend in the background.

2.3. Cities of Interest

Four cities/urban areas are selected in this work considering the availability of OCO-2 data, topography, and vegetation coverage to highlight different sources of uncertainty in the emission estimates. To explore the impact of transport model errors, three different cities are chosen according to the following criteria: (i)

distant isolation from other large anthropogenic emission sources nearby, (ii) large fossil fuel carbon emissions, (iii) low cloud cover and relatively preferable data availability, and (iv) weak contribution of biospheric signals. Cities are categorized into two different types based on the impact of local topography on dispersion: “plume cities” located in relatively flat terrain and “basin cities” located in complex terrain trapping the dispersion. In this paper, Riyadh, Saudi Arabia, with a population of 6.2 million, and Cairo, Egypt, with 18.4 million are chosen as typical “plume cities,” which are characterized by ffX_{CO_2} enhancements distributed as plumes. The Los Angeles metropolitan area (referred to as LA hereafter) with a population of more than 13 million is chosen as a characteristic “basin city.” The LA basin presents large elevation gradients from the sea surface to the top of Mount Wilson to the north. The strong enhancements in X_{CO_2} are mainly due to air masses trapped in the basin, which has been referred to as “urban dome” (Idso et al., 1998), albeit we do not adopt this terminology in this work due to the potential confusion with actual accumulation of CO_2 .

To evaluate the impact of uncertainties in biospheric fluxes on the emission estimate, the PRD region of China is selected, where an agglomeration of several cities is located, including Guangzhou, Hong Kong, Shenzhen, Zhuhai, Dongguan, and Zhongshan. The PRD region is one of the largest metropolitan areas in the world with about 45 million people. The cities are less vegetated compared to their surrounding area, leading to a distinctive contrast in the net ecosystem exchange (NEE) in the urban area and surrounding rural area (see Figure S2 in the supporting information and section 2.4.3).

2.4. Model Setup

2.4.1. Atmospheric Transport Model

The spatial heterogeneity of emissions and intense point sources (e.g., power plants) lead to complex spatial structures of urban emissions, resulting in complex ffCO_2 plume combined with local atmospheric dynamics (e.g., Feng et al., 2016). In order to capture the fine-scale variations, we simulate the ffX_{CO_2} using the Weather Research and Forecasting model (V3.6.1) with chemistry (WRF-Chem) (Grell et al., 2005; Skamarock et al., 2008), coupled to CO_2 emissions and biospheric fluxes using the passive tracer mode (Lauvaux et al., 2012). Model grids are configured with 51 vertical (eta) levels. The 6-hourly NCEP FNL (Final) Operational Global Analysis data on $0.5^\circ \times 0.5^\circ$ grids are used as the initial and boundary conditions of meteorological and land surface fields. The boundary condition of CO_2 concentration for the outermost domain is 390.0 ppm. The simulations are initiated every 4 days at 12:00 UTC with an integration time of 108 hr, including a spin-up time of 12 hr and producing hourly outputs.

One-way nested domains with resolutions of 27, 9, 3, and 1 km are used for Riyadh and Cairo, 36, 12, and 4 km for LA, and 36, 12, 4, and 1.333 km for the PRD region. The innermost domains and distributions of ffCO_2 emission for the selected cities are shown in Figure S3. For Riyadh, Cairo, and PRD, the innermost domains are used as the domains of interest to filter the OCO-2 observations. Note that for LA, the domain of interest is set to $119.0\text{--}116.3^\circ\text{W}$, $32.2\text{--}35.7^\circ\text{N}$, which is smaller than the innermost domain, and the spatial resolution is coarser than others. As reported by Feng et al. (2016), the 1.3-km run does not show significant improvement compared to the 4-km run at the surface, even though it resolves the vertical gradient of horizontal winds and planetary boundary layer (PBL) better. Given that aggregated ffX_{CO_2} along track is used to compute the scaling factor of a priori emissions, we compared the aggregated ffX_{CO_2} using 4- and 1.3-km runs, and they also present similar results owing to ffX_{CO_2} mostly being trapped within the basin during daytime. Therefore, we used 4-km resolution for LA in this study.

A summary of the simulations performed in this study is shown in Table 1. The fossil fuel CO_2 emission data and NEE data are detailed in sections 2.4.2 and 2.4.3. Note that the simulations for tracers imposed by NEE are only conducted for the PRD region. Moreover, an ensemble of modeling based on model physics parameterizations is deployed to represent the transport model errors in the simulated ffX_{CO_2} over LA (see section 2.5.1).

2.4.2. Fossil Fuel CO_2 Emissions

The Open-source Data Inventory for Atmospheric Carbon dioxide (ODIAC) Version 2015a (Oda et al., 2017; 2018; Oda & Maksyutov, 2011, 2015) is used in this paper for emissions from the cities of interest. The ODIAC emission product provides 1-km \times 1-km gridded global and monthly ffCO_2 emissions. It is developed based on country-level ffCO_2 emission estimates, fuel consumption statistics, satellite-observed nightlight data, and point source information (geographical locations and emission intensities) from the CARbon




Figure 1. Example of biogenic carbon fluxes (net ecosystem exchange, NEE) downscaling in PRD region. The top panels show the NEE from 3-hourly MsTMIP data at 12:00 LT 12 January 2010 on (a) the original $0.5^\circ \times 0.5^\circ$ grid, (b) WRF grid (1.333×1.333 km), derived by bilinear interpolation of original NEE, and (c) WRF grid (1.333×1.333 km), derived by scaling the interpolated NEE. The bottom panels show the green vegetation fraction (GVF) in January on (d) the $0.5^\circ \times 0.5^\circ$ grid, (e) WRF grid (1×1 km) by bilinear interpolation of GVF in (d), and (f) WRF grid (1×1 km). See section 2.4.3 for further details.

Monitoring for Action power plant database (Oda et al., 2018). The global nightlight data were used as a georeferenced, spatial proxy to determine the spatial extent of anthropogenic emissions from the line and diffused (area) sources (e.g., road traffic, residential, or commercial fuel consumption). The ODIAC gridded emission fields defined on a global rectangular (latitude \times longitude) coordinate are remapped to meet the grid resolutions for each simulation domain. Note that temporal variability of emissions at diurnal and weekly scales is not included in the modeling and the pseudo data experiments. We remapped the monthly emission distributions for the time periods investigated in our simulation. The ffCO_2 emission distributions are shown in Figure S3. All ffCO_2 is released at the ground surface.

2.4.3. Biogenic CO₂ Fluxes

The NEE fluxes in the PRD region are provided by the 15 different global Terrestrial Biogeochemical Models ($0.5^\circ \times 0.5^\circ$) in the Multi-scale Synthesis and Terrestrial Model Intercomparison Project (MsTMIP) (Huntzinger et al., 2013). In order to better characterize the diurnal variability and spatial distribution of biogenic fluxes, a 3-hourly data set for global biogenic fluxes (Fisher et al., 2016) is used, which is temporally downsampled from the monthly global models. Furthermore, we spatially downscale the 3-hourly NEE from the original MsTMIP grid ($0.5^\circ \times 0.5^\circ$) (e.g., Figure 1a) to the WRF domains (36-, 12-, 4-, and 1.333-km resolutions) using the green vegetation fraction (GVF), with the assumption that vegetation productivity and respiration scales linearly with canopy coverage in each grid cell. We note that using this method, besides plant productivity and respiration, soil respiration is also downsampled by GVF, which could lead to some misrepresentation of soil respiration although the impact on the results would be small. The GVF is defined as the fraction of the grid cell for which midday downward solar insolation is intercepted by a photosynthetically active green canopy. A robust relationship between canopy cover and biomass was observed in Boston,




Figure 2. (a) Local ffX_{CO_2} derived from OCO-2 data (colored dots) at about 10:00 UTC 29 December 2014 by subtracting the background concentration and the simulated ffX_{CO_2} enhancement (color shading) using ODIAC emissions, which is the truth in the OSSEs. Panels (b) and (c) show the rescaled plumes of panel (a) with wind speed error of 1.0 and -1.0 m/s. Panel (d) shows the rescaled plume with wind direction error of 5.0° .

which supports the use of GVF as a proxy for biomass, and hence as a scaling parameter for biogenic fluxes (Briber et al., 2013; Raciti et al., 2012). The NEE can be downscaled as follows:

$$E_{wrf,i,j} = (E_{blin,i,j}/GVF_{blin,i,j}) \times GVF_{wrf,i}, \quad (1)$$

where the subscripts i and j represent the coordinates of a WRF grid cell, E_{wrf} the NEE at WRF grid (e.g., Figure 1c), and E_{blin} the bilinear interpolated NEE from the original $0.5^\circ \times 0.5^\circ$ grids to WRF grid (e.g., Figure 1b). GVF_{blin} is interpolated using Moderate Resolution Imaging Spectroradiometer (MODIS) climatological GVF (e.g., Figure 1d) in the same way of deriving E_{blin} (e.g., Figure 1c), ensuring the same spatial representativeness of GVF_{blin} and E_{blin} , and GVF_{wrf} (e.g., Figure 1e) is the GVF projected to the WRF grid. The GVF data used in this study are based on MODIS climatological observations from 2001 to 2010, which is available in the geographic data since WRF v3.6. The uncertainties in biogenic X_{CO_2} are represented by the spread of simulated biogenic X_{CO_2} using the NEE from the 15-member models in MsTMIP.

2.5. Representation of Transport Model Errors

The impact of transport model errors in wind speed and wind direction on the uncertainty of emission estimates is considered in the inversions. In this section, we introduce the methods to propagate the transport model errors to the modeled ffX_{CO_2} fields across different types of cities. The method applied for plume city and basin city are explained in sections 2.5.1 and 2.5.2, respectively.

Table 2
WRF Model Configurations of the Ensemble of Simulations Conducted for Los Angeles

Ensemble member	PBL scheme	Surface layer scheme	Urban canopy model
MYJ	MYJ	Eta similarity (Janjić Eta)	None
MYJ_UCM	MYJ	Eta similarity (Janjić Eta)	Noah UCM
MYNN	MYNN	Nakanishi and Niino	None
MYNN_UCM	MYNN	Nakanishi and Niino	Noah UCM
BouLac_BEP	BouLac	Eta similarity (Janjić Eta)	BEP
BouLac_UCM	BouLac	Eta similarity (Janjić Eta)	Noah UCM

2.5.1. Plume City: Transformation of Plumes

For a “plume city,” the transport model errors are propagated by transformation of the modeled ffX_{CO_2} plume. More detail of this method is included in the supporting information (Text S1.1). The errors for the “plume city” are assumed to be unbiased based on the previous study on Indianapolis (Deng et al., 2017). For a random error (θ) in wind direction, we rotate the simulated plume $\bar{c}(x, y, t)$ at a given time (t) by an angle θ about the emission center (x_0, y_0) to get $\bar{c}_r(x_r, y_r, t)$. Note that x axis is aligned with the dominant wind direction in the domain. Then the rotated plume is transformed to incorporate random wind speed error (ε) as follows:

$$\bar{c}_r(x_r, y_r, t) = \bar{c}(x_r, y_r, t), \quad (2)$$

where

$$x_r = \frac{\bar{u} + \varepsilon}{\bar{u}}(x_r - x_0) + x_0, y_r = y_r, z_r = z_r, \bar{c}_r = \bar{c} \frac{\bar{u}}{\bar{u} + \varepsilon} \quad (3)$$

Since the ffCO_2 is confined within the PBL and well mixed during the daytime, we used the domain average wind speed within the PBL and its typical error statistics for \bar{u} and ε . The errors are assumed to follow normal distributions of $N(0, 1.0)$ (unit: m/s) for wind speed and $N(0, 15)$ (unit: °) for wind direction, respectively. The selection of the error ranges is consistent with the model error statistics in the lower troposphere (<2 km) from model evaluation studies, for example, Indianapolis (Deng et al., 2017), Los Angeles (Feng et al., 2016), and more generally over the U.S. Upper Midwest (Diaz-Issac et al., 2018).

Examples of transformed plumes using this method are shown in Figure 2 based on a simulated ffX_{CO_2} plume over Riyadh at about 10:00 UTC 29 December 2014. The impacts of positive and negative wind speed errors are represented by transformation (Figures 2b and 2c), rescaling the plume along the domain average wind direction. The impact of a wind direction error is represented by a rotation of the plume (Figure 2d).

To obtain the transport model uncertainty in the modeled ffX_{CO_2} , we transform the modeled plume by multiple times (here 10^4) with random wind speed and wind direction errors, and extract the uncertainty spread of ffX_{CO_2} by using the 25th and 75th percentiles. The transformation method is a trade-off between running a simplified model (e.g., a Gaussian plume model) and running an ensemble of simulations. With this method, we can retain the complexity of X_{CO_2} plume structures while exploring the transport error impact at low computational costs. Note that this method is derived under the assumption of spatially uniform wind errors, which is generally valid within a few tens of kilometers from the emission center.

2.5.2. Basin City: Model Physics-Based Ensemble Simulation

For LA, a typical basin city, the transport model errors are represented by an ensemble of WRF simulations with different PBL and urban canopy physics parameterizations, following Feng et al. (2016). This method is suitable for transport conditions with the dispersion of CO_2 trapped by local topography. Four different PBL parameterizations are used, that is, the Mellor-Yamada-Nakanishi-Niino 2.5 (Nakanishi & Niino, 2004) scheme, the Mellor-Yamada-Jancic scheme (Janjić, 1994), and the Bougeault and Lacarrère (BouLac) (Bougeault & Lacarrere, 1989) scheme. For the land surface processes in urban canopy, the single-layer urban canopy model (Kusaka & Kimura, 2004) and the multilayer building environment parameterization (Martilli et al., 2002) are used. The ensemble of simulations with different combinations of the model physics schemes (Table 2) can represent the model uncertainties in wind field, PBL structure, and PBL height.




Figure 3. Comparison of the modeling uncertainty and mean absolute error (MAE) of 10-m wind speeds and wind directions over 43 surface sites located in the 1.333-km resolution domain for LA. The modeling random uncertainty is calculated as the standard deviation (STD, blue scatters) and semifull range of the modeling results, that is, half of the difference between maximum and minimum values among the ensemble members (red scatters). The top two panels show results for the original six members, and the bottom two panels are for the 18 members with modeling results at ± 1 hr included. The red and blue crosses in each panel stand for average points of the scatters in the corresponding colors.

In order to evaluate the performance of the ensemble on representing the model-observation mismatches, the modeling results of wind speed and wind direction are compared with surface wind observations. Surface observations of wind speed and wind direction at 43 synoptic weather stations located within the 4-km domain covering LA were used, derived from the global hourly Integrated Surface Data and accessible at the National Centers for Environmental Information (<https://gis.ncdc.noaa.gov/geportal/catalog/search/resource/details.page?id=gov.noaa.ncdc:C00532>).

For each observation time, the mean absolute error (MAE) of modeling result is calculated to evaluate the magnitude of the model error. We compared the ensemble spread presented by two approaches: (1) the standard deviation of the ensemble results and (2) semifull range of the ensemble results (half of the difference between the maximum and minimum values). As shown in Figure 3, both of the approaches based on the simulation results of the six members exhibit a somewhat lower ensemble spread of wind speed compared to the observed MAE, suggesting an underestimation of the transport uncertainty (Figures 3a and 3b). When taking the model results at ± 1 hr relative to the observation time into account (Figures 3c and 3d with the ensemble size of 18 members), the ensemble spread is found to be enlarged, suggesting a better agreement with the observed MAE. The ensemble spreads for wind speed and wind direction both show better agreement with the MAE. It can also be notable that the semifull range yields a better representation of the model uncertainty compared to the standard deviation (Figures 3c and 3d). Hence, for LA we use the 18 ensemble members and the semifull range to estimate transport model uncertainty spread in the simulated ffX_{CO_2} .

For the systematic errors in the transport model, we compared the ensemble mean surface wind speed to the observations. The result shows a positive bias of 0.48 m/s. Similarly, Angevine et al. (2012) found a wind

speed bias of 1.1 ± 2.7 m/s. Feng et al. (2016) reported a slightly smaller bias of ~ 1.0 m/s for LA and showed larger biases near mountainous sites owing to complex topography. We note that the wind speed error is evaluated at surface in this work. The wind speed bias varies at different altitudes through the PBL, with usually larger value near the surface (Feng et al., 2016).

The wind speed bias can result in systematic error in emission estimates. In order to represent the impact of wind speed bias in the pseudo modeling data in the OSSEs, we apply a factor (k) to the simulated ffX_{CO_2} for the pseudo truth, where

$$k = \frac{\bar{u}}{\bar{u} + \Delta}, \quad (4)$$

\bar{u} is the average wind speed over the domain and Δ is the corresponding bias (in m/s). This factor assumes a single ratio over the entire domain but allows to increase/decrease the ffX_{CO_2} to represent the impact of negative/positive wind speed bias.

2.6. Emission Optimization Method

A Bayesian inversion method similar to that used by Pillai et al. (2015) is implemented, which optimizes parameters of ffCO_2 emissions with observational constraints to obtain the best consistency between modeled and observed ffX_{CO_2} enhancements. Specifically, we optimize the emissions by adjusting a scaling factor (λ) upon the a priori emissions from the entire city, without modifying the spatial distribution. The integrated ffX_{CO_2} signals along a latitudinal range of interest of OCO-2 tracks are used as the observational constraint, which can be represented as follows:

$$y_o = \int_{\text{lat1}}^{\text{lat2}} \text{ffX}_{\text{CO}_2,o}, \quad y_m = \int_{\text{lat1}}^{\text{lat2}} \text{ffX}_{\text{CO}_2,m} \quad (5)$$

where y_o and y_m are the observed and modeled results, respectively. Here, $\text{ffX}_{\text{CO}_2,o}$ is derived by subtracting the background X_{CO_2} from observations, and $\text{ffX}_{\text{CO}_2,m}$ is derived by interpolating the modeling results of the tracer tagged with fossil fuel emissions at the coincident geolocations of the observations. Compared to deriving the scaling factor with the least square error method using all the soundings, the integral ffX_{CO_2} shows less sensitivity to wind direction error (Figure S2 and Text S1.2).

For n pairs of observations and modeling results obtained with n OCO-2 tracks, the integrated enhancements can be represented as:

$$\mathbf{y}_o = \mathbf{y}_m \cdot \lambda + \boldsymbol{\varepsilon}_o \quad (6)$$

Here the state vector has been simplified as a scalar, that is, the scaling factor λ , and the Jacobian matrix that represents the sensitivity of the observations to the state vector is given by a vector \mathbf{y}_m . The term $\boldsymbol{\varepsilon}_o$ is an observational error vector, including errors in the OCO-2 measurements, forward model, and model parameters. It is assumed to follow the Gaussian distribution described with the error covariance matrix S_o . As the observation errors are assumed to be uncorrelated for different tracks, S_o is a diagonal matrix with the main diagonal entries representing the error variance of the observation (σ_o^2) for each track. As the measurement and model uncertainty are unbiased and not correlated, we estimate σ_o^2 by adding the error variances:

$$\sigma_o^2 = \sigma_{\text{measurement}}^2 + \sigma_{\text{model}}^2, \quad (7)$$

where $\sigma_{\text{measurement}}^2$ is the measurement error variance and σ_{model}^2 is the forward model error variance. The estimations of these two terms have been detailed in sections 2.1 and 2.5, respectively.

The posterior estimate of λ is derived by minimizing the cost function (J):

$$J(\lambda) = (\mathbf{y}_o - \mathbf{y}_m \lambda)^T S_o^{-1} (\mathbf{y}_o - \mathbf{y}_m \lambda) + (\lambda - \lambda_a)^2 \sigma_a^{-2} \quad (8)$$

where σ_a^2 is the error variance of the prior estimate, λ_a . The prior estimate λ_a is set to unity. The prior emission uncertainty σ_a is set up based on reported estimations in literature, as the ODIAC data product did not

provide uncertainty estimates. At annual scales, Gurney et al. (2019) investigated the difference between the ODIAC data and a high-resolution bottom-up estimate product (Hestia) in four U.S. urban areas, showing the differences of whole-city emissions ranging from -1.5% to $+20.8\%$. Oda et al. (2019) found differences of about 40% by comparing a satellite-derived annual emissions product to a gridded national inventory at 25-km resolution. However, the uncertainty would become larger at a smaller time scale. Considering the variability in anthropogenic activities, for example, different power demand on weekdays and weekends, weather-related events, domestic heating, and air conditioning, the day-to-day variability of 20 to 50% can be found in emission inventories. As most of the temporal patterns in current emission products are prescribed, and based on recent publications at different timescales, we suggest uncertainties of 50% at the daily timescale as a lower bound. As this work focuses on quantifying the whole-city emissions, we conservatively set the prior flux uncertainty σ_a to 20% for LA, since the emissions from the U.S. megacities are relatively well characterized. For Riyadh and Cairo, the prior uncertainty is set to 40%, as the fuel consumptions are expected to have higher uncertainties than over the United States.

By solving the minimum of J , the optimal estimate of scaling factor, $\hat{\lambda}$, and the posterior error variance, $\hat{\sigma}^2$ can be obtained as follows:

$$\hat{\lambda} = (\mathbf{y}_m^T \mathbf{S}_o^{-1} \mathbf{y}_m + \sigma_a^{-2})^{-1} (\mathbf{y}_m^T \mathbf{S}_o^{-1} \mathbf{y}_o + \sigma_a^{-2} \lambda_a) \quad (9)$$

$$\hat{\sigma}^2 = (\mathbf{y}_m^T \mathbf{S}_o^{-1} \mathbf{y}_m + \sigma_a^{-2})^{-1}. \quad (10)$$

2.7. OSSEs: Emission Optimization Using Multiple Tracks Under Different Meteorological Conditions

Given the limited amount of real OCO-2 overpasses and the observation geometry, OSSEs are implemented to examine the potential of using multiple OCO-2 tracks to constrain urban ffX_{CO_2} emissions in a statistical prospective, which allows examining the performances under different atmospheric transport conditions. Specifically, we examine the relation between the number of available tracks and the emission estimate uncertainty.

For each city, we use a specified and typical OCO-2 ground track locations in nadir observation mode to extract the pseudo observations and pseudo modeling data, based on the hourly modeled ffX_{CO_2} . We avoided random selection of track locations across the domain, as OCO-2 tracks repeat over time without major variations. As the OCO-2 overpasses are available only during daytime, we selected the modeling results during 09:00–15:00 LST with domain-averaged surface wind speed ≥ 2 m/s. Additionally, for Riyadh and Cairo, in order to ensure that plumes are transected by the satellite, the simulation results are also filtered with the angle from a plume axis to the typical track $\geq 10^\circ$ and $\leq 170^\circ$.

The prior emissions, that is, the ODIAC data, are set as the true emissions in the OSSEs. Hence, pseudo observations of ffX_{CO_2} are obtained by sampling from the modeled ffX_{CO_2} at the locations of the 1-s averaged soundings along the typical track. The pseudo observations are assumed to be unbiased relative to the truth. A random error per sample has been added, following a Gaussian distribution with the standard deviation of 0.2 ppm representing the lower bound of the measurement errors (section 2.1).

The pseudo modeling data of ffX_{CO_2} are obtained in different ways for different cities. For Riyadh and Cairo, the pseudo modeling data are set directly to the modeled ffX_{CO_2} sampled along the typical track and perturbed by random transport errors. Thus, the posterior emission estimate would be unbiased, and the truth of scaling factor would be unity. For LA, as the positive wind speed bias has been seen with the comparison to surface observations, we apply a factor to the simulated ffX_{CO_2} to represent the impact of wind speed bias on the pseudo modeling ffX_{CO_2} (see equation 4), which lead to a biased posterior scaling factor as will be shown in the results. The transport model uncertainty is estimated with the same methods detailed in section 2.5. We note that the uncertainty in background X_{CO_2} is not included in the observation errors.

The impact of numbers of available tracks ($N = 1, \dots, 20$) on emission estimate uncertainty is evaluated by a Monte Carlo approach. For each number of tracks (N), we randomly select N pairs of pseudo observation and modeling data. The scaling factor is derived with the same inversion method (section 2.6) as used for the real

Table 3
Inversion Results of Scaling Factors of the Whole-City Emissions for the Selected Cities Using OCO-2 X_{CO_2} Data

City	Date of OCO-2 track	Prior total emission (Mt C/month)	Prior total emission uncertainty (σ_a)	Measurement uncertainty ($\sigma_{\text{measurement}}$, units: ppm)	Transport model uncertainty (σ_{model} , units: ppm)	Scaling factor (λ) \pm posterior uncertainty ($\hat{\sigma}$)
Riyadh	27 December 2014	3.08	40%	1.34	2.80	0.92 ± 0.24
	29 December 2014	3.08		1.30	2.83	0.83 ± 0.17
Cairo	28 February 2015	2.52	40%	1.07	0.71	0.79 ± 0.15
	18 March 2015	2.70		0.55	0.31	1.18 ± 0.25
	19 May 2015	2.48		1.18	0.36	0.95 ± 0.11
	15 July 2015	2.62		1.08	0.16	0.70 ± 0.13
	16 August 2015	2.61		0.94	0.54	0.81 ± 0.10
Los Angeles	6 July 2015	4.68	20%	1.48	2.99	1.00 ± 0.15
	15 July 2015	4.68		2.11	1.62	1.84 ± 0.16
	7 August 2015	4.74		1.11	1.35	1.17 ± 0.15
	16 August 2015	4.74		1.93	3.03	1.43 ± 0.14
	10 October 2015	4.49		1.20	1.08	1.64 ± 0.15
	12 October 2015	4.49		1.40	2.70	0.66 ± 0.11

Note. The scaling factors and their posterior uncertainties are shown for each individual track, as well as for integrating the information from all the selected tracks. Uncertainty components are listed for each track, including the prior uncertainty of scaling factor, and the measurement and transport uncertainties of the integral $\text{ff}X_{CO_2}$ (the larger value between these two is shown in bold).

tracks analyses. Probability distributions of scaling factor and the associated posterior uncertainty are obtained by repeating the random selection procedure.

3. Results

3.1. Local Fossil Fuel X_{CO_2} Enhancement ($\text{ff}X_{CO_2}$)

The $\text{ff}X_{CO_2}$ enhancement, defined as the enhancement in X_{CO_2} associated with local fossil fuel emissions relative to the background concentration, is used to constrain emissions in this study. In this section, the $\text{ff}X_{CO_2}$ enhancements are shown for Riyadh, Cairo, and LA, estimated by the simulations of $\text{ff}CO_2$ tracers tagged with the ODIAC emissions. The results are compared to the observed enhancements derived from the OCO-2 X_{CO_2} data, in order to evaluate the magnitude of $\text{ff}X_{CO_2}$, that is, to assess whether the signals of local emissions are robust and detectable. Note that the whole-city emissions from these three cities in the months of the selected tracks are about 3.1, 2.5–2.7, and 4.5–4.7 Mt C/month based on the ODIAC data (see Table 3). The OCO-2 tracks shown in this section are chosen by examining the observed X_{CO_2} and the simulated plumes, to ensure that the satellite overpasses are downwind of the city with X_{CO_2} signals attributable to local emissions.

3.1.1. Comparison of Modeled and Observed $\text{ff}X_{CO_2}$

Figure 4 shows the $\text{ff}X_{CO_2}$ enhancements derived over Riyadh on 27 and 29 December 2014. Overall, the modeled enhancements over this domain are pronounced and can be up to about 6 ppm (Figures 4c and 4d), indicating signals that can be unambiguously detected from space, given the precision and accuracy of the OCO-2 observations. The modeled $\text{ff}X_{CO_2}$ distributions are characterized by elongated and non-Gaussian plume structures, mostly due to complex horizontal wind fields (Figures 4c and 4d). By examining the simulated $\text{ff}X_{CO_2}$ at one hour earlier and 1 hr later, rapidly changing fine-scale structures can be seen (Figure S4), indicating notable variations in the distributions of $\text{ff}X_{CO_2}$ over a few hours. It is also noteworthy that, to examine the impact of domain resolution on the simulated $\text{ff}X_{CO_2}$, we compared the results of 1-, 3-, and 9-km spatial resolutions. Lower and smoother peaks can be seen for the coarser resolutions (see Figure S5), which is expected given the larger aggregation errors of the emissions at the lower spatial resolution, particularly for intense point sources. This result suggests the necessity of using a high spatial resolution to reproduce the complex plume structures for the “plume cities”.

Figures 4a and 4b show the modeled and observed X_{CO_2} along the two tracks. Note that the simulated X_{CO_2} shown in these plots is derived as the sum of background line derived with the OCO-2 data and the simulated $\text{ff}X_{CO_2}$. The observed $\text{ff}X_{CO_2}$ enhancements reach ~ 1.5 and ~ 2.0 ppm for these two tracks. The magnitudes of




Figure 4. Comparisons of modeled and observed ffX_{CO_2} enhancements by the OCO-2 data on 27 and 29 December 2014 at about 10:00 UTC over Riyadh. Panels (a) and (b) show the OCO-2 X_{CO_2} (black dots) and simulated X_{CO_2} (red dotted line, sum of ffX_{CO_2} and the background concentrations) along the two tracks. The OCO-2 retrievals are filtered with quality flag ($QF = 0$) and are 1 s averaged. The blank triangles represent the data used for the derivation of background concentrations (black solid line) by linear regression with these data versus latitude. The uncertainty in the simulated ffX_{CO_2} related to transport model errors is shown by the light red shading. Panels (c) and (d) show the simulated ffX_{CO_2} and the observed ffX_{CO_2} obtained from the OCO-2 data, filtered with $QF = 0$ only. The background X_{CO_2} concentrations have been subtracted. The vectors represent 10-m wind with the reference vector standing for the wind speed of 5 m/s.

modeled ffX_{CO_2} are generally in agreement with the observations, although there is a prominent spatial displacement of $\sim 0.3^\circ$ in latitude (~ 33.4 km, Figure 4a) between the observed and simulated peaks on 27 December, and the modeled peak on 29 December is narrower than observed. The large spatial offset might be due to the satellite track transecting the edge of the plume in a nearly parallel way, so that the modeled ffX_{CO_2} values are very sensitive to errors in horizontal wind field.

For Cairo, the local ffX_{CO_2} enhancements are examined for five tracks on 28 February, 18 March, 19 May, 15 July, and 16 August 2015, respectively. As shown Figure 5, the modeled ffX_{CO_2} enhancements over Cairo are mostly < 3.0 ppm over the simulation domain, with some hot spots located close to some intense sources (Figure S3). In comparison to Riyadh, the ffX_{CO_2} enhancements are overall smaller, as expected given the lower total $ffCO_2$ emission. Compared to the simulated ffX_{CO_2} , the observed local enhancement peaks are mostly higher and narrower, especially for the tracks on 19 May and 15 July 2015, while the modeled enhancements are smoother and diffuser. Spatial displacements in the signals are also seen for the tracks on 28 February and 16 August 2015.

It is notable that the background X_{CO_2} values represented with the background lines exhibit higher latitudinal gradients for most of the selected tracks over Cairo (Figures 5a and 5b) compared to over Riyadh. The background line method used here provides likely reasonable estimations of the background X_{CO_2} , as general agreements are seen between the observed and modeled results of the integrated ffX_{CO_2} enhancements along the satellite track within the domain of interest, which can be seen in the inverse estimates of total emission scaling factors (see section 3.2). Hence, this result indicates the advantage of the background line method in deriving background X_{CO_2} for satellite observations analyzed at a spatial scale relevant to




Figure 5. Similar to Figure 4 but for the OCO-2 tracks over Cairo on (a and f) 28 February, (b and g) 18 March, (c and h) 19 May, (d and i) 15 July, and (e and j) 16 August 2015 at about 11:00 UTC.

constraining local emission sources. Neglecting the latitudinal gradients in background X_{CO_2} could lead to biases in ffX_{CO_2} , as well as in emissions derived from observations.

In respect of LA, the ffX_{CO_2} enhancements on 6 and 15 July, 7 and 16 August, and 10 and 12 October 2015 are examined with the OCO-2 data and modeled ffX_{CO_2} . As shown in Figure 6, the ffX_{CO_2} over the domain can be up to ~ 3.0 ppm and varies notably depending on meteorological conditions, as well as variations in the emissions. In comparison to the ffX_{CO_2} distributions over Riyadh and Cairo characterized by elongated plumes and rapid changing structures, the ffX_{CO_2} distributions over LA are more spread and composed of multiple plumes, mostly trapped in the basin during daytime owing to the onshore winds and local terrain. The study by Hedelius et al. (2017) demonstrated persistent differences of ~ 0.8 ppm in X_{CO_2} between two locations only 9 km apart within the LA basin induced by the steep terrain near the basin. Hence, topography plays a critical role in the distributions of ffX_{CO_2} .

Figure 7 shows the comparison of the modeled and observed X_{CO_2} over LA. Note that the OCO-2 X_{CO_2} retrievals north to the northern edge of the desert are excluded in the analysis, in order to avoid using the soundings with high warm levels owing to complex topography and high albedo of the desert area, as can be seen in Figure S6. The thresholds for latitude are $35.21^\circ N$, $35.05^\circ N$, $35.20^\circ N$, $34.90^\circ N$, $35.25^\circ N$, and $35.01^\circ N$, respectively, determined by examining the observations along with terrain height. Again, the background line has




Figure 6. Comparisons of the simulated fX_{CO_2} (color shading) and observed fX_{CO_2} enhancements (colored dots, background concentration subtracted) derived from the OCO-2 data collected at about 21:00 UTC over LA. The observation dates are labeled in each panel. The OCO-2 data are filtered by quality flag of zero (QF = 0). The vectors represent 10-m wind, with the reference vector standing for the wind speed of 5 m/s.

been added to the modeled fX_{CO_2} to present the total values; that is, the fX_{CO_2} can be seen by the increments above the background lines. Overall, the modeled fX_{CO_2} is smaller than observed, which is likely owing to the positive biases in the modeled wind speeds, and the possible underestimation of the emissions. Similar to Cairo, prominent latitudinal gradients in the background are seen for some tracks, for example, on 15 July and 12 October 2015.

3.1.2. Transport Model Uncertainty of Modeled fX_{CO_2}

The uncertainty in fX_{CO_2} related to the transport model error is an important source of uncertainty in inverse emission estimate. Here we evaluate the impact of transport model error for the above-mentioned OCO-2 tracks. Note that different error estimation methods are used for the “plume cities” and the “basin city.”

For Riyadh and Cairo, the transport model uncertainty in fX_{CO_2} has been shown with the light red shadings in Figures 4a, 4b, 5a, and 5b, estimated with the plume transformation method, as detailed in section 2.5.1. Specifically, for each track the simulated plume is perturbed 10^4 times with random errors in wind speeds and wind directions. Due to the non-Gaussian nature of the simulated plumes, as well as the nonlinearity of the sampling process, the perturbed plumes are also non-Gaussian. Hence, we used the interquartile range (difference between the 25th and the 75th percentiles) as the uncertainty spread. Note that the uncertainty spread near the domain border is not shown because of the large portion of missing data after transforming the plume, which can also be seen in Figure 2. The uncertainty in the modeled fX_{CO_2} over Cairo is up to 0.5 ppm and up to 1.2 ppm over Riyadh. Given that the same probability distribution estimations of wind speed and wind direction errors are used for both cities, the smaller transport model uncertainty in fX_{CO_2} over Cairo can be mostly attributed to its lower total emission leading to both the smaller fX_{CO_2} signals and the lower uncertainty spread. Another reason could be the relative locations of the tracks against those plumes they transect.

To evaluate the transport model errors in the modeled fX_{CO_2} over LA, an ensemble of simulations has been carried out as described in section 2.5.2. The result has been shown by the light red shadings in Figure 7.




Figure 7. Similar to Figures 4a and 4b, but for OCO-2 XCO₂ measurements over LA and the corresponding simulated results at about 21:00 UTC on (a) 6 July, (b) 15 July, (c) 7 August, (d) 16 August, (e) 10 October, and (f) 12 October 2015.

Note that this uncertainty spread is related to the random transport model error; therefore, it is independent of the systematic negative bias of the modeled ffX_{CO_2} mentioned above. As represented by the range between the minimum and maximum ffX_{CO_2} among the ensemble members, the uncertainty spread is larger where the ffX_{CO_2} is stronger, with the maximum uncertainty for each track ranging from 0.32 to 0.93 ppm. The uncertainty spread is found to be overall smaller compared to over Riyadh and comparable to over Cairo. Given that the whole-city emission is the largest for LA among these three cities, this result is likely related to the effect of local terrain trapping the $ffCO_2$ within the LA basin and making the uncertainty decrease, as well as differences in the uncertainty spreads of wind fields.

3.1.3. Integral ffX_{CO_2} Enhancement

In this work, we optimize the whole-city emissions using the latitudinal integral ffX_{CO_2} enhancements. Hence, in this section we present the modeled integral ffX_{CO_2} for each OCO-2 tracks shown in the above section and reveal how the integral ffX_{CO_2} relates to transport model errors.

Figure 8 shows the probability distributions of the simulated integral ffX_{CO_2} for the tracks over Riyadh and Cairo. The distribution for each track is derived by sampling the plumes perturbed with random transport errors 10^4 times. The distributions are mostly non-Gaussian, due to the nonlinearities in sampling transformed plumes at a specified track. To isolate the impact of transport model errors, we also show the normalized uncertainty spread, represented by the ratio between the interquartile range and the median. For Riyadh, the result shows bimodal distributions with normalized spread of 1.47 and 0.73 (Figure 8a). In comparison, the results for Cairo show narrower, less skewed, and unimodal distributions of the modeled integral ffX_{CO_2} (Figure 8b). The narrowest distribution is seen for the track on 18 March 2015 with the smallest integral ffX_{CO_2} among the five tracks. The normalized uncertainty spreads range from 0.17 to 0.38, smaller compared to Riyadh. This result is consistent with what has been shown along the tracks (Figures 4 and 5).

The large uncertainty spread over Riyadh can be attributed to the relative location of the track against the plume and structure of the plume. For example, on 27 December 2014, the large uncertainty spread is




Figure 8. Probability distributions of the modeled integral ffX_{CO_2} enhancements ($\Sigma\text{ffX}_{\text{CO}_2, \text{m}}$) for the OCO-2 tracks on 27 and 29 December 2014 over Riyadh (left) and 28 February, 28 March, 19 May, 15 July, and 16 August 2015 over Cairo (right). The total number of samples is 10^4 for each track. The distributions represent the uncertainty related to random transport model errors in wind speed and wind direction. Note that the y axis limits of the two plots are different. The numbers in the parentheses are the ratios of the interquartile range ($q_3 - q_1$) and the median (q_2).

likely due to high sensitivity of the simulated ffX_{CO_2} to transport errors, since the transaction is at the edge of the plume. For the track on 29 December, strong ffX_{CO_2} enhancements (>5.0 ppm) are found downwind of the track due to local accumulation. Those enhancements can be sampled if the wind speed is lower, corresponding to the peak centered at ~ 24 ppm in Figure 8a. Hence, the uncertainty spread also turned out to be large. Thus, it can be concluded that the transport model uncertainty in the integral ffX_{CO_2} is related to combined impacts of the magnitude of prior emissions, errors in winds, transection location relative to a plume, and the plume structure.

For LA, the modeled integral ffX_{CO_2} for each selected track is shown in Figure 9, based on the 18 ensemble members. The normalized uncertainty spread ranges between 0.06 and 0.28, smaller than the results over Riyadh and Cairo. This confirms the results shown earlier for the transport model uncertainty in ffX_{CO_2} along tracks in section 3.1.2, suggesting again that the trapped dispersion by local terrain can make the transport model uncertainty in ffX_{CO_2} smaller.

3.2. Emission Estimates and Uncertainty

In this section, we show the results of inverse emission estimates obtained by using the OCO-2 tracks shown in the above section for Riyadh, Cairo, and LA. Different sources of uncertainties are taken into consideration, including measurement errors and transport model errors. The inversions are performed to derive scaling factors of total emissions over each of the selected cities. Note that the fluxes that can be retrieved or viewed from the X_{CO_2} measurements vary from track to track, due to temporal variations in wind fields and emissions (Pillai et al., 2015). For example, if we consider an OCO-2 X_{CO_2} sounding collected at 50 km downwind of the center of an urban emission area, it takes about 5 hr for air parcels to arrive traveling with a velocity of 3 m/s. In other words, the ffX_{CO_2} obtained over or near an urban emission area can be determined by the emissions during several hours ahead of the time of the overpass (Pillai et al., 2015). Given the overpass time of the OCO-2 in early afternoon (with the equatorial crossing time at 13:30 LT), the emission estimates constrained by the OCO-2 X_{CO_2} measurements of each track could represent the emissions during morning to early afternoon on the date of the overpass. These should be taken into consideration when analyzing the results. The potential biases caused by the OCO-2 satellite observation strategy have been discussed in section 4.




Figure 9. Box plot of the modeled integral ffX_{CO_2} enhancements ($\Sigma\text{ffX}_{\text{CO}_2, \text{m}}$) for the selected OCO-2 tracks over LA on the dates (in 2015) labeled at the x axis. For each box, the central line indicates the median (q_2), and the bottom and top edges of the box indicate the 25th and 75th percentiles (q_1 and q_3), respectively. The whiskers extend to the maximum and the minimum. The numbers are the ratios of the interquartile range ($q_3 - q_1$) versus the median (q_2).

The inversions are first implemented separately for every track among the selected ones. The inverse estimates of the emission scaling factor have been shown in Table 3, as well as the measurement and transport model uncertainties of the integral ffX_{CO_2} , which are estimated with the methods described earlier in sections 2.1 and 2.5, respectively. Note that the prior scaling factor uncertainty, σ_a for LA has been set to a smaller value than those over Riyadh and Cairo, as in general there is a better knowledge of the emissions over the megacities in the United States (see also section 2.6).

The posterior scaling factors range between 0.92–0.83, 0.70–1.18, and 0.66–1.84 for the selected tracks over Riyadh, Cairo, and LA, respectively (Table 3), indicating notable temporal variations in the emission estimates from case to case. As has been explained, here the estimations represent emissions during a time period of several hours ahead of the time of overpass depending on the meteorological conditions. The posterior uncertainties, $\hat{\sigma}$, of the scaling factors for the three cities are found to range between 0.17–0.24, 0.10–0.25, and 0.11–0.16. The posterior scaling factor uncertainty has been shown overall larger for Riyadh and Cairo, compared to over LA.

With specified prior emission uncertainty for each city, the posterior uncertainty is related to both the measurement and transport model errors. The relative contribution of transport model uncertainty, σ_{model} , and measurement uncertainty, $\sigma_{\text{measurement}}$, are found to be different over the three cities. The σ_{model} for Riyadh are larger than the $\sigma_{\text{measurement}}$ by a factor of about 2 for the two tracks shown here (Table 3). By contrast, the σ_{model} for Cairo is generally smaller than the $\sigma_{\text{measurement}}$, consistent with what has been demonstrated in section 3.1.1. For LA, the relative magnitude of σ_{model} and $\sigma_{\text{measurement}}$ varies. The results show a larger σ_{model} than $\sigma_{\text{measurement}}$ for four among the six tracks analyzed here. The spread of transport model uncertainty in ffX_{CO_2} over LA is smaller compared to over Riyadh, owing to likely the trapped dispersion within the basin, which is consistent with what has been shown earlier. It can be seen that the transport model uncertainty is closely associated with the magnitude of urban emissions, relative location of plume and satellite track transection, transport model performance, and local topography. Variations of these factors lead to the temporal variability of the posterior emission uncertainty from track to track.

Although the inversion results of whole-city emissions have been demonstrated for each track separately, the sparseness of nearby OCO-2 tracks for many cities is still one of the main limitations to quantify emissions on a regular basis, as well as to track the temporal variations in emissions objectively from space. The numbers of useful tracks that have been analyzed in the above sections are 2, 5, and 6 over Riyadh, Cairo, and LA, respectively, which are chosen by examining the OCO-2 data during September 2014 to November 2015. Hence, we investigated the effect of using X_{CO_2} data from two or more tracks on the inverse emission estimates. Given that there is no correlation between the observation uncertainties for any two tracks on different days, the ffX_{CO_2} enhancement obtained from one track cannot affect the emission on another day. Therefore, in order to investigate the effect of incorporating observations on different days, we derive one single scaling factor for all the days of the useful overpasses, as shown in Table 3. Note that the resulting scaling factor for each city would represent the best estimate of emission during several hours before the general time of local overpasses.

The resulting scaling factors ($\pm 1\sigma$ uncertainty) are 0.85 ± 0.16 and 0.83 ± 0.074 for Riyadh and Cairo (see Table 3), suggesting overestimation of the prior emissions over those time periods that the observations can represent, that is, the monthly ODIAC product. While for LA, the scaling factor ($\pm 1\sigma$ uncertainty) is 1.36 ± 0.074 , indicating underestimation of the prior emission. As expected, the posterior uncertainty of scaling factor decreases substantially by using all the selected tracks over each city compared to the results with each track separately. The posterior uncertainties of the constrained whole-city emissions are reduced to about 19%, 9%, and 5% for the three cities, with the OCO-2 tracks obtained over about 1, 7, and 4 months, respectively. This result indicates the potential to obtain the emission estimate with a lower uncertainty by utilizing OCO-2 tracks over a longer time period. As explained earlier, the inversion result is closely dependent on meteorological conditions and relative location of urban plume and satellite transection. Hence, we further investigate the performance of utilizing multiple OCO-2 tracks by the OSSEs (section 3.3).

3.3. Potential of Constraining the Emissions With Multiple Tracks

In this section, we explore the effect of multiple tracks of X_{CO_2} measurements on constraining the ffCO_2 emissions. OSSEs are carried out for Riyadh, Cairo, and LA to examine the performance of OCO-2




Figure 10. Box plot of the inverse estimates of (a) whole-city emission scaling factor and (b) the posterior uncertainty for Riyadh, derived by the OSSE for different number of OCO-2 tracks (N). The number of repetition times of inversion with each specified value of N is 10^3 . For each box, the central line indicates the median, the circle represents the average, and the bottom and top edges of the box indicate the 25th and 75th percentiles (q_1 and q_3), respectively. The outliers are plotted with “x,” which are greater than $q_3 + 2 \times (q_3 - q_1)$ or less than $q_1 - 2 \times (q_3 - q_1)$. The whiskers extend to the most extreme value that is not an outlier.

truth, that is, 1, for both T01 and T02, which can be expected given that the transport model and measurement errors are assumed to be unbiased. While for LA, the median and average values are around 1.1, suggesting an overestimation by about 10% relative to the truth. This can be attributed to the overall underestimation in ffX_{CO_2} due to the positive bias in the modeled wind speed. Variability of the retrieved scaling factors for different scenes and a specified N can be seen from the height of each box, that is, the interquartile range, and the range between whiskers extending to the extreme values that are not outliers. Consistently for the three cities, the spread is found to be not obviously affected by the model errors but mainly related to the random measurement errors. The spread becomes less and less with N increasing from 2 to 20, indicating that the measurement errors tend to have less impact of biasing the emission estimate when we have more data to constrain the emission.




Figure 11. Similar to Figure 10 but for the OSSE results for Cairo.

observations from a number of tracks (N) ranging from 2 to 20. In order to take scenes with meteorological conditions into consideration, we extract the pseudo observation and pseudo modeling data using all the hourly ffX_{CO_2} modeling results in daytime for each city, rather than using only the modeling results close to the actual overpassing time. In this way we derived 217, 140, and 420 hrs of ffX_{CO_2} data in total, from which the pseudo data are generated accordingly (see section 2.7 for details). Note that the random error term used to generate pseudo observation ffX_{CO_2} represents a lower bound of measurement uncertainty, and only the times with domain-averaged surface wind speed ≥ 2 m/s are included. For Riyadh and Cairo, the pseudo data are further filtered with a criterion of the relative location of satellite transection upon ffX_{CO_2} plume (see also section 2.7).

For a specific number of tracks (N), we randomly chose N tracks from the filtered pseudo data and derive the scaling factor with these N tracks. It is assumed that these N satellite tracks were collected over a time period during which there was no bias in the prior emissions. We use a Monte Carlo method to evaluate the general performance of multiple tracks by repeating the selection and inversion process for 10^3 times. Two tests T01 and T02 are defined here. For T01, only measurement error is applied to the pseudo data, and for T02 both measurement errors and transport model errors are included.

The posterior scaling factors retrieved with different numbers of tracks (N) are shown in Figures 10a, 11a, and 12a for the three cities respectively. For Riyadh and Cairo, the median and average values are very close to the truth, that is, 1, for both T01 and T02, which can be expected given that the transport model and measurement errors are assumed to be unbiased. While for LA, the median and average values are around 1.1, suggesting an overestimation by about 10% relative to the truth. This can be attributed to the overall underestimation in ffX_{CO_2} due to the positive bias in the modeled wind speed. Variability of the retrieved scaling factors for different scenes and a specified N can be seen from the height of each box, that is, the interquartile range, and the range between whiskers extending to the extreme values that are not outliers. Consistently for the three cities, the spread is found to be not obviously affected by the model errors but mainly related to the random measurement errors. The spread becomes less and less with N increasing from 2 to 20, indicating that the measurement errors tend to have less impact of biasing the emission estimate when we have more data to constrain the emission.

Additionally, the posterior uncertainties of scaling factors are found to become smaller when the number of tracks increases (see Figures 10b, 11b, and 12b). As expected, the posterior uncertainty is overall larger for T02 in comparison to T01, suggesting the prominent contribution of the transport model errors. For each box, 75% of data are lower than the top edge, that is, the 3/4 quartile. The top whisker corresponds to about 3.275σ if the data are normally distributed, with about 99.85% coverage below it. Hence, for Riyadh, to retrieve the scaling factor with uncertainty ≤ 0.1 (i.e., 10% uncertainty of total emission) with both the measurement and transport model errors considered, 5 or 8 tracks are needed at a confidence level of 75% or 99.85% (Figure 10b). For Cairo, the numbers are found to be lower, that is, 3 or 5 at the two confidence levels (Figure 11b), owing to the smaller transport errors that have been also shown for the inversions based on real tracks. For LA, we consider a lower threshold of posterior uncertainty of 5%, given that the knowledge on prior emissions from the U.S. megacities is generally better, and the prior scaling factor uncertainty has already been set to 0.2. As shown in Figure 12b, 5 or 7 tracks are needed to obtain




Figure 12. Similar to Figure 10 but presents the OSSE results for LA.

the goal of 5% uncertainty at the two above-mentioned confidence levels. Note that in the OSSEs we have assumed no spatial correlation in the observation uncertainty. The number of tracks required to reduce the uncertainty would increase accordingly with the spatial correlation considered.

3.4. Impact of Local Biospheric CO₂ Variations on the Interpretation of Local X_{CO₂} Enhancements

Forward simulations are carried out to investigate the influence of biospheric carbon fluxes on the interpretation of local X_{CO₂} variation and the associated uncertainties. Two cases over the PRD region are analyzed using the simulation results and OCO-2 X_{CO₂} obtained on 15 January and 4 August 2015, with the results shown in Figure 13. These two cases correspond to conditions of fast atmospheric transport (i.e., high wind speeds) and weak transport (i.e., stagnant winds). The coastal circulation contrasts with the continental wind regimes, with high wind speed over the sea on 15 January and low wind speed on 4 August, opposite to the inland circulation patterns. Because the ffCO₂ emissions come from several cities located in the PRD region, the modeled ffX_{CO₂} is characterized by features of multiple overlapping plumes extending downwind from the major sources across the region.

To demonstrate the impact of local biospheric CO₂ variations on local X_{CO₂} enhancements, we derive the local X_{CO₂} enhancement (referred to as ΔX_{CO₂}) from the simulated total X_{CO₂} by subtracting the minimum value of the simulated results along a track. The constant background is valid for these simulation results, as the spatial gradient in background concentration is negligible due to a constant boundary condition used in the simulations. The observed ΔX_{CO₂} is extracted using the same method as the preceding sections. As can be seen in Figure 13, overall the simulated local enhancements including biospheric X_{CO₂} signals are larger than the simulated ffX_{CO₂}, and show a better agreement with the observed ΔX_{CO₂}. Imposed by NEE from 15 MsTMIP biospheric models, the spread of simulated ΔX_{CO₂} indicates the uncertainty in the local enhancement associated with the uncertainty in biospheric fluxes. Considering the tracks shown in Figure 13, the biogenic X_{CO₂} variability at local scales account for $\sim 32 \pm 27\%$ (1σ) and $\sim 24 \pm 18\%$ (1σ) of the latitudinally integrated local enhancement, respectively. In other words, if the biospheric signal is not separated from the local ffX_{CO₂} enhancements, the total emissions would be overestimated by about $47 \pm 37\%$ and $32 \pm 22\%$ for the two cases examined here.

For Riyadh and Cairo, the biospheric contribution is negligible compared to the local fossil fuel signals, since the local spatial gradient of NEE is much smaller than the ffCO₂ emission. Therefore, simulations for the two cities are not shown. But we note that biogenic fluxes for Cairo (Nile River delta) might be underestimated by the vegetation models. For LA, there are two reasons for not implementing a simulation like the PRD. First, the default MODIS-based GVF climatological maps in WRF show nonrealistic values compared with the real-time MODIS-based GVF maps derived by Vahmani and Ban-Weiss (2016) (Figure S7). Second, NEE is downscaled assuming a constant vegetation productivity within a NEE grid cell ($0.5^\circ \times 0.5^\circ$). However, LA has a variety of climate zones because of its proximity to the Pacific Ocean and the nearby mountain ranges, where a variety of vegetation species exist with different growth patterns (McPherson et al., 2008). Thus, it could be inappropriate to assume a constant productivity. Also, a study based on in situ flask observations in LA of ¹⁴CO₂ indicated about 25% biogenic contributions to the midday CO₂ enhancement (J. B. Miller et al., 2017), in agreement with Newman et al. (2016), but poorly simulated by vegetation model (Feng et al., 2016). More comprehensive data and method are needed to fulfill the estimation of biospheric contribution.

4. Discussion

4.1. Challenges for Other Cities

In this paper, typical cities with different local topography features are examined, which are selected following the criteria in section 2.3. The simulations over Riyadh and LA demonstrate ffX_{CO₂} enhancements




Figure 13. Simulated fX_{CO_2} over the PRD region and the 10-m wind vectors in the 1.333-km resolution domain at (a) 05:00 UTC 15 January 2015 and (b) 05:00 UTC 4 August 2015. The reference vector stands for wind speed of 5 m/s. The colored dots represent the OCO-2 data at about 05:00 UTC over this domain, filtered with quality flag of 0 (QF = 0). The background has been subtracted from the OCO-2 data. The black dots in (c) and (d) show the 1-s averaged observations in the two tracks, with the modeled fX_{CO_2} and ΔX_{CO_2} (owing to both fossil fuel and biogenic fluxes) represented by the red and blue dotted lines, respectively. The bunch of blue lines represents the results using NEE from 15 biospheric models.

overall larger than 1.0 ppm and up to about 6.0 ppm, greater than the uncertainty of retrievals over land (~1 ppm) (Eldering, O'Dell, et al., 2017). However, for Cairo the fX_{CO_2} values are mostly <3.0 ppm with some hot spots near the large emission sources. For some smaller cities, it would be even challenging to optimize their emissions from space due to limited detectability of fossil fuel imprints. The factors limiting the detectability would include (i) large cloud coverage obscuring the sensor, (ii) occasional high anthropogenic aerosol loading leading to larger measurement uncertainty, (iii) overlapping enhancements from other cities or point sources nearby, and (iv) low fX_{CO_2} emission. To obtain a bigger chance of unambiguously detecting plumes from cities, an imaging satellite with a wider swath and sufficient precision like the concept of CarbonSat (Buchwitz et al., 2013; European Space Agency, 2015) would be helpful.

Another challenge for many cities is to interpret and distinguish local X_{CO_2} variations introduced by the biospheric fluxes and fossil fuel emissions. In this work, the variations resulting from the inhomogeneity of local biospheric fluxes have been evaluated by the simulations in section 3.4. Similarly, Dayalu et al. (2017) showed equivalent magnitudes of the vegetation and the anthropogenic signal with the Vegetation, Photosynthesis, and Respiration Model (VPRM) simulations at a surface observation site in Northern China. We note that the NEE data used in this work still need further verification for regions and seasons. In addition, the downscaling of NEE data is based on the assumption of uniform local vegetation productivity, while some studies have reported impact of human interventions on urban vegetation (Hutyra et al., 2014). For example, fertilization is likely to increase both gross primary productivity and respiration of ecosystems in urban areas compared to their natural counterparts. Given the limitation in biosphere

models, many observational data have been used to extract ffCO_2 signals against the large and varying background, for example, measurements of coemitted components such as CO and NO_x (Reuter et al., 2014; Silva et al., 2013; Turnbull, Tans, et al., 2011) with efforts to determine the emission ratios accurately. Additionally, the radiocarbon content of CO_2 ($^{14}\text{CO}_2$) (Turnbull et al. 2016, 2015) has been used, although existing technology limits $^{14}\text{CO}_2$ measurement to laboratory-based analysis of individual samples at low sampling resolution. These data could provide more constraints of ffCO_2 emissions when assimilated in the inversion system jointly.

4.2. Insights From Results of the OSSEs

The performance of multiple tracks on retrieving the scaling factor of whole-city emission has been evaluated by the OSSEs, with transport model errors and measurement errors considered. The results suggest the potential of obtaining emission estimates at a lower uncertainty level and over a longer time window. At a confidence level of 99.85%, the estimated least number of tracks required to constrain the total emissions for Riyadh (<10% uncertainty), Cairo (<10%), and LA (<5%) are 8, 5, and 7, respectively. As the pseudo tracks represent different meteorological conditions, the OSSEs' results can indicate the potential of OCO-2 data in the long run. For example, we examined the number of available tracks over the three cities by counting the tracks located downwind of city and captured peaks in X_{CO_2} using OCO-2 data and surface horizontal wind data in NCEP FNL during September 2014 to November 2015, that is, 15 months in total. There are 8 (13 and 17) tracks over Riyadh (Cairo and LA) matching the criteria, corresponding to about one track per 1.75 (1.08 and 0.82) months. Therefore, the general time it takes to collect the above-mentioned number of tracks would be about 14, 5, and 6 months over the three cities. It takes the longest time for Riyadh, which can be likely owing to the larger emission than Cairo, and the less complex terrain than LA. With some other satellite missions being planned and carried out, a shorter time can be expected to retrieve emissions with uncertainty at policy-relevant level.

Positive biases in the emission scaling factors are found for LA, while the estimates are centered at the truth for Riyadh and Cairo. We suspect that the emission bias resulted from mostly the propagation of the positive wind speed bias in the atmospheric model over LA, as the wind speed bias has been considered when constructing the pseudo modeling data. In comparison, the simulations over plume cities located in flat terrain show better results of wind speed and direction; for example, wind speed errors of <1 m/s are reported for modeling results without data assimilation over Indianapolis (Deng et al., 2017) and < 0.8 m/s around Paris (Lac et al., 2013). Data assimilation systems are proven to be useful to significantly improve model performances with decreased systematic errors (<0.5 m/s) (Deng et al., 2017), which can be an effective way of reducing the wind biases for basin cities (Ware et al., 2016).

4.3. Remaining Error Components in the Inverse Emission Estimates

In addition to the uncertainties in transport model, OCO-2 measurements, and biospheric fluxes, there are several sources of errors remaining to be considered. First, the measurement errors of OCO-2 data are assumed to be noncorrelated spatially, as the correlations are yet to be characterized at high resolution at present. Additionally, the nonlinearities in the retrievals and the random components of interference errors (Connor et al., 2016) and imperfection in cloud screening especially for low clouds (Taylor et al., 2016) could introduce large errors in the retrievals. In the OSSEs, the measurement uncertainty has been evaluated using a lower bound.

For the emission inversion system, the prior emissions have been assumed to be perfectly distributed and optimized with a whole-city scaling factor. As noted by Pillai (2015), the flexibility to capture the true spatial variation of fluxes is more limited in the inversion system with one scaling factor for entire city, compared to in pixel- or parameter-wise inversions. However, the pixel-level emission uncertainties would be significantly larger than those of the whole-city emissions. Assessment of prior emissions errors in gridded field is difficult (Andres et al., 2016), which is usually done by comparing emission inventories at an aggregated spatial resolution (Hutchins et al., 2017; Oda et al., 2015). The ODIAC data have been compared to the Hestia emission product (Gurney et al., 2019), which is one of the most accurate and complete emission inventories at the scale of buildings and street segments as evaluated against in situ tower measurements (Lauvaux et al., 2016). At 1-km \times 1-km spatial scale, the result reveals a low-emission limit in ODIAC driven by saturation of the nighttime light spatial proxy, and the median difference ranging between 47% and 84%.

The largest discrepancies were found for large point sources and the on-road sector. More studies on emission comparisons would allow us to realistically constrain the emissions at sector and pixel levels.

Additionally, we note that the temporal representation of emission estimates by OCO-2 measurements can be limited given the sampling strategy. As the satellite measurements are only available in daytime clear-sky scenes, X_{CO_2} cannot be evenly sampled in time. This makes it difficult to quantify the diurnal variability in the emissions. It is suggested that X_{CO_2} retrievals must be taken into inversion modeling at the original temporal and spatial representativity (Corbin & Denning, 2006), since the clear-sky sampling biases the X_{CO_2} if they are used as daily values or averaged over a longer time period. To estimate the effect of sampling bias, we calculated the daily emission from Indianapolis by using data during daytime hours (09:00–14:00 LST) and clear-sky daytime hours, based on the hourly emission of the Hestia product (Gurney et al., 2012) for a full year and cloud cover data from a surface synoptic observation site nearby. The average daily emission for daytime hours has a +14% bias compared to the average including all hours. In addition, the bias increases to +28% when sampling only days with clear-sky conditions. Therefore, the retrieved emissions using OCO-2 measurements would be overall larger than the daily average. As another estimation, we examined the diurnal variability pattern of emissions estimated by Nassar et al. (2013). For Riyadh, Cairo, and LA, the ratios between emission at about the satellite overpassing time against the monthly average value are about 1.195, 1.127, and 1.288, corresponding to biases of about +19.5%, +12.7%, and +28.8% by only sampling in daytime compared to the monthly total emissions. More space observation missions including the Geostationary Carbon Cycle Observatory (GeoCarb) (Moore et al., 2018), OCO-3 (Eldering, Wennberg, et al., 2017), and MicroCARB (https://microcarb.cnes.fr/en/MICROCARB/GP_mission.htm) will further enhance the uniform sampling over urban areas.

5. Conclusions

In this paper, we presented the potential of using X_{CO_2} observations from OCO-2 to optimize the ffCO_2 emission from urban areas. High-resolution forward modeling of the atmospheric transport has been implemented to reproduce fine-scale structures of ffX_{CO_2} plumes, as well as to link emissions with observed X_{CO_2} . The contributions of transport model errors, measurement errors, and local variability of biospheric fluxes on the inverse estimates of whole-city emissions have been evaluated.

We used a Bayesian inversion approach to optimize the ffCO_2 emissions from three cities (Riyadh, Cairo, and LA), using the OCO-2 tracks with detectable enhancements collected between September 2014 and November 2015, namely, 15 months in total. The retrieved scaling factors ranged between 0.92–0.83, 0.70–1.18, and 0.66–1.84 for the three cities, indicating notable temporal variations in the inverse emissions from day to day. The posterior uncertainties were largest for Riyadh, mostly due to the transport model uncertainty. Prominent variability of posterior scaling factor uncertainties for the individual tracks was also due to varying meteorological conditions and locations of satellite tracks relative to city plumes. By incorporating all the selected tracks for each city, the posterior uncertainty of scaling factor was found to decrease substantially, corresponding to about 19%, 9%, and 5% uncertainty of the posterior emissions. This indicates a potential to improve current emissions estimates by utilizing OCO-2 tracks over a time period, since the frequency of nearby OCO-2 measurements is limited for each city.

We evaluated the potential of using multiple OCO-2 tracks by performing pseudo data experiments based on the high-resolution forward simulation results for the real cases analyzed above, taking the impacts of both measurement errors and transport model errors into account. For a certain number of tracks, it is assumed here that those satellite tracks were collected over a time period, during which there was no bias in the prior emissions. As revealed by the experiments, to obtain posterior uncertainty ≤ 0.1 (i.e., 10% uncertainty of total emission), five or eight (three or five) tracks are needed for Riyadh (Cairo), at a confidence level of 75% or 99.85%. For LA, we consider a lower threshold of the posterior uncertainty of 5%, and five or seven tracks are needed to achieve this goal at 75% or 99.85% confidence level.

The impact of local variability in biospheric fluxes on spatial X_{CO_2} variations is evaluated, with the uncertainty of the biospheric fluxes represented by using downscaled fluxes from the 15 biosphere models adopted in the MsTMIP intercomparison. Despite the large ffCO_2 emissions from the PRD, significant fractions, that is, $32 \pm 27\%$ (1σ) and $24 \pm 18\%$ (1σ) for the two cases shown, of the local X_{CO_2} enhancements are driven by

the local biogenic fluxes. This would lead to an overestimation of total emissions by about $47 \pm 37\%$ and $32 \pm 22\%$. For cities with biospheric fluxes of comparable magnitude but smaller fossil fuel emissions, the contribution is expected to be larger than the values shown above. Therefore, for the cities in midlatitudes and the equatorial areas with prominent local and regional biospheric fluxes, the biospheric contribution is essential for appropriate interpretation of the X_{CO_2} retrievals.

For future improvements of the quantification and monitoring of urban ffCO_2 emissions with OCO-2 data or other polar-orbit measurements, temporal and spatial correlations in prior emissions errors will likely be critical terms to be considered, which are not included in the inversions here. Given the limited satellite overpasses owing to cloud cover, retrieval issues, sampling geometry, and satellite revisiting cycle, etc., the information on prior emissions error correlations will allow us to retrieve the temporal variations and spatial structures of emissions in a more effective way, compared to using one scaling factor for the entire city. In that case, the appropriate number of tracks to constrain urban emissions will depend on the granularity, that is, the spatiotemporal resolutions of emissions in a target city, as well as the precision level required to inform policy decisions.

In addition, compared to the long-term trends in emissions that are more easily detectable, biases in retrieved emissions due to daytime-only sampling are somewhat difficult to be recovered with observations similar to OCO-2 data. Fortunately, with the continuing OCO-2 observations during its extended mission and space-based CO_2 measurement missions being deployed and planned with geostationary observations or targeting strategy over cities, the CO_2 records will be extended to potentially allow us to achieve emissions with a better temporal representativity. Our results in this work indicate a promising potential of measurements from OCO-2 or similar missions to constrain urban emissions for cities with robust ffX_{CO_2} enhancements, by using high-resolution transport modeling and the inversion approach. It can be expected that, the OCO-2 data would be more effectively used to improve the accuracy and precision of urban fossil fuel carbon fluxes, in combination with observations from other platforms to support emission reduction strategies.

Acknowledgments

This work has been funded jointly by the National Aeronautics and Space Administration (NASA) under grant NNX15AI42G, NNX15AI41G, NNX15AI40G, 80NSSC19K0092, NNX14AM76G, the National Institute for Standards and Technology (NIST) (70NANB10H245), the National Oceanic and Atmospheric Administration (NOAA) (NA13OAR4310076), and the French research program *Make Our Planet Great Again*. Emily Yang is supported by the National Science Foundation Graduate Research Fellowship (DGE 1256260). The Level 2 OCO-2 XCO_2 data used in this study are archived in permanent repository at NASA's Goddard Space Flight Center's Earth Sciences Data and Information Services Center (GES-DISC) and are also available at NASA's Jet Propulsion Laboratory (<http://co2.jpl.nasa.gov>).

References

- Ackerman, K. V., & Sundquist, E. T. (2008). Comparison of two U.S. power-plant carbon dioxide emissions data sets. *Environmental Science & Technology*, 42(15), 5688–5693. <https://doi.org/10.1021/es800221q>
- Andres, R. J., Boden, T. A., Bréon, F.-M., Ciais, P., Davis, S., Erickson, D., et al. (2012). A synthesis of carbon dioxide emissions from fossil-fuel combustion. *Biogeosciences*, 9(5), 1845–1871. <https://doi.org/10.5194/bg-9-1845-2012>
- Andres, R. J., Boden, T. A., & Higdon, D. M. (2016). Gridded uncertainty in fossil fuel carbon dioxide emission maps, a CDIAC example. *Atmospheric Chemistry and Physics*, 16(23), 14,979–14,995. <https://doi.org/10.5194/acp-16-14979-2016>
- Angevine, W. M., Eddington, L., Durkee, K., Fairall, C., Bianco, L., & Brioude, J. (2012). Meteorological model evaluation for CalNex 2010. *Monthly Weather Review*, 140(12), 3885–3906. <https://doi.org/10.1175/MWR-D-12-00042.1>
- Bougeault, P., & Lacarrere, P. (1989). Parameterization of orography-induced turbulence in a mesobeta-scale model. *Monthly Weather Review*, 117(8), 1872–1890. [https://doi.org/10.1175/1520-0493\(1989\)117<1872:POOITI>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1872:POOITI>2.0.CO;2)
- Bousquet, P. (2000). Regional changes in carbon dioxide fluxes of land and oceans since 1980. *Science*, 290(5495), 1342–1346. <https://doi.org/10.1126/science.290.5495.1342>
- Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., et al. (1999). SCIAMACHY: Mission objectives and measurement modes. *Journal of the Atmospheric Sciences*, 56(2), 127–150. [https://doi.org/10.1175/1520-0469\(1999\)056<0127:SMOAMM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<0127:SMOAMM>2.0.CO;2)
- Boxe, C. S., Worden, J. R., Bowman, K. W., Kulawik, S. S., Neu, J. L., Ford, W. C., et al. (2010). Validation of northern latitude Tropospheric Emission Spectrometer stare ozone profiles with ARC-IONS sondes during ARCTAS: Sensitivity, bias and error analysis. *Atmospheric Chemistry and Physics*, 10(20), 9901–9914. <https://doi.org/10.5194/acp-10-9901-2010>
- Bréon, F. M., Broquet, G., Puygrenier, V., Chevallier, F., Xueref-Remy, I., Ramonet, M., et al. (2015). An attempt at estimating Paris area CO_2 emissions from atmospheric concentration measurements. *Atmospheric Chemistry and Physics*, 15(4), 1707–1724. <https://doi.org/10.5194/acp-15-1707-2015>
- Briber, B., Hutyra, L., Dunn, A., Raciti, S., & Munger, J. (2013). Variations in atmospheric CO_2 mixing ratios across a Boston, MA urban to rural gradient. *Landscape*, 2(3), 304–327. <https://doi.org/10.3390/land2030304>
- Buchwitz, M., de Beek, R., Noël, S., Burrows, J. P., Bovensmann, H., Bremer, H., et al. (2005). Carbon monoxide, methane and carbon dioxide columns retrieved from SCIAMACHY by WFM-DOAS: Year 2003 initial data set. *Atmospheric Chemistry and Physics*, 5(12), 3313–3329. <https://doi.org/10.5194/acp-5-3313-2005>
- Buchwitz, M., Reuter, M., Bovensmann, H., Pillai, D., Heymann, J., Schneising, O., et al. (2013). Carbon Monitoring Satellite (CarbonSat): Assessment of atmospheric CO_2 and CH_4 retrieval errors by error parameterization. *Atmospheric Measurement Techniques*, 6(12), 3477–3500. <https://doi.org/10.5194/amt-6-3477-2013>
- Cambaliza, M. O. L., Shepson, P. B., Caulton, D. R., Stirm, B., Samarov, D., Gurney, K. R., et al. (2014). Assessment of uncertainties of an aircraft-based mass balance approach for quantifying urban greenhouse gas emissions. *Atmospheric Chemistry and Physics*, 14(17), 9029–9050. <https://doi.org/10.5194/acp-14-9029-2014>
- Ciais, P., Rayner, P., Chevallier, F., Bousquet, P., Logan, M., Peylin, P., & Ramonet, M. (2010). Atmospheric inversions for estimating CO_2 fluxes: Methods and perspectives. *Climatic Change*, 103(1–2), 69–92. <https://doi.org/10.1007/s10584-010-9909-3>

- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., et al. (2013). Carbon and other biogeochemical cycles. In Intergovernmental Panel on Climate Change (Ed.), *Climate change 2013—The Physical Science Basis*, (pp. 465–570). Cambridge: Cambridge University Press. <https://doi.org/10.1017/CBO9781107415324.015>
- Connor, B., Bösch, H., McDuffie, J., Taylor, T., Fu, D., Frankenberg, C., et al. (2016). Quantification of uncertainties in OCO-2 measurements of XCO₂: Simulations and linear error analysis. *Atmospheric Measurement Techniques*, *9*(10), 5227–5238. <https://doi.org/10.5194/amt-9-5227-2016>
- Corbin, K. D., & Denning, A. S. (2006). Using continuous data to estimate clear-sky errors in inversions of satellite CO₂ measurements. *Geophysical Research Letters*, *33*(12), L12810. <https://doi.org/10.1029/2006GL025910>
- Corbin, K. D., Denning, A. S., & Gurney, K. R. (2010). The space and time impacts on U.S. regional atmospheric CO₂ concentrations from a high resolution fossil fuel CO₂ emissions inventory. *Tellus Series B: Chemical and Physical Meteorology*, *62*(5), 506–511. <https://doi.org/10.1111/j.1600-0889.2010.00480.x>
- Crisp, D. (2008). NASA Orbiting Carbon Observatory: Measuring the column averaged carbon dioxide mole fraction from space. *Journal of Applied Remote Sensing*, *2*(1), 023508. <https://doi.org/10.1117/1.2898457>
- Crisp, D. (2015). Measuring atmospheric carbon dioxide from space with the Orbiting Carbon Observatory-2 (OCO-2). In J. J. Butler, X. Xiong, & X. Gu (Eds.), *Proceeding of SPIE 9607 Earth Observing Systems, 960702*. San Diego, California, United States. <https://doi.org/10.1117/12.2187291>
- Crisp, D., Atlas, R. M., Breon, F.-M., Brown, L. R., Burrows, J. P., Ciais, P., et al. (2004). The Orbiting Carbon Observatory (OCO) mission. *Advances in Space Research*, *34*(4), 700–709. <https://doi.org/10.1016/j.asr.2003.08.062>
- Davis, K. J., Deng, A., Lauvaux, T., Miles, N. L., Richardson, S. J., Sarmiento, D. P., et al. (2017). The Indianapolis Flux Experiment (INFLUX): A test-bed for developing urban greenhouse gas emission measurements. *Elementa: Science of the Anthropocene*, *5*(0), 21. <https://doi.org/10.1525/elementa.188>
- Dayalu, A., Munger, W., Wofsy, S. C., Wang, Y., Nehrkorn, T., Zhao, Y., et al. (2017). VPRM-CHINA: Using the vegetation, photosynthesis, and respiration model to partition contributions to CO₂ measurements in northern China during the 2005–2009 growing seasons. *Biogeosciences Discussions*, *12*, 1–29. <https://doi.org/10.5194/bg-2017-504>
- Deng, A., Lauvaux, T., Davis, K. J., Gaudet, B. J., Miles, N., Richardson, S. J., et al. (2017). Toward reduced transport errors in a high resolution urban CO₂ inversion system. *Elementa: Science of the Anthropocene*, *5*(0), 20. <https://doi.org/10.1525/elementa.133>
- Diaz-Issac, L., Lauvaux, T., Bocquet, M., & Davis, K. J. (2018). Calibration of a multi-physics ensemble for greenhouse gas atmospheric transport model uncertainty estimation. *Atmospheric Chemistry and Physics Discussions*. <https://doi.org/10.5194/acp-2018-1117>
- Duren, R. M., & Miller, C. E. (2012). Measuring the carbon emissions of megacities. *Nature Climate Change*, *2*(8), 560–562. <https://doi.org/10.1038/nclimate1629>
- Eldering, A., O'Dell, C. W., Wennberg, P. O., Crisp, D., Gunson, M. R., Viatte, C., et al. (2017). The Orbiting Carbon Observatory-2: First 18 months of science data products. *Atmospheric Measurement Techniques*, *10*(2), 549–563. <https://doi.org/10.5194/amt-10-549-2017>
- Eldering, A., Wennberg, P. O., Crisp, D., Schimel, D. S., Gunson, M. R., Chatterjee, A., et al. (2017). The Orbiting Carbon Observatory-2 early science investigations of regional carbon dioxide fluxes. *Science*, *358*(6360), eaam5745. <https://doi.org/10.1126/science.aam5745>
- European Space Agency. (2015). CarbonSat, ESA SP-1330/1 (2 volume series). Noordwijk, The Netherlands. Retrieved from http://esa-multimedia.esa.int/docs/earthobservation/sp1330-1_carbonsat.pdf
- Feng, S., Lauvaux, T., Newman, S., Rao, P., Ahmadov, R., Deng, A., et al. (2016). Los Angeles megacity: A high-resolution land-atmosphere modelling system for urban CO₂ emissions. *Atmospheric Chemistry and Physics*, *16*(14), 9019–9045. <https://doi.org/10.5194/acp-16-9019-2016>
- Fisher, J. B., Sikka, M., Huntzinger, D. N., Schwalm, C., & Liu, J. (2016). Technical note: 3-hourly temporal downscaling of monthly global terrestrial biosphere model net ecosystem exchange. *Biogeosciences*, *13*(14), 4271–4277. <https://doi.org/10.5194/bg-13-4271-2016>
- Gately, C. K., & Hutyrá, L. R. (2017). Large uncertainties in urban-scale carbon emissions. *Journal of Geophysical Research-Atmospheres*, *122*(20), 11,242–11,260. <https://doi.org/10.1002/2017JD027359>
- Gerbíg, C., Lin, J. C., Wofsy, S. C., Daube, B. C., Andrews, A. E., Stephens, B. B., ... Grainger, C. A. (2003). Toward constraining regional-scale fluxes of CO₂ with atmospheric observations over a continent: 2. Analysis of COBRA data using a receptor-oriented framework. *Journal of Geophysical Research: Atmospheres*, *108*(D24). <https://doi.org/10.1029/2003jd003770>
- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., & Eder, B. (2005). Fully coupled “online” chemistry within the WRF model. *Atmospheric Environment*, *39*(37), 6957–6975. <https://doi.org/10.1016/j.atmosenv.2005.04.027>
- Gurney, K. R., Liang, J., O’Keefe, D., Patarasuk, R., Hutchins, M., Huang, J., et al. (2019). Comparison of global downscaled versus bottom-up fossil fuel CO₂ emissions at the urban scale in four U.S. urban areas. *Journal of Geophysical Research-Atmospheres*, *124*(5), 2823–2840. <https://doi.org/10.1029/2018JD028859>
- Gurney, K. R., Razlivanov, I., Song, Y., Zhou, Y., Benes, B., & Abdul-Masih, M. (2012). Quantification of fossil fuel CO₂ emissions on the building/street scale for a large U.S. city. *Environmental Science & Technology*, *46*(21), 12,194–12,202. <https://doi.org/10.1021/es3011282>
- Gurney, K. R., Romero-Lankao, P., Seto, K. C., Hutyrá, L. R., Duren, R., Kennedy, C., et al. (2015). Climate change: Track urban emissions on a human scale. *Nature*, *525*(7568), 179–181. <https://doi.org/10.1038/525179a>
- Hakkaraínen, J., Ialongo, I., & Tamminen, J. (2016). Direct space-based observations of anthropogenic CO₂ emission areas from OCO-2. *Geophysical Research Letters*, *43*(21), 11,400–11,406. <https://doi.org/10.1002/2016GL070885>
- Hedelius, J. K., Feng, S., Roehl, C. M., Wunch, D., Hillyard, P. W., Podolske, J. R., et al. (2017). Emissions and topographic effects on column CO₂ (XCO₂) variations, with a focus on the Southern California Megacity. *Journal of Geophysical Research-Atmospheres*, *122*(13), 7200–7215. <https://doi.org/10.1002/2017JD026455>
- Hogue, S., Marland, E., Andres, R. J., Marland, G., & Woodard, D. (2016). Uncertainty in gridded CO₂ emissions estimates. *Earth’s Future*, *4*(5), 225–239. <https://doi.org/10.1002/2015EF000343>
- Houweling, S., Aben, I., Breon, F.-M., Chevallier, F., Deutscher, N., Engelen, R., et al. (2010). The importance of transport model uncertainties for the estimation of CO₂ sources and sinks using satellite measurements. *Atmospheric Chemistry and Physics*, *10*(20), 9981–9992. <https://doi.org/10.5194/acp-10-9981-2010>
- Huntzinger, D. N., Schwalm, C., Michalak, A. M., Schaefer, K., King, A. W., Wei, Y., et al. (2013). The North American carbon program multi-scale synthesis and terrestrial model Intercomparison Project—Part 1: Overview and experimental design. *Geoscientific Model Development*, *6*(6), 2121–2133. <https://doi.org/10.5194/gmd-6-2121-2013>
- Hutchins, M. G., Colby, J. D., Marland, G., & Marland, E. (2017). A comparison of five high-resolution spatially-explicit, fossil-fuel, carbon dioxide emission inventories for the United States. *Mitigation and Adaptation Strategies for Global Change*, *22*(6), 947–972. <https://doi.org/10.1007/s11027-016-9709-9>

- Hutyra, L. R., Duren, R., Gurney, K. R., Grimm, N., Kort, E. A., Larson, E., & Shrestha, G. (2014). Urbanization and the carbon cycle: Current capabilities and research outlook from the natural sciences perspective. *Earth's Future*, 2(10), 473–495. <https://doi.org/10.1002/2014EF000255>
- Idso, C. D., Idso, S. B., & Balling, R. C. Jr. (1998). The urban CO₂ dome of Phoenix, Arizona. *Physical Geography*, 19, 95–108. <https://doi.org/10.1080/02723646.1998.10642642>
- International Energy Agency (2008). World energy outlook. *International Energy Agency*. <https://doi.org/10.1787/weo-2016-en>
- IPCC. (2006). 2006 IPCC guidelines for national greenhouse gas inventories. Institute for Global Environmental Strategies. Hayama, Kanagawa, Japan. <https://doi.org/10.1007/978-1-60327-310-7>
- Janardanan, R., Maksyutov, S., Oda, T., Saito, M., Kaiser, J. W., Ganshin, A., et al. (2016). Comparing GOSAT observations of localized CO₂ enhancements by large emitters with inventory-based estimates. *Geophysical Research Letters*, 43(7), 3486–3493. <https://doi.org/10.1002/2016GL067843>
- Janjić, Z. I. (1994). The step-mountain eta coordinate model: Further developments of the convection, viscous sublayer, and turbulence closure schemes. *Monthly Weather Review*, 122(5), 927–945. [https://doi.org/10.1175/1520-0493\(1994\)122<0927:TSMECM>2.0.CO;2](https://doi.org/10.1175/1520-0493(1994)122<0927:TSMECM>2.0.CO;2)
- Janssens-Maenhout, G., Dentener, F., Van Aardenne, J., Monni, S., Pagliari, V., Orlandini, L., et al. (2012). EDGAR-HTAP: A harmonized gridded air pollution emission dataset based on national inventories. <https://doi.org/10.2788/14102>
- Keppel-Aleks, G., Wennberg, P. O., O'Dell, C. W., & Wunch, D. (2013). Towards constraints on fossil fuel emissions from total column carbon dioxide. *Atmospheric Chemistry and Physics*, 13(8), 4349–4357. <https://doi.org/10.5194/acp-13-4349-2013>
- Kort, E. A., Frankenberg, C., Miller, C. E., & Oda, T. (2012). Space-based observations of megacity carbon dioxide. *Geophysical Research Letters*, 39(17). <https://doi.org/10.1029/2012GL052738>
- Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., et al. (2013). Emissions of air pollutants and greenhouse gases over Asian regions during 2000–2008: Regional Emission inventory in ASia (REAS) version 2. *Atmospheric Chemistry and Physics*, 13(21), 11019–11058. <https://doi.org/10.5194/acp-13-11019-2013>
- Kusaka, H., & Kimura, F. (2004). Coupling a single-layer urban canopy model with a simple atmospheric model: Impact on urban heat island simulation for an idealized case. *Journal of the Meteorological Society of Japan*, 82(1), 67–80. <https://doi.org/10.2151/jmsj.82.67>
- Kuze, A., Suto, H., Nakajima, M., & Hamazaki, T. (2009). Thermal and near infrared sensor for carbon observation Fourier-transform spectrometer on the Greenhouse Gases Observing Satellite for greenhouse gases monitoring. *Applied Optics*, 48(35), 6716. <https://doi.org/10.1364/AO.48.006716>
- Lac, C., Donnelly, R. P., Masson, V., Pal, S., Riette, S., Donier, S., et al. (2013). CO₂ dispersion modelling over Paris region within the CO₂-MEGAPARIS project. *Atmospheric Chemistry and Physics*, 13(9), 4941–4961. <https://doi.org/10.5194/acp-13-4941-2013>
- Lauvaux, T., & Davis, K. J. (2014). Planetary boundary layer errors in mesoscale inversions of column-integrated CO₂ measurements. *Journal of Geophysical Research-Atmospheres*, 119(2), 490–508. <https://doi.org/10.1002/2013JD020175>
- Lauvaux, T., Miles, N. L., Deng, A., Richardson, S. J., Cambaliza, M. O., Davis, K. J., et al. (2016). High-resolution atmospheric inversion of urban CO₂ emissions during the dormant season of the Indianapolis Flux Experiment (INFLUX). *Journal of Geophysical Research-Atmospheres*, 121(10), 5213–5236. <https://doi.org/10.1002/2015JD024473>
- Lauvaux, T., Schuh, A. E., Uliasz, M., Richardson, S., Miles, N., Andrews, A. E., et al. (2012). Constraining the CO₂ budget of the corn belt: Exploring uncertainties from the assumptions in a mesoscale inverse system. *Atmospheric Chemistry and Physics*, 12(1), 337–354. <https://doi.org/10.5194/acp-12-337-2012>
- Le Quéré, C., Andrew, R., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., et al. (2018). Global carbon budget 2018. *Earth System Science Data*, 10(4), 2141–2194. <https://doi.org/10.5194/essd-10-2141-2018>
- Lin, J. C., & Gerbig, C. (2005). Accounting for the effect of transport errors on tracer inversions. *Geophysical Research Letters*, 32(1), L01802. <https://doi.org/10.1029/2004GL021127>
- Mandrake, L., O'Dell, C., Wunch, D., Wennberg, P. O., Fisher, B., Osterman, G. B., & Eldering, A. (2015). OCO-2 XCO₂ warn level, bias correction, and lite file product description. *Version 1*. Pasadena, California.
- Martilli, A., Clappier, A., & Rotach, M. W. (2002). An urban surface exchange parameterisation for mesoscale models. *Boundary-Layer Meteorology*, 104(2), 261–304. <https://doi.org/10.1023/A:1016099921195>
- McKain, K., Wofsy, S. C., Nehr Korn, T., Eluszkiewicz, J., Ehleringer, J. R., & Stephens, B. B. (2012). Assessment of ground-based atmospheric observations for verification of greenhouse gas emissions from an urban region. *Proceedings of the National Academy of Sciences*, 109(22), 8423–8428. <https://doi.org/10.1073/pnas.1116645109>
- McPherson, E. G., Simpson, J. R., Xiao, Q., & Wu, C. (2008). Los Angeles 1-million tree canopy cover assessment. *general technical report PSW-GTR-207*. Albany, CA.
- Miles, N. L., Richardson, S. J., Lauvaux, T., Davis, K. J., Balashov, N. V., Deng, A., et al. (2017). Quantification of urban atmospheric boundary layer greenhouse gas dry mole fraction enhancements in the dormant season: Results from the Indianapolis Flux Experiment (INFLUX). *Elementa: Science of the Anthropocene*, 5(0), 27. <https://doi.org/10.1525/elementa.127>
- Miller, J. B., Lehman, S., Verhulst, K. R., Yadav, V., Miller, C., Duren, R., et al. (2017). Unexpected and significant biospheric CO₂ fluxes in the Los Angeles Basin indicated by atmospheric radiocarbon (¹⁴CO₂). In *45th Global Monitoring Annual Conference*. Boulder, Colorado, USA.
- Miller, S. M., Hayek, M. N., Andrews, A. E., Fung, I., & Liu, J. (2015). Biases in atmospheric CO₂ estimates from correlated meteorology modeling errors. *Atmospheric Chemistry and Physics*, 15(5), 2903–2914. <https://doi.org/10.5194/acp-15-2903-2015>
- Moore, B., Crowell, S. M. R., Rayner, P. J., Kumer, J., O'Dell, C. W., O'Brien, D., et al. (2018). The Potential of the Geostationary Carbon Cycle Observatory (GeoCarb) to provide multi-scale constraints on the carbon cycle in the Americas. *Frontiers in Environmental Science*, 6(OCT), 1–13. <https://doi.org/10.3389/fenvs.2018.00109>
- Morino, I., Uchino, O., Inoue, M., Yoshida, Y., Yokota, T., Wennberg, P. O., et al. (2011). Preliminary validation of column-averaged volume mixing ratios of carbon dioxide and methane retrieved from GOSAT short-wavelength infrared spectra. *Atmospheric Measurement Techniques*, 4(6), 1061–1076. <https://doi.org/10.5194/amt-4-1061-2011>
- Nakanishi, M., & Niino, H. (2004). An improved Mellor–Yamada level-3 model with condensation physics: Its design and verification. *Boundary-Layer Meteorology*, 112(1), 1–31. <https://doi.org/10.1023/B:BOUN.0000020164.004146.98>
- Nassar, R., Hill, T. G., McLinden, C. A., Wunch, D., Jones, D. B. A., & Crisp, D. (2017). Quantifying CO₂ emissions from individual power plants from space. *Geophysical Research Letters*, 44(19), 10,045–10,053. <https://doi.org/10.1002/2017GL074702>
- Nassar, R., Napier-Linton, L., Gurney, K. R., Andres, R. J., Oda, T., Vogel, F. R., & Deng, F. (2013). Improving the temporal and spatial distribution of CO₂ emissions from global fossil fuel emission data sets. *Journal of Geophysical Research-Atmospheres*, 118(2), 917–933. <https://doi.org/10.1029/2012JD018196>

- Newman, S., Xu, X., Gurney, K. R., Hsu, Y. K., Li, K. F., Jiang, X., ... Yung, Y. L. (2016). Toward consistency between trends in bottom-up CO₂ emissions and top-down atmospheric measurements in the Los Angeles megacity. *Atmospheric Chemistry and Physics*, *16*(6), 3843–3863. <https://doi.org/10.5194/acp-16-3843-2016>
- Oda, T., Bun, R., Kinakh, V., Topylko, P., Haluschak, M., Marland, G., et al. (2019). Errors and uncertainties in a gridded carbon dioxide emissions inventory. *Mitigation and Adaptation Strategies for Global Change*, *24*(6), 1007–1050. <https://doi.org/10.1007/s11027-019-09877-2>
- Oda, T., Lauvaux, T., Lu, D., Rao, P., Miles, N. L., Richardson, S. J., & Gurney, K. R. (2017). On the impact of granularity of space-based urban CO₂ emissions in urban atmospheric inversions: A case study for Indianapolis, IN. *Elementa: Science of the Anthropocene*, *5*, 28. <https://doi.org/10.1525/elementa.146>
- Oda, T., & Maksyutov, S. (2011). A very high-resolution (1 km × 1 km) global fossil fuel CO₂ emission inventory derived using a point source database and satellite observations of nighttime lights. *Atmospheric Chemistry and Physics*, *11*(2), 543–556. <https://doi.org/10.5194/acp-11-543-2011>
- Oda T., & Maksyutov, S. (2015). ODIAC fossil fuel CO₂ emissions dataset (ODIAC v1.7). Center for Global Environmental Research, National Institute for Environmental Studies, Downloaded from <http://db.cger.nies.go.jp/dataset/ODIAC/>.
- Oda, T., Maksyutov, S., & Andres, R. J. (2018). The open-source data inventory for anthropogenic CO₂, version 2016 (ODIAC2016): A global monthly fossil fuel CO₂ gridded emissions data product for tracer transport simulations and surface flux inversions. *Earth System Science Data*, *10*(1), 87–107. <https://doi.org/10.5194/essd-10-87-2018>
- Oda, T., Ott, L., Topylko, P., Haluschak, M., Bun, R., Lesiv, M., et al. (2015). Uncertainty associated with fossil fuel carbon dioxide (CO₂) gridded emission datasets. In *4th International Workshop on Uncertainty in Atmospheric Emissions*, (pp. 124–129). 7–9 October 2015). Krakow, Poland: Systems Research Institute, Polish Academy of Sciences.
- O'Dell, C. W., Connor, B., Bösch, H., O'Brien, D., Frankenberg, C., Castano, R., et al. (2012). The ACOS CO₂ retrieval algorithm—Part 1: Description and validation against synthetic observations. *Atmospheric Measurement Techniques*, *5*(1), 99–121. <https://doi.org/10.5194/amt-5-99-2012>
- Pacala, S. W., Breidenich, C., Brewer, P. G., Fung, I., Gunson, M. R., Heddle, G., et al. (2010). *Verifying greenhouse gas emissions: Methods to support international climate agreements*. Washington, D.C.: National Academies Press. <https://doi.org/10.17226/12883>
- Pillai, D., Buchwitz, M., Gerbig, C., Koch, T., Reuter, M., Bovensmann, H., ... Burrows, J. P. (2016). Tracking city CO₂ emissions from space using a high-resolution inverse modelling approach: a case study for Berlin, Germany. *Atmospheric Chemistry and Physics*, *16*(15), 9591–9610. <https://doi.org/10.5194/acp-16-9591-2016>
- Raciti, S. M., Hutrya, L. R., Rao, P., & Finzi, A. C. (2012). Inconsistent definitions of “urban” result in different conclusions about the size of urban carbon and nitrogen stocks. *Ecological Applications*, *22*(3), 1015–1035. <https://doi.org/10.1890/11-1250.1>
- Reuter, M., Buchwitz, M., Hilboll, A., Richter, A., Schneising, O., Hilker, M., et al. (2014). Decreasing emissions of NO_x relative to CO₂ in East Asia inferred from satellite observations. *Nature Geoscience*, *7*(11), 792–795. <https://doi.org/10.1038/ngeo2257>
- Rotty, R. M. (1983). Distribution of and changes in industrial carbon dioxide production. *Journal of Geophysical Research*, *88*(C2), 1301. <https://doi.org/10.1029/JC088iC02p01301>
- Rypdal, K., & Winiwarter, W. (2001). Uncertainties in greenhouse gas emission inventories—Evaluation, comparability and implications. *Environmental Science & Policy*, *4*(2–3), 107–116. [https://doi.org/10.1016/S1462-9011\(00\)00113-1](https://doi.org/10.1016/S1462-9011(00)00113-1)
- Satterthwaite, D. (2008). Cities' contribution to global warming: Notes on the allocation of greenhouse gas emissions. *Environment and Urbanization*, *20*(2), 539–549. <https://doi.org/10.1177/0956247808096127>
- Schneising, O., Buchwitz, M., Burrows, J. P., Bovensmann, H., Reuter, M., Notholt, J., et al. (2008). Three years of greenhouse gas column-averaged dry air mole fractions retrieved from satellite—Part 1: Carbon dioxide. *Atmospheric Chemistry and Physics*, *8*(14), 3827–3853. <https://doi.org/10.5194/acp-8-3827-2008>
- Schneising, O., Heymann, J., Buchwitz, M., Reuter, M., Bovensmann, H., & Burrows, J. P. (2013). Anthropogenic carbon dioxide source areas observed from space: Assessment of regional enhancements and trends. *Atmospheric Chemistry and Physics*, *13*(5), 2445–2454. <https://doi.org/10.5194/acp-13-2445-2013>
- Schwandner, F. M., Gunson, M. R., Miller, C. E., Carn, S. A., Eldering, A., Krings, T., et al. (2017). Spaceborne detection of localized carbon dioxide sources. *Science*, *358*(6360). <https://doi.org/10.1126/science.aam5782>
- Silva, S. J., Arellano, A. F., & Worden, H. M. (2013). Toward anthropogenic combustion emission constraints from space-based analysis of urban CO₂/CO sensitivity. *Geophysical Research Letters*, *40*(18), 4971–4976. <https://doi.org/10.1002/grl.50954>
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., & Barker, D. M. (2008). A Description of the Advanced Research WRF Version 3. Boulder, Colorado, USA.
- Stauffer, J., Broquet, G., Bréon, F. M., Puygrenier, V., Chevallier, F., Xueref-Rémy, I., et al. (2016). The first 1-year-long estimate of the Paris region fossil fuel CO₂ emissions based on atmospheric inversion. *Atmospheric Chemistry and Physics*, *16*(22), 14703–14726. <https://doi.org/10.5194/acp-16-14703-2016>
- Super, I., Denier van der Gon, H. A. C., Visschedijk, A. J. H., Moerman, M. M., Chen, H., van der Molen, M. K., & Peters, W. (2017). Interpreting continuous in-situ observations of carbon dioxide and carbon monoxide in the urban port area of Rotterdam. *Atmospheric Pollution Research*, *8*(1), 174–187. <https://doi.org/10.1016/j.apr.2016.08.008>
- Taylor, T. E., O'Dell, C. W., Frankenberg, C., Partain, P. T., Cronk, H. Q., Savtchenko, A., et al. (2016). Orbiting Carbon Observatory-2 (OCO-2) cloud screening algorithms: Validation against collocated MODIS and CALIOP data. *Atmospheric Measurement Techniques*, *9*(3), 973–989. <https://doi.org/10.5194/amt-9-973-2016>
- Turnbull, J. C., Karion, A., Fischer, M. L., Faloona, I., Guilderson, T., Lehman, S. J., et al. (2011). Assessment of fossil fuel carbon dioxide and other anthropogenic trace gas emissions from airborne measurements over Sacramento, California in spring 2009. *Atmospheric Chemistry and Physics*, *11*(2), 705–721. <https://doi.org/10.5194/acp-11-705-2011>
- Turnbull, J. C., Keller, E. D., Norris, M. W., & Wiltshire, R. M. (2016). Independent evaluation of point source fossil fuel CO₂ emissions to better than 10%. *Proceedings of the National Academy of Sciences*, *113*(37), 10287–10291. <https://doi.org/10.1073/pnas.1602824113>
- Turnbull, J. C., Sweeney, C., Karion, A., Newberger, T., Lehman, S. J., Tans, P. P., et al. (2015). Toward quantification and source sector identification of fossil fuel CO₂ emissions from an urban area: Results from the INFLUX experiment. *Journal of Geophysical Research-Atmospheres*, *120*(1), 292–312. <https://doi.org/10.1002/2014JD022555>
- Turnbull, J. C., Tans, P. P., Lehman, S. J., Baker, D., Conway, T. J., Chung, Y. S., et al. (2011). Atmospheric observations of carbon monoxide and fossil fuel CO₂ emissions from East Asia. *Journal of Geophysical Research-Atmospheres*, *116*(D24), n/a–n/a. <https://doi.org/10.1029/2011JD016691>
- UNFCCC (2015). Adoption of the Paris Agreement. In *Report of the Conference of the Parties on its twenty-first session, Addendum* (pp. 2). Paris, France: United Nations Framework Convention on Climate Change.

- UN-Habitat. (2011). Cities and climate change: Global report on human settlements 2011. London, UK and Washington, DC, USA. ISBN: 978-92-1-132296-5
- United Nations. (1998). Kyoto protocol to the United Nations framework convention on climate change.
- Vahmani, P., & Ban-Weiss, G. A. (2016). Impact of remotely sensed albedo and vegetation fraction on simulation of urban climate in WRF-urban canopy model: A case study of the urban heat island in Los Angeles. *Journal of Geophysical Research-Atmospheres*, *121*(4), 1511–1531. <https://doi.org/10.1002/2015JD023718>
- van der Gon, H. D., Beevers, S., D'Allura, A., Finardi, S., Honoré, C., Kuenen, J., et al. (2011). Discrepancies between top-down and bottom-up emission inventories of megacities: The causes and relevance for modeling concentrations and exposure. In D. Steyn, & S. Trini Castelli (Eds.), *Air Pollution Modeling and its Application XXI. NATO Science for Peace and Security Series C: Environmental Security*, (pp. 199–204). Dordrecht: Springer. https://doi.org/10.1007/978-94-007-1359-83_4
- Verhulst, K. R., Karion, A., Kim, J., Salameh, P. K., Keeling, R. F., Newman, S., et al. (2017). Carbon dioxide and methane measurements from the Los Angeles Megacity Carbon Project—Part 1: Calibration, urban enhancements, and uncertainty estimates. *Atmospheric Chemistry and Physics*, *17*(13), 8313–8341. <https://doi.org/10.5194/acp-17-8313-2017>
- Ware, J., Kort, E. A., DeCola, P., & Duren, R. (2016). Aerosol lidar observations of atmospheric mixing in Los Angeles: Climatology and implications for greenhouse gas observations. *Journal of Geophysical Research-Atmospheres*, *121*(16), 9862–9878. <https://doi.org/10.1002/2016JD024953>
- Winker, D. (2016). CALIPSO LID L2 05kmAPro-Prov HDF file—Version3.40 [data set]. NASA Langley atmospheric science data center DAAC. https://doi.org/10.5067/caliop/calipso/lid_l2_05kmapro-prov-v3-40
- Worden, J. R., Doran, G., Kulawik, S., Eldering, A., Crisp, D., Frankenberg, C., et al. (2017). Evaluation and attribution of OCO-2 XCO₂ uncertainties. *Atmospheric Measurement Techniques*, *10*(7), 2759–2771. <https://doi.org/10.5194/amt-10-2759-2017>
- Wu, D., Lin, J. C., Oda, T., Ye, X., Lauvaux, T., Yang, E. G., & Kort, E. A. (2018). A Lagrangian approach towards extracting signals of urban CO₂ emissions from satellite observations of atmospheric column CO₂ (XCO₂): X-Stochastic Time-Inverted Lagrangian Transport model (“X-STILT v1.1”). *Geoscientific Model Development Discussion*, *11*, 4843–4871. <https://doi.org/10.5194/gmd-2018-123>
- Wunch, D., Wennberg, P. O., Osterman, G., Fisher, B., Naylor, B., Roehl, C. M., et al. (2017). Comparisons of the Orbiting Carbon Observatory-2 (OCO-2) XCO₂ measurements with TCCON. *Atmospheric Measurement Techniques*, *10*(6), 2209–2238. <https://doi.org/10.5194/amt-10-2209-2017>
- Wunch, D., Wennberg, P. O., Toon, G. C., Keppel-Aleks, G., & Yavin, Y. G. (2009). Emissions of greenhouse gases from a North American megacity. *Geophysical Research Letters*, *36*(15), L15810. <https://doi.org/10.1029/2009GL039825>