

REVIEW ARTICLE

Understanding the role of endotoxin tolerance in chronic inflammatory conditions and periodontal disease

Lena Larsson¹ | Carlos Garaicoa-Pazmino^{2,3} | Farah Asa'ad^{4,5}  | Rogerio Moraes Castilho^{6,7} 

¹Department of Periodontology Institute of Odontology, The Sahlgrenska Academy, University of Gothenburg, Göteborg, Sweden

²Department of Periodontics, University of Iowa, College of Dentistry and Dental Clinics, Iowa City, Iowa, USA

³School of Dentistry, Espíritu Santo University, Samborondon, Ecuador

⁴Department of Biomaterials, Institute of Clinical Sciences, The Sahlgrenska Academy, University of Gothenburg, Göteborg, Sweden

⁵Department of Oral Biochemistry, Institute of Odontology, The Sahlgrenska Academy, University of Gothenburg, Göteborg, Sweden

⁶Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry, Ann Arbor, Michigan, USA

⁷Laboratory of Epithelial Biology, University of Michigan School of Dentistry, Ann Arbor, Michigan, USA

Correspondence

Rogerio M. Castilho, Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry, 1011 North University Avenue, Room 2159 Commons, Ann Arbor, MI 48109-1078, USA.
Email: rcastilh@umich.edu

Funding information

EpigenCORE

Abstract

Objective: This review aims to present the current understanding of endotoxin tolerance (ET) in chronic inflammatory diseases and explores the potential connection with periodontitis.

Summary: Subsequent exposure to lipopolysaccharides (LPS) triggers ET, a phenomenon regulated by different mechanisms and pathways, including toll-like receptors (TLRs), nuclear factor kappa-light-chain enhancer of activated B-cells (NFκB), apoptosis of immune cells, epigenetics, and microRNAs (miRNAs). These mechanisms interconnect ET with chronic inflammatory diseases including periodontitis. While the direct correlation between ET and periodontal destruction has not been fully elucidated, emerging reports point towards the potential tolerization of human periodontal ligament cells (hPDLs) and gingival tissues with a significant reduction of TLR levels.

Conclusions: There is a potential link between ET and periodontal diseases. Future studies should explore the crucial role of ET in the pathogenesis of periodontal diseases, as evidence of a tolerized oral mucosa may represent an intrinsic mechanism capable of regulating the oral immune response. A clear understanding of this host immune regulatory mechanism might lead to effective and more predictable therapeutic strategies to treat chronic inflammatory diseases and periodontitis.

KEYWORDS

endotoxin shock, endotoxins, inflammation, periodontal disease, sepsis

Clinical Relevance

Scientific rationale for study: The mechanism of endotoxin tolerance (ET) could play a critical role in modulating the inflammatory response. Here we discuss the potential involvement of ET in the development and progression of periodontitis disease.

Principal findings: We present the current understanding of the mechanisms involved in the activation of ET and the emerging findings which demonstrate that low doses of LPS can induce ET, leading to reduced production of inflammatory cytokines.

Practical implications: A clear understanding of ET might lead to innovative therapeutic strategies to treat chronic inflammatory diseases and periodontitis.

1 | INTRODUCTION

The innate immune response is the first line of defence against pathogens. When the innate immune response detects pathogens through the pattern recognition receptors (PRRs), a fast, non-specific response is mounted, triggering an inflammatory response. Unregulated responses can lead to uncontrolled inflammation, which can be localized or systemic, leading to septic shock or the development of autoimmune diseases (Lopez-Collazo & del Fresno, 2013). Nonetheless, the immune system is endowed with adaptive capabilities mainly through long-term immunological memory. Similar to the adaptive properties of the immunological memory, emerging evidence points towards the ability of the innate immune system to adapt its immune response through a process known as trained immunity (Kurtz, 2005; Conrath et al., 2015). Trained immunity is defined as a higher degree of the immune response following a second pathogen challenge when compared to the first response. Such lasting re-programming of innate immune cells is mediated by epigenetic events rather than a specific transcriptional programme (Netea et al., 2019). It is important to note that erroneously trained immunity can lead to disease progression, spanning from inflammatory response to a continuous process of immunotolerance (Netea et al., 2020).

Endotoxin tolerance (ET) constitutes a functional programme from the innate immunity capable of dampening the inflammatory reaction to subsequent lipopolysaccharides (LPS) stimuli (Foster et al., 2007; del Fresno et al., 2009; Draisma et al., 2009; Novakovic et al., 2016; Vergadi et al., 2018). ET is described as a cell's reduced capacity to respond to a second endotoxin challenge (D. Liu, Cao, et al., 2019). This is the case of innate immune cells such as monocytes and macrophages, which, upon exposure to low doses of bacterial LPS, produce lower amounts of pro-inflammatory cytokines while sustaining the production of anti-inflammatory cytokine (Kopanakis et al., 2013).

Here we present the current understanding of the mechanisms involved in the activation of ET in slow, long-term inflammation (chronic inflammatory diseases) and the development of septic shock. Furthermore, we discuss the potential involvement of ET in the development and progression of periodontitis.

2 | UNDERSTANDING ENDOTOXIN TOLERANCE

2.1 | What is endotoxin tolerance?

ET constitutes a reduced response to LPS stimuli (hypo-responsiveness) upon subsequent exposures of low levels of LPS, precluding excessive innate immune response when faced with a pathogenic threat (Bauer et al., 2018) (Figure 1).

ET has been reported in many systemic inflammatory diseases, including sepsis, acting as a protective mechanism to prevent self-destruction by attenuating inflammatory pathways (tolerant response). Emerging evidence suggests that the mechanism of ET may

play a crucial function as an immune cell adaptation to continuous bacterial infection (chronic LPS exposure). Nonetheless, ET can also result in harmful effects on tissue integrity, as a hypo-responsiveness




FIGURE 1 Immune response and activation of tolerance to bacterial challenge. Graphical representation of LPS stimuli mediated by toll-like receptor (TLR) upon single stimulus or LPS re-challenge. LPS stimuli trigger NFκB signalling and enhanced inflammation (inflammatory phase). Sustained TLR activation mediated by re-challenge with LPS leads to the hypo-responsiveness of the innate immune system (immunosuppression)




FIGURE 2 Innate immune activation of naïve and endotoxin-induced tolerant cells. Graphical representation of naïve response of T-cells upon first LPS stimuli and chronic LPS exposure resulting in a tolerant response and overall hypo-responsiveness of the innate immune. The successful outcome is a result of reduced tissue damage mediated by a tolerant response

phase follows an acute pro-inflammatory response to bacterial insults as seen during sepsis (Rackov et al., 2017) (Figure 2).

The mechanism of ET was first described in the early 1900s when the injection of fever-inducing LPS was employed in the treatment of tumours and various inflammatory diseases. Interestingly, continuous administration of LPS resulted in tissue tolerization, a condition in which a gradual increase in LPS dosage was required to obtain similar therapeutic results (Bennett Jr. & Nicastrì, 1960). Similarly, rabbits receiving daily injections of bacteria demonstrated a progressive down-regulation of the immune response that lasted for a short period (Beeson & Roberts, 1947a, 1947b; Van Epps, 2006). Studies that followed have shown a significant reduction in the mortality of hosts receiving a lethal dose of endotoxin after the activation of ET (Berry & Smythe, 1965; Greisman et al., 1969; Neter, 1969; Milner, 1973). Furthermore, ET can develop after infection and/or tissue damage in which cells from the innate immune system undergo a refractory state (Lopez-Collazo & del Fresno, 2013). In vitro studies have shown that exposure to low doses of LPS results in tolerized cells capable of down-regulating inflammatory cytokines such as tumour necrosis factor- α (TNF- α), interleukin-6 (IL-6), IL-12, IL-1 β , and the mitogen-activated protein kinase kinase kinase (MAP3K). Interestingly, low doses of LPS resulted in the up-regulation of anti-inflammatory cytokines such as IL-10, transforming growth factor- β (TGF- β), and IL-1 receptor antagonist (IL-1RA). This LPS-tolerant phenotype is also characterized by inhibiting MAPK activation and impaired nuclear factor kappa-light-chain enhancer of activated B cells (NF κ B) nuclear translocation (West & Heagy, 2002; Kopanakis et al., 2013).

ET also activates high phagocytosis levels, decreases antigen presentation, reduces the expression of class II major histocompatibility complex, and impairs microbial killing (Landelle et al., 2010; Lopez-Collazo & del Fresno, 2013; Grondman et al., 2019). Patients with acute ischemic stroke present monocytes exhibiting ET and enhanced risk for pathogen colonization (Hernández-Jiménez et al., 2017). Reports suggest that around one-third of stroke patients die as a result of the potential impairment of the immune defence.

LPS from various bacteria do not have the same ability to activate toll-like receptors (TLRs) and trigger an immunological response. Thus, not all LPS have the same capacity to induce ET. When comparing LPS from *Escherichia coli* and *Porphyromonas gingivalis*, higher concentrations of *P. gingivalis* LPS are needed to induce ET (Martin et al., 2001). Macrophages also respond differently to ET signals. Cellular polarization dictates the pro-inflammatory (M1 polarization) or anti-inflammatory (M2 polarization) differentiation of macrophages and consequently its response to ET (Das et al., 2015). Macrophages of different phenotypes seem to respond differently to ET; for instance, M1 macrophages exposed to *P. gingivalis* LPS appear to be less sensitive to ET than M2 macrophages. As a matter of fact, LPS from *P. gingivalis* suppress the expression of TNF- α , IL-1 β , IL-6, and NF κ B factors from M2 macrophages (Foey & Crean, 2013; Al-Shaghдали et al., 2019). However, induction of ET using *E. coli* LPS does not interfere with TNF- α production, while IL-10 and IL-6 are suppressed in M1 but not in M2 macrophages (Al-Shaghдали et al., 2019).


2.2 | Molecular mechanisms and regulation of endotoxin tolerance

TLR-4 is a vital factor in the regulation of ET (D. Liu, Cao, et al., 2019). LPS bind to PRRs, such as TLR-4, and trigger an innate immune response, leading to the production of inflammatory cytokines (Takeda & Akira, 2015). The binding of LPS activates two signalling pathways by inducing the formation of a dimer consisting of TLR-4/MyD88 and TLR-4/TIR domain-containing adapter-inducing interferon- β (TRIF), respectively. In turn, MyD88 causes inflammatory cytokine production through the NF κ B signalling pathway, while the TRIF pathway induces interferon (IFN) type 1 via interferon regulatory factor 3 (IRF3) activation. On the other hand, NF κ B can also be activated by the TRIF/MAP3K7/inhibitor of the NF κ B kinase subunit beta (IKKB) signalling pathway (Takaesu et al., 2003; Takeda & Akira, 2015; D. Liu, Cao, et al., 2019). Other molecules such as Src homology two domain-containing inositol polyphosphate 5-phosphatase 1 (SHIP1), tumour necrosis factor- α -induced protein 3 (TNFAIP3), and interleukin-1 receptor-associated kinase-3 (IRAK3) also play a critical role in the activation of ET in different cell types (Lopez-Collazo & del Fresno, 2013; Lyroni et al., 2017). Figure 3 depicts the currently known molecular and regulatory mechanisms related to ET.

Apoptosis of immune cells has recently been suggested to be an important indicator of ET in sepsis, as the presence of bacterial endotoxin can trigger immune cell apoptosis (Rimmelé et al., 2016; van der Flier et al., 2013). This is the case of macrophages that undergo apoptosis due to the increased production of nitric oxide (NO) mediated by the presence of endotoxin in the culture media (Yamamoto et al., 1994). Furthermore, TNF- α is a cytokine suggested as an ET indicator because of its ability to induce tissue damage and cell death during sepsis. Interestingly, TNF- α has been found to activate inflammatory response as a result of cellular exposure to LPS and also be able to induce a tolerance-like state (Loosbroock & Hunter, 2014). Anti-inflammatory cytokines such as IL-10 and TGF- β can negatively regulate monocytes and activate macrophages through IRAK3 and suppressor of cytokine signalling 1 (SOCS1) signalling pathways. SOCS1 has been found to down-regulate NF κ B signalling pathway after LPS treatment (Nakagawa et al., 2002).

Recently, epigenetic modifications and microRNAs (miRNAs) have emerged as novel critical regulators of ET (Seeley & Ghosh, 2017; Vergadi et al., 2018). Epigenetic changes refer to shifts in gene expression that are not encoded in the DNA sequence, including chemical alterations of the DNA and associated proteins (i.e., histones). As a result, chromatin remodelling leads to the activation or inactivation of genes (Larsson et al., 2015). Specific epigenetic changes, such as histone modifications, are known to lead to chromatin re-organization and the control of several genes related to ET (Y. Liu, Liu, et al., 2019). One specific gene regulated by the methylation of histone H3 is IRAK3. The trimethylation of the IRAK3 promoter gene at histone H3 lysine 27 (H3K27me3) is observed in unstimulated macrophages. However, upon LPS stimuli, this silencing marker is removed, resulting in recruitment of the CCAAT/enhancer-binding protein-beta (C/EBP- β) to the promoter of IRAK-M and gene

FIGURE 3 Current molecular and regulatory mechanisms associated with ET. Graphical representation of a cellular naïve inflammatory response to LPS stimuli mediated by pattern recognition receptors (PRRs) leading the TLR-4/MyD88 and TLR-4/TRIF signalling circuitry and activation of the NF κ B pro-inflammatory signalling. Tolerant signalling prevents ubiquitination of I κ B, resulting in the inability of p65 to migrate to the nucleus



transcription (Lyroni et al., 2017). Most recently, *ex vivo* treatment with β -glucan has been shown to reverse the epigenetic immune tolerance of human macrophages while reinstating normal cytokine release levels (Novakovic et al., 2016).

Additionally, miRNAs regulate gene expression through post-transcriptional modifications. These molecules represent a group of small non-coding RNAs of about 22 bp in length that bind to the target gene's mRNA. miRNAs lead to the suppression of gene expression either by the degradation of a target mRNA or by preventing translation (Asa'ad et al., 2019). Further, the presence of differentially expressed miRNAs in periodontal diseases is known to impact the expression of TLR (Asa'ad et al., 2020) and potentially influence the activation of ET. Both non-coding RNAs (e.g., miRNAs) and long non-coding RNAs (lncRNAs) are vital regulators of ET through the TLR pathway (Vergadi et al., 2018). The LPS-inducible miRNA-155 and miRNA-146 α have been identified to be coordinately regulated via gene colocalization and transcription factor binding (Doxaki et al., 2015). While miRNA-146 α targets *IRAK1* and TNF receptor-associated factor-6 (*TRAF6*), miRNA-155 inhibits *SHIP1* and *SOCS1* (Nahid et al., 2011; X. Sun et al., 2018). Thus, these miRNAs influence critical components of TLR signalling. Additionally, several TLR-inducible lncRNAs can limit the excessive inflammatory responses by negatively regulating TLR signalling (Vergadi et al., 2018). Myocardial infarction associated transcript 2 (Mirt2) (Du et al., 2017), TNF- α

heterogenous nuclear ribonucleoprotein L (THRIL) (Li et al., 2014), metastasis associated lung adenocarcinoma transcript 1 (MALAT1) (Zhao et al., 2016), NF κ B interacting lncRNA (NKILA) (B. Liu et al., 2015), long intergenic non-coding RNA p21 (lincRNA-21) (Zhou et al., 2016), and lymphotoxin/TNF- α locus lncRNA "Sense Transcript" (SeT) (Stratigi et al., 2015) are responsible molecules to suppress the expression of pro-inflammatory mediation, such as TNF- α , and thereby contributing to ET.

2.3 | Endotoxin tolerance and sepsis

Perhaps one of the most notable examples of ET is sepsis. Sepsis is a systemic inflammatory response to an infection that can trigger a cascade of events leading to multi-organ dysfunction and organ failure. According to a recent definition, sepsis is considered a life-threatening organ dysfunction caused by a dysregulated host response to infection (Singer et al., 2016). Moreover, a septic shock is caused by gram-negative bacteria in about 70% of cases, with a mortality rate of approximately 40% (Touyz, 2013; Hernandez et al., 2019).

During sepsis, there is a deregulation of the innate immune response upon systemic bacterial infection, leading to LPS-mediated stimulation of TLR and activation of ET. The following features have been proposed as hallmarks of sepsis: (1) a decrease in lymphocyte

proliferation in response to antigen stimulation, (2) a decrease in production of IL-2 and IFN- γ by peripheral blood mononuclear cells, (3) a diminished monocyte expression of human leukocyte antigen-DR isotype (HLA-DR) and CD86, and of lymphocyte CD28, (4) an increase in cytotoxic T-lymphocyte antigen-4 (CTLA4) expression, and (5) a decrease in monocyte capacity to release pro-inflammatory cytokines in response to LPS (Monneret et al., 2008). Furthermore, ET's clinical features during sepsis comprise two main phases: an inflammatory stage, followed by a tolerance phenotype phase of the host immune cells. This results in a high risk of secondary infections, leading to an increased mortality rate (Lopez-Collazo & del Fresno, 2013). ET is often correlated with the late stage of sepsis when there is a pronounced hypo-inflammatory response due to decreased TLR-4/MyD88 signalling and M2 macrophage polarization. This explains the greater risk of secondary infections (Monneret et al., 2008).

3 | ENDOTOXIN TOLERANCE AND CHRONIC INFLAMMATION

3.1 | The inter-relationship between endotoxin tolerance and chronic inflammatory diseases

The presence and activation of ET have been reported in diseases and conditions such as trauma, surgery, pancreatitis, and cystic fibrosis. Of interest, cystic fibrosis is characterized by repeated infections leading to the inflammation of the lungs and progressive tissue destruction, which result in a compensatory anti-inflammatory response syndrome (CARS), a condition often referred to as the clinical manifestation of ET (Biswas & Lopez-Collazo, 2009; Lopez-Collazo & del Fresno, 2013). CARS is an autoimmune suppression state observed during significant insults to the human body, such as extensive burns, large tissue injuries, and sepsis (Ward et al., 2008). Like CARS, ET is defined as a hyporesponsiveness disease presenting a reduced response of myeloid cells to inflammatory stimuli, particularly those initiated by bacterial LPS (Vergadi et al., 2018). Compromised epithelial barrier has a severe impact on the maintenance of chronic inflammatory diseases. Loss of epithelial protection often results in endotoxemia, that is, the accumulation of LPS in the blood, leading to chronic stimulation of myeloid cells and potential association with ET (Carron et al., 2019). One example of endotoxemia is found during chronic inflammation of end-stage renal disease (ESRD), where the loss of intestinal epithelium integrity and the consequent bacterial presence in the bloodstream result in the binding of LPS to LPS-binding proteins (LBPs), activation of MyD88 and TRIF, and the production of pro-inflammatory cytokines, leading to chronic inflammation (Carron et al., 2019). The connection between endotoxemia and the development of ET has been demonstrated in patients with acute respiratory distress syndrome (ARDS). The development and severity of ARDS are associated with the endotoxemia levels and the development of tolerance as judged by the reduced ability of monocytes to release cytokines (Buttenschoen et al., 2008).

The role of circulating LPS and their influence in ET and chronic systemic diseases have been discussed (van Lier et al., 2019). From

experimental human models of endotoxemia, intravenous administration of LPS results in neutrophilia, mono- and lymphocytopenia after 2–4 h, associated with a marked up-regulation of reactive oxygen species (ROS) (Pillay et al., 2010; Kiers et al., 2017). Plasma cytokine levels (e.g., TNF- α , IL-1 β , IL-6, IL-10) have a distinct, dose-dependent, and highly reproducible time course that returns to baseline values 6–8 h after the LPS challenge (Kiers et al., 2017). TNF- α leads the LPS-induced inflammatory response by peaking at 90–120 min followed by IL-1 β , IL-6, and IL-10 and reaching maximum plasma concentrations at approximately 3 h (van Lier et al., 2019). This experimental model has been used to induce short-lived, well-tolerated, and controlled inflammatory responses, like those observed in sepsis, to explore the role of endotoxemia-induced, organ-specific effects in the immune system, serum, cardiovascular system, kidneys, lungs, respiratory muscle, and the gastrointestinal tract.

Conversely, sub-clinical serum LPS may be a risk factor for conditions characterized by a non-resolving inflammation as observed in atherosclerosis (Geng et al., 2016). Interestingly, epithelial cells of the nasal cavity and airway present a tolerant phenotype in healthy individuals. A second signal stimulating the Fc-gamma receptor III (Fc γ RIII) and LPS stimulation of TLR-4 was needed in these cells to break this immune tolerance (Golebski et al., 2019).

4 | ENDOTOXIN TOLERANCE AND PERIODONTAL DISEASE

4.1 | Periodontal disease is a form of chronic inflammatory disease

Periodontal disease development involves an intricate relationship between disease susceptibility, bacterial colonization, and host response. Although oral hygiene and the microbial community are critical factors in developing periodontitis, the imbalance in the host susceptibility is the final element for the disease. Recent findings suggest that the host immune response may play an integral role in chronic periodontitis susceptibility (Hajishengallis, 2015). Whole blood cell cultures obtained from periodontitis-affected and non-periodontitis patients were stimulated with *P. gingivalis* LPS, revealing increased IFN- γ levels only in the former (Nogueira-Filho et al., 2014). These data support the notion that periodontitis alters the systemic immune cell response. LPS from other periodontal pathogens are also known to show similar effects. The low dose of bloodstream LPS (endotoxemia), similar to that found in periodontitis patients, appears to be sufficient to prime monocytes capable of affecting immune and inflammatory cells (Nakamura et al., 2004). This study involved pre-treating blood samples obtained from seven systemically and periodontally healthy individuals with or without *Aggregatibacter actinomycetemcomitans* LPS, followed by further LPS stimulation. The pre-treatment with *A. actinomycetemcomitans* LPS significantly enhanced the production of IL-1 β and IL-6 from whole blood, suggesting the role of trained innate immunity (Nakamura et al., 2004).

4.2 | The link between endotoxin tolerance and periodontal disease

For years, it was suspected that LPS led to the exaggerated innate immune response, resulting in tissue destruction observed with periodontitis (Bainbridge et al., 2002). Currently, little is known on the mechanism by which ET regulates periodontal homeostasis by providing sufficient tolerance to the presence of a healthy and balanced oral microbiota (Figure 4). However, the unbalance of the protective effects of ET poses a question on the potential mechanisms by which ET becomes dysfunctional. First, the presence of a dysbiotic oral microbiota may fail to activate the mechanism of ET, leading to the exacerbation of the inflammatory response and consequently tissue destruction. Conversely, excessive activation of ET mediated by a dysbiotic microbiota or the loss of epithelial barrier and consequently the development of endotoxemia may either prevent the recruitment of inflammatory cells or reduce the ability of monocytes to release cytokines, resulting in uncontrolled bacteria-induced tissue destruction (Figure 4). It remains unknown whether the progression from a healthy periodontium to periodontitis is driven by an overly robust or insufficient immune response. Figure 4 illustrates our view of the potential role of ET in the pathogenesis of periodontal disease.

Despite the fact that the clearance of periodontal pathogens through an inflammatory response relies heavily on the recognition of LPS, it has been shown that low doses of LPS can induce ET, leading to reduced production of inflammatory cytokines (Rios et al., 2016; Seeley & Ghosh, 2017). In periodontitis, biofilm can disrupt the gingival tissues, resulting in a compromised epithelial barrier, and periodontal pockets followed by colonization with gram-negative bacteria (Touyz, 2013). Subsequently, this can further lead to endothelial damage and a leak of bacterial products into the bloodstream (endotoxemia), which supports the link between periodontitis and septic shock syndrome (Touyz, 2013).

As noted in vitro, isolated gingival fibroblasts have demonstrated a marked difference in LPS tolerance between healthy and inflamed gingiva (Kang et al., 2016). The induction of pro-inflammatory cytokines (i.e., IL-6, IL-8, IL-1 β , and TNF- α) by *P. gingivalis* LPS occurred at significantly lower levels among fibroblasts isolated from inflamed gingiva affected with periodontitis compared with fibroblasts obtained from healthy patients, and thus suggesting increased tolerance in disease. Conversely, monocytes exposed to either *P. gingivalis* or *E. coli* LPS are down-regulated for TLR-2 and TLR-4 at the mRNA and protein levels and reduce IL-1 β mRNA and TNF- α secretion (Muthukuru et al., 2005).

Monocyte cultures can be tolerized upon exposure to LPS from *P. gingivalis*. The conditioned media from tolerized monocytes have a direct negative impact on the migration and apoptotic process of neutrophils while simultaneously increasing the concentration levels of ROS (Zhu et al., 2016). As such, the role of ET becomes more crucial in periodontal diseases, as neutrophil apoptosis has been suggested to influence the progression of periodontitis. Using an ET model, the exposure of human periodontal ligament cells (hPDLs) to *P. gingivalis* LPS showed that they could not develop ET (Blufstein et al., 2018). In contrast, a significant reduction of TLR-4 was observed in pre-treated cells compared to non-pre-treated cells. The pre-treatment with *P. gingivalis* LPS did not decrease IL-6, IL-8, and monocyte chemoattractant protein-1 (MCP-1), thus indicating that these cells may play an important role in sustaining the inflammation in periodontal disease. Another study using a similar model reported that hPDLs differ in pro-inflammatory cytokines and TLRs after the first and second dose of the same LPS (Wu et al., 2015). The authors hypothesized that since the oral cavity is continuously exposed to a large amount and a diversity of periodontal pathogens, hPDLs are exposed to a “negative regulation of immune response” due to this repeated exposure to pathogens. These findings indicate ET's induction by LPS, leading to down-regulation of TLRs and the production of inflammatory cytokines.




FIGURE 4 Graphical representation of ET in periodontal health and disease. ET plays a crucial role in maintaining the periodontium tissue integrity (healthy periodontium). Upon dysbiosis, LPS fail to trigger ET, resulting in the increased inflammatory response and tissue destruction. Alternatively, dysbiosis led to excessive ET function and reduced inflammatory response along with enhanced bacteria-induced tissue destruction

Moreover, periodontal disease is characterized by rampant tissue destruction mediated by matrix metalloproteinases (MMPs). A reduction in MMPs would be expected in LPS-tolerized individuals; however, this is not the case, as one study identified the secretion of MMP-9 was not affected by LPS tolerance (Muthukuru & Cutler, 2015). Conversely, tissue inhibitor of metalloproteinase-1 (TIMP-1), the main antagonist of MMP-9, is greatly up-regulated in monocytes that have been induced to a tolerant state.

Compared to LPS from other gram-negative bacteria such as *E. coli*, LPS from *P. gingivalis* induce much lower levels of pro-inflammatory cytokines, including IL-1 β , IL-6, and TNF- α (Martin et al., 2001). *P. gingivalis* LPS are also known to trigger a state of imbalanced immune tolerance through continuous secretion of IL-8 and a decreased production of TNF- α (Zaric et al., 2010). These events can lead to enhanced migration of neutrophils to the site of infection and reduced apoptosis. The low toxicity of *P. gingivalis* LPS compared to those of other gram-negative bacteria and the pattern of pro-inflammatory cytokines induced after primary and secondary stimulation play a critical role in the chronic inflammatory state seen in periodontal disease.

In addition to the down-regulation of host response, modifications of LPS signalling by *P. gingivalis* may affect the immune response. Patients with chronic forms of periodontitis are well known to have increased TLR-2 and TLR-4 positive cells and IL-1 β levels within the inflamed gingival tissues. Interestingly, the numbers of TLR-2 cells increase linearly with inflammation. In a study using peripheral blood monocytes, the first stimulus of LPS resulted in the up-regulation of TLR-2 and TLR-4 mRNA. In contrast, a second stimulus down-regulated both receptors at the mRNA and protein levels.

Along with the TLR receptor, mRNA levels of IL-1 β were also found to be reduced and TNF- α secretion levels were decreased by 10-folds. However, cytokines such as IL-6, IL-8, and IL-10 were also negatively impacted by the re-stimulation at a lesser degree than TNF- α (Muthukuru et al., 2005). Additionally, TLR-4 response to LPS varies significantly depending on the phosphorylation patterns of the lipid A component. As a clear example, 1'-phosphorylated lipid A demonstrated more pronounced TLR-4 activation than 4'-phosphorylated lipid A (Coats et al., 2009).

The exposure of THP-1 cells to *P. gingivalis* LPS stimulated NF κ B activation and TNF- α release upon re-stimulation with the same LPS (Hajishengallis et al., 2002). Conversely, NF κ B-dependent transcription inhibition was observed in pre-treated cell lines with *E. coli* LPS and re-stimulated with *P. gingivalis* LPS. Although *P. gingivalis* LPS-treated cells could not become tolerant to subsequent exposure to *E. coli* LPS, *P. gingivalis* LPS inhibited the *E. coli* LPS-induced TNF- α and IL-6 release when added simultaneously into the same cell line. The re-stimulation of THP-1 monocyte cells using *P. gingivalis* and *E. coli* LPS resulted in decreased TNF- α and IL-1 β but increased IL-10 levels (Y. Sun et al., 2014). Cells treated with LPS following re-treatment with *P. gingivalis* LPS showed down-regulation of TLR-2. In contrast, when re-stimulated with *E. coli* LPS, a decrease in TLR-4 was observed and associated with the irregular expression of IRAK3 and SOCS1 genes. In a similar study, the pre-treatment of THP-1 cells with *E. coli* LPS resulted in a decrease of TLR-4, TNF- α , IL-1 β , and IL-6 levels without

changes in the expression of CD14 (Martin et al., 2001). Nonetheless, cells pre-treated with a similar concentration of *P. gingivalis* LPS displayed an increase in CD14 and TLR-2 associated with a decrease in IL-1 β . It is important to point out that *P. gingivalis* LPS suppressed *P. gingivalis* FimA protein-induced NF κ B-dependent transcription in a 3E10/huTLR4 cell line not expressing TLR-2 (Hajishengallis et al., 2002). Thus, it suggests that *P. gingivalis* and *E. coli* LPS do not compete for common signalling intermediates and that *P. gingivalis* may target components in the TLR-4 receptor complex.

Observations from ligature-induced periodontitis models have shown that endotoxin-tolerant animals have significantly reduced immune system protein mannose-binding lectin levels (Nowotny & Sanavi, 1983). While the direct correlation between ET and diminished periodontal destruction has not been extensively investigated, limited reports have explored the immunological response after exposure to bacteria. TLR-2 and TLR-4 positive cells were present in more significant numbers among gingival tissue biopsies from periodontitis patients (Muthukuru et al., 2005).

The question remains open as to how ET can drive periodontal disease progression while reducing the host's pro-inflammatory response. The study from the Fujihashi's lab sheds light on this conundrum by demonstrating that the presence of CD4+ T-cells in chronic periodontitis shifts from a pro-inflammatory state characterized by the production of IL-6 and TNF- α to an IL-10-producing phenotype in an experimental periodontitis model (Kobayashi et al., 2011).

5 | CONCLUSIONS

There is still a lack of knowledge on the regulatory machinery involved in the activation of ET and a clear understanding of the role of tolerized cells in disease progression or tissue defence against bacterial insults. Future research should elucidate the crucial role of ET in the pathogenesis of periodontal diseases and the intricate process that balances the host-bacteria interactions during health. It is also important to determine the potential imbalance of the ET machinery in patients presenting increased susceptibility to periodontitis and unresponsive to traditional periodontal therapy. A clear understanding of this host immune regulatory mechanism can lead to the development of effective and more predictable therapeutic strategies to treat chronic inflammatory diseases and periodontitis.

ACKNOWLEDGEMENTS

This work was supported by EpigenCORE (a consortium of researchers tackling the effects of epigenetic modifications in disease), a non-profit initiative resulting from a collaboration between researchers from the University of Iowa, Iowa City, IA, the University of Michigan, Ann Arbor, MI, and the University of Gothenburg, Göteborg, Sweden.

CONFLICT OF INTEREST

The authors do not have any financial interests, either directly or indirectly, in the products or information identified in the paper. None of the authors has any competing interests in the manuscript.

AUTHOR CONTRIBUTIONS

Rogério M. Castilho and Lena Larsson conceived the main idea for this review. Lena Larsson, Carlos Garaicoa-Pazmino, Farah Asa'ad, and Rogério M. Castilho contributed to the conception, design, writing, and critical review of the manuscript.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analysed in this study.

ORCID

Farah Asa'ad  <https://orcid.org/0000-0001-5186-2645>

Rogério Moraes Castilho  <https://orcid.org/0000-0001-5358-612X>

REFERENCES

- Al-Shaghda, K., Durante, B., Hayward, C., Beal, J., & Foey, A. (2019). Macrophage subsets exhibit distinct *E. coli*-LPS tolerable cytokines associated with the negative regulators, Irak-M and Tollip. *PLoS One*, 14(5), e0214681. <https://doi.org/10.1371/journal.pone.0214681>
- Asa'ad, F., Garaicoa-Pazmino, C., Dahlin, C., & Larsson, L. (2020). Expression of MicroRNAs in periodontal and Peri-implant diseases: A systematic review and meta-analysis. *International Journal of Molecular Sciences*, 21(11), 4147–4160. <https://doi.org/10.3390/ijms21114147>
- Asa'ad, F., Monje, A., & Larsson, L. (2019). Role of epigenetics in alveolar bone resorption and regeneration around periodontal and peri-implant tissues. *European Journal of Oral Sciences*, 127(6), 477–493. <https://doi.org/10.1111/eos.12657>
- Bainbridge, B. W., Coats, S. R., & Darveau, R. P. (2002). *Porphyromonas gingivalis* lipopolysaccharide displays functionally diverse interactions with the innate host defense system. *Annals of Periodontology*, 7(1), 29–37. <https://doi.org/10.1902/annals.2002.7.1.29>
- Bauer, M., Weis, S., Netea, M. G., & Wetzker, R. (2018). Remembering pathogen dose: Long-term adaptation in innate immunity. *Trends in Immunology*, 39(6), 438–445. <https://doi.org/10.1016/j.it.2018.04.001>
- Beeson, P. B., & Roberts, E. (1947a). Tolerance to bacterial pyrogens: I. factors influencing its development. *The Journal of Experimental Medicine*, 86(1), 29–38. <https://doi.org/10.1084/jem.86.1.29>
- Beeson, P. B., & Roberts, E. (1947b). Tolerance to bacterial pyrogens: II. Role of the Reticulo-endothelial system. *The Journal of Experimental Medicine*, 86(1), 39–44. <https://doi.org/10.1084/jem.86.1.39>
- Bennett, I. L., Jr., & Nicastrì, A. (1960). Fever as a mechanism of resistance. *Bacteriological Reviews*, 24(1), 16–34.
- Berry, L. J., & Smythe, D. S. (1965). Some metabolic aspects of tolerance to bacterial endotoxin. *Journal of Bacteriology*, 90(4), 970–977. <https://doi.org/10.1128/JB.90.4.970-977.1965>
- Biswas, S. K., & Lopez-Collazo, E. (2009). Endotoxin tolerance: New mechanisms, molecules and clinical significance. *Trends in Immunology*, 30(10), 475–487. <https://doi.org/10.1016/j.it.2009.07.009>
- Blufstein, A., Behm, C., Nguyen, P. Q., Rausch-Fan, X., & Andrukhov, O. (2018). Human periodontal ligament cells exhibit no endotoxin tolerance upon stimulation with *Porphyromonas gingivalis* lipopolysaccharide. *Journal of Periodontal Research*, 53(4), 589–597. <https://doi.org/10.1111/jre.12549>
- Buttenschoen, K., Kornmann, M., Berger, D., Leder, G., Beger, H. G., & Vasilescu, C. (2008). Endotoxemia and endotoxin tolerance in patients with ARDS. *Langenbeck's Archives of Surgery*, 393(4), 473–478. <https://doi.org/10.1007/s00423-008-0317-3>
- Carron, C., Pais de Barros, J. P., Gaiffe, E., Deckert, V., Adda-Rezig, H., Roubiou, C., Laheurte, C., Masson, D., Simula-Faivre, D., Louvat, P., Moulin, B., Frimat, L., Rieu, P., Mousson, C., Durrbach, A., Heng, A. E., Saas, P., Ducloux, D., Lagrost, L., & Bamouid, J. (2019). End-stage renal disease-associated gut bacterial translocation: Evolution and impact on chronic inflammation and acute rejection after renal transplantation. *Frontiers in Immunology*, 10, 1630. <https://doi.org/10.3389/fimmu.2019.01630>
- Coats, S. R., To, T. T., Jain, S., Braham, P. H., & Darveau, R. P. (2009). *Porphyromonas gingivalis* resistance to polymyxin B is determined by the lipid a 4'-phosphatase, PGN_0524. *International Journal of Oral Science*, 1(3), 126–135. <https://doi.org/10.4248/IJOS.09062>
- Conrath, U., Beckers, G. J., Langenbach, C. J., & Jaskiewicz, M. R. (2015). Priming for enhanced defense. *Annual Review of Phytopathology*, 53, 97–119. <https://doi.org/10.1146/annurev-phyto-080614-120132>
- Das, A., Sinha, M., Datta, S., Abas, M., Chaffee, S., Sen, C. K., & Roy, S. (2015). Monocyte and macrophage plasticity in tissue repair and regeneration. *The American Journal of Pathology*, 185(10), 2596–2606. <https://doi.org/10.1016/j.ajpath.2015.06.001>
- del Fresno, C., García-Río, F., Gómez-Piña, V., Soares-Schanoski, A., Fernández-Ruiz, I., Jurado, T., Kajiji, T., Shu, C., Marín, E., Gutierrez del Arroyo, A., Prados, C., Arnalich, F., Fuentes-Prior, P., Biswas, S. K., & López-Collazo, E. (2009). Potent phagocytic activity with impaired antigen presentation identifying lipopolysaccharide-tolerant human monocytes: Demonstration in isolated monocytes from cystic fibrosis patients. *Journal of Immunology*, 182(10), 6494–6507. <https://doi.org/10.4049/jimmunol.0803350>
- Doxaki, C., Kampranis, S. C., Eliopoulos, A. G., Spilianakis, C., & Tsatsanis, C. (2015). Coordinated regulation of miR-155 and miR-146a genes during induction of endotoxin tolerance in macrophages. *Journal of Immunology*, 195(12), 5750–5761. <https://doi.org/10.4049/jimmunol.1500615>
- Draisma, A., Pickkers, P., Bouw, M. P., & van der Hoeven, J. G. (2009). Development of endotoxin tolerance in humans in vivo. *Critical Care Medicine*, 37(4), 1261–1267. <https://doi.org/10.1097/CCM.0b013e31819c3c67>
- Du, M., Yuan, L., Tan, X., Huang, D., Wang, X., Zheng, Z., Mao, X., Li, X., Yang, L., Huang, K., Zhang, F., Wang, Y., Luo, X., Huang, D., & Huang, K. (2017). The LPS-inducible lncRNA Mirt2 is a negative regulator of inflammation. *Nature Communications*, 8(1), 2049. <https://doi.org/10.1038/s41467-017-02229-1>
- Foey, A. D., & Crean, S. (2013). Macrophage subset sensitivity to endotoxin toleration by *Porphyromonas gingivalis*. *PLoS One*, 8(7), e67955. <https://doi.org/10.1371/journal.pone.0067955>
- Foster, S. L., Hargreaves, D. C., & Medzhitov, R. (2007). Gene-specific control of inflammation by TLR-induced chromatin modifications. *Nature*, 447(7147), 972–978. <https://doi.org/10.1038/nature05836>
- Geng, S., Chen, K., Yuan, R., Peng, L., Maitra, U., Diao, N., Chen, C., Zhang, Y., Hu, Y., Qi, C. F., Pierce, S., Ling, W., Xiong, H., & Li, L. (2016). The persistence of low-grade inflammatory monocytes contributes to aggravated atherosclerosis. *Nature Communications*, 7, 13436. <https://doi.org/10.1038/ncomms13436>
- Golebski, K., Hoepel, W., van Egmond, D., de Groot, E. J., Amatngalim, G. D., Beekman, J. M., Fokkens, W. J., van Drunen, C. M., & den Dunnen, J. (2019). FcγRIII stimulation breaks the tolerance of human nasal epithelial cells to bacteria through cross-talk with TLR4. *Mucosal Immunology*, 12(2), 425–433. <https://doi.org/10.1038/s41385-018-0129-x>
- Greisman, S. E., Young, E. J., & Carozza, F. A., Jr. (1969). Mechanisms of endotoxin tolerance. V. Specificity of the early and late phases of pyrogenic tolerance. *Journal of Immunology*, 103(6), 1223–1236.
- Grondman, I., Arts, R. J. W., Koch, R. M., Leijte, G. P., Gerretsen, J., Bruse, N., Kempkes, R. W. M., ter Horst, R., Kox, M., Pickkers, P., Netea, M. G., & Gresnigt, M. S. (2019). Frontline science: Endotoxin-induced immunotolerance is associated with loss of monocyte metabolic plasticity and reduction of oxidative burst. *Journal of Leukocyte Biology*, 106(1), 11–25. <https://doi.org/10.1002/JLB.5HI0119-018R>
- Hajishengallis, G. (2015). Periodontitis: From microbial immune subversion to systemic inflammation. *Nature Reviews. Immunology*, 15(1), 30–44. <https://doi.org/10.1038/nri3785>

- Hajshengallis, G., Martin, M., Schifferle, R. E., & Genco, R. J. (2002). Counteracting interactions between lipopolysaccharide molecules with differential activation of toll-like receptors. *Infection and Immunity*, 70(12), 6658–6664. <https://doi.org/10.1128/IAI.70.12.6658-6664.2002>
- Hernandez, A., Patil, N. K., Stothers, C. L., Luan, L., McBride, M. A., Owen, A. M., Burelbach, K. R., Williams, D. L., Sherwood, E. R., & Bohannon, J. K. (2019). Immunobiology and application of toll-like receptor 4 agonists to augment host resistance to infection. *Pharmacological Research*, 150, 104502. <https://doi.org/10.1016/j.phrs.2019.104502>
- Hernández-Jiménez, E., Gutierrez-Fernández, M., Cubillos-Zapata, C., Otero-Ortega, L., Rodríguez-Frutos, B., Toledano, V., Martínez-Sánchez, P., Fuentes, B., Varela-Serrano, A., Avendaño-Ortiz, J., Blázquez, A., Mangas-Guijarro, M. Á., Díez-Tejedor, E., & López-Collazo, E. (2017). Circulating monocytes exhibit an endotoxin tolerance status after acute ischemic stroke: Mitochondrial DNA as a putative explanation for Poststroke infections. *Journal of Immunology*, 198(5), 2038–2046. <https://doi.org/10.4049/jimmunol.1601594>
- Kang, W., Hu, Z., & Ge, S. (2016). Healthy and inflamed gingival fibroblasts differ in their inflammatory response to *Porphyromonas gingivalis* lipopolysaccharide. *Inflammation*, 39(5), 1842–1852. <https://doi.org/10.1007/s10753-016-0421-4>
- Kiers, D., Koch, R. M., Hamers, L., Gerretsen, J., Thijs, E. J., van Ede, L., Riksen, N. P., Kox, M., & Pickkers, P. (2017). Characterization of a model of systemic inflammation in humans in vivo elicited by continuous infusion of endotoxin. *Scientific Reports*, 7, 40149. <https://doi.org/10.1038/srep40149>
- Kobayashi, R., Kono, T., Bolerjack, B. A., Fukuyama, Y., Gilbert, R. S., Fujihashi, K., Ruby, J., Kataoka, K., Wada, M., Yamamoto, M., & Fujihashi, K. (2011). Induction of IL-10-producing CD4+ T-cells in chronic periodontitis. *Journal of Dental Research*, 90(5), 653–658. <https://doi.org/10.1177/0022034510397838>
- Kopanakis, K., Tzepi, I. M., Pistiki, A., Carrer, D. P., Netea, M. G., Georgitsi, M., Lymperi, M., Droggiti, D. I., Liakakos, T., Machairas, A., & Giamarellos-Bourboulis, E. J. (2013). Pre-treatment with low-dose endotoxin prolongs survival from experimental lethal endotoxin shock: Benefit for lethal peritonitis by *Escherichia coli*. *Cytokine*, 62(3), 382–388. <https://doi.org/10.1016/j.cyto.2013.03.028>
- Kurtz, J. (2005). Specific memory within innate immune systems. *Trends in Immunology*, 26(4), 186–192. <https://doi.org/10.1016/j.it.2005.02.001>
- Landelle, C., Lepape, A., Voirin, N., Tognet, E., Venet, F., Bohé, J., Vanhems, P., & Monneret, G. (2010). Low monocyte human leukocyte antigen-DR is independently associated with nosocomial infections after septic shock. *Intensive Care Medicine*, 36(11), 1859–1866. <https://doi.org/10.1007/s00134-010-1962-x>
- Larsson, L., Castilho, R. M., & Giannobile, W. V. (2015). Epigenetics and its role in periodontal diseases: A state-of-the-art review. *Journal of Periodontology*, 86(4), 556–568. <https://doi.org/10.1902/jop.2014.140559>
- Li, Z., Chao, T. C., Chang, K. Y., Lin, N., Patil, V. S., Shimizu, C., Head, S. R., Burns, J. C., & Rana, T. M. (2014). The long noncoding RNA THRIL regulates TNF α expression through its interaction with hnRNPL. *Proceedings of the National Academy of Sciences of the United States of America*, 111(3), 1002–1007. <https://doi.org/10.1073/pnas.1313768111>
- Liu, B., Sun, L., Liu, Q., Gong, C., Yao, Y., Lv, X., Lin, L., Yao, H., Su, F., Li, D., Zeng, M., & Song, E. (2015). A cytoplasmic NF- κ B interacting long noncoding RNA blocks I κ B phosphorylation and suppresses breast cancer metastasis. *Cancer Cell*, 27(3), 370–381. <https://doi.org/10.1016/j.ccr.2015.02.004>
- Liu, D., Cao, S., Zhou, Y., & Xiong, Y. (2019). Recent advances in endotoxin tolerance. *Journal of Cellular Biochemistry*, 120(1), 56–70. <https://doi.org/10.1002/jcb.27547>
- Liu, Y., Liu, C., Zhang, A., Yin, S., Wang, T., Wang, Y., Wang, M., Liu, Y., Ying, Q., Sun, J., Wei, F., Liu, D., Wang, C., & Ge, S. (2019). Down-regulation of long non-coding RNA MEG3 suppresses osteogenic differentiation of periodontal ligament stem cells (PDLSCs) through miR-27a-3p/IGF1 axis in periodontitis. *Aging (Albany NY)*, 11(15), 5334–5350. <https://doi.org/10.18632/aging.102105>
- Loosbroock, C., & Hunter, K. W. (2014). Inhibiting TNF- α signaling does not attenuate induction of endotoxin tolerance. *Journal of Inflammation Research*, 7, 159–167. <https://doi.org/10.2147/JIR.S75037>
- Lopez-Collazo, E., & del Fresno, C. (2013). Pathophysiology of endotoxin tolerance: Mechanisms and clinical consequences. *Critical Care*, 17(6), 242. <https://doi.org/10.1186/cc13110>
- Lyrioni, K., Patsalos, A., Daskalaki, M. G., Doxaki, C., Soennichsen, B., Helms, M., Liapis, I., Zacharioudaki, V., Kampranis, S. C., & Tsatsanis, C. (2017). Epigenetic and transcriptional regulation of Irak-M expression in macrophages. *Journal of Immunology*, 198(3), 1297–1307. <https://doi.org/10.4049/jimmunol.1600009>
- Martin, M., Katz, J., Vogel, S. N., & Michalek, S. M. (2001). Differential induction of endotoxin tolerance by lipopolysaccharides derived from *Porphyromonas gingivalis* and *Escherichia coli*. *Journal of Immunology*, 167(9), 5278–5285. <https://doi.org/10.4049/jimmunol.167.9.5278>
- Milner, K. C. (1973). Patterns of tolerance to endotoxin. *The Journal of Infectious Diseases*, 128(Suppl), 237–245. https://doi.org/10.1093/infdis/128.supplement_1.s237
- Monneret, G., Venet, F., Pachot, A., & Lepape, A. (2008). Monitoring immune dysfunctions in the septic patient: A new skin for the old ceremony. *Molecular Medicine*, 14(1–2), 64–78. <https://doi.org/10.2119/2007-00102.Monneret>
- Muthukuru, M., & Cutler, C. W. (2015). Resistance of MMP9 and TIMP1 to endotoxin tolerance. *Pathogens and Disease*, 73(5), ftu003. <https://doi.org/10.1093/femspd/ftu003>
- Muthukuru, M., Jotwani, R., & Cutler, C. W. (2005). Oral mucosal endotoxin tolerance induction in chronic periodontitis. *Infection and Immunity*, 73(2), 687–694. <https://doi.org/10.1128/IAI.73.2.687-694.2005>
- Nahid, M. A., Satoh, M., & Chan, E. K. (2011). Mechanistic role of microRNA-146a in endotoxin-induced differential cross-regulation of TLR signaling. *Journal of Immunology*, 186(3), 1723–1734. <https://doi.org/10.4049/jimmunol.1002311>
- Nakagawa, R., Naka, T., Tsutsui, H., Fujimoto, M., Kimura, A., Abe, T., Seki, E., Sato, S., Takeuchi, O., Takeda, K., Akira, S., Yamanishi, K., Kawase, I., Nakanishi, K., & Kishimoto, T. (2002). SOCS-1 participates in negative regulation of LPS responses. *Immunity*, 17(5), 677–687. [https://doi.org/10.1016/s1074-7613\(02\)00449-1](https://doi.org/10.1016/s1074-7613(02)00449-1)
- Nakamura, T., Nitta, H., & Ishikawa, I. (2004). Effect of low dose *Actinobacillus actinomycetemcomitans* lipopolysaccharide pre-treatment on cytokine production by human whole blood. *Journal of Periodontal Research*, 39(2), 129–135. <https://doi.org/10.1111/j.1600-0765.2004.00717.x>
- Netea, M. G., Dominguez-Andrés, J., Barreiro, L. B., Chavakis, T., Divangahi, M., Fuchs, E., Joosten, L. A. B., van der Meer, J. W. M., Mhlanga, M. M., Mulder, W. J. M., Riksen, N. P., Schlitzer, A., Schultze, J. L., Stabel Benn, C., Sun, J. C., Xavier, R. J., & Latz, E. (2020). Defining trained immunity and its role in health and disease. *Nature Reviews Immunology*, 20(6), 375–388. <https://doi.org/10.1038/s41577-020-0285-6>
- Netea, M. G., Schlitzer, A., Placek, K., Joosten, L. A. B., & Schultze, J. L. (2019). Innate and adaptive immune memory: An evolutionary continuum in the host's response to pathogens. *Cell Host & Microbe*, 25(1), 13–26. <https://doi.org/10.1016/j.chom.2018.12.006>
- Neter, E. (1969). Endotoxins and the immune response. *Current Topics in Microbiology and Immunology*, 47, 82–124. https://doi.org/10.1007/978-3-642-46160-6_5
- Nogueira-Filho, G., Rosa, B. T., Santos, P. F., Tunes, U. R., Freire, S. M., Meyer, R., & Darveau, R. P. (2014). Whole-blood cultures from patients with chronic periodontitis respond differently to *Porphyromonas gingivalis* but not *Escherichia coli* lipopolysaccharide. *Journal of Periodontology*, 85(2), e18–23. <https://doi.org/10.1902/jop.2013.120735>

- Novakovic, B., Habibi, E., Wang, S. Y., Arts, R. J. W., Davar, R., Megchelenbrink, W., Kim, B., Kuznetsova, T., Kox, M., Zwaag, J., Matarese, F., van Heeringen, S. J., Janssen-Megens, E. M., Sharifi, N., Wang, C., Keramati, F., Schoonenberg, V., Flicek, P., Clarke, L., ... Stunnenberg, H. G. (2016). Beta-glucan reverses the epigenetic state of LPS-induced immunological tolerance. *Cell*, 167(5), 1354–1368. e1314. <https://doi.org/10.1016/j.cell.2016.09.034>
- Nowotny, A., & Sanavi, F. (1983). Induction of nonspecific tolerance to endotoxins reduces the alveolar bone resorption in ligature-treated rats. *Infection and Immunity*, 39(2), 873–878. <https://doi.org/10.1128/IAI.39.2.873-878.1983>
- Pillay, J., Ramakers, B. P., Kamp, V. M., Loi, A. L., Lam, S. W., Hietbrink, F., Leenen, L. P., Tool, A. T., Pickkers, P., & Koenderman, L. (2010). Functional heterogeneity and differential priming of circulating neutrophils in human experimental endotoxemia. *Journal of Leukocyte Biology*, 88(1), 211–220. <https://doi.org/10.1189/jlb.1209793>
- Rackov, G., Shokri, R., De Mon, M. A., Martinez, A. C., & Balomenos, D. (2017). The role of IFN-beta during the course of sepsis progression and its therapeutic potential. *Frontiers in Immunology*, 8, 493. <https://doi.org/10.3389/fimmu.2017.00493>
- Rimmelé, T., Payen, D., Cantaluppi, V., Marshall, J., Gomez, H., Gomez, A., Murray, P., & Kellum, J. A. (2016). Immune cell phenotype and function in sepsis. *Shock*, 45(3), 282–291. <https://doi.org/10.1097/SHK.0000000000000495>
- Rios, E. C., de Lima, T. M., Moretti, A. I., & Soriano, F. G. (2016). The role of nitric oxide in the epigenetic regulation of THP-1 induced by lipopolysaccharide. *Life Sciences*, 147, 110–116. <https://doi.org/10.1016/j.lfs.2016.01.041>
- Seeley, J. J., & Ghosh, S. (2017). Molecular mechanisms of innate memory and tolerance to LPS. *Journal of Leukocyte Biology*, 101(1), 107–119. <https://doi.org/10.1189/jlb.3MR0316-118RR>
- Singer, M., Deutschman, C. S., Seymour, C. W., Shankar-Hari, M., Annane, D., Bauer, M., Bellomo, R., Bernard, G. R., Chiche, J. D., Cooper-Smith, C. M., Hotchkiss, R. S., Levy, M. M., Marshall, J. C., Martin, G. S., Opal, S. M., Rubenfeld, G. D., van der Poll, T., Vincent, J. L., & Angus, D. C. (2016). The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA*, 315(8), 801–810. <https://doi.org/10.1001/jama.2016.0287>
- Stratigi, K., Kapsetaki, M., Aivaliotis, M., Town, T., Flavell, R. A., & Spilianakis, C. G. (2015). Spatial proximity of homologous alleles and long noncoding RNAs regulate a switch in allelic gene expression. *Proceedings of the National Academy of Sciences of the United States of America*, 112(13), E1577–E1586. <https://doi.org/10.1073/pnas.1502182112>
- Sun, X., Sun, J., Shao, X., Feng, J., Yan, J., & Qin, Y. (2018). Inhibition of microRNA-155 modulates endotoxin tolerance by upregulating suppressor of cytokine signaling 1 in microglia. *Experimental and Therapeutic Medicine*, 15(6), 4709–4716. <https://doi.org/10.3892/etm.2018.6032>
- Sun, Y., Li, H., Sun, M. J., Zheng, Y. Y., Gong, D. J., & Xu, Y. (2014). Endotoxin tolerance induced by lipopolysaccharides derived from *Porphyromonas gingivalis* and *Escherichia coli*: Alternations in toll-like receptor 2 and 4 signaling pathway. *Inflammation*, 37(1), 268–276. <https://doi.org/10.1007/s10753-013-9737-5>
- Takaesu, G., Surabhi, R. M., Park, K. J., Ninomiya-Tsuji, J., Matsumoto, K., & Gaynor, R. B. (2003). TAK1 is critical for I κ B kinase-mediated activation of the NF- κ B pathway. *Journal of Molecular Biology*, 326(1), 105–115. [https://doi.org/10.1016/s0022-2836\(02\)01404-3](https://doi.org/10.1016/s0022-2836(02)01404-3)
- Takeda, K., & Akira, S. (2015). Toll-like receptors. *Current Protocols in Immunology*, 109, 14.12.1–14.12.10. <https://doi.org/10.1002/0471142735.im1412s109>
- Touyz, L. Z. (2013). Periodontitis contributes to initiation, progress and aggravation of septic shock; a feasible hypothesis. *Medical Hypotheses*, 81(4), 650–652. <https://doi.org/10.1016/j.mehy.2013.06.033>
- van der Flier, M., Sharma, D. B., Estevão, S., Emonts, M., Rook, D., Hazelzet, J. A., van Goudoever, J. B., & Hartwig, N. G. (2013). Increased CD4(+) T cell co-inhibitory immune receptor CEACAM1 in neonatal sepsis and soluble-CEACAM1 in meningococcal sepsis: A role in sepsis-associated immune suppression? *PLoS One*, 8(7), e68294. <https://doi.org/10.1371/journal.pone.0068294>
- Van Epps, H. L. (2006). Ignoring endotoxin. *The Journal of Experimental Medicine*, 203(5), 1137. <https://doi.org/10.1084/jem.2035fta>
- van Lier, D., Geven, C., Leijte, G. P., & Pickkers, P. (2019). Experimental human endotoxemia as a model of systemic inflammation. *Biochimie*, 159, 99–106. <https://doi.org/10.1016/j.biochi.2018.06.014>
- Vergadi, E., Vaporidi, K., & Tsatsanis, C. (2018). Regulation of endotoxin tolerance and compensatory anti-inflammatory response syndrome by non-coding RNAs. *Frontiers in Immunology*, 9, 2705. <https://doi.org/10.3389/fimmu.2018.02705>
- Ward, N. S., Casserly, B., & Ayala, A. (2008). The compensatory anti-inflammatory response syndrome (CARS) in critically ill patients. *Clinics in Chest Medicine*, 29(4), 617–625, viii. <https://doi.org/10.1016/j.ccm.2008.06.010>
- West, M. A., & Heagy, W. (2002). Endotoxin tolerance: A review. *Critical Care Medicine*, 30(1 Supp), S64–S73.
- Wu, X., Zhang, G., Wang, X., Zhang, P., & Tan, Y. (2015). Endotoxin tolerance induction in human periodontal ligament fibroblasts stimulated with different bacterial lipopolysaccharides. *Archives of Oral Biology*, 60(3), 463–470. <https://doi.org/10.1016/j.archoralbio.2014.10.005>
- Yamamoto, Y., He, P., Klein, T. W., & Friedman, H. (1994). Endotoxin induced cytotoxicity of macrophages is due to apoptosis caused by nitric oxide production. *Journal of Endotoxin Research*, 1(3), 181–187. <https://doi.org/10.1177/096805199400100307>
- Zaric, S., Shelburne, C., Darveau, R., Quinn, D. J., Weldon, S., Taggart, C. C., & Coulter, W. A. (2010). Impaired immune tolerance to *Porphyromonas gingivalis* lipopolysaccharide promotes neutrophil migration and decreased apoptosis. *Infection and Immunity*, 78(10), 4151–4156. <https://doi.org/10.1128/IAI.00600-10>
- Zhao, G., Su, Z., Song, D., Mao, Y., & Mao, X. (2016). The long noncoding RNA MALAT1 regulates the lipopolysaccharide-induced inflammatory response through its interaction with NF- κ B. *FEBS Letters*, 590(17), 2884–2895. <https://doi.org/10.1002/1873-3468.12315>
- Zhou, W. Q., Wang, P., Shao, Q. P., & Wang, J. (2016). Lipopolysaccharide promotes pulmonary fibrosis in acute respiratory distress syndrome (ARDS) via lincRNA-p21 induced inhibition of Thy-1 expression. *Molecular and Cellular Biochemistry*, 419(1–2), 19–28. <https://doi.org/10.1007/s11010-016-2745-7>
- Zhu, X. Q., Lu, W., Chen, Y., Cheng, X. F., Qiu, J. Y., Xu, Y., & Sun, Y. (2016). Effects of *Porphyromonas gingivalis* lipopolysaccharide tolerized monocytes on inflammatory responses in neutrophils. *PLoS One*, 11(8), e0161482. <https://doi.org/10.1371/journal.pone.0161482>

How to cite this article: Larsson, L., Garaicoa-Pazmino, C., Asa'ad, F., & Castilho, R. M. (2022). Understanding the role of endotoxin tolerance in chronic inflammatory conditions and periodontal disease. *Journal of Clinical Periodontology*, 49(3), 270–279. <https://doi.org/10.1111/jcpe.13591>