

Stephen E. Kesler · John T. Chesley
John N. Christensen · Richard D. Hagni
Wouter Heijlen · J. Richard Kyle · Phillipe Muechez
Kula C. Misra · Rob van der Voo

Discussion of “Tectonic controls of Mississippi Valley-type lead-zinc mineralization in orogenic forelands” by D.C. Bradley and D.L. Leach

Received: 22 March 2004 / Accepted: 22 March 2004 / Published online: 8 July 2004
© Springer-Verlag 2004

The interesting and useful study by Bradley and Leach (2003) of tectonic controls on Mississippi Valley-type (MVT) mineralization in orogenic forelands is marred by a surprising disregard for published evidence for the age of these deposits. This problem is of major importance, because the ages of the deposits control their relation to tectonic features. Here are three examples of the problem, showing how they relate to the suggested tectonic models.

Editorial handling: L. Meinert

S. E. Kesler (✉)
Department of Geological Sciences,
University of Michigan, Ann Arbor, MI 48109, USA
E-mail: skesler@umich.edu

J. T. Chesley
Department of Geosciences,
University of Arizona, Tucson, AZ 85721, USA

J. N. Christensen
Center for Isotope Geochemistry,
Lawrence Berkeley National Laboratory, Berkeley,
CA 94720-8179, USA

R. D. Hagni
Department of Geology and Geophysics,
University of Missouri-Rolla, Rolla,
MO 65409-0410, USA

W. Heijlen · P. Muechez
Fysico-chemische Geologie, Katholieke Universiteit,
3001 Leuven, Belgium

J. R. Kyle
Department of Geological Sciences,
University of Texas at Austin,
Austin, TX 78712-0254, USA

K. C. Misra
Department of Earth and Planetary Sciences,
University of Tennessee, Knoxville,
TN 37996-1410, USA

R. van der Voo
Department of Geological Sciences,
University of Michigan, Ann Arbor,
MI 48109, USA

1. Bradley and Leach (2003, p. 657 and p. 663) cite the Canadian Rockies as an example of MVT deposits formed in an Andean-type orogen based on Laramide-age paleomagnetic poles for Pine Point, Kicking Horse and Robb Lake. They ignore Devonian Rb/Sr ages for sphalerite from Pine Point because they “may date late Devonian clays entrapped in the much younger sphalerite.” This statement is referenced to Symons et al. (1998b), which contains nothing about Pine Point. Perhaps they meant to refer to Symons et al. (1998a), which states (p. 79) that “Garven and Sverjensky (1994) noted that the (Rb-Sr) method is prone to contamination by colloidal clay particles entrapped in sphalerite....” However, Garven and Sverjensky (1994, p. 1,150) say only that “Other workers dated early-stage sphalerite at Pine Point as Devonian using a Rb-Sr method, but this type of analysis may be prone to contamination by clay particles incorporated during rapid precipitation of colloform aggregates of metal sulfides.” None of these studies contains any information on clay minerals at Pine Point. Nakai et al. (1990, 1993), which were omitted by Bradley and Leach (2003), deal with this issue specifically and show that inclusions of clay minerals are unlikely to account for Rb-Sr compositions of MVT sphalerite. If Cordilleran MVT deposits are Devonian in age, they could have formed when the western margin of North America was “...dominated by long-lived regional extension” (Nelson et al. 2002).
2. Bradley and Leach (2003, p. 657 and Fig. 5) cite the Cevennes and Maestrat areas as examples of MVT deposits formed in an inversion-type orogen of Santonian to Miocene-age in the Pyrenees. In support of this interpretation, they refer to U-Pb dating by Grandia et al. (2000), isotopic ages on fluorite in Leach et al. (2001), and paleomagnetic measurements by Lewchuk et al. (1998). Sample and analytical data on the fluorite dated by Leach et al. (2001) are, to our knowledge, not published or available for evaluation. However, the U-Pb isochron age of about 62.6 Ma

for calcite and galena from the AVECILLA mine in the Maestrat basin has been interpreted to indicate that mineralization took place during a rifting or post-rifting interval and before Oligocene-age inversion to form the Pyrenees (Grandia et al. 2000, 2003).

- Bradley and Leach (2003, p. 662) suggest East Tennessee as an example of MVT mineralization that could have formed after thrust faulting. They note that this would require that the famous “sphalerite sands” (Fig. 1), which indicate that ore formed before folding (Kendall 1960; Matlock and Misra 1983), “actually formed by grain-by-grain replacement of pre-tectonic carbonate sands.” In support of this revised origin for the sphalerite sands, they cite a paleomagnetic study that contains no information about the sands and says only that “...epigenetic sphalerite could easily replace a clastic carbonate grain...” (Symons and Stratakos 2000, p. 376), as does a longer version of the same study (Symons and Stratakos 2003). Thus, Bradley and Leach (2003) and references therein provide no information to support the contention that “the timing of mineralization with respect to thrusting in East Tennessee is debatable” and that the Devonian Rb-Sr ages for sphalerite in the district (Nakai et al. 1990, 1993) should be ignored.

These conclusions and others by Bradley and Leach (2003) about tectonic controls of MVT mineralization in orogenic forelands reflect a clear preference for MVT-age estimates based on paleomagnetic measurements over those based on geologic observations or isotopic age analyses (unless they agree with the paleomagnetic ages). However, there is no proof that remnant magnetism measured in MVT deposits is actually associated with the ore minerals. The carrier for remnant magne-

tization in most deposits is magnetite or pyrrhotite, both of which are extremely rare in MVT ore. In the Viburnum Trend, which is commonly cited in support of the coexistence of magnetite and MVT ore minerals, only a few magnetite grains have been observed in thousands of polished sections, and its paragenetic relation to other minerals is uncertain (Hagni 1986, p. 123). In East Tennessee, micron-size magnetite is associated with authigenic feldspar that formed after MVT mineralization (Suk et al. 1990a, b; Aleinikoff et al. 1993). Post-ore magnetite and pyrrhotite of this type probably form when small amounts of iron are released from minerals such as dolomite or sphalerite during alteration caused by new pore fluids or deeper burial. Because so little magnetite or pyrrhotite is needed to produce a magnetic signature in carbonate rocks, this alteration is very difficult to recognize.

Evidence that paleomagnetic ages for MVT deposits are susceptible to alteration and resetting is seen in their relation to isotopic ages. In East Tennessee, Pine Point, and Silesia, paleomagnetic ages are younger than Rb-Sr ages on sphalerite; only at Polaris do the two methods yield the same age (Nakai et al. 1990, 1993; Symons and Sangster 1992; Symons et al. 1993, 1995; Christensen et al. 1996; Christensen et al. 1995; Symons and Stratakos 2000, 2003; Heijlen et al. 2003). Where paleomagnetic measurements can be compared to Sm-Nd ages on fluorite in the Illinois-Kentucky fluorite district, the paleomagnetic age is much younger and the Sm-Nd age agrees with Rb-Sr ages on sphalerite from the possibly related Upper Mississippi Valley district (Brannon et al. 1992; Chesley et al. 1994; Symons 1994). Finally, in Central Tennessee, the age indicated by paleomagnetic poles is the same as that obtained from Th-Pb isotopes in ore-stage calcite (Brannon et al. 1996; Lewchuk and Symons 1995). In all cases, the paleomagnetic age is the same as or younger than the isotopic age, suggesting that paleomagnetic measurements frequently reflect later fluid events. (It is important to note that this relation is the same for minerals such as fluorite and calcite in which the crystallographic position of the radioactive isotope is well understood and sphalerite, where its exact crystallographic setting remains unclear.) A few recent paleomagnetic studies have found evidence for multiple events in complex orogens (Weil and Van der Voo 2002), but most studies lack this resolution and reflect only late events. The fact that paleomagnetic age estimates are older than some orogenic events that did not form MVT deposits (Leach et al. 2002), tells us only that paleomagnetic signatures can survive some events, not that they provide an age for MVT mineralization.

In their closing statement, Bradley and Leach (2003, p. 664) indicate that “...other tectonic models are needed for MVT genesis...” outside obvious convergent-margin settings with well-defined foreland basins. In particular, their selective rejection of isotopic measurements and heavy reliance on questionable paleomagnetic age constraints appears to have obscured the number and distribution of MVT deposits that formed in an extensional




Fig. 1 Hand-samples of “sphalerite sand” from the Mascot-Jefferson City district showing relation between sphalerite-bearing clasts and dolomite sands. Note that large clasts consist only partly of sphalerite (S) and some are mantled by sparry dolomite (D) of post-sphalerite age, making a replacement origin particularly unlikely

setting. In addition to the Nanisivik and Canning Basin deposits that were noted by Bradley and Leach (2003) and those of the Canadian Cordillera and Pyrenees mentioned above, Rb-Sr ages for MVT mineralization in the Upper Silesian district of Poland indicate that it formed "...in response to Early Cretaceous crustal extension preceding the opening of the northern Atlantic Ocean" (Heijlen et al. 2003), and a similar extensional setting is thought to have prevailed for Mesozoic-age MVT mineralization in the Verviers-Aachen MVT district in Belgium (Heijlen et al. 2001). If MVT deposits in extensional settings are as widespread as this growing list suggests, then tectonic and fluid flow models related to orogenic forelands will have to be revised. Answers to this important question will require a more balanced evaluation of all data that constrain the age of MVT deposits.

References

- Aleinikoff JN, Walter M, Kunk MJ, Hearn PP Jr. (1993) Do ages of authigenic K-feldspar date the formation of Mississippi Valley-type Pb-Zn deposits, central and southeastern United States?: Pb isotopic evidence. *Geology* 21:73–76
- Bradley DC, Leach DL (2003) Tectonic controls of Mississippi Valley-type lead-zinc mineralization in orogenic forelands. *Miner Deposita* 38:652–667
- Brannon JC, Podosek FA, McLimans RK (1992) Alleghanian age of the Upper Mississippi Valley zinc-lead deposit determined by Rb-Sr dating of sphalerite. *Nature* 356:509–511
- Brannon JC, Podosek FA, Cole SC (1996) Radiometric dating of Mississippi Valley-type ore deposits. Society of Economic Geologists, Spec. Publ. 4, College Station, TX, pp 536–545
- Chesley JT, Halliday AN, Kyser TK, Spry PG (1994) Direct dating of Mississippi Valley-type mineralization: use of Sm-Nd in fluorite. *Econ Geol* 89:1192–1199
- Christensen JN, Halliday AN, Kesler SE, Leigh KE, Randell RN (1995) Direct dating of sulfides by Rb-Sr: a critical test using the Polaris Mississippi Valley-type Zn-Pb deposit. *Geochim Cosmochim Acta* 59:5191–5197
- Christensen JN, Halliday AN, Kesler SE (1996) Rb-Sr dating of sphalerite and the ages of Mississippi Valley-type Pb-Zn deposits. Society of Economic Geologists, Spec. Publ. 4, College Station, TX, pp 527–5335
- Garven G, Sverjensky DA (1994) Paleohydrogeology of the Canadian Rockies and origins of brines, Pb-Zn deposits and dolomitization in the Western Canada sedimentary basin: comment. *Geology* 22:1149–1150
- Grandia F, Asmerom Y, Getty S, Cardellach E, Canals Á (2000) U-Pb dating of MVT ore-stage calcite: implications for fluid flow in a Mesozoic extensional basin from Iberian Peninsula. *J Geochem Explor* 69–70:377–380
- Grandia F, Cardellach E, Canals Á, Banks DA (2003) Geochemistry of the fluids related to epigenetic carbonate-hosted deposits in the Maestrat Basin, eastern Spain: fluid inclusion and isotope (Cl C O S Sr) evidence. *Econ Geol* 98:933–954
- Hagni RD (1986) Mineral paragenetic sequence of the lead-zinc-copper-cobalt-nickel ores of the Southeast Missouri Lead District, USA. In: Mineral parageneses. Theophrastus. Athens, Greece, pp 93–132
- Heijlen W, Muechez P, Banks DA (2001) Origin and evolution of high-salinity Zn-Pb mineralizing fluids in the Variscides of Belgium. *Miner Deposita* 36:165–176
- Heijlen W, Muechez P, Banks DA, Schneider J, Kucha H, Keppens E (2003) Carbonate-hosted Zn-Pb deposits in Upper Silesia, Poland: origin and evolution of mineralizing fluids and constraints on genetic models. *Econ Geol* 98:911–932
- Kendall DL (1960) Ore deposits and sedimentary features, Jefferson City Mine, Tennessee. *Econ Geol* 55:985–1003
- Leach DL, Bradley D, Lewchuk MT, Symons DTA, de Marsily G, Brannon J (2001) Mississippi Valley-type lead-zinc deposits through geological time: implications from recent age-dating research. *Miner Deposita* 36:711–740
- Leach DL, Bradley D, Lewchuk MT, Symons DTA, de Marsily G, Brannon J (2002) Discussion of Mississippi Valley-type lead-zinc deposits through geological time: implications from recent age-dating research, reply. *Miner Deposita* 37:803–805
- Lewchuk MT, Symons DTA (1995) Age and duration of Mississippi Valley-type ore mineralizing events. *Geology* 23:233–236
- Lewchuk MT, Rouvier H, Henry B, Macquar J-C, Leach DL (1998) Paleomagnetism of Mississippi Valley-type mineralization in southern France and Cenozoic orogenesis. In: European Geophysical Society 23 Gen Assoc, part 1. Soc. Symp. Solid Earth Geophysics and Geodesy, 20–24 April 1998, Nice
- Matlock JF, Misra KC (1993) Sphalerite-bearing detrital "sand" bodies in Mississippi Valley-type zinc deposits, Mascot-Jefferson City district, Tennessee: implications for the age of mineralization. *Miner Deposita* 28:344–353
- Nakai S, Halliday AN, Kesler SE, Jones HD (1990) Rb-Sr dating of sphalerite and genesis of MVT deposits. *Nature* 346:354–357
- Nakai S, Halliday AN, Kesler SE, Jones HD, Kyle JR, Lane TE (1993) Rb-Sr dating of sphalerites from Mississippi Valley-type (MVT) ore deposits. *Geochim Cosmochim Acta* 57:417–427
- Nelson J, Paradis S, Christensen J, Gabites J (2002) Canadian Cordilleran Mississippi Valley-type deposits: a case for Devonian-Mississippian back-arc hydrothermal origin. *Econ Geol* 97:1013–1036
- Suk D, Peacor D R, Van der Voo R (1990a) Replacement of pyrite framboids by magnetite in limestones and implication for paleomagnetism. *Nature* 345:611–613
- Suk D, Van der Voo R, Peacor DR (1990b) Scanning and transmission electron microscope observations of magnetite and other iron phases in Ordovician carbonates from East Tennessee. *J Geophys Res* 95:12327–12336
- Symons DTA (1994) Paleomagnetism and the late Jurassic genesis of the Illinois-Kentucky fluorspar deposits. *Econ Geol* 89:438–449
- Symons DTA, Sangster DF (1992) Late Devonian paleomagnetic age for the Polaris Mississippi Valley-type Zn-Pb deposit, Canadian Arctic Archipelago. *Can J Earth Sci* 29:15–25
- Symons DTA, Stratakis KK (2000) Paleomagnetic dating of dolomitization and Mississippi Valley-type mineralization in the Mascot-Jefferson City district of eastern Tennessee: a preliminary analysis. *J Geochem Explor* 69–70:373–376
- Symons DTA, Stratakis KK (2003) Paleomagnetic dating of Alleghanian orogenesis and mineralisation in the Mascot-Jefferson City zinc district or East Tennessee, USA. *Tectonophysics* 348:51–72
- Symons DTA, Pan H, Sangster DF, Jowett EC (1993) Paleomagnetism of the Pine Point Zn-Pb-deposits. *Can J Earth Sci* 30:1028–1036
- Symons DTA, Sangster DF, Leach DL (1995) A Tertiary age from paleomagnetism for Mississippi Valley-type zinc-lead mineralization in Upper Silesia, Poland. *Econ Geol* 90:782–794
- Symons DTA, Lewchuk M, Sangster DF (1998a) Laramide orogenic fluid flow into the Western Canada Sedimentary Basin: evidence from paleomagnetic dating of the Kicking Horse Mississippi Valley-type ore deposit. *Econ Geol* 93:68–83
- Symons DTA, Lewchuk M, Leach DF (1998b) Age and duration of the Mississippi Valley-type mineralizing fluid flow events in the Viburnum Trend, southeast Missouri, USA, from paleomagnetism. *J Geol Soc* 144:27–39
- Weil AB, Van der Voo R (2002) Insights into the mechanism for orogen-related carbonate remagnetization from growth of authigenic Fe-oxide: a SEM and rock magnetic study of Devonian carbonates from northern Spain. *J Geophys Res* 107(B4):2063–2066