also exhibited by birds in an experiment on simulated distasteful Batesian mimicry¹² and is probably indicative of a conflict of behavioural tendencies in the animal. This slowing down of attack would give a real mimic more time to escape. Behaviour of this type was never shown towards the non-mimetic (green) seed by the experimental group, nor to any colour of seed by the control group.

The formation of an avoidance image² by the birds was due to the simulated escape mechanism and not to some feature of the apparatus, as the control group fed readily on all seeds from it and the experimental group went for the green seed without hesitation throughout the experiment. Avoidance image formation was therefore through 'frustration learning' which can be as effective as pain learning¹³, the mechanism operative in discriminating distasteful from palatable prey. It would seem then, that 'an efficient escape mechanism' as observed by Lindroth³ could be as powerful as distastefulness in qualifying its possessor as a model for Batesian mimicry. Further observations in the field may show the phenomenon to be less rare than is now supposed.

Table 1 Model-mimic/non-mimetic ratio for each bird during the learning period, retention period and both periods combined

Bird	Learning		Retention		Both periods combined (model/mimic complex)	
	B/G	R/G	B/G	R/G	B/G	R/G
B	2.7	4.1	4.0	7.8	3.5	6.0
M	6.8	10.7	4.0	4.5	5.5	7.8
WR	3.6	5.8	4.8	4.9	4.2	5.4
WL	2.5	3.5	2.1	2.8	2.4	3.3
All	3.1	4.7	3.8	5.7	3.3	5.0

B, Blue seed, (cryptic model/mimic); G, green seed, (non-mimetic); R, red seed, (aposematic model/mimic).

These findings also have implications for the phenomenon of balanced polymorphism in a prey population with aposematic morphs. Birds feed in a frequency dependent manner, the more common the prey the stronger the searching image¹⁴ formed. The selection of apostates (non-mimetic polymorphs) enables a prey species to occur in an area at a greater density than if it were monomorphic. Some of these morphs may appear aposematic and will create a searching image more readily than a cryptic morph. These aposematic forms, if in a species acceptable to a predator, will be found at a very low frequency in the population through apostatic selection mechanisms. *P. spumarius marginella* occurs in an area studied by Thompson⁴ at a level above that of other aposematic morphs, but below that of the cryptic morphs. As all four morphs have the escape mechanism the only difference is that *marginella* is strikingly coloured. In the experiment reported here, learning to avoid the aposematic model was more rapid and this learning was retained longer than for the cryptic model, though they had equally efficient escape mechanisms. In a natural situation the escape mechanisms would presumably be less than 100% effective but, as shown by the experiment, a predator would remember an encounter with the aposematic morph more readily than one with a cryptic morph³. If in a dimorphic species the aposematic form is, say, four times as easy to detect as the cryptic form then the strength of an avoidance or searching image formed would equal that formed by four of the cryptic form. In this simplified hypothetical situation with two morphs, a balanced polymorphic level would be reached when the cryptic morph was four times as common as the aposematic. If the aposematic morph becomes more frequent in the population 'the selective disadvantages of being conspicuous'¹⁴ and predation will increase until the aposematic morph is reduced to the balanced polymorphic level again.

I thank Mr John Lazarus for his reading and criticisms of the manuscript.

DIANNE O. GIBSON

Department of Zoology,
University of Newcastle-upon-Tyne,
Newcastle-upon-Tyne, UK

Received December 31, 1973; revised February 14, 1974.

- ¹ Cott, H. B., *Adaptive Coloration in Animals*, 256-258 (Methuen, London, 1940).
- ² Clarke, B., *Systematics Association Publications No. 4 Taxonomy & Geography*, 47-70 (1962).
- ³ Lindroth, C. H., *Ent. Scand.*, **2**, 41-48 (1972).
- ⁴ Thompson, V., *Nature*, **242**, 126-128 (1973).
- ⁵ Jones, F. M., *Trans. R. ent. Soc. Lond.*, **79**, 345-385 (1932).
- ⁶ Sommeron, V. G. L., and Jackson, T. H. E., *J. Lepit. Soc.*, **13**, 121-150 (1960).
- ⁷ Hespdenheide, H. A., *J. Ent. Ser. A.*, **43**, 49-55 (1973).
- ⁸ Rothschild, The Hon. M., in *Ecological Genetics & Evolution* (Edit. by Creed, R.) 202-223 (Blackwell, Oxford, 1971).
- ⁹ Pumphrey, R. J., *Ibis*, **90**, 171-190 (1948).
- ¹⁰ Klopfer, P. M., in *Behavioural Aspects of Ecology*, 10-34 (Prentice-Hall, New Jersey, 1962).
- ¹¹ Siegel, S., in *Non-parametric Statistics for the Biological Sciences*, 184-193 (McGraw-Hill, 1956).
- ¹² Reiskind, J., *Anim. Behav.*, **13**, 466-469 (1965).
- ¹³ Sutherland, N. S., and Mackintosh, N. J., in *Mechanisms of Animal Discrimination Learning*, 338-403 (Academic Press, 1971).
- ¹⁴ Tinbergen, L., *Archs neerl. Zool.*, **13**, 265-336 (1960).

Temperature effects on hearing in two species of *Amphisbaenia*

SEVERAL recent studies have documented diverse auditory responses in lizards and turtles as a function of body temperature. Previous work was with iguanid lizards, which are diurnal, and can regulate body temperature by basking; and with the aquatic turtle *Pseudemys*. We have worked with amphisbaenians, which are highly modified burrowing reptiles.

Werner (refs 1 and 2, and personal communication) showed that the ears of six species of iguanid lizards were most sensitive in terms of the electrical potentials of the cochlea at temperatures corresponding either to the 'eccritic' or to the similar 'preferred' temperature. At this optimum temperature the species-specific auditory sensitivity curve is regular and has a characteristic zone of maximum sensitivity². Temperature shifts produce fairly predictable species-specific variations. A temperature drop impairs the sensitivity, especially for the high tones, and makes the function highly irregular, with many sharp variations in frequency. A moderate rise improves the response to high tones while reducing it for low tones. A further elevation is likely to have little effect on high-tone sensitivity, but that to low tones is seriously impaired. In the region of best sensitivity the change often approaches 40 dB. Thus a torpid lizard is likely to have poor perception of sound, even though this may signal a predator or other sort of danger.

A lesser temperature effect was found in the turtle, *Pseudemys scripta elegans*³. For a head temperature range of 14° to 33° C the response to a constant tone of 330 or 630 Hz varied up to 4.3 dB, showing two peaks of sensitivity, respectively at 19° and 29.5° C. According to Patterson *et al.*, temperature has two effects: it acts on the energy source for the generation of potentials and involves either a conductive mechanism or blood pressure.

An earlier study⁴ reported limited observations on a single specimen of the African amphisbaenid *Chirindia langi*. Cochlear potentials differed up to 18 dB in the region around 3,000 Hz between 24.4° and 29.4° C.

Observations on two other species of amphisbaenians now demonstrate strikingly different behaviour. The animals were

four *Blanus cinereus cinereus* (Vandelli) from the vicinity of Jerez de la Frontera, Spain, and three *Diplometopon zarudnyi* Nikolski: one from the vicinity of Damman, Saudi Arabia, and two from the vicinity of Kuwait. *Blanus cinereus* is probably the most primitive living amphisbaenian, and *Diplometopon zarudnyi* is a highly modified trogonophid.

Specimens were anaesthetised with ethyl carbamate (0.012 ml per g body weight, 20% in physiological saline), often supplemented. The cochlear potentials were recorded with a needle electrode in contact with the perilymph of the saccule while stimulating with various tones over the range of 100 to 10,000 Hz. Temperature was monitored by a thermal probe deep in the cloaca and controlled within 1° C by means of an electric blanket draped over the animal. For details see ref. 4.

Fig. 1 Cochlear potentials of the left ear of *Blanus cinereus* (F868) taken at 25°, 31°, and 34° C in that sequence.

Figure 1 shows sensitivity curves (sound pressure in decibels relative to a pressure of 10^{-5} N m $^{-2}$, required at various frequencies to produce a standard potential of 0.1 μ V (that is, the lower the curve, the greater the sensitivity) for a specimen of *Blanus cinereus* at body temperatures of 25°, 31°, and 34° C. The curves differ, especially for the medium low and middle tones, but lack a clear trend as a function of temperature. Figure 2 shows similar observations on specimens of *Diplometopon zarudnyi* at four temperatures between 25° C and 40° C. All curves have the same general form.

A second specimen of *Diplometopon zarudnyi* tested at 37° C and then at 28° C yielded closely similar curves that cross one another several times. There was no obvious temperature effect, despite differences of 5–8 dB at two or three frequencies. The temperature was next set at 43° C, but the animal became hyperactive and a further test run had to be interrupted. After a period of time at a temperature of 35° C, a final series of measurements indicated reduced sensitivity. Possibly the high temperature of 43° C impaired the cochlear response.

These results reveal large differences among reptiles in the effects of temperature upon the response of the ear to sound. *Blanus* and *Diplometopon* maintain a remarkable constancy over a temperature range of as much as 15° C in sharp contrast to the lizards tested by Werner. *Pseudemys* and *Chirindia* evidently are intermediate.

It is as yet impossible to identify the process or processes involved in the observed temperature effects. Werner's observations seem definitely to exclude the sound conductive pathways, and point to the cochlear processes, and probably the hair cells themselves. The temperature effect may depend upon the magnitude of the polarisation potential set up at the surface membrane of the hair cell, and reflects the metabolic processes and membrane permeabilities by which this polarisation is established. Such possibilities should be tested and interspecies differences more broadly assayed. Do the responses reflect such environmental parameters as the range across

Fig. 2 Cochlear potentials of the left ear of *Diplometopon zarudnyi* (F869) taken at temperatures of 25°, 31°, 36°, and 40° C as indicated.

which a species normally thermoregulates? Can these amphisbaenians control the temperature of the otic capsule independent of general body temperature?

The sensitivity curves represented above show considerable variability tentatively attributed to the activity of middle ear muscles connecting to the terminal disk of the columella and probably affecting sound transmission to the inner ear⁵. Their activity may explain instability in the meter readings during presentation of a steady tone.

Efforts in one experiment to reduce this instability by the use of curare (*d*-tubocurarine) yielded uncertain results, perhaps because we were overcautious in the dosage used. Further experiments are needed on a more readily available amphisbaenian.

We thank M. Moya, J. Mandaville, A. d'A. Bellairs, and S. Howe who made these animals available for study. The work of C. G. is supported by a grant from the National Science Foundation, and that of E. G. W. by a grant from the National Institutes of Health.

CARL GANS

Department of Zoology,
The University of Michigan,
Ann Arbor, Michigan 48104

ERNEST GLEN WEVER

Auditory Research Laboratories,
Princeton University,
Princeton, New Jersey

Received October 29, 1973; revised March 25, 1974.

¹ Werner, Y. L., *J. Herp.*, **2**, 178 (1968).

² Werner, Y. L., *J. Herp.*, **6**, 147 (1972).

³ Patterson, W. C., *J. aud. Res.*, **6**, 453 (1966).

⁴ Gans, C., and Wever, E. G., *J. exp. Zool.*, **179**, 17 (1972).

⁵ Wever, E. G., and Gans, C., *Proc. natn. Acad. Sci. U.S.A.*, **69**, 2714 (1972).

Green cones of the piñon pine stimulate late summer breeding in the piñon jay

REPRODUCTION in birds is influenced by a variety of environmental factors, usually classified as ultimate or proximate¹. Ultimate factors determine efficiency of breeding, for example food supply for the young both before and after they become independent of their parents. Proximate factors, on the other hand, are the timers or predictors which initiate breeding at the appropriate time. Photoperiod is the most reliable and