

Characterization of a *Bacillus anthracis* spore coat-surface protein that influences coat-surface morphology

Michael Mallozzi¹, Joel Bozue², Rebecca Giorno¹, Krishna-Sulayman Moody², Alex Slack¹, Christopher Cote², Dengli Qiu³, Rong Wang³, Peter McKenney⁴, Erh-Min Lai⁵, Janine R. Maddock⁶, Arthur Friedlander⁷, Susan Welkos², Patrick Eichenberger⁴ & Adam Driks¹

¹Department of Microbiology and Immunology, Loyola University Medical Center, Maywood, IL, USA; ²Bacteriology Division, United States Army Medical Research Institute of Infectious Diseases, Frederick, MD, USA; ³Department of Biological, Chemical, and Physical Sciences, Illinois Institute of Technology, Chicago, IL, USA; ⁴Department of Biology and Center for Genomics and Systems Biology, New York University, NY, USA; ⁵Institute of Plant and Microbial Biology, Academia Sinica, Taipei, Taiwan, Republic of China; ⁶Department of Molecular, Cellular and Developmental Biology, University of Michigan, Ann Arbor, MI, USA; and ⁷Headquarters, United States Army Medical Research Institute of Infectious Diseases, Frederick, MD, USA

Correspondence: Adam Driks, Department of Microbiology and Immunology, Loyola University Medical Center, 2160 South First Avenue, Bldg. 105, Rm. 3820, Maywood, IL 60153, USA. Tel.: +1 708 216 3706; fax: +1 708 216 9574; e-mail: adriks@lumc.edu

Received 14 August 2008; accepted 15 September 2008.
First published online October 2008.

DOI:10.1111/j.1574-6968.2008.01380.x

Editor: Ezio Ricca

Keywords

Bacillus anthracis; coat; germination; spore assembly; anthrax.

Introduction

Bacterial spores can resist extremes of temperature, pH, toxic chemicals and predation by other microorganisms (Klobutcher *et al.*, 2006; Setlow, 2006). Studies in *Bacillus subtilis* demonstrate that spore resistance properties are derived from concentrically organized shells that encase the spore (Aronson & Fitz-James, 1976; Driks, 1999; Henriques *et al.*, 2004). A critical protective structure is the coat, a protein shell that encases bacterial spores of all species. In some species, such as *B. subtilis*, the coat is the external surface of the spore. Unlike *B. subtilis*, *Bacillus anthracis* and *Bacillus cereus* possess an additional balloon-like structure, called the exosporium, that encases the entire spore (Holt & Leadbetter, 1969; Aronson & Fitz-James, 1976). The *B. subtilis* coat is composed of 60–80 protein species which

Abstract

Bacterial spores are encased in a multilayered proteinaceous shell, called the coat. In many *Bacillus* spp., the coat protects against environmental assault and facilitates germination. In *Bacillus anthracis*, the spore is the etiological agent of anthrax, and the functions of the coat likely contribute to virulence. Here, we characterize a *B. anthracis* spore protein, called Cot β , which is encoded only in the genomes of the *Bacillus cereus* group. We found that Cot β is synthesized specifically during sporulation and is assembled onto the spore coat surface. Our analysis of a *cot β* null mutant in the Sterne strain reveals that Cot β has a role in determining coat-surface morphology but does not detectably affect germination. In the fully virulent Ames strain, a *cot β* null mutation has no effect on virulence in a murine model of *B. anthracis* infection.

are organized into distinctly staining inner and outer layers as visualized using thin-section electron microscopy (TEM) (Warth *et al.*, 1963; Kim *et al.*, 2006). The *B. anthracis* coat also has multiple layers, but these are more difficult to visualize because the coat is relatively thin (Holt & Leadbetter, 1969; Aronson & Fitz-James, 1976). Analysis of the composition and assembly of spore structures are critical steps towards understanding the molecular basis of spore resistance.

In addition to its protective roles, the *B. subtilis* coat also facilitates germination (Aronson & Fitz-James, 1976; Setlow, 2006). In *B. anthracis*, the etiological agent of anthrax, the spore is the infectious form of the bacterium, and the protective and germination-regulating functions of the coat (Kim *et al.*, 2004; Giorno *et al.*, 2007) are very likely to be important to virulence. During germination, the spore rehydrates and swells. The coat accommodates these changes

in volume by relaxing a series of folds on the dormant spore coat surface (Chada *et al.*, 2003; Plomp *et al.*, 2005, 2007). Soon after rehydration begins, the coat is shed, permitting further expansion of cell volume and completion of the conversion to the vegetative state (Santo & Doi, 1974).

The spore is formed during a multistage developmental process called sporulation. An early step in sporulation is the division of the cell into two unequally sized compartments by an asymmetrically positioned septum. Next, the smaller forespore compartment, which will become the spore, is engulfed by the larger mother cell. Ultimately, the mature spore is released into the milieu after mother cell lysis. In *B. subtilis*, spore coat assembly begins with synthesis of the spore coat proteins in the mother cell, just after the appearance of the sporulation septum (Zheng & Losick, 1990).

In this study, we characterize the coat protein Cot β , which is present in the genomes of *B. anthracis*, *B. cereus*, *Bacillus thuringiensis*, *Bacillus mycoides*, *Bacillus pseudomycooides* and *Bacillus weihenstephanensis* (members of the so-called *B. cereus* group) but not in any other as yet sequenced species, and which we identified in a search for novel *B. anthracis* spore proteins. We show that, in the Sterne strain of *B. anthracis*, Cot β is at the spore coat surface, and has a major effect on the topography of the coat surface. We further demonstrate that, in the fully virulent Ames strain, *cot* β is not necessary for infection in an animal model.

Materials and methods

General methods

Escherichia coli strains were cultured in Luria–Bertani medium and antibiotics were used at standard concentrations (Sambrook *et al.*, 1989). All *B. anthracis* strains are congenic with either the Ames or Sterne 34F2 strains. For the Ames and Sterne derivatives of *B. anthracis*, we used erythromycin at 5 $\mu\text{g mL}^{-1}$ and kanamycin at 20 $\mu\text{g mL}^{-1}$, and for Sterne derivatives, we used tetracycline at 5 $\mu\text{g mL}^{-1}$. Routine molecular biological techniques were performed using standard methodologies (Sambrook *et al.*, 1989). For atomic force-, fluorescence-, and immunofluorescence microscopy (IFM) experiments, Sterne strain spores were prepared by exhaustion sporulation in liquid Difco sporulation medium (DSM) (Harwood & Cutting, 1990). Spores judged to be dormant using phase contrast microscopy were collected and washed 5–10 times in sterile deionized water and analyzed immediately, as described below. For sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), spores of the Sterne and Ames strains were generated by culturing in DSM medium at 37 °C as described (Giorno *et al.*, 2007). For immunoblot analysis, reverse transcriptase (RT)-PCR analysis and animal

challenge assays, spores of the Sterne and Ames strains (and derivatives) were prepared in Leighton–Doi broth (Leighton & Doi, 1971) either at room temperature (for immunoblot) or at 37 °C (for RT-PCR and challenge assays), and then purified by centrifugation through a Hypaque-76 gradient (Nycomed Inc.). MS was performed as described previously (Lai *et al.*, 2003). To monitor *cot* β transcription in the Ames strain, mRNA was purified between 2 and 4 h after the onset of sporulation and RT-PCR analysis was performed as described previously (Bozue *et al.*, 2005). The plasmids used in this study to generate recombinant *B. anthracis* strains are listed in Table 1 and their construction is described in Supporting Information. Recombinant *B. anthracis* Sterne strains were generated by electroporation using unmethylated plasmid DNA (Koehler *et al.*, 1994). Recombinant Ames strains were generated using a protocol described previously (Mendelson *et al.*, 2004). A strain in which *cot* β was inactivated, Ames-JAB-7, was obtained by screening for kanamycin resistance and erythromycin sensitivity (encoded by plasmid pEO-3). The disruption of the *cot* β gene in the chromosome was confirmed using PCR. Spore proteins were extracted as described previously (Little & Driks, 2001), and were fractionated on 12% bis-Tris Gels (Invitrogen) and stained with either the Gel Code Blue system (Pierce) or Coomassie Brilliant Blue, or were electrotransferred to a polyvinylidene fluoride (PVDF) membrane in a 20% methanol, 3.7% SDS, Nupage Transfer Buffer (Invitrogen), in an Owl Panther transfer unit (Owl Separation Systems), and then stained with India Ink in a phosphate-buffered saline (PBS) solution (0.4% Tween-20, 0.1% India Ink) or for immunoblotting, were blocked in a 5% nonfat milk solution and then incubated with rabbit anti-green fluorescent protein (GFP) antibodies (Molecular Probes) at a dilution of 1 : 2000. After washing, membranes were incubated with anti-rabbit secondary antibodies conjugated to horseradish peroxidase (BioRad) at a dilution of 1 : 4000. Bands were visualized using the 1-Step 4CN reagent (Pierce).

Results and discussion

Roles of Cot β in structure and resistance

We have calculated, that *c.* 10–20 *B. anthracis*-specific spore proteins remain to be identified (Kim *et al.*, 2006; Giorno *et al.*, 2007). Because these proteins are likely to play roles in species-specific spore properties, we seek to identify these proteins using SDS-PAGE fractionation of spore extracts followed using MS analysis of excised bands (Lai *et al.*, 2003). Using this approach, we identified a peptide from a *B. anthracis* spore protein with a predicted mass of 8.4 kDa that migrates as a *c.* 10 kDa species (currently annotated as BAS1956 in the Sterne strain and BA2104 in the Ames

Table 1. Strains and plasmids

Strain	Genotype/description	Sources
Strains		
<i>E. coli</i>		
DH5 α	Cloning host	Lab stock
GM2163	<i>dam dcm</i> ; for transformation of <i>B. anthracis</i>	Lab stock
GM1684	<i>dam</i>	T. Koehler
Sterne strains		
34F2	pXO1 ⁺ pXO2 ⁻ , wild type	P. Jackson
RG61	pXO1 ⁺ pXO2 ⁻ , <i>cotβ::km</i>	This study
RG134	pXO1 ⁺ pXO2 ⁻ , <i>cotβΩpAS5</i>	This study
MGM68	pXO1 ⁺ pXO2 ⁻ , <i>cotβ::km</i>	This study
RG56	pXO1 ⁺ pXO2 ⁻ , <i>cotE::km</i>	Giorno et al. (2007)
MGM37	pXO1 ⁺ pXO2 ⁻ , <i>cotE::km, cotβΩpAS5</i>	This study
Ames strains		
Ames	pXO1 ⁺ pXO2 ⁺ , wild type	Little & Knudson (1986)
Ames-JAB-7	pXO1 ⁺ pXO2 ⁺ , <i>cotβ::km</i>	This study
Ames-JAB-10	pXO1 ⁺ pXO2 ⁺ , <i>cotβΩpEO3-cotβ-gfp</i>	This study
Name	Description	Sources
Plasmids		
pMR1	Used to introduce <i>kan^r</i> into the chromosome by marker replacement	Giorno et al. (2007)
pMR2	Used to replace <i>cotβ</i> with <i>kan^r</i>	This study
pAS1	Used to introduce <i>gfp</i> -fusion genes into the chromosome	Lab stock
pAS5	pAS1 bearing 3' fragment of <i>cotβ</i>	This study
pEO3	<i>E. coli/Bacillus</i> shuttle vector	Mendelson et al. (2004)
pJRS182	Origin of the Ωkm-2 cassette	Perez-Casal et al. (1991)
pJRS100.2	Source of the Ωkm-2 cassette	M. Caparon
pEO3- <i>cotβ</i>	pEO3+ 1.3 kb fragment containing <i>cotβ</i>	This study
pEO3- <i>cotβ</i> -Ωkm-2	pEO3- <i>cotβ</i> bearing Ωkm-2	This study
pEO3- <i>cotβ</i> - <i>gfp</i>	pEO3 containing <i>cotβ-gfp</i>	This study

strain) which we designate Cotβ (Fig. 1) [see Kim *et al.* (2004) for precedent for the use of Greek letters to name *B. anthracis* coat proteins]. This peptide does not match any sequences other than Cotβ and its orthologues in publicly available databases.

To analyze the role of *cotβ* in spore function and assembly, we inactivated it by replacing the gene with a kanamycin resistance cassette in the Sterne and Ames strains of *B. anthracis*, resulting in strains MGM68 and Ames-JAB-7, respectively. These mutations are unlikely to affect the transcription of the surrounding genes for three reasons: (1) the kanamycin resistance cassette is flanked by transcriptional termination sequences (Perez-Casal *et al.*, 1991), (2) RT-PCR analysis shows that *cotβ* is encoded by a separate mRNA transcript than that of the immediately upstream ORF (data not shown), and (3) the downstream ORF is oriented in the opposite direction from *cotβ*.

Both mutant strains sporulated normally and produced spores that were indistinguishable from wild type, based on phase-contrast light microscopy and TEM (data not shown).



Fig. 1. Identification of a novel spore protein in *Bacillus anthracis*. *Bacillus anthracis* coat proteins from the wild-type Sterne strain were extracted from spores, subjected to 16% SDS-PAGE and stained with Coomassie Blue G-250. The band indicated by an arrowhead was cut out and analyzed using matrix-assisted laser desorption/ionization time-of-flight MS. The peptide sequence is aligned with the best-fit ORF (BAS1956) from the annotated genome of the Sterne strain (inset). Molecular weight markers are indicated in kDa.

To assess general spore resistance properties, we subjected the Sterne strain *cot β* mutant (MGM68) to bleach, lysozyme, and heat challenges (using the methods described in Harwood & Cutting, 1990; Young & Setlow, 2003). We detected no significant differences between the mutant and wild-type strains (data not shown). Additionally, there were no readily detectable differences in the protein compositions of the extractable portion of the spore coat proteins from wild-type and mutant spores, as judged using one-dimensional SDS-PAGE (data not shown). We visualized bands in these gels both by staining with Coomassie Brilliant Blue, and by India ink staining after electrotransfer to a PVDF membrane, which allows detection of bands not stained by Coomassie (data not shown). In *B. subtilis*, the presence of multiple spore proteins in a single band is well documented (Kuwana *et al.*, 2002). We infer, therefore, that a protein other than Cot β is also present at this location in the gel (Fig. 1).

To determine whether Cot β has a role in directing coat-surface topography, we examined the Sterne strain *cot β* mutant spore surface using atomic force microscopy (AFM), which produces topographical relief maps. We have found that AFM can detect morphological defects not revealed using TEM (Chada *et al.*, 2003; McPherson *et al.*, 2005; Giorno *et al.*, 2007). Previously, we showed that the *B. anthracis* spore coat surface is spanned by ridges that extend, for the most part, from pole-to-pole along the long axis of the spore (Chada *et al.*, 2003; Wang *et al.*, 2007). In *c.* 85% of *B. anthracis* spores, the ridges appear in pairs, in close apposition (Fig. 2a), as opposed to well-separated single ridges for the case of *B. subtilis* (Chada *et al.*, 2003; Plomp *et al.*, 2005; Wang *et al.*, 2007). In contrast to wild-type *B. anthracis* spores, < 7% of *cot β* mutant Sterne strain spores (of 150 spores examined) possessed clear ridges. Instead, the coat surface has a smooth appearance, but with a cleft extending across the length of the spore (Fig. 2b and c). We note that this phenotype is very unlikely to be due to a mutation in an unlinked gene as the *cot β*

null mutation (in strain MGM68) was introduced by transduction from a strain (RG61) in which it was originally created (see Supporting Information). The *B. subtilis* coat-surface protein CotB also controls coat topography (Donovan *et al.*, 1987; Istatico *et al.*, 2001; Chada *et al.*, 2003). Possibly, a role in coat-surface morphology is typical of coat-surface proteins in all species.

Physiological roles of Cot β

Because defects in the coat can affect germination (Henriques & Moran, 2006), we analyzed the ability of *cot β* mutant spores to germinate using a variety of assays (see Supporting Information) that measure distinct steps in germination (Harwood & Cutting, 1990). However, we found that *cot β* mutant spores germinated identically to the wild-type strains (data not shown), suggesting that Cot β does not have a role in germination.

To detect a role for Cot β in pathogenesis, we infected female BALB/c mice intranasally (Lyons *et al.*, 2004) with either *c.* 1×10^6 wild-type Ames or *cot β* mutant (Ames-JAB-7) spores. Infection with wild-type spores resulted in a typical survival curve, with 93.4% of animals (out of 15 animals) having succumbed by day 6 postinfection. The survival curves of animals challenged with *cot β* spores or wild-type spores were indistinguishable (data not shown).

Timing of Cot β synthesis and subcellular localization

The observation that Cot β affects coat-surface morphology focused our attention on its subcellular location. To study this, we introduced *cot β -gfp* at the *cot β* locus (generating strain RG134) via a Campbell-type single reciprocal recombination, resulting in a chromosomal copy of the fusion gene at the endogenous locus. The endogenous, wild-type, copy of *cot β* is very likely to be inactive in RG134 because its





Fig. 2. Effect of *cot β* on spore coat-surface topography. Wild-type Sterne (a and a₁) and *cot β* mutant (strain MGM68) spores (b, b₁, and c) were analyzed by AFM in the tapping mode. Spore outlines and spore coat ridges imaged in (a) and (b) are indicated by white-dotted lines and arrows in (a₁) and (b₁), respectively. Scale bars represent 0.5 μ m in (a), (b) and (c).

promoter is separated from the coding sequence by the inserted plasmid backbone. We then used fluorescence microscopy to localize Cot β -GFP in sporulating cells and spores. To estimate the stage of development, we imaged the sporangial cell envelopes with the lipophilic counter stain FM4-64 (Pogliano *et al.*, 1999) as well as using phase-contrast microscopy (data not shown). Because sporulation had to be carried out at room temperature to visualize GFP-localization (see Supporting Information), the progression through sporulation was much slower than the *c.* 8 h required at 37 °C (see below).



We first detected Cot β -GFP when the forespore was engulfed (stage III) (Losick *et al.*, 1986) (at about 24 h of culturing, Fig. 3a), consistent with σ^E -control of *cot* β transcription (Hilbert & Piggot, 2004). Sequences resembling the -10 and -35 RNA polymerase-binding sites of the *B. subtilis* σ^E consensus sequence (Eichenberger *et al.*, 2003) are present upstream of *cot* β (at nucleotide positions -16 and -39 , relative to the beginning of the ORF). At stage III, Cot β -GFP was dispersed throughout the mother cell cytoplasm. When about 90% of the cells in the culture had reached stage IV, when forespores dehydrate and become refractile, we detected caps or rings of fluorescence around the forespores, as well as dispersed fluorescence in the mother cell cytoplasm (Fig. 3a). Released spores were surrounded by rings or caps of fluorescence. These rings were more pronounced after deconvolution (Fig. 3b). We argue that this fluorescence pattern is indicative of a coat localization for Cot β , as we have shown previously that GFP alone does not target any particular location in sporulating cells of *B. subtilis* (Little & Driks, 2001), and a cap or ring-like localization pattern is consistent with that of a coat or exosporium protein as demonstrated by several laboratories (Pogliano *et al.*, 1995; Bauer *et al.*, 1999; Asai *et al.*, 2001; Ozin *et al.*, 2001; Eichenberger *et al.*, 2003; van Ooij *et al.*, 2004; Kim *et al.*, 2006; R. Giorno, J. Bozue, M. Mallozzi *et al.*, unpublished data). However, when viewed in the context of the temporal progression of sporulation, this localization pattern differs in an interesting way from previous studies, in that there is a distinct stage during

Fig. 3. Fluorescence microscopic and Western blot analysis of Cot β -GFP in Ames and Sterne strain spores and sporulating cells. (a) Fluorescence microscopic localization of Cot β -GFP in strain RG134. Cells were grown at room temperature and analyzed 24 h (corresponding to stage III), 28 h (corresponding to stage IV), and 48 h (after spore release) after inoculation. Membrane staining (MEM) is shown in the upper panels, GFP fluorescence (GFP) is shown in the middle panels, and the merged (MERGE) image is shown in the lower panels. Forespore membranes are not stained due to the exclusion of the dye by the coat. The sporangia pictured in the merged images are cartooned below for clarity; spores and forespores are indicated by dark grey ovals, and mother cells are indicated by white rounded rectangles. Released spores in (a) were prepared by water washing. Fluorescence of Hypaque-purified released spores was indistinguishable from that of water-washed spores (data not shown). (b) Deconvolved micrograph from a spore from strain MGM37. (c) Spore coat extracts of *cot* β -gfp fusion bearing spores, from strains Ames-JAB-10 and RG134 (lanes 1 and 2, respectively), or from the wild-type Ames strain spores (lane 3) were fractionated using SDS-PAGE and transferred to a PVDF membrane, and then probed with anti-GFP antibodies. The asterisk (*) indicates a cross-reacting species, and arrowheads indicate the Cot β -GFP-specific doublet that is present in both fusion-bearing strains. (d) After sporulation for 2 days at room temperature, spores bearing *cot* β -gfp (from strain RG134, panels 1 and 2), or bearing *cot* β -gfp and a *cotE* mutation (from strain MGM37, panels 3 and 4) were collected and imaged by fluorescence (β -GFP) or IFM (α -GFP).

which Cot β is present but not yet deposited around the forespore. It would appear, therefore, that an as yet unidentified signal acts prior to stage IV to trigger Cot β deposition. This dynamic localization pattern is consistent with observations of other *B. anthracis* spore proteins (R. Giorno, J. Bozue, M. Mallozzi *et al.*, unpublished data).

Unexpectedly, we did not detect fluorescent rings in a version of the Ames strain bearing *cot β -gfp* (Ames-JAB-10) (data not shown). We confirmed that the *cot β* gene was transcribed in the Ames strain using RT-PCR analysis of mRNA collected from sporangia (data not shown). To confirm that Cot β -GFP was synthesized and present on spores, we extracted spore proteins from Ames-JAB-10 spores (as well as from RG134 spores), fractionated them using SDS-PAGE and analyzed them by Western blot analysis using anti-GFP antibodies. We detected Cot β -GFP-specific bands in both fusion-bearing strains (Fig. 3c, lanes 1 and 2). These bands migrate as a doublet of about 37 kDa, consistent with the expected size of the fusion protein (37.5 kDa). However, the intensity of the doublet bands from strain Ames-JAB-10 was substantially less than strain RG134, and three additional bands were present in the RG134 extracts (migrating between 45 and 50 kDa) which were not detected in Ames-JAB-10. These data suggest that, in the Ames strain, Cot β -GFP is made (albeit at a level below the limit of detection using fluorescence microscopy). The reason for the lower abundance of Cot β -GFP in cells from the Ames strain background is unknown. We note, however, that the adenine at position -13 (relative to the beginning of the ORF) in the Sterne strain is missing in the Ames strain, altering a putative σ^E consensus sequence. Possibly, this alteration reduces *cot β* transcription (and presumably Cot β abundance). The identities of the 45–50 kDa RG134-specific bands are unknown, although they apparently contain at least part of the fusion protein. A reasonable speculation is that they represent posttranslationally modified Cot β -GFP.

To determine the location of Cot β within the spore more precisely, we first performed IFM on Cot β -GFP-bearing spores (from strain RG134) using anti-GFP antibodies. While we could detect GFP fluorescence (Fig. 3d, panel 1), we did not detect a signal by IFM (Fig. 3d, panel 2) presumably because antibodies are prevented from binding by the presence of the exosporium. We then carried out IFM on cells bearing *cot β -gfp*, but where the exosporium is absent due to a mutation in *cotE* (in strain MGM37) (Giorno *et al.*, 2007). IFM analysis of strain MGM37 revealed rings of fluorescence. These rings, as well as those due to GFP-fluorescence, were interrupted at several positions around the spore (Fig. 3d, panels 3 and 4). We infer that Cot β -GFP is present beneath the exosporium surface (although it could also be present elsewhere in the exosporium) and, most likely, on the

coat surface. Interestingly, GFP-fluorescence in MGM37 spores was reduced relative to wild type (Fig. 3d, compare panels 1 and 3). Therefore, although some Cot β -GFP is assembled in the absence of CotE, a significant amount of Cot β -GFP likely assembles in a CotE-dependent manner.

Antigens specific to *B. cereus*-group spores (such as Cot β) could be used in future technologies to discriminate these spores from nonharmful spores, to improve medical diagnostics, food safety, and possibly biodefense. However, because Cot β is not required for disease in the model we analyzed, it is not an optimal marker for detecting a biological weapon, as it could be removed by a technically capable enemy. Nonetheless, because Cot β is conserved in food-borne disease causing strains of *B. cereus* it could serve as a marker for pathogens in the food supply.

In summary, we have characterized Cot β , a *B. cereus* group-specific protein in *B. anthracis* present on the spore coat surface. The abundance of Cot β in Sterne strain spores is significantly greater than in Ames strain spores, providing evidence that spore composition varies among *B. anthracis* strains. While Cot β does not detectably contribute to spore resistance, germination, or virulence (as measured by a murine intranasal challenge assay), it has a significant role in coat-surface architecture, similar to the *B. subtilis* coat-surface protein CotB (Chada *et al.*, 2003).

Acknowledgements

This work was supported by grants GM53989 and AI53365 from the National Institutes of Health (NIH) (A.D.), Department of the Army Award Number W81XWH-04-01-0307 (P.E.), and the United States Army Medical Research and Materiel Command under Projects 02-4-5C-018 (J.A.B.), 02-4-5C-023 (S.W.) and 04-0-IL-002 (S.W.). Research was conducted in compliance with the Animal Welfare Act, and other federal statutes and regulations relating to animals and experiments involving animals, and adheres to principles stated in the Guide for the Care and Use of Laboratory Animals, National Research Council, 1996. The facility where this research was conducted is fully accredited by the Association for Assessment and Accreditation of Laboratory Animal Care International. Opinions, interpretations, conclusions, and recommendations are those of the author and are not necessarily endorsed by the United States Army. We thank the Michigan Proteome Consortium for the identification of Cot β .

References

Aronson AI & Fitz-James P (1976) Structure and morphogenesis of the bacterial spore coat. *Bacteriol Rev* **40**: 360–402.

- Asai K, Takamatsu H, Iwano M, Kodama T, Watabe K & Ogasawara N (2001) The *Bacillus subtilis yabQ* gene is essential for formation of the spore cortex. *Microbiology* **147**: 919–927.
- Bauer T, Little S, Stover AG & Driks A (1999) Functional regions of the *Bacillus subtilis* spore coat morphogenetic protein CotE. *J Bacteriol* **181**: 7043–7051.
- Bozue JA, Parthasarathy N, Phillips LR, Cote CK, Fellows PF, Mendelson I, Shafferman A & Friedlander AM (2005) Construction of a rhamnose mutation in *Bacillus anthracis* affects adherence to macrophages but not virulence in guinea pigs. *Microb Pathogenesis* **38**: 1–12.
- Chada VG, Sanstad EA, Wang R & Driks A (2003) Morphogenesis of *Bacillus* spore surfaces. *J Bacteriol* **185**: 6255–6261.
- Donovan W, Zheng LB, Sandman K & Losick R (1987) Genes encoding spore coat polypeptides from *Bacillus subtilis*. *J Mol Biol* **196**: 1–10.
- Driks A (1999) *Bacillus subtilis* spore coat. *Microbiol Mol Biol R* **63**: 1–20.
- Eichenberger P, Jensen ST, Conlon EM *et al.* (2003) The sigmaE regulon and the identification of additional sporulation genes in *Bacillus subtilis*. *J Mol Biol* **327**: 945–972.
- Giorno R, Bozue J, Cote C *et al.* (2007) Morphogenesis of the *Bacillus anthracis* spore. *J Bacteriol* **189**: 691–705.
- Harwood CR & Cutting S (1990) *Molecular Biological Methods for Bacillus*. John Wiley and Sons, Sussex, UK.
- Henriques AO & Moran CP Jr (2006) Structure, assembly, and function of the spore surface layers. *Annu Rev Microbiol* **61**: 555–558.
- Henriques AO, Costa TV, Martins LO & Zilhao R (2004) The functional architecture and assembly of the coat. *Bacterial Spore Formers: Probiotics and Emerging Applications* (Ricca E, Henriques AO & Cutting SM, eds), pp. 65–86. Horizon Biosciences, Norfolk.
- Hilbert DW & Piggot PJ (2004) Compartmentalization of gene expression during *Bacillus subtilis* spore formation. *Microbiol Mol Biol R* **68**: 234–262.
- Holt SC & Leadbetter ER (1969) Comparative ultrastructure of selected aerobic spore-forming bacteria: a freeze-etching study. *Bacteriol Rev* **33**: 346–378.
- Isticato R, Cangiano G, Tran HT, Ciabattini A, Medagliani D, Oggioni MR, De Felice M, Pozzi G & Ricca E (2001) Surface display of recombinant proteins on *Bacillus subtilis* spores. *J Bacteriol* **183**: 6294–6301.
- Kim H, Hahn M, Grabowski P, McPherson DC, Otte MM, Wang R, Ferguson CC, Eichenberger P & Driks A (2006) The *Bacillus subtilis* spore coat protein interaction network. *Mol Microbiol* **59**: 487–502.
- Kim HS, Sherman D, Johnson F & Aronson AI (2004) Characterization of a major *Bacillus anthracis* spore coat protein and its role in spore inactivation. *J Bacteriol* **186**: 2413–2417.
- Klobutcher LA, Ragkousi K & Setlow P (2006) The *Bacillus subtilis* spore coat provides “eat resistance” during phagocytic predation by the protozoan *Tetrahymena thermophila*. *Proc Natl Acad Sci USA* **103**: 165–170.
- Koehler TM, Dai Z & Kaufman-Yarbray M (1994) Regulation of the *Bacillus anthracis* protective antigen gene: CO₂ and a *trans*-acting element activate transcription from one of two promoters. *J Bacteriol* **176**: 586–595.
- Kuwana R, Kasahara Y, Fujibayashi M, Takamatsu H, Ogasawara N & Watabe K (2002) Proteomics characterization of novel spore proteins of *Bacillus subtilis*. *Microbiology* **148**: 3971–3982.
- Lai EM, Phadke ND, Kachman MT, Giorno R, Vazquez S, Vazquez JA, Maddock JR & Driks A (2003) Proteomic analysis of the spore coats of *Bacillus subtilis* and *Bacillus anthracis*. *J Bacteriol* **185**: 1443–1454.
- Leighton TJ & Doi RH (1971) The stability of messenger ribonucleic acid during sporulation in *Bacillus subtilis*. *J Biol Chem* **246**: 3189–3195.
- Little S & Driks A (2001) Functional analysis of the *Bacillus subtilis* morphogenetic spore coat protein CotE. *Mol Microbiol* **42**: 1107–1120.
- Little SF & Knudson GB (1986) Comparative efficacy of *Bacillus anthracis* live spore vaccine and protective antigen vaccine against anthrax in the guinea pig. *Infect Immun* **52**: 509–512.
- Losick R, Youngman P & Piggot PJ (1986) Genetics of endospore formation in *Bacillus subtilis*. *Annu Rev Genet* **20**: 625–669.
- Lyons CR, Lovchik J, Hutt J, Lipscomb MF, Wang E, Heninger S, Berliba L & Garrison K (2004) Murine model of pulmonary anthrax: kinetics of dissemination, histopathology, and mouse strain susceptibility. *Infect Immun* **72**: 4801–4809.
- McPherson DC, Kim H, Hahn M, Wang R, Grabowski P, Eichenberger P & Driks A (2005) Characterization of the *Bacillus subtilis* spore morphogenetic coat protein CotO. *J Bacteriol* **187**: 8278–8290.
- Mendelson I, Tobery S, Scorpio A, Bozue J, Shafferman A & Friedlander AM (2004) The NheA component of the non-hemolytic enterotoxin of *Bacillus cereus* is produced by *Bacillus anthracis* but is not required for virulence. *Microb Pathogenesis* **37**: 149–154.
- Ozin AJ, Samford CS, Henriques AO & Moran CP Jr (2001) SpoVID guides SafA to the spore coat in *Bacillus subtilis*. *J Bacteriol* **183**: 3041–3049.
- Perez-Casal J, Caparon MG & Scott JR (1991) Mry, a *trans*-acting positive regulator of the M protein gene of *Streptococcus pyogenes* with similarity to the receptor proteins of two-component regulatory systems. *J Bacteriol* **173**: 2617–2624.
- Plomp M, Leighton TJ, Wheeler KE & Malkin AJ (2005) The high-resolution architecture and structural dynamics of *Bacillus* spores. *Biophys J* **88**: 603–608.
- Plomp M, Leighton TJ, Wheeler KE, Hill HD & Malkin AJ (2007) *In vitro* high-resolution structural dynamics of single germinating bacterial spores. *Proc Natl Acad Sci USA* **104**: 9644–9649.
- Pogliano J, Osborne N, Sharp MD, Abanes-De Mello A, Perez A, Sun YL & Pogliano K (1999) A vital stain for studying membrane dynamics in bacteria: a novel mechanism controlling septation during *Bacillus subtilis* sporulation. *Mol Microbiol* **31**: 1149–1159.

- Pogliano K, Harry E & Losick R (1995) Visualization of the subcellular location of sporulation proteins in *Bacillus subtilis* using immunofluorescence microscopy. *Mol Microbiol* **18**: 459–470.
- Sambrook J, Fritsch EF & Maniatis T (1989) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Santo LY & Doi RH (1974) Ultrastructural analysis during germination and outgrowth of *Bacillus subtilis* spores. *J Bacteriol* **120**: 475–481.
- Setlow P (2006) Spores of *Bacillus subtilis*: their resistance to and killing by radiation, heat and chemicals. *J Appl Microbiol* **101**: 514–525.
- van Ooij C, Eichenberger P & Losick R (2004) Dynamic patterns of subcellular protein localization during spore coat morphogenesis in *Bacillus subtilis*. *J Bacteriol* **186**: 4441–4448.
- Wang R, Krishnamurthy SN, Jeong J-S, Driks A, Mehta M & Gingras BA (2007) Finger-printing species and strains of *Bacillus* spores by distinctive coat surface morphology. *Langmuir* **23**: 10230–10234.
- Warth AD, Ohye DF & Murrell WG (1963) The composition and structure of bacterial spores. *J Cell Biol* **16**: 579–592.
- Young SB & Setlow P (2003) Mechanisms of killing of *Bacillus subtilis* spores by hypochlorite and chlorine dioxide. *J Appl Microbiol* **95**: 54–67.
- Zheng LB & Losick R (1990) Cascade regulation of spore coat gene expression in *Bacillus subtilis*. *J Mol Biol* **212**: 645–660.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1. Primers.

Appendix S1. Generation of recombinant Sterne strains.

Please note: Wiley-Blackwell is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.