

Table I. Plasmids.

plasmid	description	Source
pLG Δ 312S Δ SS	<i>CYC1-lacZ URA3</i> 2 μ	Bowdish <i>et al</i> , 1993
pKB144	T ₄ C- <i>CYC1-lacZ URA3</i> 2 μ	Bowdish <i>et al</i> , 1995
pKB148	URS1(<i>IME2,3'</i> →5' ^(a))- <i>CYC1-lacZ URA3</i> 2 μ	Bowdish <i>et al</i> , 1995
pKB150	[URS1(<i>IME2,3'</i> →5' ^(a))] ₂ - <i>CYC1-lacZ URA3</i> 2 μ	Bowdish <i>et al</i> , 1995
pKB160	T ₄ C-URS1(<i>IME2,5'</i> →3')- <i>CYC1-lacZ URA3</i> 2 μ	Bowdish <i>et al</i> , 1995
pKBmtURS1	T ₄ C-mtURS1- <i>CYC1-lacZ URA3</i> 2 μ	this study
pLG(URS1)-52	[URS1(<i>IME2,1x3'</i> →5'+5x5' <i>→3'</i>)] ₆ - <i>CYC1-lacZ URA3</i> 2 μ	this study
pRS314	<i>TRP1 CEN</i>	Sikorski and Hieter, 1989
pRS314- <i>HAC1</i> ⁱ	HA- <i>HAC1</i> ⁱ <i>TRP1 CEN</i>	this study
pRS316	<i>URA3 CEN</i>	Sikorski and Hieter, 1989
pRS316- <i>HAC1</i> ⁱ	HA- <i>HAC1</i> ⁱ <i>URA3 CEN</i>	Schröder <i>et al</i> , 2000
p2UG	GRE ₃ - <i>CYC1 URA3</i> 2 μ	Schena <i>et al</i> , 1991
pG-N795	<i>GPD-N795</i> ^(b) <i>TRP1</i> 2 μ	Schena <i>et al</i> , 1991
p2UG-HA- <i>HAC1</i> ⁱ	GRE ₃ - <i>CYC1-HA-HAC1</i> ⁱ <i>URA3</i> 2 μ	this study

(a) Orientation was designated on the basis of the consensus URS1 sequence.

(b) N795 denotes the 795 amino acid rat glucocorticoid receptor (Schena *et al*, 1991).