

# Isotopes of nitrogen on Mars: Atmospheric measurements by Curiosity's mass spectrometer

Michael H. Wong,<sup>1,2</sup> Sushil K. Atreya,<sup>1</sup> Paul N. Mahaffy,<sup>3</sup> Heather B. Franz,<sup>3</sup> Charles Malespin,<sup>3</sup> Melissa G. Trainer,<sup>3</sup> Jennifer C. Stern,<sup>3</sup> Pamela G. Conrad,<sup>3</sup> Heidi L. K. Manning,<sup>4</sup> Robert O. Pepin,<sup>5</sup> Richard H. Becker,<sup>5</sup> Christopher P. McKay,<sup>6</sup> Tobias C. Owen,<sup>7</sup> Rafael Navarro-González,<sup>8</sup> John H. Jones,<sup>9</sup> Bruce M. Jakosky,<sup>10</sup> and Andrew Steele<sup>11</sup>

Received 3 September 2013; revised 18 November 2013; accepted 18 November 2013; published 9 December 2013.

[1] The Sample Analysis at Mars (SAM) instrument suite on the Mars Science Laboratory (MSL) measured a Mars atmospheric  $^{14}\text{N}/^{15}\text{N}$  ratio of  $173 \pm 11$  on sol 341 of the mission, agreeing with Viking's measurement of  $168 \pm 17$ . The MSL/SAM value was based on Quadrupole Mass Spectrometer measurements of an enriched atmospheric sample, with  $\text{CO}_2$  and  $\text{H}_2\text{O}$  removed. Doubly ionized nitrogen data at  $m/z$  14 and 14.5 had the highest signal/background ratio, with results confirmed by  $m/z$  28 and 29 data. Gases in SNC meteorite glasses have been interpreted as mixtures containing a Martian atmospheric component, based partly on distinctive  $^{14}\text{N}/^{15}\text{N}$  and  $^{40}\text{Ar}/^{14}\text{N}$  ratios. Recent MSL/SAM measurements of the  $^{40}\text{Ar}/^{14}\text{N}$  ratio ( $0.51 \pm 0.01$ ) are incompatible with the Viking ratio ( $0.35 \pm 0.08$ ). The meteorite mixing line is more consistent with the atmospheric composition measured by Viking than by MSL. **Citation:** Wong, M. H., et al. (2013), Isotopes of nitrogen on Mars: Atmospheric measurements by Curiosity's mass spectrometer, *Geophys. Res. Lett.*, 40, 6033–6037, doi:10.1002/2013GL057840.

## 1. Introduction

[2] Isotope ratios are excellent tracers of atmospheric source and loss processes over planetary timescales. In the case of Martian oxygen and carbon, atmospheric evolution models must account for the exchange between atmospheric and surface/subsurface reservoirs [Jakosky, 1991; Webster et al., 2013]. The potential for exchange of nitrogen between the atmosphere and surface of Mars is not currently understood, but such exchange would be very important to the study of Martian habitability that is the focus of the Mars Science Laboratory (MSL) mission [Grotzinger et al., 2012].

[3] The atmospheric nitrogen isotopic ratio is difficult to measure. Very weak absorption lines are produced by the dipole forbidden vibrational-rotational transitions of ground state  $\text{N}_2$ , so the spectroscopic techniques used to measure isotope ratios in other molecules cannot be applied to molecular nitrogen. Of the 10 science instruments of the MSL payload [Grotzinger et al., 2012], Sample Analysis at Mars's (SAM's) mass spectrometer provides the best capability to measure the nitrogen isotopic ratio [Mahaffy et al., 2012]. Independent measurements of  $^{14}\text{N}/^{15}\text{N}$  were conducted by mass spectrometers on Viking Lander 2 [Owen et al., 1977; Owen, 1992] and the two Viking lander aeroshells [Nier and McElroy, 1977]. The surface and descent measurements found  $^{14}\text{N}/^{15}\text{N}$  of  $170 \pm 15$  and  $168 \pm 17$ , respectively.

## 2. Experiments

[4] SAM is a highly capable and configurable instrument suite for analyzing gases either directly sampled from the atmosphere or evolved from solid samples [Mahaffy et al., 2012]. SAM/Quadrupole Mass Spectrometer (QMS) determines isotopic ratios based on ratios of detector counting rates. In direct atmospheric experiments,  $\text{CO}^+$  ions (from  $\text{CO}_2$ ) interfere at  $m/z$  28 and 29 (where the  $\text{N}_2^+$  parent ions are detected). But there is no  $\text{CO}^{++}$  contribution at  $m/z$  14, so we can derive  $^{14}\text{N}/^{15}\text{N}$  from direct atmospheric experiments using the  $m/z$  14/14.5 count ratio. In enrichment experiments, chemical scrubbers remove  $\text{CO}_2$ ,  $\text{H}_2\text{O}$ , and other species with chemical affinity to the scrubber material. The enrichment process allows determination of the isotopic ratio using  $m/z$  28/29 and 14/29 count ratios. We further boost signal level (compared to the normal dynamic mode) using a semistatic mode, where the pumping rate out of the QMS is reduced. Figure 1 shows a mass spectrum from

Additional supporting information may be found in the online version of this article.

<sup>1</sup>Department of Atmospheric, Oceanic, and Space Sciences, University of Michigan, Ann Arbor, Michigan, USA.

Corresponding author: M. H. Wong, Department of Atmospheric, Oceanic, and Space Sciences, University of Michigan, Space Research Bldg., 2455 Hayward St., Ann Arbor, MI 48109-2143, USA. (mike.wong@umich.edu)

©2013 The Authors. *Geophysical Research Letters* published by Wiley on behalf of the American Geophysical Union.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made. 0094-8276/13/10.1002/2013GL057840

<sup>2</sup>Astronomy Department, University of California, Berkeley, California, USA.

<sup>3</sup>Code 699, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA.

<sup>4</sup>Department of Physics, Concordia College, Moorhead, Minnesota, USA.

<sup>5</sup>School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota, USA.

<sup>6</sup>NASA Ames Research Center, Moffet Field, California, USA.


<sup>7</sup>Institute for Astronomy, University of Hawai'i at Mānoa, Honolulu, Hawaii, USA.

<sup>8</sup>Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, México City, México.

<sup>9</sup>NASA Johnson Space Center, Houston, Texas, USA.

<sup>10</sup>LASP, University of Colorado Boulder, Colorado, USA.

<sup>11</sup>Geophysical Laboratory, Carnegie Institution of Washington, Washington, District of Columbia, USA.



**Figure 1.** High-resolution mass spectrum from the enrichment experiment on sol 232. Sample spectrum is an average of data from enrichment cycle 9 (see Figure 2). Nitrogen ions are present at  $m/z$  14, 14.5, 28, and 29. The singly ionized  $N_2$  “parent ion” appears in the Mars data at  $m/z$  28 (for  $^{14}N^{14}N$ ) and at  $m/z$  29 (for  $^{15}N^{14}N$ ). A third parent ion,  $^{15}N^{15}N$ , is so rare as to produce only a small signal below the background level (difference between red and blue curves). For a comparison of the  $m/z$  14–15 range in the direct atmospheric experiment, see *Wong et al.* [2013].

an enrichment experiment in dynamic mode, spanning the relevant  $m/z$  range for nitrogen analysis.


[5] On Jupiter, the nitrogen isotopic ratio was determined using Galileo Probe Mass Spectrometer measurements of  $NH_3^+$ , to avoid interferences at  $m/z$  17 and 18 from water [Owen *et al.*, 2001; Wong *et al.*, 2004]. We use a similar approach in the MSL/SAM analysis, although there is an additional complication because the count ratio of  $m/z$  14/14.5 includes contributions from two nitrogen daughter ions,  $N^+$  and  $N_2^+$ .

[6] To extract the nitrogen isotopic ratio, we first measure a splitting fraction (derived from SAM/QMS preflight cali-

bration data) that gives the relative contributions from these two daughter ions:

$$\beta = \frac{N^+}{N^+ + N_2^+} = \frac{2 \left[ \frac{m14}{m14.5} \right] - {}^{14}N/{}^{15}N}{2 \left[ \frac{m14}{m14.5} \right] + 1}, \quad (1)$$

where the count rates at a mass/charge ratio of  $m/z$   $X$  are represented as  $mX$ , and  $^{14}N/{}^{15}N$  is independently measured in the calibration gas. We find  $\beta = 0.404 \pm 0.033$  based on five experiments [Wong *et al.*, 2013]. The  $^{14}N/{}^{15}N$  isotopic ratio is then derived from the splitting fraction



**Figure 2.** Visual narrative of the sol 232 enrichment experiment. Parent and daughter nitrogen ions are shown as colored points. Background measurements (shaded) are used to characterize the background signal that must be subtracted from the data before calculating count ratios. Detector corrections have been applied to the data (see supporting information). In each of 10 enrichment cycles, an atmospheric sample was introduced, and  $CO_2$  and  $H_2O$  were removed by chemical scrubbers [Mahaffy *et al.*, 2012]. The enriched sample from one cycle was then combined with a new injection of Mars atmosphere and scrubbed to produce the next sample. Signal at  $m/z$  28 decreases in enrichment cycle 2 and later due to detector saturation at high source pressures.

**Table 1.** Mars Nitrogen Isotopic Ratios Measured in Atmospheric N<sub>2</sub>

MSL Sol <sup>a</sup>	<sup>14</sup> N/ <sup>15</sup> N	δ <sup>15</sup> N(‰) <sup>b</sup>	Count Ratio	Limiting S/BG <sup>c</sup>	Experiment Type
341 (night)	173 ± 11	572 ± 82	m14/m14.5	≤ 388	Enrichment (MSL/SAM, semistatic mode)
341 (night)	175 ± 21	554 ± 187	m28/m29	3.4	Enrichment (MSL/SAM, semistatic mode)
232 (night)	178 ± 12	528 ± 103	m14/m14.5	≤ 97	Enrichment (MSL/SAM, dynamic mode)
232 (night)	179 ± 11	519 ± 91	m28/m29	≤ 13	Enrichment (MSL/SAM, dynamic mode)
19 (night)	150 ± 60	813 ± 725	m14/m14.5	2	Direct atmospheric (MSL/SAM)
45 (night)	151 ± 27	798 ± 324	m14/m14.5	9	Direct atmospheric (MSL/SAM)
77 (night)	169 ± 19	611 ± 183	m14/m14.5	8	Direct atmospheric (MSL/SAM)
278 (day)	173 ± 27	572 ± 245	m14/m14.5	14	Direct atmospheric (MSL/SAM)
284 (night)	172 ± 38	582 ± 345	m14/m14.5	17	Direct atmospheric (MSL/SAM)
292 (night)	177 ± 14	537 ± 123	m14/m14.5	20	Direct atmospheric (MSL/SAM)
321 (day)	167 ± 14	628 ± 140	m14/m14.5	12	Direct atmospheric (MSL/SAM)
Testbed <sup>d</sup>	270 ± 28	8 ± 104	m14/m14.5	≤ 33	Testbed enrichment TID 50765
Comparison measurements					
Viking aeroshell mass spec	168 ± 17	619 ± 182	m28/m29	< 50	Viking, 125 km [Nier and McElroy, 1977]
Viking lander mass spec	165 ± 17	649 ± 183	m28/m29 <sup>e</sup>	? <sup>e</sup>	Viking, surface [Owen et al., 1977]
Viking lander mass spec	170 ± 15	600 ± 155	m28/m29 <sup>e</sup>	? <sup>e</sup>	Viking, surface [Owen, 1992]

<sup>a</sup>MSL landing occurred on sol 0 at 15:03 local mean solar time or UTC 2012-08-06 05:17.

<sup>b</sup>δ<sup>15</sup>N (‰) = 1000(R/R<sub>AIR</sub> - 1), where R = <sup>15</sup>N/<sup>14</sup>N in the sample, and R<sub>AIR</sub> = 0.003676 [Coplen et al., 2002].

<sup>c</sup>Signal to background level (S/BG) is “limiting” because it refers to the count rate with the poorest S/BG (e.g., m/z 14 in the case of the m/z 14/14.5 ratio). The “≤” symbol means that the S/BG level changes over the course of the experiment, and the best S/BG ratio over the full experiment range is reported.

<sup>d</sup>Testbed experiment is listed to confirm SAM accuracy with a gas sample of known isotopic composition (δ<sup>15</sup>N = 0‰, <sup>14</sup>N/<sup>15</sup>N = 272). Analysis method is the same as for flight data.

<sup>e</sup>Viking lander mass spectrometer publications do not describe count ratios used for isotopic determination or S/BG levels in the data. Experiments used enriched atmospheric samples.

β and the flight m/z 14/14.5 count ratio (see supporting information):

$$^{14}\text{N}/^{15}\text{N} = 2(1 - \beta) \left[ \frac{\text{m}14}{\text{m}14.5} \right] - \beta. \quad (2)$$

[7] Additional count ratios can be used to measure nitrogen isotopes in the enrichment experiments. Measurement of the <sup>14</sup>N/<sup>15</sup>N ratio directly from the parent ions is difficult because the signal level is very high at m/z 28 over most of the enrichment experiment; in fact, Figure 2 shows that the m/z 28 signal is near saturation in the first enrichment cycle (labeled 1a and 1b). The m/z 28 counting rate decreases in enrichment cycles 2–4 due to saturation, with 0 counts/s in the subsequent cycles. The isotope ratio can also be derived from m/z 14/29 data, but different voltage frequencies used above and below m/z 20 [Mahaffy et al., 2012] introduce complications that have not yet been fully resolved.

### 3. Results

[8] Table 1 lists nitrogen isotopic results from SAM up through MSL sol 360 (Planetary Data System release 4). Uncertainties include all sources of error (see supporting information). Specifically, we include uncertainties from detector corrections, background subtraction, statistical noise, and splitting fractions. SAM background corrections are significant, so the limiting S/BG level is listed as a measure of the confidence in each measurement. Where known, S/BG levels from other instruments are also given for comparison.

[9] The most reliable MSL/SAM measurement of <sup>14</sup>N/<sup>15</sup>N comes from m/z 14/14.5 in the semistatic enrichment experiment, where a S/BG ratio of 388 was achieved. Results at low S/BG have much more scatter than results at S/BG > 50 (see supporting information), suggesting that imperfect background corrections dominate the uncertainty in the


direct atmospheric measurements. The largest single source of uncertainty in the isotopic ratio is the uncertainty in β (equation (1) and Figure S1). The range of <sup>14</sup>N/<sup>15</sup>N ratios derived from MSL/SAM data falls completely within the error bars of Viking measurements.

### 4. Discussion

[10] Nonterrestrial nitrogen isotopic ratios in Martian meteorite samples have been interpreted as tracers of atmospheric gas, trapped within the meteorites at their time of ejection from Mars [Bogard and Johnson, 1983]. Figure 3 compares laboratory analyses of gases evolved from meteorite samples to in situ measurements of the Mars atmosphere by Viking and MSL. Selected step-heating gas releases in the terrestrial laboratory experiments can be used to define mixing lines between the compositional measurements. The choice of steps for analysis (black symbols and references in figure caption) was intended to eliminate contributions from terrestrial air or organic contamination. For comparison, we also show meteorite data (red symbols) and linear fits based on the total gas released in each experiment. These total gas release data are free from uncertainties introduced by subjective interpretation of stepped heating data. The mixing line slope is not significantly affected by uncertainties in cosmic ray exposure ages [Nyquist et al., 2001; Schwenzner et al., 2007].

[11] There is a clear disagreement between the Ar/N measurement by MSL/SAM of 0.51 ± 0.01 [Mahaffy et al., 2013] and by multiple Viking instruments of 0.35 ± 0.08 [Oyama and Berdahl, 1977]. All of the meteorite mixing lines are consistent with Viking composition and inconsistent with MSL.

[12] Three possible explanations for the disagreements can be considered: that the MSL composition is wrong, that the Viking composition is wrong, or that the Martian atmospheric composition is variable. Internal and peer reviews



**Figure 3.** Pseudo-three-isotope plots at two different magnifications (comparing  $^{14}\text{N}$ ,  $^{15}\text{N}$ , and  $^{40}\text{Ar}$ ) showing Mars meteorite and atmosphere measurements and linear fits. The meteorite data suggest mixing between at least two components: Martian atmosphere and terrestrial atmosphere. Black meteorite data include only selected temperature steps in stepped heating gas release experiments. Red data points include the total gas released at all temperatures. Meteorite data are from *Becker and Pepin* [1984, EETA 79001A samples C1 and C2], *Wiens et al.* [1986, sample C3], *Marti et al.* [1995, Zagami], *Chennaoui Aoudjehane et al.* [2012, Tissint], and *Avicé* [2012, Tissint]. Viking Ar/N ratio is from *Oyama and Berdahl* [1977], and Viking  $\delta^{15}\text{N}$  is from *Nier and McElroy* [1977]. MSL values are from SAM/QMS measurements, with  $\delta^{15}\text{N}$  described in this work, and Ar/N is from *Mahaffy et al.* [2013]. Linear fits included uncertainties in both Ar/N and  $\delta^{15}\text{N}$ , following the method of *York et al.* [2004], assuming uncorrelated errors.

of the MSL/SAM data argue against the first possibility at this time [*Mahaffy et al.*, 2013]. Similarly, the Viking composition seems highly robust, with good agreement among multiple instruments. The MSL  $^{14}\text{N}/^{15}\text{N}$  ratio falls easily within the error bars of the Viking measurements by both lander and aeroshell mass spectrometer experiments. So the primary disagreement is with the Ar/N ratio. This ratio was repeatedly measured by both the Viking lander mass spectrometer [*Owen et al.*, 1977] and the gas exchange experiment [*Oyama and Berdahl*, 1977]. Unfortunately, Viking flight data are no longer available, so we cannot repeat the same validation we applied to the MSL data. Therefore,

the possibility that Viking Ar/N measurements were erroneous can only be speculation.

[13] There is currently no satisfactory mechanism that could explain a variable Ar/N ratio in the Martian atmosphere. Adsorbed  $\text{CO}_2$  and  $\text{H}_2\text{O}$  in the regolith make up a large volatile reservoir that exchanges over time with the atmosphere and polar caps [e.g., *Fanale et al.*, 1982; *Fanale and Jakosky*, 1982]. Nitrogen and argon are not thought to participate in this cycle, although these gases do adsorb onto Martian soil [*Ballou et al.*, 1978]. A previously unknown surface-atmosphere exchange mechanism would need to involve very large column density changes: a release of up to  $1 \text{ kg m}^{-2}$  Ar or sequestration of up to  $0.7 \text{ kg m}^{-2}$   $\text{N}_2$ , prior to 2012. Column density changes might be thermally driven by global warming of 0.65 K since Viking [*Fenton et al.*, 2007], or related to a reduction of polar cap mass [*Malin et al.*, 2001], but MSL measurements from early spring to late summer on Mars do not support any variation in Ar/N [*Atreya et al.*, 2013; *Trainer et al.*, 2013].

[14] Heavy nitrogen in the atmosphere of Mars is enriched relative to both the protosolar isotopic ratio and the ratio in the terrestrial and Venusian atmospheres (Table 2), as well as to the primordial Mars atmosphere as suggested by the meteorite ALH84001 with  $\delta^{15}\text{N} = 4 \pm 2 \text{ ‰}$  [*Marti and Mathew*, 2000]. The enrichment of  $^{15}\text{N}$  is due to preferential escape of  $^{14}\text{N}$  from the Martian atmosphere [e.g., *McElroy et al.*, 1976; *Fox and Dalgarno*, 1983; *Fox and Hać*, 1997; *Chassefière and Leblanc*, 2004]. Some models of sputtering and photochemical escape of nitrogen to space suggest that exchange with surface/interior reservoirs is needed to buffer the nitrogen against excessive fractionation due to escape [*Jakosky et al.*, 1994; *Zent et al.*, 1994], leading to a loss of  $\sim 90\%$  of atmospheric nitrogen over time [*Jakosky and Phillips*, 2001]. The detailed history and mechanisms of exchange between atmosphere and surface/interior are poorly understood for nitrogen, compared to similar processes that fractionate atmospheric carbon and oxygen [*Webster et al.*, 2013], via exchange between polar ices, carbonates, and regolith adsorption [*Jakosky*, 1991; *Fanale et al.*, 1992].

[15] SAM may provide insight into these processes by investigating nitrogen-containing compounds in solid samples. Reduced and oxidized compounds have tentatively been identified in evolved gases, including N, NO, HCN, CN, and  $\text{CH}_3\text{CN}$  [*Stern et al.*, 2013; *Navarro-González et al.*, 2013]. The origin of the detected nitrogen could include terrestrial sources within the SAM instrument or the Curiosity sample chain, exogenous nitrogen delivered by meteorites, or indigenous Martian inorganic and organic matter. Exchange between solid and atmospheric reservoirs

**Table 2.** Solar System Nitrogen Isotopic Ratios

Measurement	$^{14}\text{N}/^{15}\text{N}$	$\delta^{15}\text{N}$ (‰)	Reference
Protosolar/primordial			
Sun (Genesis)	$441 \pm 5$	$-383 \pm 8$	<i>Marty et al.</i> [2011]
Jupiter	$434 \pm 7.5$	$-373 \pm 11$	<i>Owen et al.</i> [2001]
Interstellar medium	$450 \pm 98$	$-395 \pm 168$	<i>Dahmen et al.</i> [1995]
Secondary atmospheres			
Mars (MSL)	$173 \pm 11$	$572 \pm 82$	(This work)
Titan	$183 \pm 5$	$487 \pm 42$	<i>Niemann et al.</i> [2005]
Venus	$272 \pm 54$	$0 \pm 250$	<i>Hoffman et al.</i> [1979]
Earth	272	0	

could occur as organic surface nitrogen is photolytically or radiolytically decomposed to N<sub>2</sub> or oxidized to inorganic nitrogen in the form of nitrates, which are also converted to N<sub>2</sub> by the current impact flux [Manning et al., 2008].

[16] **Acknowledgments.** The authors thank the MSL Team for successful operation of the mission, Jane Fox and Richard Quinn for thorough and helpful reviews, and Bernard Marty and Guillaume Avice for insightful comments on the analysis of Tissint. This research was supported by the NASA Mars Science Laboratory Project.

[17] The Editor thanks Richard Quinn and Jane Fox for their assistance in evaluating this manuscript.

## References

- Atreya, S. K., S. W. Squyres, P. R. Mahaffy, L. A. Leshin, H. B. Franz, M. G. Trainer, M. H. Wong, C. P. McKay, R. Navarro-González, and MSL Science Team (2013), MSL/SAM measurements of non-condensable volatiles in the atmosphere of Mars—Possibility of seasonal variations, *LPI Contrib.*, 1719, 2130.
- Avice, G. (2012), Analyse géochimique d'une nouvelle météorite martienne: Tissint, Master's thesis, École Nationale Supérieure de Géologie, Vandœuvre-lès-Nancy, France.
- Ballou, E. V., P. C. Wood, T. Wydeven, M. E. Lehwalt, and R. E. Mack (1978), Chemical interpretation of Viking Lander 1 life detection experiment, *Nature*, 271, 644–645.
- Becker, R. H., and R. O. Pepin (1984), The case for a Martian origin of the shergottites—Nitrogen and noble gases in EETA 79001, *Earth Planet. Sci. Lett.*, 69, 225–242.
- Bogard, D. D., and P. Johnson (1983), Martian gases in an Antarctic meteorite?, *Science*, 221, 651–654.
- Chassefière, E., and F. Leblanc (2004), Mars atmospheric escape and evolution; interaction with the solar wind, *Planet. Space Sci.*, 52, 1039–1058.
- Chennaoui Aoudjehane, H., et al. (2012), Tissint Martian meteorite: A fresh look at the interior, surface, and atmosphere of Mars, *Science*, 338, 785–788.
- Coplen, T. B., et al. (2002), Isotope-abundance variations of selected elements (IUPAC technical report), *Pure Appl. Chem.*, 74, 1987–2017.
- Dahmen, G., T. L. Wilson, and F. Matteucci (1995), The nitrogen isotope abundance in the galaxy. I: The galactic disk gradient, *Astron. Astrophys.*, 295, 194–198.
- Fanale, F. P., and B. M. Jakosky (1982), Regolith-atmosphere exchange of water and carbon dioxide on Mars—Effects on atmospheric history and climate change, *Planet. Space Sci.*, 30, 819–831.
- Fanale, F. P., S. E. Postawko, J. B. Pollack, M. H. Carr, and R. O. Pepin (1992), Mars—Epochal climate change and volatile history, in *Mars*, edited by H. H. Kieffer et al., pp. 1135–1179, Univ. of Ariz. Press, Tucson, Ariz.
- Fanale, F. P., W. B. Banerdt, R. S. Saunders, L. A. Johansen, and J. R. Salvail (1982), Seasonal carbon dioxide exchange between the regolith and atmosphere of Mars: Experimental and theoretical studies, *J. Geophys. Res.*, 87, 10,215–10,225.
- Fox, J. L., and A. Hać (1997), The <sup>15</sup>N/<sup>14</sup>N isotope fractionation in dissociative recombination of N<sub>2</sub><sup>+</sup>, *J. Geophys. Res.*, 102, 9191–9204.
- Fox, J. L., and A. Dalgarno (1983), Nitrogen escape from Mars, *J. Geophys. Res.*, 88, 9027–9032.
- Fenton, L. K., P. E. Geissler, and R. M. Haberle (2007), Global warming and climate forcing by recent albedo changes on Mars, *Nature*, 446, 646–649.
- Grotzinger, J. P., et al. (2012), Mars Science Laboratory mission and science investigation, *Space Sci. Rev.*, 170, 5–56.
- Hoffman, J. H., R. R. Hodges, M. B. McElroy, T. M. Donahue, and M. Kolpin (1979), Composition and structure of the Venus atmosphere—Results from Pioneer Venus, *Science*, 205, 49–52.
- Jakosky, B. M. (1991), Mars volatile evolution: Evidence from stable isotopes, *Icarus*, 94, 14–31.
- Jakosky, B. M., and R. J. Phillips (2001), Mars' volatile and climate history, *Nature*, 412, 237–244.
- Jakosky, B. M., R. O. Pepin, R. E. Johnson, and J. L. Fox (1994), Mars atmospheric loss and isotopic fractionation by solar-wind-induced sputtering and photochemical escape, *Icarus*, 111, 271–288.
- Mahaffy, P. R., et al. (2012), The sample analysis at Mars investigation and instrument suite, *Space Sci. Rev.*, 170, 401–478.
- Mahaffy, P. R., et al. (2013), Abundance and isotopic composition of gases in the Martian atmosphere from the Curiosity rover, *Science*, 341(263), 263–266.
- Malin, M. C., M. A. Caplinger, and S. D. Davis (2001), Observational evidence for an active surface reservoir of solid carbon dioxide on Mars, *Science*, 294, 2146–2148.
- Manning, C. V., C. P. McKay, and K. J. Zahnle (2008), The nitrogen cycle on Mars: Impact decomposition of near-surface nitrates as a source for a nitrogen steady state, *Icarus*, 197, 60–64.
- Marti, K., and K. J. Mathew (2000), Ancient Martian nitrogen, *Geophys. Res. Lett.*, 27, 1463–1466.
- Marti, K., J. S. Kim, A. N. Thakur, T. J. McCoy, and K. Keil (1995), Signatures of the Martian atmosphere in glass of the Zagami meteorite, *Science*, 267, 1981–1984.
- Marty, B., M. Chaussidon, R. C. Wiens, A. J. G. Jurewicz, and D. S. Burnett (2011), A <sup>15</sup>N-poor isotopic composition for the solar system as shown by Genesis solar wind samples, *Science*, 332, 1533–1536.
- McElroy, M. B., Y. L. Yung, and A. O. Nier (1976), Isotopic composition of nitrogen: Implications for the past history of Mars atmosphere, *Science*, 194, 70–72.
- Navarro-González, R., et al. (2013), Possible detection of nitrates on Mars by the Sample Analysis at Mars (SAM) instrument, *LPI Contrib.*, 1719, 2648.
- Nier, A. O., and M. B. McElroy (1977), Composition and structure of Mars' upper atmosphere: Results from the neutral mass spectrometers on Viking 1 and 2, *J. Geophys. Res.*, 82, 4341–4349.
- Niemann, H. B., et al. (2005), The abundances of constituents of Titan's atmosphere from the GCMS instrument on the Huygens probe, *Nature*, 438, 779–784.
- Nyquist, L. E., D. D. Bogard, C.-Y. Shih, A. Greshake, D. Stöffler, and O. Eugster (2001), Ages and geologic histories of Martian meteorites, *Space Sci. Rev.*, 96, 105–164.
- Owen, T. C. (1992), The composition and early history of the atmosphere of Mars, in *Mars*, edited by H. H. Kieffer et al., pp. 818–834, Univ. of Ariz. Press, Tucson, Ariz.
- Owen, T., K. Biemann, D. R. Rushneck, J. E. Biller, D. W. Howarth, and A. L. Lafleur (1977), The composition of the atmosphere at the surface of Mars, *J. Geophys. Res.*, 82, 4635–4639.
- Owen, T., P. R. Mahaffy, H. B. Niemann, S. Atreya, and M. H. Wong (2001), Protosolar nitrogen, *Astrophys. J.*, 553, L77–L79.
- Oyama, V. I., and B. J. Berdahl (1977), The Viking gas exchange experiment results from Chryse and Utopia surface samples, *J. Geophys. Res.*, 82, 4669–4676.
- Schwenzer, S. P., S. Herrmann, R. K. Mohapatra, and U. Ott (2007), Noble gases in mineral separates from three shergottites: Shergotty, Zagami, and EETA79001, *Meteorit. Planet. Sci.*, 42, 387–412.
- Stern, J. C., et al. (2013), Detection of reduced nitrogen compounds at rock-vent using the Sample Analysis at Mars (SAM) instrument on the Mars Science Laboratory (MSL), *LPI Contrib.*, 1719, 2790.
- Trainer, M. G., et al. (2013), Change in the <sup>40</sup>Ar/N of the Mars Atmosphere from Viking to MSL: A possible indication of climate change on Mars, presented at 2013 Fall Meeting, P31C-03, AGU, San Francisco, Calif.
- Webster, C. R. et al. (2013), Isotope ratios of H, C, and O in CO<sub>2</sub> and H<sub>2</sub>O of the Martian atmosphere, *Science*, 341, 260–263.
- Wiens, R. C., R. H. Becker, and R. O. Pepin (1986), The case for a Martian origin of the shergottites. II. Trapped and indigenous gas components in EETA 79001 glass, *Earth Planet. Sci. Lett.*, 77, 149–158.
- Wong, M. H., P. R. Mahaffy, S. K. Atreya, H. B. Niemann, and T. C. Owen (2004), Updated Galileo probe mass spectrometer measurements of carbon, oxygen, nitrogen, and sulfur on Jupiter, *Icarus*, 171, 153–170.
- Wong, M. H., et al. (2013), MSL/SAM measurements of nitrogen and argon isotopes in the Mars atmosphere, *LPI Contrib.*, 1719, 1712. LPSC E-Poster at <http://www.lpi.usra.edu/meetings/lpsc2013/eposter/1712.pdf>.
- York, D., N. M. Evensen, and M. L. Martínez (2004), Unified equations for the slope, intercept, and standard errors of the best straight line, *Am. J. Phys.*, 72, 367–375.
- Zent, A. P., R. C. Quinn, and B. M. Jakosky (1994), Fractionation of nitrogen isotopes on Mars: The role of the regolith as a buffer, *Icarus*, 112, 537–540.