

HEALTH POLICY

Adolescent vulnerability, sexual health, and the NP's role in health advocacy

Elizabeth K. Kuzma, DNP, FNP-BC (Clinical Instructor)¹ & Rosalind M. Peters, PhD, RN, FAAN (Associate Professor)²

¹School of Nursing, University of Michigan, Ann Arbor, Michigan

²College of Nursing, Wayne State University, Detroit, Michigan

Keywords

Adolescent; youth-friendly; vulnerable; health policy; role of NPs; advocacy.

Correspondence

Elizabeth K. Kuzma, DNP, FNP-BC, School of Nursing, University of Michigan, 400 N. Ingalls Room 3173, Ann Arbor, MI 48109.
Tel: 734-647-0149; Fax: 734-647-0351;
E-mail: ekuzma@med.umich.edu

Received: 28 May 2015;

accepted: 12 October 2015

doi: 10.1002/2327-6924.12331

Abstract

Purpose: To describe adolescents as a vulnerable population with unique healthcare needs, especially in relation to their sexual risk-taking behaviors; and to explicate the necessity of nurse practitioners (NPs) advocating for youth-friendly services and policies to meet adolescents' sexual and reproductive healthcare needs.

Data sources: CINAHL, Medline, PsychInfo, and PsychArticle databases were searched to identify theoretical and empirical literature regarding adolescence, vulnerability, sexual health outcomes, barriers to accessing reproductive health services, what it means to be youth friendly, and health advocacy to meet the health needs of adolescents.

Conclusions: Adolescents' health needs may not be fully met in traditional healthcare settings. Lack of access to youth-friendly sexual and reproductive health services increases adolescents' risks for poor health outcomes including sexually transmitted disease and unplanned pregnancy. Clinic, state, and national policies can create barriers for adolescents in obtaining sexual health services.

Implications for practice: NPs are philosophically and educationally prepared to be leaders in improving adolescent health outcomes. NPs can directly provide youth-friendly care as well as advocate for youth-friendly practices within their health system. In addition, NPs are well positioned to be leaders in advocating for state and national policies that improve adolescents' access to appropriate sexual and reproductive health care.

Introduction

Adolescence is a critical period in the transition from childhood into adulthood during which young people, 12–18 years of age, experience significant physical, psychological, social, and emotional changes. Adjusting to their sexually maturing bodies, developing new perspectives on human relationships, and developing new coping and decision-making skills are among the major tasks of this developmental stage (Erikson & Erikson, 1998; Havighurst, 1956). Risk taking is part of normal healthy development as adolescents move toward independence. However, risky behaviors, especially sexual risk-taking behaviors, have consequences that may significantly impact the youth's future health and well-being. Although adolescents are at risk for many adverse outcomes from sexual risk-taking behaviors, they are less likely to

seek reproductive health services because of a variety of barriers. To help adolescents transition into a healthy adulthood, nurse practitioners (NPs) need to address the unique needs of youth, especially those related to their sexual expression (Lim, Chhabra, Rosen, Racine, & Alderman, 2012). The purpose of this article is to address the role of NPs in improving adolescent health by understanding why adolescents are a vulnerable group, directly providing youth-friendly health care (i.e., care that is safe, confidential, and age appropriate), and by advocating for youth-friendly policies at their practice sites, as well as advocating for policy change at the state and national level.

Adolescent vulnerability

Within health care, vulnerable populations are those that are not well integrated into the healthcare system,

putting them at risk for serious health problems (Urban Institute, 2010). Typically, vulnerable groups are those at risk because of racial, ethnic, economic, geographic, or health characteristics (e.g., disabilities), but age is also a potential source of vulnerability (Office of Minority Health, 2014). Adolescents are a vulnerable population because of their psychosocial developmental stage and its associated challenges, as well as the structural and functional changes occurring in the brain (World Health Organization: Department of Maternal, Newborn, Child and Adolescent Health, 2012).

Psychosocially, adolescents are increasing their level of autonomy and independence, yet are still dependent on their families for food, housing, financial support, and safety, including access to health care (Bailey, 2012). Risk taking in adolescence is a part of identity development (Erikson & Erikson, 1998). Many of the risks taken by adolescents are those associated with a sense of invincibility: the belief that consequences from risky behaviors will not happen to them.

Risk-taking behaviors may reflect not only psychosocial changes, but also developmental changes occurring in the brain. Researchers have suggested two processes of brain maturation resulting in development of the social-emotional and cognitive control systems (Steinberg, 2007, 2008). The social-emotional system, involving the amygdala, matures more rapidly than the prefrontal cortex which acts as the cognitive control system. This imbalance in maturation predisposes adolescents to risk-taking behaviors. While the brain is maturing, the adolescent may be making decisions guided by the amygdala, the portion of the brain responsible for emotions and emotional behaviors, leading to increased reward seeking, especially in the presence of peers (Steinberg, 2008). It takes longer for the prefrontal cortex, the portion of the brain responsible for decision making, impulse control, and the ability to envision long-term consequences, to mature. Thus, the adolescent prefrontal cortex may not provide the self-regulatory capacity needed to adequately assess and control impulsivity and risk-taking behavior that is driven by social-emotional processes (Blakemore, 2012; Steinberg, 2007, 2008).

Adolescent vulnerability and sexual health

As part of normal physical and biological development, it is natural for adolescents to begin to explore their sexual identity and sexual feelings through dating and sexual experimentation (Mannheim, Zieve, Eltz, Slon, & Wang, 2013). When paired with increased reward-seeking behaviors and poor impulse control, this stage of development makes adolescents especially vulnerable to problems related to sexual behaviors.

Adolescents are susceptible to sexual risk taking because of social pressures, mixed messages about sexuality, as well as having limited resources, or support to protect themselves from unsafe sex, sexually transmitted infections (STIs), or pregnancy (Bailey, 2012). In addition, the use of tobacco, alcohol, or illicit drugs, even casually, increases the chance that an adolescent will engage in high-risk sexual behaviors (Centers for Disease Control and Prevention [CDC], 2015a).

In 2013, nearly 50% of all high school students in the United States reported engaging in some form of sexual activity (oral, vaginal, or anal) at least once. Among those reported engaging in sexual activity in the past 3 months, 40.9% did not use a condom. Fifteen percent of the students surveyed reported having four or more sexual partners while still in high school (CDC, 2014). Specific consequences of sexual risk-taking behaviors include teen pregnancy, STIs, and human immunodeficiency virus (HIV; CDC, 2015b). The consequences of adolescent sexual risk-taking behaviors are associated with significant costs for both the individual and society as a whole (Blank, Baxter, Payne, Guillaume, & Pilgrim, 2010).

Consequences and costs

While older adolescents (ages 15–19) and young adults (ages 20–24) represent only 14.1% of the population in the United States (U.S. Census Bureau, 2011), in 2013 they accounted for 56% of new diagnoses of gonorrhea, 67% of new chlamydia cases (CDC, 2014), and those ages 13–24 accounted for 26% of new HIV infections (CDC, 2013b). The CDC estimates these infections account for nearly \$16 billion in direct medical costs (CDC, 2013a).

Teen birth is another major health consequence of adolescent sexual risk behavior. Although the U.S. teen birth rate has decreased by 40% over the last 20 years, in 2013 there were still more than a quarter million babies born to adolescent girls aged 15–19 (Hamilton, Martin, Osterman, & Curtin, 2014). Teen pregnancy has negative consequences for teen parents, their children, and society. Children born to teenage mothers are at increased risk of being born prematurely and having low birthweight resulting in complications requiring additional medical interventions (March of Dimes, 2012). Teen pregnancy and childbirth is estimated to cost U.S. taxpayers \$9.4–\$28 billion annually. These costs are due to increased healthcare and foster care costs, increased incarceration rates among children of teen parents, and lost tax revenue because of lower educational attainment and income among teen mothers (National Campaign, 2013; U.S. HHS, 2015).

Cleland, Peipert, Westhoff, Spear, and Trussell (2011) estimated that for every \$1 spent on family planning services the U.S. taxpayers could save \$3.74 in

pregnancy-related healthcare costs. In addition, the CDC (2015b) stresses the urgency of improving access to STI screening, treatment, and primary prevention services to decrease the burden of STIs. Yet, teens often do not avail themselves of sexual health services. Identifying and overcoming the barriers to sexual health care is critical in improving health outcomes for this vulnerable population.

Barriers to sexual health services for adolescents

Adolescents in the United States experience a number of barriers to sexual and reproductive healthcare services. A review of relevant literature revealed that the major barriers to adolescents obtaining appropriate sexual health care included access, communication, and trust in provider, with concerns about confidentiality as an overarching issue influencing the other barriers. State and federal policies can create additional barriers to care by limiting youths' ability to access necessary sexual and reproductive health services.

Confidentiality

Concern for confidentiality is seen as a significant barrier for adolescents in accessing healthcare services. Adolescents may be concerned about being recognized in the waiting room by family, friends, or other people in the community who may know their parents. They may also be concerned that their provider may have social interactions with parents in which the provider might purposefully or accidentally disclose that the teen was seen by them (World Health Organization: Department of Maternal, Newborn, Child and Adolescent Health, 2012). Having parents learn about services sought is a profound fear of adolescents. One study found that 83% of the adolescent participants would stop accessing sexual health services if their parents were notified, while only 1% of the participants would abstain from sex (Alford, 2009).

Access. Access to sexual health care is another major barrier facing adolescents. Access can be impeded in multiple ways including: time and space available for services, access to a provider who will give them time alone for appointments and will talk about sexual issues, access to confidential services that are affordable, and access to transportation for appointments (World Health Organization: Department of Maternal, Newborn, Child and Adolescent Health, 2012).

Adolescents may have a healthcare provider they see regularly, but may lack access to sexual health services because the provider does not see them privately without a parent being present. Additionally, some healthcare providers will either choose not to ask adolescents about

sexual health issues, or will do so in front of a parent, which inadvertently limits adolescents' ability to access necessary services.

Access may also be affected by the nature of the clinic itself. Many clinics do not have late, weekend, or adolescent-specific hours. Further, an adolescent may feel uncomfortable addressing sexual concerns in a pediatric clinic, yet may not know where or how to access an adult clinic, let alone one that is youth-friendly.

Cost creates another access issue for adolescents as well as a potential breach in confidentiality (Tebb et al., 2014). If adolescents are insured under their parents' plan and the services are billed to that insurance, an explanation of benefits letter is often sent home. Such a letter would alert parents that the adolescent had sought health services, threatening the confidentiality of the services provided. To avoid this breach in confidentiality, adolescents may elect to not have the insurance billed, which then requires adolescents to pay out of pocket for what could be expensive clinic and laboratory services.

Access to transportation for confidential health appointments may create an additional barrier (World Health Organization: Department of Maternal, Newborn, Child and Adolescent Health, 2012). Clinics may not be conveniently located near public transportation routes and adolescents may not have the financial means to afford cab service or other types of transportation to obtain health care without their parents' knowledge.

Communication/trust. Effective communication skills are essential for establishing trust with adolescent patients (Ambresin, Bennett, Patton, Sanci, & Sawyer, 2013). Adolescents have expressed a willingness and desire to discuss sensitive topics with their providers, but the providers' communication style is a determining factor as to what information the adolescent will share (Martyn et al., 2013). Informing adolescents they have a right to confidentiality protected by law helps promote a culture of openness and disclosure. A significant barrier can be created if a healthcare provider unintentionally disrespects, ignores, or judges an adolescent (Alford, 2009). This can happen when providers fear their guidance is not being accepted by the adolescent. This fear may lead to frustration impeding effective communication and potentially jeopardizing the adolescents' return for needed follow-up care.

Health policy as a barrier

Public health policies are made in the legislative arena with the presumed goal of achieving specific health objectives (World Health Organization, 2015). However, health policies are influenced by the beliefs and values of those who develop and enact them (Leavitt, 2009), and thus

may reflect political rather than health-oriented goals. Adolescent sexual and reproductive health is a very controversial political issue. Many view allowing adolescents to access confidential reproductive health services as a threat to parental rights (Herrman, 2013). Currently, minors aged 13–17 have the legal right to access confidential health services in all states, but some policymakers want to restrict the confidential services youth can access by requiring parental notification (American Civil Liberty Union [ACLU], 2015). While parental involvement is vital to the reproductive health of their children, consent laws could create a severe barrier and put their teens' health at risk. Research has shown that mandated parental notification does not deter adolescents from engaging in sexual activity, as adolescents reported they would continue having sex even if they could not access confidential health services (Goodwin et al., 2012).

Clinical implications—youth-friendly health care

Adolescents are generally healthy and the major health problems they face are because of preventable conditions arising from risky behaviors. Meeting the sexual and reproductive healthcare needs of adolescents requires a rethinking of the clinical services provided to this at-risk, vulnerable population. As identified by World Health Organization: Department of Maternal, Newborn, Child and Adolescent Health (2012), the major clinical implication to improve care for adolescents is the need to establish youth-friendly practices and policies that reduce or remove identified barriers to care. Because of their philosophical approach to practice, NPs are well suited to providing care that is sensitive to adolescents' cognitive, biological, and psychological stage of development. NPs can play an important role in reducing adolescents' barriers to care by directly providing youth-friendly services, advocating for the creation of youth-friendly environments, and addressing state and federal policy issues that could help overcome barriers to adolescent care.

Youth-friendly care

Being youth-friendly means meeting the needs of youth by providing a safe, comfortable, confidential environment for youth (World Health Organization: Department of Maternal, Newborn, Child and Adolescent Health, 2012). Youth-friendly clinics have policies and services that attract and retain adolescent patients. Characteristics of a youth-friendly clinic include having providers who are specially trained to work with adolescents and who understand the need for protecting privacy and confidentiality, offer services at an affordable cost, and have flexible, convenient hours for youth.

NPs directly providing youth-friendly health care

To provide youth-friendly services, NPs may have to undergo a paradigm shift to recognize adolescents as a vulnerable population with needs very different from children or adult patients. Barriers are reduced when NPs are committed to confidentiality, therapeutic communication, and clearly understand minor consent laws.

Commitment to confidentiality. A major way in which NPs can reduce barriers to care for adolescents is by recognizing the need for confidentiality and ways in which it can be broken, even inadvertently. NPs must communicate their commitment to maintain confidentiality to the adolescent, while also being honest about the limits to confidentiality, such as ensuring patient safety, and the laws related to reporting communicable diseases (World Health Organization: Department of Maternal, Newborn, Child and Adolescent Health, 2012).

NP communication. NPs should be prepared to talk to adolescents about sexual health in a nonjudgmental, respectful manner. Discussions should address: the youth's emotional readiness for sex, the benefits of abstinence, strategies for maintaining abstinence, and safe sexual practices for sexually active teens including how to prevent pregnancy and/or STIs. NPs must provide accurate information about the options, safety, and side effects of multiple contraceptive methods, including condoms, as well as information regarding where contraceptives are available for free or low cost (Martyn et al., 2013). NPs must also be willing and able to discuss pregnancy options for an adolescent, including choosing to keep the pregnancy, considering adoption, or termination of the pregnancy. NPs must know state laws regarding a minor's ability to consent to adoption or access abortion services if desired. To ensure patients can access information to which they are legally entitled, NPs also need to know local resources to refer an adolescent if not comfortable discussing all pregnancy options, including abortion.

What are the laws? Adolescents are not legally adults, thus it is crucial for NPs to be knowledgeable about laws that affect adolescents' access to health care (Riley, Ranalli, Lane, & Sohikian, 2014). Nationally, minors are legally able to access confidential reproductive and sexual health services (Guttmacher, 2015). A minor's right to access contraception services is protected by the Supreme Court ruling in *Carey vs. Population Services International* (1977, 431 U.S. 678). Additionally, minors aged 13 and older are ensured access to confidential reproductive health services through requirements in federally funded programs such as Medicaid and Title X (U.S. HHS, 2014). In most states, minors can be tested and receive treatment for STIs, including HIV, receive testing for pregnancy and prenatal care, and obtain contraception

Table 1 Health policy resources: state laws and information to locate elected officials

Website	Highlights
http://www.reproductiverights.org	Center for Reproductive Law & Policy ■ Information regarding state laws for a minor's ability to access contraception, consent for an abortion, and consent for an adoption
http://www.guttmacher.org	Guttmacher Institute ■ Information regarding state laws for a minor's ability to access contraception, consent for an abortion, and consent for an adoption
https://www.opencongress.org/people/zipcodelookup	Open Congress ■ This site lists Federal Senators and Representatives by zip code.
http://openstates.org/find_your legislator/	Open States ■ This site lists State Senators and Representatives by zip code.
http://www.statelocalgov.net/	State & Local Government ■ This site lists local elected officials by zip code.

without parental consent. Many states require minors to have parental consent for adoption or an abortion. The laws vary by state requiring healthcare providers working with adolescents to be aware of the specific minor consent law in the state(s) in which they practice. Information on state laws affecting adolescent services can be found in Table 1.

NPs creating a youth-friendly care environment

NPs can advocate for environmental and policy changes at their clinic or health system level. Clinic level changes to improve access for adolescents could include having specific clinic times dedicated to adolescent services, allowing walk-in or evening appointments. Having specific time designated for adolescents and allowing more flexibility when they arrive late could reduce the barriers they face in accessing care. Youth-friendly clinics would have exam rooms that are appealing and inviting to adolescents by having posters that clearly list confidential and sensitive services to encourage them to discuss risky behaviors. All office staff need to be trained in serving youth, including minor consent laws, documentation, and billing prac-

tices for confidential services because staff attitudes, values, and beliefs can greatly impact the clinic's ability to provide youth-friendly care (World Health Organization: Department of Maternal, Newborn, Child and Adolescent Health, 2012). Any breach in confidentiality or perceived disrespect could prevent an adolescent from seeking services again (Alford, 2009).

NPs role in advocacy for youth-friendly health policy

Nurses have a long history of patient advocacy at both the individual and population level (Selanders & Crane, 2012). NPs can continue this tradition by using their clinical expertise in a health advocacy role. Health advocacy encompasses direct services at the individual and family level, but also includes being involved in the political processes that affect access to health care for communities and the general public.

As described above, there are numerous ways in which NPs can advocate for individually focused youth-friendly services within their own practice sites. Policy advocacy efforts at the population level need to recognize the varying opinions and preferences about how tax dollars should be spent. Further, there are public concerns that investing in the health of adolescents is an enormous expenditure with limited return. Thus, much of the NP advocacy role may be related to educating policymakers and the public about the costs of *not* meeting the sexual and reproductive health needs of adolescents. Areas for advocacy efforts could include: augmenting services in existing youth-friendly models of care, improving funding to clinics that provide comprehensive sexual and reproductive health services, removal of unnecessary and burdensome requirements for clinics that provide abortion services, and broader over-the-counter (OTC) access for emergency contraception.

Advocating for enhanced services in existing youth-friendly models of care. NPs could advocate for policy changes that would allow established youth-friendly clinics, such as school-based health centers (SBHCs), to provide comprehensive sexual and reproductive health services on-site. When SBHCs are able to provide comprehensive services, there is a decreased prevalence of sexual risk-taking behaviors and decreased incidence of STIs among patients (Gonzales, 2011). Multiple studies endorse the benefits of SBHCs providing contraception on-site, including improved consistency of contraception use and increased chance of SBHC patients using hormonal or emergency contraception at their last sexual encounter (Blank et al., 2010; Ethier et al., 2011). Despite the research demonstrating high-quality outcomes of SBHCs, less than half of the 1900 SBHCs nationwide have the legal ability to prescribe or dispense contraception

on site, thus limiting their ability to fully meet the needs of the youth they serve (Keeton, Soleimanpour, & Brindis, 2012). The services provided by SBHCs may be restricted at either the state or local community level, so NPs must be prepared to advocate for policy change at all levels.

Advocating for improved funding to clinics that provide comprehensive reproductive health services. NPs also could advocate for sustained funding to programs that provide vital sexual and reproductive services to youth. Title X is the only federal grant program dedicated solely to providing individuals with comprehensive family planning and related preventive health services. Programs funded by Title X offer a broad range of contraceptive methods approved by the Food and Drug Administration and related counseling, as well as breast and cervical cancer screening, pregnancy testing, screening and treatment for STIs including HIV, and other patient education and referrals (U.S. HHS, 2014). By law, Title X funds cannot be used to pay for abortion. Despite the fact that Title X centers serve over 4.5 million clients annually, for the fifth year in a row, the U.S. House of Representatives has proposed ending the program, and the U.S. Senate is recommending a major reduction in funding. Such policy decisions disproportionately affect adolescents and disadvantaged women who do not have resources to obtain reproductive health care elsewhere (Guttmacher, 2015). As patient advocates, NPs must work to ensure that patients have access to essential reproductive healthcare services to achieve and maintain optimal health.

Advocating for the removal of unnecessary and burdensome requirements for clinics that provide abortion services. Although abortion remains a legal right for women in the United States, a number of policy and funding decisions have created major barriers for women to access abortion services. There has been almost a 50% reduction in clinics providing abortion services since the 1980s (Guttmacher, 2014). This has occurred as states have enacted laws that severely limit who provides and where abortion services can be offered. The ability of adolescents to obtain abortions also varies by state. Some states require parental notification while others require either parental or other adult family member consent. Some states have judicial bypass laws allowing a teen to petition the court to obtain an abortion without parental consent. NPs working with adolescents considering abortion need to be aware of the laws in their state. Table 2 lists a website that offers an overview of state abortion laws.

Advocating for broader OTC access to emergency contraception. OTC access to emergency contraception is a safe, effective method for preventing pregnancy, but only branded medications (e.g., Plan B)

are available without age restrictions (Munro, Dulin, & Kuzma, 2015). Generic formulations of emergency contraception still have restrictions on availability, some require a prescription, and others are only available OTC for youth 17 years and older. Many pharmacies keep emergency contraceptives behind the counter, available for purchase without a prescription, which requires teens to have confidence in requesting the medication and the financial means to pay for it. Such restrictions result in cost and access barriers for adolescents.

Steps in health advocacy. There are multiple ways for NPs to be involved in advocating for health policies that support adolescents' access to comprehensive sexual and reproductive health services. The level of involvement depends on the level of investment an NP is willing to make. The simplest level of involvement is to become a member of an adolescent health organization that endorses advocacy efforts that an NP supports (HealthPAC, 2015). Membership dues help maintain the organization's ability to continue advocacy efforts, and donating to the organization's political action fund (PAC) helps the organization to be active and effective in the political arena.

Another method of involvement that requires minimal time commitment and effort is signing up for listservs, e-mail alerts, or following organizations social media page(s) in order to be notified when action on important issues is necessary. These action efforts usually involve easily signed prewritten electronic messages sent to policymakers to express support for an important issue and take only a few minutes to complete.

Much of advocacy is interpersonal in nature, thus establishing a relationship with lawmakers is important. The most valuable advocacy effort requires NPs to personally connect with their local, state, and federal elected officials (Table 2 provides website links for finding elected officials by zip code). Communication can be achieved by sending e-mails, speaking with office staff, and/or scheduling face-to-face meetings. It is essential that in these communications the NP presents factual information along with related patient stories to counter any misinformation regarding adolescents' ability to access comprehensive sexual and reproductive health services. In order to be effective in policy advocacy, NPs must be informed of the issues surrounding adolescent access to comprehensive sexual and reproductive health care. Table 2 presents numerous resources to help NPs be able to respond effectively to questions about adolescent sexual and reproductive health. As part of the most trusted profession in the United States, NPs are in a unique position to use their knowledge, interpersonal skills, and clinical expertise to influence health policy.

Table 2 Resources on adolescent sexual and reproductive health for nurse practitioners

Website	Highlights
http://www.guideline.gov/content.aspx?id=47031&search=abortion+and+adolescent	2013 UK national guideline sexual history taking
http://umhs-adolescenthealth.org	Adolescent Health Initiative ■ Provider resources, trainings, annual conference
http://www.ahwg.net/	Adolescent Health Working Group ■ Training sessions, modules, online toolkits
http://www.advocatesforyouth.org	Advocates for Youth ■ Publications, resources, professional education, and patient resources
http://www.aap.org	American Academy of Pediatrics ■ Publications, resources, professional education, and patient resources
http://www.acog.org	American College of Obstetricians and Gynecologists ■ Publications, resources, professional education, and patient resources
https://www.arhp.org/	Association of Reproductive Health Professionals ■ Publications, resources, professional education, and patient resources
https://brightfutures.aap.org/materials-and-tools/tool-and-resource-kit/Pages/default.aspx	Bright Futures: Promoting Healthy Sexual Development and Sexuality ■ Toolkit for providers
http://leah.mchtraining.net/	Leadership Education in Adolescent Health ■ Leadership training program
https://www.napnap.org/	National Association of Pediatric Nurse Practitioners ■ National conference offers sessions on adolescent sexual and reproductive health and other online resources
http://www.ncsddc.org/	National Coalition of STD Directors, promoting sexual health through STD prevention ■ Resources and publications
http://www.naspag.org/	North American Society for Pediatric and Adolescent Gynecology ■ Publications, resources, professional education, and patient resources
http://prh.org/teen-reproductive-health/arshep-downloads/	Physicians for Reproductive Health ■ Many power point presentations and other helpful resources for providers
http://www.plannedparenthood.org	Planned Parenthood Federation of America ■ Publications, resources, professional education, and patient resources
http://www.cdc.gov/mmwr/preview/mmwrhtml/rr6304a1.htm?s_cid=rr6304a1_w	Providing Quality Family Planning Services: Recommendations of CDC and the U.S. Office of Population Affairs: Recommendations and Reports April 25, 2014/63(RR04);1-29
http://www.reproductiveaccess.org/	The Reproductive Health Access Project ■ Publications, resources, professional education, and patient resources
http://www.adolescenthealth.org/	Society for Adolescent Health and Medicine ■ Offers fellowships and trainings annually, webinars, clinical care guidelines, annual conference
http://www.hhs.gov/ash/oah/	U.S. Office of Adolescent Health ■ Publications, resources, professional education, patient resources, health initiatives, state data

NPs as leaders in youth-friendly initiatives and care

Adolescents are a vulnerable population at increased risk of serious health problems because of their developmental state. NPs advocacy efforts can help improve adolescent health outcomes. Advocacy by a particularly skilled and knowledgeable group with a common philosophy, such as NPs, brings the strength of numbers of like-minded peo-

ple advocating for youth-friendly practices and policies to overcome barriers to care faced by adolescents. NPs can effect change, not only in their own practice, but can influence and implement change at the clinic, healthcare system, and community level, as well as advocate for policy change at the state and national level to ensure the healthcare needs of adolescents are met. NPs have an obligation to advocate for their adolescent patients, serving as champions for change.

Acknowledgments

Drs. Elizabeth K. Kuzma and Rosalind M. Peters provided substantial contributions to the conception and design of the manuscript. Dr. Elizabeth K. Kuzma wrote the initial draft. Both authors revised it critically for important intellectual content, made a final approval of the version to be published, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy and integrity of any part of the work are appropriately investigated and resolved.

References

- Alford, S. (2009). Best practices for youth-friendly services. *Advocates for Youth*. Retrieved from <http://www.advocatesforyouth.org/component/content/article/1347-best-practices-for-youth-friendly-clinical-services>
- Ambresin, A. E., Bennett, K., Patton, G. C., Sanci, L. A., & Sawyer, S. M. (2013). Assessment of youth-friendly healthcare: A systematic review of indicators drawn from young people's perspectives. *Journal of Adolescent Health, 52*(2013), 670–681.
- American Civil Liberty Union (ACLU). (2015). *Preventing teenagers from getting contraceptives unless they tell a parent puts teens at risk*. Retrieved from <https://www.aclu.org/preventing-teenagers-getting-contraceptives-unless-they-tell-parent-puts-teens-risk>
- Bailey, L. D. (2012). Adolescent girls: A vulnerable population. *Advances in Neonatal Care, 12*(2), 102–106.
- Blakemore, S. J. (2012). Development of the social brain in adolescence. *Journal of the Royal Society of Medicine, 105*(10), 111–116.
- Blank, L., Baxter, S. K., Payne, N., Guillaume, L. R., & Pilgrim, H. (2010). Systematic review and narrative synthesis of the effectiveness of contraception service interventions for young people, delivered in educational settings. *Journal of Pediatric Adolescent Gynecology, 23*(2010), 341–351.
- Centers for Disease Control and Prevention. (2013a). *CDC fact sheet: Incidence, prevalence, and cost of sexually transmitted infections in the United States*. Retrieved from <http://www.cdc.gov/std/stats/sti-estimates-fact-sheet-feb-2013.pdf>
- Centers for Disease Control and Prevention. (2013b). *HIV incidence*. Retrieved from <http://www.cdc.gov/hiv/statistics/surveillance/incidence/>
- Centers for Disease Control and Prevention. (2014). *2013 Sexually transmitted diseases surveillance*. Retrieved from <http://www.cdc.gov/std/stats13/adol.htm>
- Centers for Disease Control and Prevention. (2015a). *HIV and substance use in the United States*. Retrieved from <http://www.cdc.gov/hiv/riskbehaviors/substanceuse.html>
- Centers for Disease Control and Prevention. (2015b). *Sexual risk behavior: HIV, STD, & teen pregnancy prevention*. Retrieved from <http://www.cdc.gov/healthyouth/sexualbehaviors/>
- Cleland, K., Peipert, J. F., Westhoff, C., Spear, S., & Trussell, J. (2011). Family planning as a cost-saving preventive health service. *New England Journal of Medicine, 364*(37), 1–3.
- Erikson, E. H., & Erikson, J. M. (1998). *The life cycle completed*. New York, NY: W. W. Norton & Company.
- Ethier, K. A., Dittus, P. J., DeRosa, C. J., Chung, E. Q., Martinez, E., & Kerndt, P. R. (2011). School-based health center access, reproductive health care, and contraception use among sexually experienced high school students. *Journal of Adolescent Health, 48*(6), 562–565.
- Gonzales, M. (2011). *Position statement: Providing reproductive health services in Colorado school-based health centers*. Denver: Colorado Association for School-Based Health Care Retrieved from <http://www.casbhc.org/Publications/Policy%20Briefs/Reproductive%20Health%20Position%20Statement.pdf>
- Goodwin, K. D., Taylor, M. M., Brown, E. C., Winscott, M., Scanlon, M., Hodge, Jr, J. G., ... England, B. (2012). Protecting adolescents' right to seek treatment for sexually transmitted diseases without parental consent: The Arizona experience with Senate Bill 1309. *Public Health Reports, 127*(3), 253–258.
- Guttmacher. (2014). *Trends in abortion in the United States, 1973–2011*. Retrieved from <https://www.guttmacher.org/presentations/trends.pdf>
- Guttmacher. (2015). *State policies in brief: An overview of minor's consent laws*. Retrieved from <http://www.guttmacher.org/statecenter/spibs/spib'OMCL.pdf>
- Hamilton, B. E., Martin, J. A., Osterman, M. J. K., & Curtin, S. C. (2014). Births: Preliminary data for 2013. *National Center for Health Statistics*. Retrieved from <http://www.cdc.gov/nchs/data/nvsr/nvsr63/nvsr63'02.pdf>
- Havighurst, R. J. (1956). Research on the developmental-task concept. *School Review, 64*(5), 215–223.
- HealthPAC. (2015). *Healthcare advocacy*. Retrieved from <http://www.healthpaonline.net/health-care-advocacy.htm>
- Herrman, J. W. (2013). *Delaware stakeholder perceptions of the provision of reproductive health services by school-based health centers*. Retrieved from <http://dethrives.com/wp-content/uploads/2013/06/perceptions-2.pdf>
- Keeton, V., Soleimanpour, S., & Brindis, C. D. (2012). School-based health centers in an era of healthcare reform: Building on history. *Current Problems in Pediatric and Adolescent Healthcare, 12*(7), 132–156.
- Leavitt, J. K. (2009). Leaders in health policy: A critical role for nursing. *Nursing Outlook, 2009*(57), 73–77.
- Lim, S. W., Chhabra, R., Rosen, A., Racine, A. D., & Alderman, E. M. (2012). Adolescents' views on barriers to healthcare: A pilot study. *Journal of Primary Care & Community Health, 3*(2), 99–103.
- Mannheim, J. K., Zieve, D., Eltz, R. D., Slon, S., & Wang, N. (2013). *U. S. national library of medicine, medline plus: Adolescent development*. Retrieved from <http://www.nlm.nih.gov/medlineplus/ency/article/002003.htm>
- March of Dimes. (2012). *Teenage pregnancy*. Retrieved from <http://www.marchofdimes.org/materials/teenage-pregnancy.pdf>
- Martyn, K. K., Munro, M. L., Darling-Fisher, C. S., Ronis, D. L., Villarruel, A. M., Pardee, M., ... Fava, N. M. (2013). Patient-centered communication and health assessment with youth. *Nursing Research, 62*(6), 383–393.
- Munro, M. L., Dulin, A. C., & Kuzma, E. (2015). Update on Plan B emergency contraception and the nurse's role. *Nursing for Women's Health, 19*(2), 142–153.
- Riley, M., Ranalli, L., Lane, J., & Sohikian, V. (2014). *Adolescent centered environment toolkit*. Ann Arbor, MI: The Adolescent Health Initiative—The University of Michigan Health System.
- Selanders, L., & Crane, P. (January 31, 2012). The Voice of Florence Nightingale on Advocacy. *OJIN: The Online Journal of Issues in Nursing, 17*(1), 1. doi: 10.3912/OJIN.Vol17No01Man01
- Steinberg, L. (2007). Risk taking in adolescence: New perspectives from brain and behavioral science. *Current Directions in Psychological Science, 16*, 55–59.
- Steinberg, L. (2008). A social neuroscience perspective on adolescent risk-taking. *Developmental Review, 28*, 78–106.
- Tebb, K. P., Sedlander, E., Pica, G., Diaz, A., Peake, K., & Brindis, C. D. (2014). *Protecting adolescent confidentiality under healthcare reform: The special case of explanation of benefits (EOBs)*. Retrieved from http://healthpolicy.ucsf.edu/sites/healthpolicy.ucsf.edu/files/documents/EOB%20Policy%20Brief_FINAL.pdf
- The National Campaign to Prevent Teen and Unplanned Pregnancy. (2013). *Counting it up: The public costs of teen childbearing: Key data*. Washington, DC: The National Campaign to Prevent Teen and Unplanned Pregnancy. Retrieved from <http://thenationalcampaign.org/sites/default/files/resource-primary-download/counting-it-up-key-data-2013-update.pdf>
- United States Supreme Court. (06/09/1977). *Care versus Population Services International (75–443) 431 U.S. 678 (1977)*.
- United States [U.S.] Census Bureau. (2011). *Age and sex composition: 2010*. Retrieved from <http://www.census.gov/prod/cen2010/briefs/c2010br-03.df>
- U.S. Department of Health and Human Services [U.S. HHS]: Office of Adolescent Health. (2015). *Strategies & approaches*. Retrieved from <http://www.hhs.gov/ash/oah/adolescent-health-topics/reproductive-health/teen-pregnancy/strategies-and-approaches.html>
- U.S. Department of Health and Human Services [U.S. HHS]: Office of Population Affairs. (2014). *Title X 90 family planning*. Retrieved from <http://www.hhs.gov/opa/title-x-family-planning/>

United States [U.S.] Office of Minority Health. (2014). *Other at-risk populations*. Retrieved from <http://www.cdc.gov/minorityhealth/populations/atrisk.html>

United States Supreme Court. (1977). *Carey vs. Population Services International* (75-443) 431 U.S. 678.

Urban Institute. (2010). *Vulnerable populations*. Retrieved from <http://www.urban.org/health-policy/vulnerable-populations/>

World Health Organization. (2015). *Health policy*. Retrieved from <http://www.who.int/topics/health-policy/en/>

World Health Organization: Department of Maternal, Newborn, Child and Adolescent Health. (2012). *Making health services adolescent friendly: Developing national quality standards for adolescent friendly health services*. Retrieved from <http://apps.who.int/iris/bitstream/10665/75217/1/9789241503594-eng.pdf>