

# Novel Roles for CRF-Binding Protein and CRF Receptor 2 in Binge Drinking

Kyle D. Ketchesin and Audrey F. Seasholtz

**S**TRESS IS A key environmental factor in alcohol addiction, and the stress system has been linked to alcohol use prior to dependence, after dependence, and in relapse to drinking after abstinence in both humans and rodent model systems (reviewed in Lowery and Thiele, 2010; Phillips et al., 2015; Ray, 2011). The key central nervous system regulator of the mammalian stress response is corticotropin-releasing factor (CRF). This peptide mediates its primary effects through binding to G-protein-coupled receptors, CRF receptor 1 (CRF-R1) and CRF receptor 2 (CRF-R2), and its activity is modulated by the CRF-binding protein (CRF-BP). Dysregulation of the CRF system has been observed in a number of rodent models of binge drinking and alcohol dependence, with most studies focusing on the roles of CRF and CRF receptors, particularly CRF-R1 (reviewed in Lowery and Thiele, 2010; Phillips et al., 2015). CRF-R1 antagonists decreased binge drinking and self-administration in alcohol-dependent animals (Lowery and Thiele, 2010), suggesting potential therapeutic uses for CRF-R1 antagonists. However, fewer studies have investigated the roles of CRF-R2 and CRF-BP in binge drinking and alcohol dependence.

In the recent study by Albrechet-Souza and colleagues (2015), a variety of drinking paradigms and pharmacological tools were utilized to examine the roles of CRF-BP and CRF-R2 in the ventral tegmental area (VTA) and central nucleus of the amygdala (CeA) in binge drinking and alcohol dependence. To model binge drinking, the authors used the drinking in the dark (DID) paradigm, in which mice were given limited access to 20% ethanol in the dark phase of their circadian cycle, resulting in drinking to intoxication and pharmacologically relevant blood ethanol concentration. Two variations of the DID paradigm were employed, both

1-bottle and 2-bottle choice (Rhodes et al., 2007). Alcohol dependence was modeled using a 4-week 2-bottle choice, intermittent access to alcohol (IAA) paradigm. To investigate the role of CRF-BP in each of these paradigms, the CRF-BP ligand inhibitor CRF<sub>6-33</sub> was utilized (Sutton et al., 1995). The authors found that CRF-BP in the VTA, but not CeA, regulates 1-bottle choice binge drinking, but not dependence-induced alcohol consumption. To investigate the role of CRF-R2 in the VTA in binge drinking, the CRF-R2 selective antagonist astressin 2B (A2B) was used. A2B injected into the VTA at a high dose led to a decrease in ethanol intake in 2-bottle choice DID, while low dose A2B had no effect. Together, these exciting results reveal novel roles for CRF-BP and CRF-R2 in the VTA in binge drinking.

## ROLES FOR CRF-BINDING PROTEIN

The CRF-BP is a 37-kDa secreted glycoprotein that binds CRF and the CRF-like ligand, urocortin 1 (Ucn1), with very high affinity. It is co-localized with CRF or CRF receptors at numerous sites (i.e., amygdala and bed nucleus of the stria terminalis), suggesting potential sites of interaction in stress-reward pathways (reviewed in Westphal and Seasholtz, 2006). In humans, it is estimated that 40 to 60% of CRF in the brain is bound by CRF-BP, and *CRHBP* single-nucleotide polymorphisms have been associated with alcohol use disorder and stress-induced alcohol craving (Ray, 2011), suggesting a role for the CRF-BP in susceptibility to alcohol use and addiction.

The CRF-BP has been studied for over 20 years with many postulated roles. In an inhibitory role, CRF-BP reduces CRF receptor activation, likely by sequestering CRF or Ucn1 and/or targeting them for degradation. Consistent with this model, purified CRF-BP reduces CRF-mediated adrenocorticotrophic hormone release from anterior pituitary cultures or AtT-20 cells (Cortright et al., 1995; Potter et al., 1991; Sutton et al., 1995). Recombinant CRF-BP also attenuates CRF-R1-mediated increases in cAMP (Boorse et al., 2006). Similarly, CRF-BP-deficient mice show increased baseline anxiety-like behavior and slowed return to homeostasis after lipopolysaccharide stress, consistent with elevated free levels of CRF in the absence of an inhibitory CRF-BP (Karolyi et al., 1999). In contrast, the study by Albrechet-Souza and colleagues (2015) suggests an enhancing or facilitatory role. In this role, CRF-BP may bind CRF

From the Neuroscience Graduate Program (KDK, AFS), University of Michigan, Ann Arbor, Michigan; Molecular and Behavioral Neuroscience Institute (KDK, AFS), University of Michigan, Ann Arbor, Michigan; and Department of Biological Chemistry (AFS), University of Michigan, Ann Arbor, Michigan.

Received for publication August 26, 2015; accepted September 2, 2015.

Reprint requests: Audrey F. Seasholtz, PhD, University of Michigan, 5035 Biomedical Science Research Building, 109 Zina Pitcher Place, Ann Arbor, MI 48109; Tel.: 734-936-2072; Fax: 734-936-2690; E-mail: aseashol@umich.edu

Copyright © 2015 by the Research Society on Alcoholism.

DOI: 10.1111/acer.12897

and increase CRF signaling, likely by delivering CRF to the receptor, extending the half-life of CRF, or modulating its interaction with the receptor. Consistent with this hypothesis, VTA CRF-BP may normally facilitate alcohol consumption, as administration of CRF<sub>6-33</sub> into VTA reduced binge drinking in mice (Albrechet-Souza et al., 2015). Similarly, intra-VTA administration of CRF<sub>6-33</sub> attenuated CRF-induced relapse to cocaine seeking in rats (Wang et al., 2007). Altogether, these results suggest that CRF-BP may have different roles depending on the specific cell type or context in which it is expressed, consistent with ultrastructural localization studies (Peto et al., 1999). Finally, the CRF-BP may exert ligand and/or receptor independent actions. Intracerebroventricular (i.c.v.) administration of CRF<sub>6-33</sub> elicits c-fos activation not only in a subset of CRF receptor-expressing cells, but also in CRF-BP-expressing cells that do not express ligand or CRF receptors, suggesting additional actions of CRF<sub>6-33</sub> and/or CRF-BP (Chan et al., 2000). Hence, further studies are needed to elucidate the multiple roles of CRF-BP in vivo, as the actions of CRF-BP are likely brain region- and cell type-dependent.

### CRF RECEPTORS IN BINGE DRINKING AND DEPENDENCE

The CRF receptors are highly homologous G-protein-coupled receptors with similar, yet distinct ligand binding profiles. CRF binds with very high affinity to CRF-R1, but binds to CRF-R2 with about 30-fold lower affinity. Both CRF receptors couple predominantly to G $\alpha_s$ , activating adenylyl cyclase and increasing cAMP levels and protein kinase A activity. However, they have also been shown to act through other G-protein subtypes, including G $\alpha_q/11$ , G $\alpha_o$ , and G $\alpha_i$ , in certain cellular contexts (reviewed in Hillhouse and Grammatopoulos, 2006).

There is a large body of literature implicating the role of CRF-R1 in excessive alcohol drinking (reviewed in Lowery and Thiele, 2010; Phillips et al., 2015). Altered CRF-R1 expression has been observed in rats with a history of alcohol dependence, and peripheral administration of CRF-R1 antagonists reduced both binge drinking and dependence-induced alcohol consumption (Lowery and Thiele, 2010). Furthermore, CRF receptor regulation of alcohol consumption may be brain region-specific, with CeA and VTA as major sites of action. Intra-CeA (but not basolateral amygdala) administration of a CRF-R1 antagonist decreased binge drinking (Lowery-Gionta et al., 2012). Similarly, intra-VTA CRF-R1 antagonist administration decreased binge drinking in a DID paradigm (Sparta et al., 2013) and decreased excessive drinking during the 2-bottle choice IAA paradigm (Hwa et al., 2013). Although less studied, CRF-R2 has also been implicated in excessive alcohol drinking, but often in an inverse fashion to CRF-R1. For example, i.c.v. administration of the CRF-R2-selective agonist urocortin 3 (Ucn3) dose-dependently decreased binge drinking, and intra-CeA administration of Ucn3 resulted in decreased

alcohol self-administration in alcohol-dependent rats (reviewed in Lowery and Thiele, 2010; Phillips et al., 2015). Hence, CRF-R1 antagonists and CRF-R2 agonists often elicit similar responses. In contrast, in the recent study by Albrechet-Souza and colleagues (2015), intra-VTA CRF-R2 antagonist (A2B) administration decreased alcohol drinking in the 2-bottle DID paradigm, suggesting that VTA CRF-R2 is facilitating drinking. Additional studies that directly compare the effects of CRF-R1 and CRF-R2, via genetic approaches and/or receptor-selective agonists and antagonists, in the VTA, CeA, and other brain regions will be required to fully elucidate the roles of these 2 receptors in alcohol binge drinking and dependence.

### POTENTIAL CRF-BP/CRF RECEPTOR INTERACTIONS AND FUTURE DIRECTIONS

The CRF system is unique in that very few neuropeptides have a high-affinity binding protein that is totally distinct from its receptors. CRF and Ucn1 bind to CRF-BP with very high affinity ( $K_d < 1$  nM), equal to or greater than the affinity of these ligands for CRF-R1 and/or CRF-R2. Interestingly, the CRF-BP is widely expressed in brain, localizing to sites of CRF or Ucn1 expression or release, sites of CRF-R1 or CRF-R2 expression, and sites where neither the receptors nor ligands are detected (Chan et al., 2000; Potter et al., 1992). CRF-BP synthesis and release are also highly responsive to environmental cues, as stress, CRF, glucocorticoids, and interleukin-6 all regulate CRF-BP expression (Westphal and Seasholtz, 2006). Yet we still know little about the molecular mechanisms of its actions. Purified CRF-BP attenuates CRF or Ucn1 activity at endogenous CRF receptors in cultured cells (Westphal and Seasholtz, 2006). In contrast, in vivo or slice studies suggest a facilitatory role for CRF-BP in VTA, with CRF<sub>6-33</sub> administration into the VTA decreasing ethanol consumption, CRF-induced relapse to cocaine, and CRF-mediated potentiation of N-methyl-D-aspartate excitatory postsynaptic currents on dopamine neurons (Albrechet-Souza et al., 2015; Ungless et al., 2003; Wang et al., 2007). Strikingly, many of the facilitatory effects of CRF-BP in VTA appear to be associated with CRF-R2 (Ungless et al., 2003; Wang et al., 2007). In fact, Ungless and colleagues (2003) have proposed that CRF-BP is required for CRF activation of CRF-R2 and the downstream phospholipase C/protein kinase C, but not protein kinase A, signaling pathway in VTA dopamine neurons. The study by Albrechet-Souza and colleagues (2015) begins to address this VTA CRF-BP/CRF-R2 interaction in the 2-bottle choice DID paradigm, but does not test CRF<sub>6-33</sub> alone in this protocol, making it difficult to ascertain whether CRF-BP and CRF-R2 are interacting.

Hence, while this interesting study by Albrechet-Souza and colleagues (2015) reveals new roles for the CRF system, especially CRF-BP and CRF-R2, in alcohol binge drinking, many questions remain. We propose that CRF-BP plays multiple roles in stress and alcohol addiction,

likely in a brain region- and cell type-specific manner, with CRF receptor subtype and signaling pathway as potential determinants. Clearly, additional *in vitro* and *in vivo* studies are needed to define the mechanisms of CRF-BP/CRF/CRF receptor interactions. Future studies should also include viral and genetic approaches allowing cell-type-specific overexpression or knockdown of CRF-BP, and/or CRF receptors, in various brain regions within the stress-reward pathway, as the transition from predependent to dependent states in alcohol addiction involves different sites and circuits. Together, these multidisciplinary approaches will allow us to better understand the multiple roles of the CRF-BP in inhibiting and facilitating CRF and/or Ucn1 interactions with CRF-R1 and CRF-R2 and the functional outcomes of these interactions in stress and addiction.

### ACKNOWLEDGMENTS

The authors thank Dr. Gwen Stinnett for critical reading of the manuscript. This work was supported by NIAAA INIA U01 AA013641 (AFS) and NSF Graduate Research Fellowship F031543 (KDK). The authors declare no conflict of interest.

### REFERENCES

- Albrechet-Souza L, Hwa LS, Han X, Zhang EY, Debold JF, Miczek KA (2015) Corticotropin releasing factor binding protein and CRF receptors in the ventral tegmental area: modulation of ethanol binge drinking in C57BL/6J mice. *Alcohol Clin Exp Res* 39: 1609–1618.
- Boorse GC, Kholdani CA, Seasholtz AF, Denver RJ (2006) Corticotropin-releasing factor is cytoprotective in *Xenopus* tadpole tail: coordination of ligand, receptor, and binding protein in tail muscle cell survival. *Endocrinology* 147:1498–1507.
- Chan RK, Vale WW, Sawchenko PE (2000) Paradoxical activational effects of a corticotropin-releasing factor-binding protein “ligand inhibitor” in rat brain. *Neuroscience* 101:115–129.
- Cortright DN, Nicoletti A, Seasholtz AF (1995) Molecular and biochemical characterization of the mouse brain corticotropin-releasing hormone-binding protein. *Mol Cell Endocrinol* 111:147–157.
- Hillhouse EW, Grammatopoulos DK (2006) The molecular mechanisms underlying the regulation of the biological activity of corticotropin-releasing hormone receptors: implications for physiology and pathophysiology. *Endocr Rev* 27:260–286.
- Hwa LS, Debold JF, Miczek KA (2013) Alcohol in excess: CRF(1) receptors in the rat and mouse VTA and DRN. *Psychopharmacology* 225:313–327.
- Karolyi IJ, Burrows HL, Ramesh TM, Nakajima M, Lesh JS, Seong E, Camper SA, Seasholtz AF (1999) Altered anxiety and weight gain in corticotropin-releasing hormone-binding protein-deficient mice. *Proc Natl Acad Sci U S A* 96:11595–11600.
- Lowery EG, Thiele TE (2010) Pre-clinical evidence that corticotropin-releasing factor (CRF) receptor antagonists are promising targets for pharmacological treatment of alcoholism. *CNS Neurol Disord Drug Targets* 9:77–86.
- Lowery-Gionta EG, Navarro M, Li C, Pleil KE, Rinker JA, Cox BR, Sprow GM, Kash TL, Thiele TE (2012) Corticotropin releasing factor signaling in the central amygdala is recruited during binge-like ethanol consumption in C57BL/6J mice. *J Neurosci* 32:3405–3413.
- Peto CA, Arias C, Vale WW, Sawchenko PE (1999) Ultrastructural localization of the CRF-binding protein in rat brain and pituitary. *J Comp Neurol* 413:241–254.
- Phillips TJ, Reed C, Pastor R (2015) Preclinical evidence implicating corticotropin-releasing factor signaling in ethanol consumption and neuroadaptation. *Genes Brain Behav* 14:98–135.
- Potter E, Behan DP, Fischer WH, Linton EA, Lowry PJ, Vale WW (1991) Cloning and characterization of the cDNAs for human and rat corticotropin releasing factor-binding proteins. *Nature* 349:423–426.
- Potter E, Behan DP, Linton EA, Lowry PJ, Sawchenko PE, Vale WW (1992) The central distribution of a corticotropin-releasing factor (CRF)-binding protein predicts multiple sites and modes of interaction with CRF. *Proc Natl Acad Sci U S A* 89:4192–4196.
- Ray LA (2011) Stress-induced and cue-induced craving for alcohol in heavy drinkers: preliminary evidence of genetic moderation by the OPRM1 and CRH-BP genes. *Alcohol Clin Exp Res* 35:166–174.
- Rhodes JS, Ford MM, Yu CH, Brown LL, Finn DA, Garland T Jr, Crabbe JC (2007) Mouse inbred strain differences in ethanol drinking to intoxication. *Genes Brain Behav* 6:1–18.
- Sparta DR, Hopf FW, Gibb SL, Cho SL, Stuber GD, Messing RO, Ron D, Bonci A (2013) Binge ethanol-drinking potentiates corticotropin releasing factor R1 receptor activity in the ventral tegmental area. *Alcohol Clin Exp Res* 37:1680–1687.
- Sutton SW, Behan DP, Lahrichi SL, Kaiser R, Corrigan A, Lowry P, Potter E, Perrin MH, Rivier J, Vale WW (1995) Ligand requirements of the human corticotropin-releasing factor-binding protein. *Endocrinology* 136:1097–1102.
- Ungless MA, Singh V, Crowder TL, Yaka R, Ron D, Bonci A (2003) CRF requires CRF-binding protein to potentiate NMDA receptors via CRF receptor 2 in dopamine neurons. *Neuron* 39:401–407.
- Wang B, You ZB, Rice KC, Wise RA (2007) Stress-induced relapse to cocaine seeking: roles for the CRF(2) receptor and CRF-binding protein in the ventral tegmental area of the rat. *Psychopharmacology* 193:283–294.
- Westphal NJ, Seasholtz AF (2006) CRH-BP: the regulation and function of a phylogenetically conserved binding protein. *Front Biosci* 11:1878–1891.