

RESEARCH

Open Access

Smoking decreases the response of human lung macrophages to double-stranded RNA by reducing TLR3 expression

Jill C Todt¹, Christine M Freeman^{1,2}, Jeanette P Brown¹, Joanne Sonstein¹, Theresa M Ames¹, Alexandra L McCubbrey³, Fernando J Martinez¹, Stephen W Chensue^{3,4,5}, James M Beck^{1,6} and Jeffrey L Curtis^{1,3,6*}

Abstract

Background: Cigarette smoking is associated with increased frequency and duration of viral respiratory infections, but the underlying mechanisms are incompletely defined. We investigated whether smoking reduces expression by human lung macrophages (M ϕ) of receptors for viral nucleic acids and, if so, the effect on CXCL10 production.

Methods: We collected alveolar macrophages (AM ϕ) by bronchoalveolar lavage of radiographically-normal lungs of subjects undergoing bronchoscopies for solitary nodules ($n = 16$) and of volunteers who were current or former smokers ($n = 7$) or never-smokers ($n = 13$). We measured expression of mRNA transcripts for viral nucleic acid receptors by real-time PCR in those AM ϕ and in the human M ϕ cell line THP-1 following phorbol myristate acetate/vitamin D3 differentiation and exposure to cigarette smoke extract, and determined TLR3 protein expression using flow cytometry and immunohistochemistry. We also used flow cytometry to examine TLR3 expression in total lung M ϕ from subjects undergoing clinically-indicated lung resections ($n = 25$). Of these, seven had normal FEV1 and FEV1/FVC ratio (three former smokers, four current smokers); the remaining 18 subjects (14 former smokers; four current smokers) had COPD of GOLD stages I-IV. We measured AM ϕ production of CXCL10 in response to stimulation with the dsRNA analogue poly(I:C) using Luminex assay.

Results: Relative to AM ϕ of never-smokers, AM ϕ of smokers demonstrated reduced protein expression of TLR3 and decreased mRNA for TLR3 but not TLR7, TLR8, TLR9, RIG-I, MDA-5 or PKR. Identical changes in TLR3 gene expression were induced in differentiated THP-1 cells exposed to cigarette smoke-extract in vitro for 4 hours. Among total lung M ϕ , the percentage of TLR3-positive cells correlated inversely with active smoking but not with COPD diagnosis, FEV1% predicted, sex, age or pack-years. Compared to AM ϕ of never-smokers, poly(I:C)-stimulated production of CXCL10 was significantly reduced in AM ϕ of smokers.

Conclusions: Active smoking, independent of COPD stage or smoking duration, reduces both the percent of human lung M ϕ expressing TLR3, and dsRNA-induced CXCL10 production, without altering other endosomal or cytoplasmic receptors for microbial nucleic acids. This effect provides one possible mechanism for increased frequency and duration of viral lower respiratory tract infections in smokers.

Trial registration: ClinicalTrials.gov NCT00281190, NCT00281203 and NCT00281229.

Keywords: Lung, Cigarette smoking, Effects, Toll-like receptors, Macrophages, Alveolar

* Correspondence: jlc Curtis@umich.edu

¹Division of Pulmonary & Critical Care Medicine, Department of Internal Medicine, University of Michigan Health Care System, Ann Arbor, MI 48109-2399, USA

³Graduate Program in Immunology, University of Michigan Health Care System, Ann Arbor, MI 48109-2399, USA

Full list of author information is available at the end of the article

Background

Viral respiratory infections are major risk factors for exacerbations of chronic obstructive pulmonary disease (COPD) and asthma [1,2], contributing to enormous societal costs due to healthcare utilization and reduced productivity. The frequency and duration of viral respiratory infections is increased by cigarette smoking [3-6] through multiple mechanisms, including reduced mucociliary clearance, impaired production of epithelial defensins, and decreased neutrophil chemotaxis {reviewed in [7,8]}. Smoking also negatively impacts host defense by alveolar macrophages (AM \emptyset), a key phagocyte and source of inflammatory mediators. Acting via both nicotinic cholinergic receptors [9] and uncharacterized mechanisms [10], smoking decreases pro-inflammatory cytokine release by human AM \emptyset [11-13]. Cigarette smoke extract (CSE) also blunts the ability of AM \emptyset to be activated by IFN- γ produced by T cells and natural killer (NK) cells [14]. Whether smoking has additional effects predisposing to viral respiratory infections is unknown.

We hypothesized that cigarette smoke could also reduce the ability of AM \emptyset to detect and activate innate responses to respiratory viruses. Cells recognize viruses by sensing double-stranded (ds) RNA using TLR3, which is expressed on endosomes and in some cell types, on the cell surface [15,16]. The other endosomal microbe-associated molecular pattern (MAMP) receptors, TLR7, TLR8 and TLR9, are also important for recognizing single stranded RNA and bacterial DNA with unmethylated CpG dinucleotides, respectively. Alternatively, dsRNA can also be recognized via cytosolic sensors, such as double-stranded RNA-dependent protein kinase R (PKR), retinoic acid-inducible gene I (RIG-I), or melanoma differentiation-associated gene 5 (MDA-5) [17].

Experimentally, the response to dsRNA is measured using the synthetic analogue polyinosinic acid:cytidylic acid (poly(I:C)), which stimulates secretion of the chemokine CXCL10 (formerly IP-10) [18-20], in some cell types in an entirely TLR3-dependent fashion [21]. CXCL10 is important to recruit and activate neutrophils, lymphocytes and NK cells [22-24], which are crucial to limit viral replication. At concentrations 100-fold higher than needed for chemotactic activity, CXCL10 also has defensin-like antimicrobial activities [25] and can enhance M \emptyset killing of intracellular *Leishmania* [26].

The purpose of this study was to determine the effect of smoking on endosomal and cytoplasmic receptors for nucleic acids. Our results demonstrate the susceptibility in human AM \emptyset of TLR3, but not of other receptors tested, to smoke-induced down-regulation.

Methods

Research subjects

Studies and consent procedures were performed in accordance with the Declaration of Helsinki at the VA Ann Arbor Healthcare System (all bronchoscopies and some clinically-indicated lung resections) and the University of Michigan Health System (some clinically-indicated lung resections, including all lung transplants and lung volume reduction surgeries) and were approved by their Institutional Review Boards (FWA 00000348 & FWA 00004969, respectively). All subjects understood the purpose of the study and gave written consent before any research procedures. All subjects underwent a complete history and physical examination by a Pulmonologist, spirometry, chest imaging, prospective collection of medication history, and complete blood count with differential, coagulation studies and chemistry panel.

Bronchoalveolar lavage (BAL) cohort

We recruited 13 healthy never-smoker volunteers and 23 subjects with a smoking history who either were scheduled to undergo bronchoscopy in the evaluation of solitary pulmonary nodules ($n = 16$) or who volunteered ($n = 7$). Among the 23 subjects who had ever smoked, 18 were active smokers (<six months since quitting) and 5 were former smokers (>six months since quitting). Of those with any smoking history, subjects ($n = 10$) with ≥ 10 pack years, a ratio of forced expiratory volume in 1 second to forced vital capacity (FEV₁/FVC) >0.7, normal spirometry, and no clinical diagnosis of COPD represent control smokers. Subjects ($n = 13$) with a smoking history, FEV₁/FVC <0.7 and abnormal spirometry were considered to have COPD. All subjects were without evidence of lung infection, interstitial lung disease or collagen vascular disease.

Importantly, not all types of experiments were performed on cells from every subject in this cohort, and conversely, some subjects were used for more than one type of experiment. The characteristics of the BAL subjects used in each type of experiments are summarized in Tables in the Results sections; demographic and clinical data for this entire cohort are shown in Additional file 1: Table S1.

Lung tissue cohort

Lung tissue was collected from consented subjects undergoing clinically-indicated resections for pulmonary nodules, lung volume reduction surgery, or lung transplantation ($n = 25$). Using the same definitions as in the BAL cohort, this cohort comprised seven control smokers and 18 subjects with COPD. This cohort was used exclusively for flow cytometric analysis of total lung M \emptyset . Characteristics of this cohort are summarized in the

Results section, and demographic and clinical data for individual subjects are shown in Additional file 2: Table S2.

BAL procedure and cell preparation

BAL was performed in the right middle lobe and lingula of the volunteers and in whichever of these sites was contralateral to the nodule in the clinically-indicated bronchoscopies, in which the research BAL was performed as early as feasible, and always before any biopsies or brushings. We instilled 100 ml normal saline per site, using a 30 ml syringe and gentle manual suction. BAL fluid was filtered through sterile gauze to remove mucous, and cells were washed thrice with PBS, with centrifugation between washes.

BAL cells (>95% AM \emptyset by Wright-Giemsa-stained cytopins) were either immediately processed for flow cytometry and immunohistochemistry or were cultured briefly to purify AM \emptyset by adherence. For this purpose, cells were resuspended in complete medium (RPMI 1640 containing 25 mM HEPES, 2 mM l-glutamine, 1 mM pyruvate, 100 U/ml penicillin/streptomycin, 10% heat-inactivated AB human serum (all from Invitrogen, Carlsbad, CA), and 55 μ M 2-ME (Sigma Chemical, St. Louis, MO), and were plated at 2×10^5 cells/well in sterile 24-well plates. Cells were incubated for 1.5 h in 5% CO₂ at 37°C, washed to remove nonadherent cells, then incubated in AIM-V serum-free medium (Invitrogen) under experimental conditions. For TLR3 stimulation, adherence-purified AM \emptyset were cultured in AIM-V alone or with poly(I:C) at 50 μ g/ml for 24 h.

Lung sample preparation

Lung tissue was dissected free of any areas containing nodules, cancers or evidence of infection by a Pathologist before being released to the study. Lung sections weighing approximately 3 g were dispersed using a Waring blender in a biosafety cabinet without enzyme treatments, which we have previously shown produces single cell suspensions of high viability and functional capacity [27,28]. Cells were filtered through a 40 μ m strainer to remove debris and were resuspended in staining buffer (2% FBS in PBS) for flow cytometry.

Differentiation and treatment of THP-1 cells

THP-1 cells obtained from ATCC (Manassas, VA) were grown according to the supplier's guidelines. Cells were plated at 4×10^5 cells/well in 24-well plates, and after adherence, were differentiated by exposure to 150 ng/ml phorbol myristate acetate (PMA) and 0.01 μ M vitamin D3 for 48 hrs, and then were washed with fresh complete medium before exposure to various concentrations of cigarette smoke-extract (CSE) for 4 hr. Fresh CSE was prepared by bubbling the mainstream smoke of two University of Kentucky Tobacco Health Research

Institute standardized cigarettes (lot 2R4F), using a Jaeger-Baumgartner Cigarette Smoking Machine (C.H. Technologies, New Jersey) driven by dry compressed air, into 20 ml RPMI 1640; the result was defined as 10% CSE. CSE was filter-sterilized using a 0.22 μ m membrane and was used the same day.

Quantitative real-time PCR

RNA was isolated using RiboPure kits (Applied Biosystems, Foster City, CA), TURBO DNA-free (Applied Biosystems) to remove genomic DNA, and Retroscript kits (Applied Biosystems) for reverse-transcription. We performed quantitative real-time PCR using the Stratagene Mx3000P (LaJolla, CA), with human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) as the endogenous reference, and commercial primer-probe sets (Taqman chemistry, Applied Biosystems). Transcript levels were reported relative to a calibrator of unstimulated THP cells, or AM \emptyset from never smokers, as appropriate, as calculated on the thermocycler.

Flow cytometry

BAL cells were resuspended in 100 μ L staining buffer {1% FA Buffer (BD Bioscience, San Jose, CA), 1% FCS, 0.01% sodium azide} per tube. We used antibodies against the following antigens (clones in parentheses): CD45 (2D1) (eBioscience, San Diego, CA), TLR3 (40C1285.6), TLR7 (polyclonal), TLR9 (26C593.2) (Imgenex Corp., San Diego, CA), and TLR8 (303F1.14) (Dendritics, Lyon, France). To detect intracellular receptor expression, cells were treated with Fixation and Permeabilization Buffers (eBioscience). In all experiments, we used isotype-matched controls, analyzed cells on an LSR II flow cytometer (BD Bioscience), and collected a minimum of 10,000 CD45+ events per sample, using FACS Diva software with automatic compensation, and FlowJo analysis software (Tree Star, Inc., Ashland, OR). Details of instrument setup have been described recently [29].

Immunohistochemistry

Cytopins were immersed in cold acetone followed by methanol plus hydrogen peroxide. The slides were stained with biotinylated anti-human TLR3 (Imgenex) or isotype control antibodies, followed by Universal ABC peroxidase complex and 3-amino-9-ethylcarbazole peroxidase substrate. Slides were counter-stained with hematoxylin and photographed using an Olympus BX51 digital camera.

CXCL10 analysis

CXCL10 levels were determined using Biosource Multiplex Assays (Invitrogen) and a Luminex 200™ (Luminex Corporation, Austin, TX).

Statistical analyses

Statistical analysis was performed using GraphPad Prism (GraphPad, La Jolla, CA) except as specified. We used the nonparametric unpaired Mann–Whitney test to determine differences between two groups (smokers versus never-smokers; healthy subjects versus COPD subjects) in mRNA and flow cytometry data; the nonparametric Kruskal-Wallis one-way ANOVA to compare multiple conditions on THP cells in vitro; and ANOVA with Bonferroni's Multiple Comparison post-hoc testing to determine differences between smokers and never-smokers in stimulated cytokine secretion. We used Spearman non-parametric analysis to correlate TLR3 mRNA and protein expression with clinical variables and performed linear regression using SPSS (IBM Corp.; Armonk, NY). A two-tailed p value of < 0.05 was considered significant.

Results

TLR3 expression is selectively decreased in AM \emptyset of smokers

To test whether cigarette smoking impacts expression by human AM \emptyset of TLRs implicated in defense against respiratory viruses, we first analyzed adherence-purified AM \emptyset obtained by BAL of 11 current or former smoking subjects and six never-smokers (Table 1) using quantitative real-time PCR. Results showed significantly decreased TLR3 mRNA transcripts in smokers compared with never-smokers ($p = 0.0015$; Mann–Whitney test) (Figure 1A), but no significant differences in transcripts for TLR7, TLR8 or TLR9, which are also found in the endosome (Figure 1A). We also found no significant differences between smokers and never-smokers in mRNA transcripts for the cytoplasmic dsRNA receptors RIG-I, MDA-5 and PKR (Additional file 3: Figure S1).

These findings were supported by two measures of AM \emptyset protein expression. Flow cytometry (Additional file 4:

Figure S2) permitted identification of specific staining for TLR3 and TLR9 amongst AM \emptyset of never-smokers, although expression of TLR7 was very low (Additional file 4: Figure S2, middle panel) in both groups, and was not analyzed further. Comparing BAL samples from smokers ($n = 13$), and never-smokers ($n = 10$) (Table 2), we found a significantly decreased percentage ($p < 0.0001$, Mann–Whitney test) of AM \emptyset positive for intracellular TLR3 in AM \emptyset of smokers (Figure 1B). Note that these experiments contain some of the same subjects studied in Table 1, as well as other BAL subjects, as described in *Methods*; for additional details about individual subjects, see Additional file 1: Table S1). TLR3 was not detected on the surface of unpermeabilized AM \emptyset of either group (not shown). There were no differences between smokers ($n = 6$) and never-smokers ($n = 5$) in the percentage of AM \emptyset expressing intracellular TLR9 (Figure 1B) (or TLR7 and TLR8, data not shown). Similarly, immunohistochemical staining of BAL cytopins (Additional file 1: Table S1) demonstrated greatly reduced TLR3 expression in the AM \emptyset from smokers, relative to AM \emptyset from the never-smokers (Figure 1C). These independent data indicate that reduction in TLR3 protein expression measured by flow cytometry did not result simply from greater autofluorescence in AM \emptyset of smokers.

In univariate analyses including both smokers and never-smokers, TLR3 RNA transcripts correlated inversely with both FEV $_1$ % predicted and subject age (Figure 2A-B). Considering only those with a history of smoking, there was no correlation with pack-years (Figure 2C). However, in a linear regression, none of the variables (smoker versus never-smoker, age, sex, FEV $_1$ % predicted) reached statistical significance for mRNA transcripts. AM \emptyset positivity for TLR3 protein did not correlate in univariate analyses with FEV $_1$ % predicted (Figure 2D) or pack-years of smoking exposure (Figure 2F), but did correlate with subject age (Figure 2E). In a linear regression, smoking status (smoker vs. never-smoker) was strongly associated with TLR3+ AM \emptyset ($p < 0.0001$) and subject age was also significant ($p = 0.035$), but sex and FEV $_1$ % predicted remained insignificant. When we analyzed smokers with COPD versus smokers without COPD, no significant differences were seen in TLR3 mRNA transcripts (COPD, 0.23 ± 0.16 vs. no COPD, 0.49 ± 0.49 , mean \pm SEM dRn; $p = 0.52$, unpaired t test) or flow cytometric results (COPD, 12.2 ± 12.2 vs. non-COPD, 10.4 ± 5.2 ; mean \pm SEM % TLR3-positive AM \emptyset , $p = 0.87$, unpaired t test). Despite the use of inhaled corticosteroids (ICS) by ~50% of the smoking subjects, the decrease in the fraction of TLR3+ AM \emptyset of smokers was not a result of steroid usage ($p = 0.79$, determined by t -test comparing TLR3 mRNA transcript expression between ICS users and non-users; $p = 0.92$ when comparing TLR3 protein measurements by flow cytometry).

Table 1 Characteristics of BAL subjects used for RNA studies

Group	Smokers	Never-smokers	p value
Subjects, n	11	6	
Sex ratio, M/F	11/0	2/4	0.003
Age, years (SD)	65.5 (7.2)	37.7 (12)	0.002
Smoking, pack-years (SD)	65.5 (23.1)	0 (0)	0.0008
Smoking status (Former/Current)	2/9	0/0	0.003
FEV $_1$, % pred. (SD)	57 (22.1)	101 (8.7)	0.002
FEV $_1$ /FVC% (SD)	58.5 (15.6)	86.8 (4.9)	0.0007
ICS use (yes/no)	7/4	0/6	0.02

ICS Inhaled Corticosteroid Use, M Male, F Female. Data are represented as mean (SD), ratio of current smokers to former smokers, or fraction of ICS users (yes/no). All FEV $_1$ values are pre-bronchodilator. p values calculated using Mann Whitney test.




Figure 1 AMø of current smokers show reduced expression of TLR3 mRNA transcripts and intracellular protein relative to AMø of never-smokers. **A.** mRNA transcripts. RNA was isolated from AMø, depleted of contaminating genomic DNA, reverse-transcribed and analyzed by quantitative real-time RT-PCR using Taqman chemistry and specific primer-probe sets, normalized to GAPDH transcripts. Data are expressed on the horizontal axis as mean \pm SEM for relative quantity (dRn), calculated in comparison to a single never-smoker who was arbitrarily designated the reference sample. Never-smokers ($n = 6$), red bars; smokers ($n = 11$), blue bars. *, $p < 0.05$ by Mann-Whitney test. **B.** Intracellular TLR3 (top panel) and TLR9 (bottom panel) expression by flow cytometry. AMø were permeabilized, stained for TLR3 and TLR9 expression and analyzed by flow cytometry. Data are expressed as TLR-positive AMø as mean \pm SEM, never-smokers ($n = 10$ for TLR3, $n = 5$ for TLR9), red circles; smokers ($n = 13$ for TLR3, $n = 6$ for TLR9), blue circles. *, $p < 0.05$ by Mann-Whitney test. **C.** Immunohistochemical staining. Cytospins from a representative smoker (right) and a never-smoker (left), stained for intracellular TLR3 expression using AEC (red product) and hematoxylin (original 20 X). Top row, isotype control staining, bottom row, specific TLR3 staining. Representative of three experiments with similar results.

Thus, these secondary analyses supported a significant effect of smoking on AMø expression of TLR3. However, due to the distribution of our subjects, we were unable to exclude the possibility that our findings resulted from the preponderance of women in the never-smoker group and of men among the smokers in this cohort, or from the significant difference in ages between the volunteer and clinical subjects.

Reduced TLR3 expression in lung Mø correlates with smoking status but not COPD stage

In apparent contradiction to our findings, a recent paper primarily using immunohistochemistry in lung tissue sections reported that the percentage of TLR3-positive lung Mø were significantly increased in smokers compared with never-smokers [30]. To address this discrepancy and the questions about the possible effects of age and sex in

Table 2 Characteristics of BAL subjects used for flow cytometry studies

Group	Smokers	Never-smokers	<i>p</i> value
Subjects, <i>n</i>	13	10	
Sex ratio, M/F	11/2	1/9	0.0006
Age, years (SD)	60.3 (8.9)	42.2 (13.7)	0.002
Smoking, pack-years (SD)	47.4 (33.7)	0 (0)	0.0001
Smoking status (Former/Current)	3/10	0/0	0.0001
FEV ₁ , % pred. (SD)	83.5 (23.6)	100.7 (10.5)	0.077
FEV1/FVC% (SD)	72.5 (14.3)	84.2 (5.7)	0.0249
ICS use (yes/no)	4/9	0/10	N.S.

ICS Inhaled Corticosteroid Use; M Male, F Female. Data are represented as mean (SD), ratio of current smokers to former smokers, or fraction of ICS users (yes/no). All FEV₁ values are pre-bronchodilator. *p* values calculated using Mann Whitney test.

our BAL subjects, we studied a different cohort of subjects (*n* = 25) undergoing clinically-indicated lung resection procedures (Table 3). This cohort also had the advantage of having more nearly balanced ratios of male and female subjects, as well as a wide range of spirometry values. We used mechanical disaggregation of tumor-free lung parenchyma to produce a single cell suspension of high viability containing both AM ϕ and interstitial lung M ϕ , which we identified as autofluorescent, CD45+, high side scatter cells.

Flow cytometric analysis permitted objective quantification of the percentage of individual lung M ϕ s positive for intracellular TLR3, relative to simultaneously analyzed isotype-control monoclonal antibodies (Figure 3A, B). We found the percentage of TLR3-positive lung M ϕ s from current smokers was significantly reduced (*p* = 0.006) compared to the percentage of TLR3-positive lung M ϕ s from former smokers (Figure 3C). By contrast, considering these same individuals, there were no significant differences in percentages of lung M ϕ expressing TLR3 between those with normal pulmonary function and those with COPD (Figure 3D). Nor was there a correlation between the percentage of TLR3-positive lung M ϕ by flow cytometry and either FEV₁ % predicted or duration of smoking in pack-years (Additional file 5: Figures S3A & B). In a linear regression, smoking status (current vs. former) was significantly associated with TLR3 positivity (*p* = 0.016) and was not affected by age, FEV₁ % predicted, sex, or pack-years. These independent data agree with and extend results of our BAL experiments and collectively show that active smoking reduces TLR3 expression by resident human lung M ϕ .

Exposure to CSE specifically reduces expression of TLR3 mRNA transcripts by differentiated human THP-1 cells

To explore the effect of acute smoke-exposure in vitro, we used the well-characterized system of differentiating


the human M ϕ cell line THP-1 by treatment with PMA plus vitamin D3. In four independent experiments, we found that CSE at concentrations of 1.25% and 2.5% significantly decreased expression of TLR3 transcripts (Figure 4A), but had no effect on transcripts of TLR7, TLR8 or TLR9 (Figures 4B-D). CSE at these concentrations had no effect on viability as assessed by trypan blue exclusion (not shown). CSE also had no significant effect on mRNA transcripts for RIG-I, MDA-5 or PKR at concentrations up to 2.5% CSE (Figure 4E-G). Thus, the effect of smoke exposure on TLR3 mRNA levels can be induced within as little as four hours in THP-1 cells differentiated to a mature M ϕ phenotype. These findings indicate that smoking reduces M ϕ TLR3 expression directly, and not secondarily due to an effect on another lung cell type or on the lung microbiome.

AM ϕ of smokers show reduced CXCL10 production in response to poly(I:C) stimulation in vitro

Finally, we investigated whether decreased TLR3 expression on the AM ϕ of smokers would affect CXCL10 production following poly(I:C) stimulation. These experiments were performed using AM ϕ from subjects in the BAL cohort (Additional file 2: Table S2). CXCL10 concentrations were close to the level of detection in unstimulated cells from both smokers (*n* = 5) and never-smokers (*n* = 4) (Table 4) and did not differ significantly (Figure 5). Following poly(I:C) stimulation, AM ϕ from never-smokers showed a 1000-fold increase in CXCL10 production, and differed significantly from the response of AM ϕ from smokers, which showed only a 10-fold increase from unstimulated levels (Figure 5). These results illustrate a functional consequence of the difference in TLR3 expression between the two groups.

Discussion

We demonstrate that smoking reduces expression by human lung M ϕ of TLR3, a receptor for microbial dsRNA and endogenous danger signals. Relative to AM ϕ of never-smokers, AM ϕ of current smokers showed reduced TLR3 mRNA and protein, and lower secretion of CXCL10 in response to the viral dsRNA analogue poly(I:C). Smoking did not significantly reduce AM ϕ expression of other receptors for viral nucleic acids in the cytoplasm (RIG-I, MDA-5 or PKR) or endosome (TLR7, TLR8 or TLR9). In a separate cohort, TLR3 protein expression by total lung M ϕ was also reduced in active smokers, relative to former smokers. TLR3 protein expressed as the percentage of positive lung M ϕ did not correlate significantly in either cohort with a diagnosis of COPD, FEV₁% predicted or total history of smoking expressed as pack-years. The direct nature of the smoking effect was shown in vitro using differentiated cells of the human M ϕ line THP-1. These findings identify a



new mechanism by which smoking impairs anti-viral lung host defenses.

Our findings address the intersection of two highly prevalent clinical issues, host defense against respiratory viruses and smoking. Respiratory viruses are implicated, alone or as mixed infections, in most COPD exacerbations, including those of greatest severity [31,32]. Viral respiratory infections are also the leading cause of asthma exacerbations in adults and children [33,34], including those requiring hospitalization [35]. Although

only some smokers develop COPD, those who continue to smoke sustain more frequent respiratory infections and faster lung function decline [36]. Smoking is also a key contributor to morbidity in asthma [37]. Asthmatic smokers experience worse symptom control, greater risk of hospitalization, and increased resistance to the therapeutic effects of inhaled corticosteroids [38-41]. Finally, even in individuals with normal spirometry, smoking is an independent risk factor for increased number and severity of respiratory infections [3]. Hence, these results

Table 3 Characteristics of surgical lung tissue subjects


Group	Smokers with COPD	Smokers without COPD	<i>p</i> value
Subjects, <i>n</i>	18	7	
Sex ratio, M/F	10/8	5/2	0.49
Age, years (SD)	61.8 (9.4)	54.4 (9.1)	0.74
Smoking, pack-years (SD)	60.3 (41.4)	44.5 (23.9)	0.47
Smoking status (Former/Current)	4/14	3/4	0.11
FEV ₁ , % pred. (SD)	41.8 (27.7)	95.4 (11.2)	0.0008
FEV1/FVC% (SD)	42.5 (18.6)	76.0 (5.2)	0.0002
ICS use (yes/no)	12/6	1/6	0.02

ICS Inhaled Corticosteroid Use, M Male, F Female. Data are represented as mean (SD), ratio of current smokers to former smokers, or fraction of ICS users (yes/no). All FEV₁ values are pre-bronchodilator. *p* values calculated using Mann Whitney test.

provide another rationale to urge all smokers to quit smoking absolutely, regardless of pulmonary function or specific diagnosis.

Optimal defense against respiratory viruses requires coordination of multiple elements of innate and adaptive immunity [42], especially prompt production by epithelial

cells of types I (alpha/beta) and III (lambda) IFNs [43-45]. Appropriate contributions by AM ϕ are likely also important [42]. In response to viruses, AM ϕ produce measurable amounts of type I IFNs [46] and of the type III IFN lambda 1 (IL-29) [47]. Human monocyte-derived M ϕ (MDM ϕ) produce mRNA for type III IFNs in response to






Figure 4 Exposure to CSE specifically and rapidly reduces TLR3 gene expression in a differentiated human M ϕ cell line. THP-1 cells differentiated using PMA & vitamin D3 for 48 hours were cultured in complete medium alone or with various concentration of CSE for an additional 4 hr, then analyzed by quantitative real-time RT-PCR. Data are mean \pm SEM of four independent experiments, expressed as dRn (relative to differentiated THP-1 cells cultures in complete medium alone). **A**, TLR3; **B**, TLR7; **C**, TLR8; **D**, TLR9; **E**, RIG-1; **F**, MDA-5; **G**, PKR. Kruskal-Wallis one-way ANOVA was used to compare groups. *, $p < 0.05$, compared to cells that did not receive CSE exposure.

stimulation via TLR3 or TLR4 [48] or to viruses [47,49]. A specific anti-viral role of AM ϕ has been confirmed experimentally by cellular depletion in two animal models [50,51], but in a murine model of respiratory syncytial virus pneumonia, AM ϕ depletion reduced early antiviral responses, increased viral load, but had no effect on late outcomes [52]. Hence, the significance of the current findings may vary with the particular viral pathogen. Additional studies will be needed to determine whether the effect we found on TLR3 expression translates into defective responses by the AM ϕ of smokers to intact respiratory viruses.

These results expand the known effects of cigarette smoking on AM ϕ function, which are clearly complex. Two studies using Affymetrix Human Genome U133 Plus 2.0 GeneChips found AM ϕ from human smokers to have distinctive, globally altered gene expression profiles relative to non-smokers [53,54]. Interestingly, TLR3 is not mentioned as a differentially expressed gene in either study. Disparity from our results likely relates to the difference in methodology; importantly, we showed very reduced TLR3 protein expression. The more pronounced reduction we found in TLR3 protein relative to mRNA implies the possibility of post-transcriptional

Table 4 Characteristics of BAL subjects used for in vitro stimulation studies

Group	Smokers	Never-smokers	p value
Subjects, n	5	4	
Sex ratio, M/F	5/0	2/2	0.13
Age, years (SD)	60 (11)	39.5 (12.1)	0.06
Smoking, pack-years (SD)	47.4 (35.2)	0 (0)	0.02
Smoking status (Former/Current)	0/5	0/0	0.007
FEV ₁ , % pred. (SD)	66.6 (30.6)	106.5 (12)	0.06
FEV1/FVC% (SD)	61.8 (19)	81.0 (7.5)	0.10
ICS use (yes/no)	3/2	0/4	0.10

ICS Inhaled Corticosteroid Use, M Male, F Female. Data are represented as mean (SD), ratio of current smokers to former smokers, or fraction of ICS users (yes/no). All FEV₁ values are pre-bronchodilator. p values calculated using Mann Whitney test.

smoking-induced downregulation, an important issue for future studies.

AM \emptyset of human smokers are often described as being “activated” [55-57], but both they and AM \emptyset from COPD patients show impaired bacterial uptake and killing [58,59]. Production of TNF- α and IL-6 by AM \emptyset of smokers (and COPD patients, which often mixed current and former smokers) equaled or exceeded that of non-smokers, but was strikingly less inhibitable by steroids [60,61]. Increased secretion of IL-8 and other neutrophil-

attracting chemokines by AM \emptyset of human smokers has been shown in most [62,63] but not all [64] studies. Thus, smoking appears to make AM \emptyset less likely to respond efficiently to pathogens but more likely to enhance sustained inflammation.

TLR3 has been shown experimentally to be essential for optimal protection against some but not all viruses, even those with a dsRNA genome [17]. Moreover, under some circumstances TLR3 may be responsible for exaggerated inflammation [65], and anti-TLR3 blocking antibody is being developed for therapeutic use [66]. Importantly, TLR3 is a receptor not only for microbial dsRNA, but also for dsRNA from necrotic host cells [67], an example of an endogenous danger-associated molecular pattern. This finding assumes increased significance based on the recognition of increased apoptotic cell death in emphysema [68] and decreased clearance of apoptotic cells by AM \emptyset of smokers [69-71], a defect postulated to foster lung injury and emphysema progression [71,72]. Hence, the putative detrimental effect of reduced TLR3 expression by AM \emptyset of smokers on antiviral defenses might be partially offset by reduced responsiveness to necrosis-induced lung inflammation, an intriguing possibility that will require considerably greater investigation. Our finding that the percentage of TLR3-positive AM \emptyset was increased in former smokers when compared to active smokers agrees with the partial resolution of other smoking-induced AM \emptyset defects on smoking cessation [73]. These data are also congruent with the observation that smoking but not COPD decreases AM \emptyset expression of TLR2, but not of TLR4 [74]. The molecular mechanisms for down regulation by active smoking of TLR3 (this study) and TLR2 [74] remain unclear.

Our data partially differ from those of Koarai and colleagues, who recently reported increased TLR3 expression by AM \emptyset of smokers that correlated positively with smoking history and inversely with DLCO but not FEV₁ % predicted [30]. Our results agree in finding no differences in AM \emptyset TLR3 expression among smokers based on the presence or absence of spirometrically-defined COPD. That study examined paraffin-fixed sections using a polyclonal anti-TLR3 antibody and immunohistochemistry, whereas we analyzed freshly isolated cells using the same monoclonal antibody in both of our cohorts. Hence, the disparity regarding TLR3 expression results might relate to antigen availability. Our studies differ in the cell type used for in vitro exposure to CSE (in that study MDM \emptyset), incubation time (24 hr vs. 4 hr in the current study) and CSE concentration (5% vs. a significant effect at 1.25% in the current study). These disparities make comparison difficult, as do multiple differences in design and endpoints of the experiments testing poly(I:C)-stimulated mediator production. One other study found that CSE reduced




Figure 5 Production of CXCL10 by AM \emptyset of smokers in response to poly(I:C) stimulation is impaired. AM \emptyset from never-smokers (n = 4) (white and red columns) and smokers (n = 5) (light and dark blue columns) were cultured for 24 h in either complete medium alone or with poly(I:C). Supernatants were collected and CXCL10 protein was measured by Luminex assay; results are expressed as the fold-increase in CXCL10 over the unstimulated control condition; note the logarithmic scale of the vertical axis. Data are mean \pm SEM. One-way ANOVA with Bonferroni's Multiple Comparison post-hoc testing was used to compare differences among groups, *, p < 0.01.

mRNA expression by MDM \emptyset of TLR8 and MDA5 and that MDM \emptyset of smokers had reduced mRNA expression of RIG-I and MDA-5, relative to MDM \emptyset of never-smokers [75]. That study does not mention TLR3 expression, which presumably was not altered in either case. Differences from our results likely reflect disparity both in cell type studied and possibly in methodology, as those results were obtained by Affimetrix assay.

The current findings are interesting in regard to two recent murine studies. Gaschler and colleagues found attenuated *in vitro* secretion of TNF- α , IL-6 and CCL5 in response to poly(I:C) by AM \emptyset from smoke-exposed mice, but did not find decreased TLR3 expression [76]. That disparity from our results may reflect a species difference. Kang and colleagues found that cigarette smoke selectively augmented airway and alveolar inflammation induced by viral MAMPs and influenza virus, as measured by induction of type I IFN, IFN- γ , IL-18 and IL-12/IL-23 p40 and PKR activation [77]. Further analysis using knockout mice indicated a role for both acute TLR3-dependent and chronic TLR3-independent pathways, as well as a pathway dependent on mitochondrial antiviral signaling protein (MAVS), IL-18R α , IFN- γ and PKR [77]. That important study, however, did not measure TLR3 expression by AM \emptyset , and its findings likely reflect the interaction of multiple cell types *in vivo*.

Considerable evidence indicates the importance of CXCL10 in antiviral host defense but also in development or prevention of certain lung diseases [78,79]. CXCL10 is one of three chemokines, along with CXCL9 and CXCL11, that are highly induced by IFN- γ , as well as by types I and III IFNs [49,80]. Of note, however, CXCL10 production is also induced directly by infection of human AM \emptyset by rhinovirus, of murine AM \emptyset by RSV and of human MDM \emptyset by influenza or HIV [19,20,81-83], making this cytokine a suitable endpoint to test the functional importance of the observed TLR3 downregulation. A key action of CXCL10 is inflammatory cell recruitment, acting via CXCR3, which is found on M \emptyset , CD8 $^+$ T cells, activated CD4 $^+$ T cells (especially T $_{H1}$ cells), NK cells and plasmacytoid dendritic cells [84]. CXCR3 is essential for recruitment of antigen-specific CD4 $^+$ T cells to the lungs in a murine model of parainfluenza pneumonia [85]. In a model of Coxsackievirus infection, CXCL10 was shown to limit viral replication via NK cell recruitment [24].

Markedly reduced induction of CXCL10 in AM \emptyset from smokers is of interest in the context of other recent human studies. CXCL10 is one of several M1-associated genes down-regulated in AM \emptyset of healthy smokers and even more so, in smokers with COPD, relative to healthy never-smokers [54]. Although significantly increased CXCL10 concentrations have been found in the sputum of COPD patients, when compared with nonsmokers but not with smokers without obstruction [86], the

origin of that chemokine is unclear, as it is also produced by epithelial cells that significantly outnumber M \emptyset in the airways. Both by showing the role of reduced TLR3 expression and by studying AM \emptyset , we extend a study of gene expression by MDM \emptyset , which found reduced expression of CXCL10 at baseline in smokers relative to never-smokers, and also that CSE reduced IFN- γ -induced CXCL10 production by MDM \emptyset of never-smokers [75].

Our study has several limitations. One is the absence of never-smokers from our surgical cohort, who were recruited before clinically-indicated resections. Another is the predominance in our BAL cohort of female never-smokers (11 of 13 subjects) versus a predominance of male current smokers with preserved lung function (eight of nine subjects) and COPD patients (11 of 11 subjects). Concerns that differences in sex or age could confound our BAL results should be reduced by the congruent effect of smoking in our surgical cohort, which comprised 40% female subjects and which showed no significant effect of age on TLR3 expression by lung M \emptyset . Moreover, at equivalent levels of tobacco exposure, women appear to be at greater risk of lung function impairment [reviewed in [87]], the opposite of what would be expected if the current finding were a male-specific effect. Although we cannot formally exclude the possibility that differences in subject age explain the reduced expression of CXCL10 in response to TLR3, we consider that possibility unlikely, as serum levels of CXCL10 increase with age [88,89]. Other limitations are the absence of DLCO measurements, which precludes comparison of our results with those of Koarai and colleagues [30] on this point, and that because post-bronchodilator FEV1 values were not available on all subjects, we have presented and analyzed exclusively pre-bronchodilator FEV1 values.

Conclusions

Active smoking reduces expression of TLR3 by human lung M \emptyset , assayed both as AM \emptyset harvested by BAL and as a mixture of AM \emptyset and interstitial lung M \emptyset from surgical tissue. This effect occurs via both down-regulation of mRNA transcripts, which can be induced *in vitro* within four hours, and from a more pronounced effect on TLR3 protein expression, which appears to be at least partially reversible on smoking cessation. Reduction in lung M \emptyset TLR3 expression may be one mechanism contributing to the increased incidence of viral respiratory infections in smokers and to viral induction of acute exacerbations of COPD. Defining the effect of smoking on the phenotype of human AM \emptyset and other lung cell types is a crucial step in the translation of basic science into therapies.

Endnotes

Subjects were recruited via observational studies registered with ClinicalTrials.gov as NCT00281190, NCT00281203 and NCT00281229. These data were presented in part at the International Scientific Conference of the American Thoracic Society, May 17, 2010 in New Orleans, LA, and have been published in abstract form *Am J Respir Crit Care Med* 2010; 181: A3875.

Supported by R01 HL082480 (JLC, FJM), R01 HL056309 (JLC) and U01 HL098961 (JMB, JLC) from the USPHS; a Career Development Award (CMF) and a Merit Review Award (CMF) and a Research Enhancement Award Program (SWC, JMB, JLC) from the Biomedical Laboratory Research & Development Service, Department of Veterans Affairs. These investigations were also supported in part by the Tissue Procurement Core of the University of Michigan Comprehensive Cancer Center, Grant P30 CA46952, and by the Lung Tissue Research Consortium (Clinical Centers), Grant N01 HR046162.

Current e-mail addresses: James M. Beck, M.D.: james.beck@denver.edu.

Additional files

Additional file 1: Table S1. Summary of all BAL subjects demographics, smoking histories, spirometry & current smoking status. ICS, inhaled corticosteroid use; FC, flow cytometry; IHC, immunohistochemistry. Summarized data are shown in bold and are represented as mean (SD), ratio of current smokers to former smokers, or fraction of ICS users (yes/no). All FEV1 values are pre-bronchodilator.

Additional file 2: Table S2. Summary of all lung tissue subject demographics, smoking histories, spirometry & current smoking status. ICS, inhaled corticosteroid use. M, Male; F, Female. Summarized data are shown in bold and are represented as mean (SD), ratio of current smokers to former smokers, or fraction of ICS users (yes/no). All FEV1 values are pre-bronchodilator.

Additional file 3: Figure S1. AM ϕ of current smokers show no reduction in mRNA expression of cytoplasmic dsRNA receptors, relative to AM ϕ of never-smokers. RNA from AM ϕ was isolated, depleted of contaminating genomic DNA, reverse-transcribed and analyzed by quantitative real-time RT-PCR using Taqman chemistry and specific primer-probe sets, normalized to GAPDH transcripts. Data are expressed on the horizontal axis as mean \pm SEM for relative quantity (dRn), calculated in comparison to a single never-smoker who was arbitrarily designated the reference sample. Never-smokers ($n = 6$), red bars; smokers ($n = 11$), blue bars. The Mann-Whitney test was used to calculate statistical significance.

Additional file 4: Figure S2. Representative flow cytometry results. AM ϕ were permeabilized, stained for expression of TLR3 (left-hand panels), TLR 7 (middle panel) and/or TLR9 (right-hand panels), and analyzed by flow cytometry, gating on AM ϕ (CD45+, high side scatter cells). A,B; specific staining (red line), isotype control staining (blue line). A, never-smoker, B, smoker.

Additional file 5: Figure S3. Lack of correlation of spirometry or total smoking history with lung AM ϕ TLR3 expression in surgical cohort. Total lung M ϕ were harvested from excess lung tissue removed surgically for clinical indications as described in the legend to Figure 3. Lung M ϕ were permeabilized, stained for TLR3 expression and analyzed by flow cytometry, gating on CD45+, high side-scatter cells. Data are shown as the percentage of TLR3-positive lung M ϕ on the vertical axis versus A,

FEV1 % predicted; B, smoking history in pack-years. In panel A, circles represent current smokers and "x" represents former smokers. In panel B, all subjects are shown as inverted triangles, regardless of smoking status (active vs. former); in both panels, $n = 25$.

Abbreviations

AM ϕ : Alveolar macrophage(s); BAL: Bronchoalveolar lavage; COPD: Chronic obstructive pulmonary disease; CSE: Cigarette smoke extract; Ds: Double-stranded; FEV₁: Forced expiratory volume in 1 second; FVC: Forced vital capacity; GAPDH: Glyceraldehyde-3-phosphate dehydrogenase; M ϕ : Macrophage(s); MDA-5: Melanoma differentiation-associated gene 5; MDM ϕ : Monocyte-derived macrophage(s); NK: Natural killer; PKR: Double-stranded RNA-dependent protein kinase; PMA: Phorbol myristate acetate; poly(I:C): Polyinosinic acid:cytidylic acid; RIG-I: Retinoic acid-inducible gene I; TLR3: Toll-like receptor 3.

Competing interests

Jill C Todt, Jeanette P Brown, Joanne Sonstein, Theresa M Ames, Alexandra McCubbrey and Stephen W Chensue have no competing interests to declare. Christine M Freeman, Fernando J Martinez, James M Beck and Jeffrey L Curtis were supported by research grants as outlined in the Endnotes, but have no other competing interests to declare.

Authors' contributions

JCT: Designed and performed experiments, analyzed data, produced graphs and tables, wrote the initial draft of manuscript, reviewed and approved final manuscript; CMF: designed and performed experiments, analyzed data and produced graphs and tables, wrote the initial draft of the revised manuscript, reviewed and approved final manuscript; JPB designed and performed experiments, analyzed data and produced graphs and tables, reviewed and approved final manuscript; JS: performed and analyzed flow cytometry experiments, reviewed and approved final manuscript; TMA: performed and photographed immunocytochemistry experiments, participated in performance of other experiments, reviewed and approved final manuscript; ALM: designed and performed experiments, reviewed and approved final manuscript; FJM: secured research funding, participated in study design, analyzed data, participated in manuscript generation, reviewed and approved final manuscript; SWC: performed dissection of surgical lung specimens, reviewed and approved final manuscript; JMB: secured research funding, participated in study design, analyzed data, participated in manuscript generation, reviewed and approved final manuscript; JLC: secured research funding, oversaw study design, performed all subject-related activities including bronchoscopies, analyzed data, generated the final manuscript, and takes responsibility for the scientific integrity of the overall project.

Acknowledgements

The authors thank Drs. Peter Mancuso, Michal A Olszewski and Martin R Stämpfli for providing helpful discussions and suggestions; Catherine Meldrum, R.N. for human subjects recruitment; Lisa McCloskey, R.R.T. and Tricia Geal, R.R.T. for human subjects recruitment and assistance in bronchoscopies; Bradley T Todd, and Angela M Preston, M.P.H. for generating cigarette smoke extract; Catherine Spino, Sc.D. and Glen Feak for database support; and Mary Freer, M.B.A., Tameka Lewis, Joyce O'Brien, Rebecca Weeks and Sara Whisenant for administrative support.

Author details

¹Division of Pulmonary & Critical Care Medicine, Department of Internal Medicine, University of Michigan Health Care System, Ann Arbor, MI 48109-2399, USA. ²Research Service, Department of Veterans Affairs Health Care System, Ann Arbor, MI 48105-2303, USA. ³Graduate Program in Immunology, University of Michigan Health Care System, Ann Arbor, MI 48109-2399, USA. ⁴Department of Pathology, University of Michigan Health Care System, Ann Arbor, MI 48109-5602, USA. ⁵Pathology & Laboratory Medicine Service, Department of Veterans Affairs Health Care System, Ann Arbor, MI 48105-2303, USA. ⁶Pulmonary & Critical Care Medicine Section, Medical Service, Department of Veterans Affairs Health Care System, 2215 Fuller Road, Ann Arbor, MI 48105-2303, USA.

Received: 3 May 2012 Accepted: 22 February 2013
Published: 9 March 2013

References

- Johnston SL: Overview of virus-induced airway disease. *Proc Am Thorac Soc* 2005, **2**:150–156.
- Proud D, Chow CW: Role of viral infections in asthma and chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol* 2006, **35**:513–518.
- Arcavi L, Benowitz NL: Cigarette smoking and infection. *Arch Intern Med* 2004, **164**:2206–2216.
- Aronson MD, Weiss ST, Ben RL, Komaroff AL: Association between cigarette smoking and acute respiratory tract illness in young adults. *JAMA* 1982, **248**:181–183.
- Bensenor IM, Cook NR, Lee IM, Chown MJ, Hennekens CH, Buring JE, Manson JE: Active and passive smoking and risk of colds in women. *Ann Epidemiol* 2001, **11**:225–231.
- Marcy TW, Merrill WW: Cigarette smoking and respiratory tract infection. *Clin Chest Med* 1987, **8**:381–391.
- Sopori M: Effects of cigarette smoke on the immune system. *Nat Rev Immunol* 2002, **2**:372–377.
- Mehta H, Nazzari K, Sadikot RT: Cigarette smoking and innate immunity. *Inflamm Res* 2008, **57**:497–503.
- Matsunaga K, Klein TW, Friedman H, Yamamoto Y: Involvement of nicotinic acetylcholine receptors in suppression of antimicrobial activity and cytokine responses of alveolar macrophages to *Legionella pneumophila* infection by nicotine. *J Immunol* 2001, **167**:6518–6524.
- Ouyang Y, Virasch N, Hao P, Aubrey MT, Mukerjee N, Bierer BE, Freed BM: Suppression of human IL-1beta, IL-2, IFN-gamma, and TNF-alpha production by cigarette smoke extracts. *J Allergy Clin Immunol* 2000, **106**:280–287.
- Wewers MD, Diaz PT, Wewers ME, Lowe MP, Nagaraja HN, Clanton TL: Cigarette smoking in HIV infection induces a suppressive inflammatory environment in the lung. *Am J Respir Crit Care Med* 1998, **158**:1543–1549.
- Chen H, Cowan MJ, Hasday JD, Vogel SN, Medvedev AE: Tobacco smoking inhibits expression of proinflammatory cytokines and activation of IL-1R-associated kinase, p38, and NF-kappaB in alveolar macrophages stimulated with TLR2 and TLR4 agonists. *J Immunol* 2007, **179**:6097–6106.
- Birrell MA, Wong S, Catley MC, Belvisi MG: Impact of tobacco-smoke on key signaling pathways in the innate immune response in lung macrophages. *J Cell Physiol* 2008, **214**:27–37.
- Edwards K, Braun KM, Evans G, Sureka AO, Fan S: Mainstream and sidestream cigarette smoke condensates suppress macrophage responsiveness to interferon gamma. *Hum Exp Toxicol* 1999, **18**:233–240.
- Matsumoto M, Funami K, Tanabe M, Oshiumi H, Shingai M, Seto Y, Yamamoto A, Seya T: Subcellular localization of Toll-like receptor 3 in human dendritic cells. *J Immunol* 2003, **171**:3154–3162.
- Orinska Z, Bulanova E, Budagian V, Metz M, Maurer M, Bulfone-Paus S: TLR3-induced activation of mast cells modulates CD8+ T-cell recruitment. *Blood* 2005, **106**:978–987.
- Meylan E, Tschopp J: Toll-like receptors and RNA helicases: two parallel ways to trigger antiviral responses. *Mol Cell* 2006, **22**:561–569.
- Matsukura S, Kokubu F, Kurokawa M, Kawaguchi M, Ieki K, Kuga H, Odaka M, Suzuki S, Watanabe S, Homma T, et al: Role of RIG-I, MDA-5, and PKR on the expression of inflammatory chemokines induced by synthetic dsRNA in airway epithelial cells. *Int Arch Allergy Immunol* 2007, **143**(Suppl 1):80–83.
- Hui KP, Lee SM, Cheung CY, Ng IH, Poon LL, Guan Y, Ip NY, Lau AS, Peiris JS: Induction of proinflammatory cytokines in primary human macrophages by influenza A virus (H5N1) is selectively regulated by IFN regulatory factor 3 and p38 MAPK. *J Immunol* 2009, **182**:1088–1098.
- Dhillon N, Zhu X, Peng F, Yao H, Williams R, Qiu J, Callen S, Ladner AO, Buch S: Molecular mechanism(s) involved in the synergistic induction of CXCL10 by human immunodeficiency virus type 1 Tat and interferon-gamma in macrophages. *J Neurovirol* 2008, **14**:196–204.
- Livengood AJ, Wu CC, Carson DA: Opposing roles of RNA receptors TLR3 and RIG-I in the inflammatory response to double-stranded RNA in a Kaposi's sarcoma cell line. *Cell Immunol* 2007, **249**:55–62.
- Michalec L, Choudhury BK, Postlethwait E, Wild JS, Alam R, Lett-Brown M, Sur S: CCL7 and CXCL10 orchestrate oxidative stress-induced neutrophilic lung inflammation. *J Immunol* 2002, **168**:846–852.
- Agostini C, Calabrese F, Poletti V, Marcer G, Facco M, Miorin M, Cabrelle A, Baesso I, Zambello R, Trentin L, Semenzato G: CXCR3/CXCL10 interactions in the development of hypersensitivity pneumonitis. *Respir Res* 2005, **6**:20.
- Yuan J, Liu Z, Lim T, Zhang H, He J, Walker E, Shier C, Wang Y, Su Y, Sall A, et al: CXCL10 inhibits viral replication through recruitment of natural killer cells in coxsackievirus B3-induced myocarditis. *Circ Res* 2009, **104**:628–638.
- Cole AM, Ganz T, Liese AM, Burdick MD, Liu L, Strieter RM: Cutting edge: IFN-inducible ELR- CXC chemokines display defensin-like antimicrobial activity. *J Immunol* 2001, **167**:623–627.
- Vasquez RE, Xin L, Soong L: Effects of CXCL10 on dendritic cell and CD4+ T-cell functions during *Leishmania amazonensis* infection. *Infect Immun* 2008, **76**:161–169.
- Freeman CM, Curtis JL, Chensue SW: CC chemokine receptor 5 and CXC chemokine receptor 6 expression by lung CD8+ cells correlates with chronic obstructive pulmonary disease severity. *Am J Pathol* 2007, **171**:767–776.
- Freeman CM, Han MK, Martinez FJ, Murray S, Liu LX, Chensue SW, Polak TJ, Sonstein J, Todt JC, Ames TM, et al: Cytotoxic potential of lung CD8(+) T cells increases with chronic obstructive pulmonary disease severity and with in vitro stimulation by IL-18 or IL-15. *J Immunol* 2010, **184**:6504–6513.
- McCubbrey AL, Sonstein J, Ames TM, Freeman CM, Curtis JL: Glucocorticoids relieve collectin-driven suppression of apoptotic cell uptake in murine alveolar macrophages through downregulation of SIRPalpha. *J Immunol* 2012, **189**:112–119.
- Koara A, Yanagisawa S, Sugiura H, Ichikawa T, Akamatsu K, Hirano T, Nakanishi M, Matsunaga K, Minakata Y, Ichinose M: Cigarette smoke augments the expression and responses of toll-like receptor 3 in human macrophages. *Respirology* 2012, **17**:1018–1025.
- Seemungal T, Harper-Owen R, Bhowmic A, Moric I, Sanderson G, Message S, MacCallum P, Meade T, Jeffries D, Johnston S, Wedzicha J: Respiratory viruses, symptoms, and inflammatory markers in acute exacerbations and stable chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001, **164**:1618–1623.
- Papi A, Bellettato C, Braccioni F, Romagnoli M, Casolari P, Caramori G, Fabbri L, Johnston S: Infections and airway inflammation in chronic obstructive pulmonary disease severe exacerbations. *Am J Respir Crit Care Med* 2006, **173**:1114–1121.
- Nicholson KG, Kent J, Ireland DC: Respiratory viruses and exacerbations of asthma in adults. *BMJ* 1993, **307**:982–986.
- Johnston SL, Pattermore PK, Sanderson G, Smith S, Lampe F, Josephs L, Symington P, O'Toole S, Myint SH, Tyrrell DA, et al: Community study of role of viral infections in exacerbations of asthma in 9–11 year old children. *BMJ* 1995, **310**:1225–1229.
- Johnston SL, Pattermore PK, Sanderson G, Smith S, Campbell MJ, Josephs LK, Cunningham A, Robinson BS, Myint SH, Ward ME, et al: The relationship between upper respiratory infections and hospital admissions for asthma: a time-trend analysis. *Am J Respir Crit Care Med* 1996, **154**:654–660.
- Kanner RE, Anthonisen NR, Connett JE: Lower respiratory illnesses promote FEV(1) decline in current smokers but not ex-smokers with mild chronic obstructive pulmonary disease: results from the lung health study. *Am J Respir Crit Care Med* 2001, **164**:358–364.
- Thomson NC, Chaudhuri R: Asthma in smokers: challenges and opportunities. *Curr Opin Pulm Med* 2009, **15**:39–45.
- Eisner MD, Iribarren C: The influence of cigarette smoking on adult asthma outcomes. *Nicotine Tob Res* 2007, **9**:53–56.
- Sippel JM, Pedula KL, Vollmer WM, Buist AS, Osborne ML: Associations of smoking with hospital-based care and quality of life in patients with obstructive airway disease. *Chest* 1999, **115**:691–696.
- Siroux V, Pin I, Oryszczyn MP, LeMoual N, Kauffmann F: Relationships of active smoking to asthma and asthma severity in the EGEA study. Epidemiological study on the Genetics and Environment of Asthma. *Eur Respir J* 2000, **15**:470–477.
- Lazarus SC, Chinchilli VM, Rollings NJ, Boushey HA, Cherniack R, Craig TJ, Deykin A, DiMango E, Fish JE, Ford JG, et al: Smoking affects response to inhaled corticosteroids or leukotriene receptor antagonists in asthma. *Am J Respir Crit Care Med* 2007, **175**:783–790.
- Kohlmeier JE, Woodland DL: Immunity to respiratory viruses. *Annu Rev Immunol* 2009, **27**:61–82.
- Khaitov MR, Laza-Stanca V, Edwards MR, Walton RP, Rohde G, Contoli M, Papi A, Stanciu LA, Kotenko SV, Johnston SL: Respiratory virus induction of alpha-, beta- and lambda-interferons in bronchial epithelial cells and peripheral blood mononuclear cells. *Allergy* 2009, **64**:375–386.
- Contoli M, Message SD, Laza-Stanca V, Edwards MR, Wark PA, Bartlett NW, Kebabdzic T, Mallia P, Stanciu LA, Parker HL, et al: Role of deficient type III

- interferon-lambda production in asthma exacerbations. *Nat Med* 2006, **12**:1023–1026.
45. Cakebread JA, Xu Y, Grainge C, Kehagia V, Howarth PH, Holgate ST, Davies DE: Exogenous IFN-beta has antiviral and anti-inflammatory properties in primary bronchial epithelial cells from asthmatic subjects exposed to rhinovirus. *J Allergy Clin Immunol* 2011, **127**:1148–1154. e1149.
 46. Laza-Stanca V, Stanciu LA, Message SD, Edwards MR, Gern JE, Johnston SL: Rhinovirus replication in human macrophages induces NF-kappaB-dependent tumor necrosis factor alpha production. *J Virol* 2006, **80**:8248–8258.
 47. Wang J, Oberley-Deegan R, Wang S, Nikrad M, Funk CJ, Hartshorn KL, Mason RJ: Differentiated human alveolar type II cells secrete antiviral IL-29 (IFN-lambda 1) in response to influenza A infection. *J Immunol* 2009, **182**:1296–1304.
 48. Siren J, Pirhonen J, Julkunen I, Matikainen S: IFN-alpha regulates TLR-dependent gene expression of IFN-alpha, IFN-beta, IL-28, and IL-29. *J Immunol* 2005, **174**:1932–1937.
 49. Sheppard P, Kindsvogel W, Xu W, Henderson K, Schlutsmeyer S, Whitmore TE, Kuestner R, Garrigues U, Birks C, Roraback J, et al: IL-28, IL-29 and their class II cytokine receptor IL-28R. *Nat Immunol* 2003, **4**:63–68.
 50. Kim HM, Lee YW, Lee KJ, Kim HS, Cho SW, Van Rooijen N, Guan Y, Seo SH: Alveolar macrophages are indispensable for controlling influenza viruses in lungs of pigs. *J Virol* 2008, **82**:4265–4274.
 51. Kumagai Y, Takeuchi O, Kato H, Kumar H, Matsui K, Morii E, Aozasa K, Kawai T, Akira S: Alveolar macrophages are the primary interferon-alpha producer in pulmonary infection with RNA viruses. *Immunity* 2007, **27**:240–252.
 52. Pribul PK, Harker J, Wang B, Wang H, Tregoning JS, Schwarze J, Openshaw PJ: Alveolar macrophages are a major determinant of early responses to viral lung infection but do not influence subsequent disease development. *J Virol* 2008, **82**:4441–4448.
 53. Woodruff PG, Koth LL, Yang YH, Rodriguez MW, Favoreto S, Dolganov GM, Paquet AC, Erle DJ: A distinctive alveolar macrophage activation state induced by cigarette smoking. *Am J Respir Crit Care Med* 2005, **172**:1383–1392.
 54. Shaykhiyev R, Krause A, Salit J, Strulovici-Barel Y, Harvey BG, O'Connor TP, Crystal RG: Smoking-dependent reprogramming of alveolar macrophage polarization: implication for pathogenesis of chronic obstructive pulmonary disease. *J Immunol* 2009, **183**:2867–2883.
 55. Hunninghake GW, Gadek JE, Kawanami O, Ferrans VJ, Crysta I RG: Inflammatory and immune processes in the human lung in health and disease. Evaluation by bronchoalveolar lavage. *Am J Pathol* 1979, **97**:149–206.
 56. Richards SW, Peterson PK, Verbrugh HA, Nelson RD, Hammerschmidt DE, Hoidal JR: Chemotactic and phagocytic responses of human alveolar macrophages to activated complement components. *Infect Immun* 1984, **43**:775–778.
 57. Holt PG: Immune and inflammatory function in cigarette smokers. *Thorax* 1987, **42**:241–249.
 58. Marti-Llites P, Regueiro V, Morey P, Hood DW, Saus C, Sauleda J, Agusti AG, Bengoechea JA, Garmendia J: Nontypeable *Haemophilus influenzae* clearance by alveolar macrophages is impaired by exposure to cigarette smoke. *Infect Immun* 2009, **77**:4232–4242.
 59. King TE Jr, Savici D, Campbell PA: Phagocytosis and killing of *Listeria monocytogenes* by alveolar macrophages: smokers versus nonsmokers. *J Infect Dis* 1988, **158**:1309–1316.
 60. Cosio BG, Tsaprouni L, Ito K, Jazrawi E, Adcock IM, Barnes PJ: Theophylline restores histone deacetylase activity and steroid responses in COPD macrophages. *J Exp Med* 2004, **200**:689–695.
 61. Cosio MG, Saetta M, Agusti A: Immunologic aspects of chronic obstructive pulmonary disease. *N Engl J Med* 2009, **360**:2445–2454.
 62. McCrea KA, Ensor JE, Nall K, Bleecker ER, Hasday JD: Altered cytokine regulation in the lungs of cigarette smokers. *Am J Respir Crit Care Med* 1994, **150**:696–703.
 63. Morrison D, Strieter RM, Donnelly SC, Burdick MD, Kunkel SL, MacNee W: Neutrophil chemokines in bronchoalveolar lavage fluid and leukocyte-conditioned medium from nonsmokers and smokers. *Eur Respir J* 1998, **12**:1067–1072.
 64. Ohta T, Yamashita N, Maruyama M, Sugiyama E, Kobayashi M: Cigarette smoking decreases interleukin-8 secretion by human alveolar macrophages. *Respir Med* 1998, **92**:922–927.
 65. Hutchens M, Luker KE, Sottile P, Sonstein J, Lukacs NW, Nunez G, Curtis JL, Luker GD: TLR3 increases disease morbidity and mortality from vaccinia infection. *J Immunol* 2008, **180**:483–491.
 66. Bunting RA, Duffy KE, Lamb RJ, San Mateo LR, Smalley K, Raymond H, Liu X, Petley T, Fisher J, Beck H, et al: Novel antagonist antibody to TLR3 blocks poly(I:C)-induced inflammation in vivo and in vitro. *Cell Immunol* 2011, **267**:9–16.
 67. Cavassani KA, Ishii M, Wen H, Schaller MA, Lincoln PM, Lukacs NW, Hogaboam CM, Kunkel SL: TLR3 is an endogenous sensor of tissue necrosis during acute inflammatory events. *J Exp Med* 2008, **205**:2609–2621.
 68. Henson PM, Tuder RM: Apoptosis in the lung: induction, clearance and detection. *Am J Physiol Lung Cell Mol Physiol* 2008, **294**:L601–L611.
 69. Hodge S, Hodge G, Brozyna S, Jersmann H, Holmes M, Reynolds PN: Azithromycin increases phagocytosis of apoptotic bronchial epithelial cells by alveolar macrophages. *Eur Respir J* 2006, **28**:486–495.
 70. Hodge S, Hodge G, Scicchitano R, Reynolds PN, Holmes M: Alveolar macrophages from subjects with chronic obstructive pulmonary disease are deficient in their ability to phagocytose apoptotic airway epithelial cells. *Immunol Cell Biol* 2003, **81**:289–296.
 71. Richens TR, Linderman DJ, Horstmann SA, Lambert C, Xiao YQ, Keith RL, Boe DM, Morimoto K, Bowler RP, Day BJ, et al: Cigarette smoke impairs clearance of apoptotic cells through oxidant-dependent activation of RhoA. *Am J Respir Crit Care Med* 2009, **179**:1011–1021.
 72. Vandivier RW, Henson PM, Douglas IS: Burying the dead: the impact of failed apoptotic cell removal (efferocytosis) on chronic inflammatory lung disease. *Chest* 2006, **129**:1673–1682.
 73. Hodge S, Hodge G, Ahern J, Jersmann H, Holmes M, Reynolds PN: Smoking alters alveolar macrophage recognition and phagocytic ability: implications in chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol* 2007, **37**:748–755.
 74. Drömann D, Goldmann T, Tiedje T, Zabel P, Dalhoff K, Schaub B: Toll-like receptor 2 expression is decreased on alveolar macrophages in cigarette smokers and COPD patients. *Respir Res* 2005, **6**:68.
 75. Doyle I, Ratcliffe M, Walding A, Vanden Bon E, Dymond M, Tomlinson W, Tilley D, Shelton P, Dougall I: Differential gene expression analysis in human monocyte-derived macrophages: impact of cigarette smoke on host defence. *Mol Immunol* 2010, **47**:1058–1065.
 76. Gaschler GJ, Zavitz CC, Bauer CM, Skrtic M, Lindahl M, Robbins CS, Chen B, Stampfli MR: Cigarette smoke exposure attenuates cytokine production by mouse alveolar macrophages. *Am J Respir Cell Mol Biol* 2008, **38**:218–226.
 77. Kang MJ, Lee CG, Lee JY, Dela Cruz CS, Chen ZJ, Enelow R, Elias JA: Cigarette smoke selectively enhances viral PAMP- and virus-induced pulmonary innate immune and remodeling responses in mice. *J Clin Invest* 2008, **118**:2771–2784.
 78. Bonocchi R, Galliera E, Borroni EM, Corsi MM, Locati M, Mantovani A: Chemokines and chemokine receptors: an overview. *Front Biosci* 2009, **14**:540–551.
 79. Jiang D, Liang J, Campanella GS, Guo R, Yu S, Xie T, Liu N, Jung Y, Homer R, Meltzer EB, et al: Inhibition of pulmonary fibrosis in mice by CXCL10 requires glycosaminoglycan binding and syndecan-4. *J Clin Invest* 2010, **120**:2049–2057.
 80. Pekarek V, Srinivas S, Eskdale J, Gallagher G: Interferon lambda-1 (IFN-lambda1/IL-29) induces ELR(-) CXC chemokine mRNA in human peripheral blood mononuclear cells, in an IFN-gamma-independent manner. *Genes Immun* 2007, **8**:177–180.
 81. Korpi-Steiner NL, Bates ME, Lee WM, Hall DJ, Bertics PJ: Human rhinovirus induces robust IP-10 release by monocytic cells, which is independent of viral replication but linked to type I interferon receptor ligation and STAT1 activation. *J Leukoc Biol* 2006, **80**:1364–1374.
 82. Miller AL, Bowlin TL, Lukacs NW: Respiratory syncytial virus-induced chemokine production: linking viral replication to chemokine production in vitro and in vivo. *J Infect Dis* 2004, **189**:1419–1430.
 83. Proost P, Vynckier AK, Mahieu F, Put W, Grillet B, Struyf S, Wuyts A, Opendakker G, Van Damme J: Microbial Toll-like receptor ligands differentially regulate CXCL10/IP-10 expression in fibroblasts and mononuclear leukocytes in synergy with IFN-gamma and provide a mechanism for enhanced synovial chemokine levels in septic arthritis. *Eur J Immunol* 2003, **33**:3146–3153.
 84. D'Ambrosio D, Mariani M, Panina-Bordignon P, Sinigaglia F: Chemokines and their receptors guiding T lymphocyte recruitment in lung inflammation. *Am J Respir Crit Care Med* 2001, **164**:1266–1275.
 85. Kohlmeier JE, Cookenham T, Miller SC, Roberts AD, Christensen JP, Thomsen AR, Woodland DL: CXCR3 directs antigen-specific effector CD4+ T cell migration to the lung during parainfluenza virus infection. *J Immunol* 2009, **183**:4378–4384.

86. Costa C, Rufino R, Traves SL, Lapa ESJR, Barnes PJ, Donnelly LE: **CXCR3 and CCR5 chemokines in induced sputum from patients with COPD.** *Chest* 2008, **133**:26–33.
87. Han MK, Postma D, Mannino DM, Giardino ND, Buist S, Curtis JL, Martinez FJ: **Gender and chronic obstructive pulmonary disease: why it matters.** *Am J Respir Crit Care Med* 2007, **176**:1179–1184.
88. Antonelli A, Rotondi M, Fallahi P, Ferrari SM, Paolicchi A, Romagnani P, Serio M, Ferrannini E: **Increase of CXC chemokine CXCL10 and CC chemokine CCL2 serum levels in normal ageing.** *Cytokine* 2006, **34**:32–38.
89. Antonelli A, Rotondi M, Fallahi P, Romagnani P, Ferrari SM, Ferrannini E, Serio M: **Age-dependent changes in CXC chemokine ligand 10 serum levels in euthyroid subjects.** *J Interferon Cytokine Res* 2005, **25**:547–552.

doi:10.1186/1465-9921-14-33

Cite this article as: Todt et al.: Smoking decreases the response of human lung macrophages to double-stranded RNA by reducing TLR3 expression. *Respiratory Research* 2013 **14**:33.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

