

CHEMISTRY

A European Journal

Supporting Information

High Surface Area, Thermally Stable, Hydrophobic, Microporous, Rigid Gels Generated at Ambient from MeSi(OEt)₃/ (EtO)₃SiCH₂CH₂Si(OEt)₃ Mixtures by F⁻-Catalyzed Hydrolysis

Joseph C. Furgal,^{*[a, e]} Honami Yamane,^[b] Timothy R. Odykirk,^[c] Eongyu Yi,^[c] Yoshiki Chujo,^[b] and Richard M. Laine^{*[c, d]}

chem_201704941_sm_miscellaneous_information.pdf

Supporting Information




Figure S1. FTIR studies of dried gels from Table 1.




Figure S2. FTIR of gels 12-14 in Table 3, blue (12), red (13), purple (14).




Figure S3. FTIR of gel (15), Table 3.




Figure S3. FTIR of gels 17 (blue) and 18 (red), Table 3.




Figure S4. FTIR of gels 16 (blue), and 19 (red), Table 3.




Figure S5. FTIR of gels 21 (red), and 22 (blue), Table 3.




Figure S6. FTIR of gels 20 (purple), 23 (red), and 24 (blue), Table 3.




Figure S7. FTIR of gel (26), Table 3.




Figure S8. FTIR of gel (27), Table 3.




Figure S9. FTIR of gel (28), Table 3.