

Development of Quality Metrics for Ambulatory Pediatric Cardiology – Transposition of the Great Arteries after Arterial Switch Operation

Running Title: Quality Metrics and TGA/ASO

Carissa M Baker-Smith, MD, MPH<sup>\*</sup>, Karina Carlson, MD<sup>†</sup>, Jose Etedgui, MD<sup>‡</sup>, Takeshi Tsuda, MD<sup>×</sup>, K. Anitha Jayakumar, MD<sup>§</sup>, Matthew Park, MD<sup>||</sup>, Nikola Tede, MD<sup>#</sup>, Karen Uzark, PhD PNP<sup>\*\*</sup>, Craig Fleishman, MD<sup>††</sup>, David Connuck, MD<sup>‡‡</sup>, Maggie Likes, MD<sup>xx</sup>, Daniel J Penny, MD, PhD<sup>ss</sup>

<sup>\*</sup> University of Maryland School of Medicine, Baltimore, MD

<sup>†</sup> Children's Mercy Hospital, Kansas City, MO

<sup>‡</sup> University of Florida, Jacksonville, FL

<sup>×</sup> Alfred I. DuPont Hospital for Children, Wilmington, DC

<sup>§</sup> University of Virginia, Charlottesville, VA

<sup>||</sup> Pediatrix Medical Group - Northwest Children's Heart Care, Tacoma, WA

<sup>#</sup> California Pacific Medical Center, San Francisco, CA

<sup>\*\*</sup> University of Michigan Congenital Heart Center, Ann Arbor, MI

<sup>††</sup> The Heart Center at Arnold Palmer Hospital for Children, Orlando, FL

<sup>‡‡</sup> Janet Weis Children's Hospital, Danville, PA

<sup>xx</sup> Seattle Children's Hospital, University of Washington, Seattle, WA

<sup>ss</sup> Baylor College of Medicine and Texas Children's Hospital, Houston, TX

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version record](#). Please cite this article as [doi:10.1111/chd.12540](https://doi.org/10.1111/chd.12540).

Corresponding Author:

Carissa M. Baker-Smith, MD, MPH, F.A.A.P., F.A.H.A.

University of Maryland School of Medicine

110 S. Paca Street

7<sup>th</sup> floor

Baltimore, Maryland 21201

Phone: 410-328-4348

Fax: 410-328-8670

E-mail: cbaker-smith@peds.umaryland.edu

Conflicts of Interest by the Authors: None

Disclosure: No funding was provided for this study

Figures: 0; Tables 2

#### ABSTRACT:

**Objective:** To develop quality metrics (QMs) for the ambulatory care of patients with transposition of the great arteries following arterial switch operation (TGA/ASO).

**Design:** Under the auspices of the American College of Cardiology Adult Congenital and Pediatric Cardiology (ACPC) Steering committee, the TGA/ASO team generated candidate QMs related to TGA/ASO ambulatory care. Candidate QMs were submitted to the ACPC Steering Committee and were reviewed for validity and feasibility using individual expert panel member scoring according to the RAND-UCLA methodology. QMs were then made available for review by the entire ACC ACPC during an “open comment period.” Final approval of each QM was provided by a vote of the ACC ACPC Council.

Patients: Patients with TGA who had undergone an ASO were included. Patients with complex transposition were excluded.

Results: Twelve candidate QMs were generated. Seven metrics passed the RAND-UCLA process. Four passed the “open comment period” and were ultimately approved by the Council.

These included: (1) at least 1 echocardiogram performed during the first year of life reporting on the function, aortic dimension, degree of neoaortic valve insufficiency, the patency of the systemic and pulmonary outflows, the patency of the branch pulmonary arteries and coronary arteries, (b) neurodevelopmental (ND) assessment after ASO; (c) lipid profile by age 11 years; and (d) documentation of a transition of care plan to an adult congenital heart disease (CHD) provider by 18 years of age.

Conclusion(s): Application of the RAND-UCLA methodology and linkage of this methodology to the ACPC approval process led to successful generation of 4 QMs relevant to the care of TGA/ASO pediatric patients in the ambulatory setting. These metrics have now been incorporated into the ACPC Quality Network providing guidance for the care of TGA/ASO patients across 30 CHD centers.

Keywords: transposition, arterial switch operation, quality, pediatrics

## Introduction

The goal of quality metric (QM) development is to create self assessment measures that can be easily utilized by providers and health care systems. These metrics should address key issues related to patient management and allow for evidence based care, greater consistency in care and ultimately, when utilized, lead to improvements in clinical care and patient outcome.<sup>i</sup> Historically, the outpatient management of children with congenital heart disease (CHD) has varied according to center and provider preference.<sup>ii</sup> Consistent and evidence based clinical practice has been limited by the paucity of established guidelines and the challenge of creating evidence based recommendations given heterogeneity within the CHD population, even for a single diagnosis and intervention.

While mortality rates associated with transposition of the great arteries (TGA) following arterial switch operation (ASO) have improved over time, patients with TGA/ASO remain at risk for significant mid-term and long-term complications including *myocardial ischemia, left ventricular dysfunction, arrhythmia and sudden cardiac death (SCD)*,<sup>ii iii iv v vi vii viii ix x xi ,xii xiii</sup> *neoaortic valve regurgitation (NAR), aortic root dilation (ARD), supraaortic pulmonary stenosis (SPS), supraaortic aortic stenosis (SAS)*,<sup>ii xiv xv xvi xvii xviii</sup> *impaired neurodevelopmental (ND) outcome*,<sup>xix xx xxi xxii</sup> *and the development of risk factors for premature atherosclerosis, including unhealthy weight, elevated blood pressure and high cholesterol*.<sup>xxvi</sup> Similar to many CHD patients, teenagers with TGA/ASO are vulnerable to delays in transition of care to an adult provider and lapses in care, as a result.<sup>xxvii xxviii</sup>

To date there have been no established guidelines offering metrics for quality improvement in the ambulatory care setting for children with TGA/ASO.<sup>ii xxv xxix xxx xxxi</sup> Recognizing the need to develop valid and feasible quality metrics (QM) in 5 key areas of pediatric and congenital disease care, including children with TGA/ASO, the American College of Cardiology (ACC)

Adult Congenital and Pediatric Cardiology Council (ACPC) proposed use of a 2 step review process, incorporating the modified RAND-UCLA methodology, for the creation of QMs relevant to the ambulatory care of TGA/ASO pediatric patients.<sup>xxxii</sup> This paper outlines the process utilized to generate QMs, reviews the candidate QMs and describes the QMs that were ultimately selected for TGA/ASO patients.

## Methods

Members of the TGA/ASO quality metrics group were members of the ACC and were selected on a volunteer basis. There were 12 members of the TGA/ASO group. Members were expected to have content expertise and to participate in bi-weekly conference calls. Two to three person groups were created among the members in order to divide up the work of evaluating the existing literature and in order to create candidate QMs to be reviewed by the entire team. Each 2-3 person group generated candidate QMs. Given that the goal was to generate metrics for quality improvement in ambulatory practice, proposed candidate QMs included a description of the proposed metric, a numerator (e.g., what was to be measured from a patient's chart), a denominator (e.g., the target population), exclusions, and a summary of the key references (see Appendix 1). The QMs were designed for ease of ascertainment of data from a patient's electronic medical record (EMR). All resulting QMs were discussed over a series of conference calls involving the entire TGA/ASO team.

Only QMs relevant to the outpatient management of TGA/ASO recipients and that addressed risk factors for mid and long term morbidity and mortality following the ASO procedure were created. We excluded patients with complex TGA, including those with a hemodynamically significant ventricular septal defect (VSD), aortic arch obstruction, or congenitally-corrected transposition, in order to analyze a more homogenous population. We also excluded TGA patients who had undergone Mustard or Senning operations.

Candidate QMs were then submitted to the ACC ACPC Steering Committee for expert panel review. Employing the RAND-UCLA process,<sup>xxxiii</sup> candidate QMs were first individually scored for validity and feasibility by a single member of the expert panel and then independently scored by each member of the expert panel during an in-person meeting of the Steering Committee and expert panel on October 3<sup>rd</sup> and 4<sup>th</sup>, 2013 at the ACC Heart House. Members discussed and refined the QMs before a final score was obtained from each member of the expert panel. Candidate metrics were scored for validity and feasibility on a scale to 1 to 9, where 9 represented the most valid or most feasible. In order to be recommended for “open comment”, a candidate metric required a mean validity score of 7 to 9 and a mean feasibility score of 4 to 9. Consensus among expert panel members was not required for approval during either round of scoring.

QMs recommended following the RAND- UCLA 2-step process were posted on the ACC website for “open comment” from existing ACC ACPC members. Comments received during this process were reviewed and discussed by the Steering Committee and were used to modify the language of the QM. Final approval of the QMs was determined by a vote of the ACC ACPC Council.<sup>xxxii</sup>

## Results

### *Literature Review*

Morbidity and mortality related outcomes post ASO among “simple” TGA patients are generally quite good.<sup>iii xiii xviii</sup> However, patients remain at risk for arrhythmia. They are also at risk for myocardial ischemia secondary to problems that occur during coronary artery transfer and secondary to more complex coronary artery anatomy as well as to the translocation process itself.<sup>xiii</sup> Common mid and late-term complications following ASO include aortic root dilation,<sup>xvi</sup>

neoaortic valve insufficiency, supraaortic stenosis (e.g., aortic and pulmonary) and branch pulmonary artery stenosis.<sup>iii</sup> Children who undergo ASO are also at risk for abnormal neurodevelopment<sup>xx</sup> as well as acquired cardiovascular disease (CVD).<sup>xxvi</sup> Finally, transition of care to an adult provider represents a particularly vulnerable time for the child post ASO, providing an avenue for poor outcome.<sup>xxxiv</sup>

● A formal systematic review was beyond the scope of this project. However, a thorough review of the existing literature was carried out in order to generate candidate metrics. We also searched and reviewed existing guidelines.<sup>ii xxxi xxxv</sup> Existing guidelines regarding the management of TGA patients following ASO include the ACC/AHA 2008 Guidelines for the Management of Adults with Congenital Heart Disease<sup>xxxv</sup> and a single center study providing general recommendations for outpatient management post ASO.<sup>ii</sup>

#### *Metric#1-3: Electrocardiogram (ECG), Holter Monitoring and Exercise Stress Testing post ASO*

The ASO procedure and the translocation of the coronary arteries poses a potential for immediate as well as mid-term and long-term complications related to compromised coronary artery perfusion.<sup>xxxvi</sup> Children with complex coronary artery anatomy (e.g., intramural course) are at particularly high risk for coronary artery stenosis and subsequent myocardial ischemia.<sup>xxxvii</sup>

Sudden cardiac death post surgery is rare and has been reported in 0.3 to 0.5% of all TGA/ASO patients.<sup>ix xxxviii</sup> Children are also at risk for arrhythmia.<sup>vi x lix</sup> However, rates of arrhythmia are low and 98% of ASO/TGA patients have normal sinus rhythm by holter monitoring with occasional premature beats reported in approximately 50%. Rarely reported are episodes of supraventricular tachycardia or ventricular tachycardia following ASO.<sup>liii</sup>

Exercise capacity following ASO is typically normal. Exercise testing has been considered useful for detecting exercise induced ischemia or arrhythmia.<sup>xxxix</sup>

#### Metric #4-#5: Echocardiogram Evaluation

Neo-aortic regurgitation (NAR), aortic root dilation (ARD), supra-aortic pulmonary stenosis (SPS), and supra-aortic aortic stenosis (SAS) are commonly reported mid and long-term complications post the ASO procedure.<sup>ii xiv xviii xl xli</sup> NAR occurs early post ASO, is most commonly mild, but can progress especially among individuals with a history of size discrepancy between the pulmonary artery and aorta.<sup>xlii</sup> Freedom from moderate (or more) NAR is 98% at 1 year and 93% at 10 years.<sup>xli</sup>

Aortic root dilation has been reported to occur in at least two-thirds of TGA/ASO patients<sup>xli</sup> with a freedom from ARD at 10 years of 51%.<sup>xli</sup> Furthermore, 10-15% of TGA/ASO patients develop supra-aortic pulmonary stenosis by 20 years after ASO.<sup>x</sup> Branch pulmonary artery stenosis is the most frequent reason for re-intervention.<sup>xliii</sup>

Typically ventricular function post ASO is normal, however, mild left ventricular dysfunction can develop (e.g., EF 30-45%) and may occur secondary to coronary artery stenosis.<sup>xvii</sup> Given these findings, echocardiographic monitoring has been proposed as a reasonable method of screening for these complications.

#### Metric#6: Advanced Imaging

It was the opinion of the TGA/ASO team that there are times when echocardiogram images are limited and additional imaging modalities are necessary to evaluate for the presence of common mid and long term complications post ASO. Computed tomography (CT)<sup>xliv</sup>, magnetic resonance imaging (MRI) and cardiac catheterization with angiography<sup>xlv</sup> were proposed. Studies have shown that CT is a safe and feasible imaging modality for evaluating the coronary arteries in children 5- 6 years of age.<sup>xliv</sup> Unfortunately, advanced imaging is costly and invasive (e.g., coronary angiography) and it remains unclear what percentage of TGA/ASO patients will ultimately require such testing for such a recommendation to be considered a QM.



## Metric#7: Neurodevelopmental (ND) Assessment

The ASO procedure and the circulatory arrest required to carry out the ASO procedure, places patients at a greater risk of neurocognitive delay with 27% and 40% greater prevalence of neurologic and speech impairments among TGA/ASO, respectively.<sup>xxi</sup> Heart surgery performed with circulatory arrest as the predominant support strategy is associated with a higher risk of delayed motor development and neurologic abnormalities at the age of one year.<sup>xx</sup>

Transient postoperative seizures following the ASO procedure also place children at greater risk for worse ND outcomes at ages 1 to 2 1/2 years.<sup>xlvi</sup> Among children status post TGA/ASO, weaknesses have been noted in expressive language, visual-motor integration, motor planning and organization, and oromotor control.<sup>xxii xxiii xxiv</sup> Given the prevalence of these complications, neurocognitive testing was proposed as a QM.

## Metric#8-#10: Screening for acquired CVD risk factors

It is well known that children with congenital heart disease are at risk for acquired heart disease, deconditioning and metabolic disease.<sup>xlvi</sup> Studies suggest that children with TGA/ASO do not meet the recommended guidelines for physical activity<sup>xlvi</sup> and many children with CHD have modifiable risk factors for premature CVD, including diets rich in high fat and sugar and poor in the recommended intake of fresh fruit, vegetables and lean meat.<sup>xlix</sup> Furthermore, children who have undergo surgical manipulation of the coronary arteries, such as TGA/ASO patients, have been shown to have abnormalities in coronary function in later childhood and may be at increased risk for atherosclerosis as adults.<sup>ii xxvi xlvii</sup> Measurement of body mass index (BMI), blood pressure, and lipid screening are important and recommended according to existing guidelines.<sup>xxxi</sup>

## Metric#11-#12: Transition Plan

The AHA currently recommends that all adolescents have a written transition of care from the pediatric to the adult provider.<sup>xxxv</sup> The AAP recommends that this transition of care take place between 18 and 21 years of age.<sup>xxx</sup> The TGA/ASO team also recognized the importance of transition of care as well as reproductive health counseling.<sup>xxviii li lii liii</sup>

*Quality Metrics:*

Final recommendations from the TGA/ASO group submitted to the Steering Committee are shown in Table 1. Twelve QMS were proposed by the TGA/ASO group (See Appendix 1). Of the 12 candidate QMs, 7 passed the RAND process and were recommended for review by ACC members during the “open comment period.” Four QMs passed the “open comment period.”<sup>xxxii</sup> Selected QMs included: (a) the performance of at least 1 echocardiogram during the first year of life, after ASO, that reports the left ventricular function, aortic root dimension, the degree of neo-aortic valve insufficiency, and the patency of systemic and pulmonary outflow tracts, the branch pulmonary arteries and the coronary arteries; (b) ND assessment after ASO; (c) assessment of lipid profile by age 11 years; and (d) documentation of a transition of care plan to an adult congenital heart disease (ACHD) provider within the prior 2 years for ASO recipients 18 years of age or older.

Candidate QMs that did not pass the RAND process included QMs related to reproductive health counseling, arrhythmia screening, stress testing and use of additional, more expensive imaging modalities; these candidate QMs did not pass due to low validity scores (i.e., mean validity score of 6 during round 1 and mean validity score of 4 to 5 during round 2) and a lack of sufficient evidence. In particular, the following QMs did not pass the RAND process: (a) patients after ASO should be provided with age appropriate reproductive health counseling on sexual health, contraception, and pregnancy beginning early adolescence; (b) a patient after ASO should undergo periodic ECGs; (c) a patient after ASO should undergo periodic Holter

monitoring after the ASO; (d) a patient after ASO should have at least 1 stress test by age 11 years; and (e) advanced imaging (MRI, CT, cath) after ASO between 7 and 11 years of age.

Candidate QMs that passed the RAND process but that were not ultimately selected by the ACC ACPC Council following the “open comment period” included: (a) periodic echocardiogram monitoring beyond infancy following the ASO; (b) a patient after ASO should undergo regular surveillance of BMI and arterial pressure after ASO; and (c) provision of information outlining exercise recommendations after ASO. Although some single center studies suggested echocardiograms be performed every 1 to 2 years following ASO,<sup>ii</sup> comments received during the “open comment period” stated that the proposed frequency of echocardiography after infancy among ASO recipients was too aggressive. Several comments were received with suggestions to include additional imaging modalities, beyond echocardiography alone and a recommendation to decrease the frequency of echocardiograms to every 3-5 years. A recommendation for annual visits with the pediatric cardiologist were also found to be excessive. Although it is recommended that BMI and BP be assessed yearly among all children,<sup>xxxii</sup> not all children following ASO require annual visits with a pediatric cardiologist. Lipid levels on the other hand can typically be checked much less frequently and often with a single measurement between 9 and 11 years of age.<sup>xxxii</sup> Finally, a recommendation for providing information outlining exercise recommendations was considered too difficult to measure as data would be difficult to collect from the EMR.<sup>xxxii</sup>

## Discussion

Since the first ASO procedure in 1975, associated mortality rates have declined and efforts now focus on ensuring appropriate and data driven mid-term and long-term follow-up. The current study is timely and unique in that it identifies methods for improving the care of children with TGA/ASO. To date, existing guidelines regarding the management of individuals with

TGA/ASO have focused on adult populations or single center studies.<sup>ii xxxv</sup> The QMs created are now being utilized as a tool for data collection and quality improvement by 30 centers, participating in the Adult Congenital and Pediatric Quality Network- a national network of congenital heart disease (CHD) centers dedicated to collecting and sharing data with the goal of providing best practice for CHD patients.<sup>liv</sup>

- Twelve candidate QMs were proposed and 4 were approved. The approved QMs provide valid and feasible methods for assessing quality and improving the care of “simple” TGA/ASO patients. The metrics are not intended to address the highest risk TGA patients, including those with complex TGA who are more likely to require re-intervention or individuals with variant coronary artery anatomy who are at greatest risk for arrhythmia and sudden death.<sup>lv lvi lvii</sup>

The selected QMs address, however, the most common complications and risks for poor outcome in the simple TGA/ASO patient population. The selected metrics include use of echocardiograms to identify anatomic problems, followed by identifying risk factors associated with acquired heart disease (i.e., lipid screening), ND screening and transition of care. Metrics addressing less common complications, such as arrhythmia and coronary artery stenosis, were not selected due to lack of evidence and significantly lower reported rates of such complications in the non-high risk TGA/ASO population.<sup>xxi</sup> More costly imaging modalities such as MRI, CT and angiography were not selected as these tests may be of low yield in the majority of TGA/ASO patients.<sup>lviii</sup>

While not selected, screening BMI and arterial blood pressure should be performed at all ambulatory care visits regardless of the presence of TGA/ASO. Furthermore, even though a formal discussion of reproductive issues was not selected as a metric, it is quite possible that these discussions take place during the transition process.

Potential challenges with implementation of these metrics include variations in practice style, preference and limited resource availability. While echocardiograms are routinely performed and while lipid panels can be performed in the majority of children, ND and adult congenital resources may be limited. Future challenges will include not only making ND and adult congenital resources more readily available, but identifying appropriate quality metrics for complex cases of TGA. The current metrics do not take into consideration special circumstances, such as variant coronary artery anatomy, the presence of a large VSD or the presence of risk factors for early mortality including left ventricular outflow tract obstruction, arrhythmia, weight less than 2.5kg at time of ASO, or the need for extracorporeal membrane oxygenation.<sup>lix</sup>

Standardized practices for addressing less critical complications following the ASO are also needed. For example, metrics that address how best to manage supravalvar stenosis, branch pulmonary artery stenosis, and ARD are needed. Future efforts may include a potential role for biomarkers and the inclusion of studies that address risk stratification processes for those at greatest risk of SCD following ASO.

**Acknowledgements:**

The authors would like to thank Devyani Chowdhury, MD for her help and input

**Author Contributions:**

Carissa M Baker-Smith is a member of the transposition committee. She participated in content/design of the paper. She also drafted and critically reviewed the paper.

Karina Carlson and Jose Etedgui are members of the transposition committee, participated in content/design of the paper and critically reviewed the paper.

Takeshi Tsuda, K. Anitha Jayakumar, Matthew Park, Nikola Tede, Karen Uzark, Craig Fleishman, David Connuck and Maggie Likes were members of the transposition committee, assisted with generating content for the manuscript and approved the content of the paper.

Daniel J Penny was the team lead for the TGA/ASO committee and contributed to the content/design of the paper and critically reviewed the paper.

**REFERENCES:**

---

<sup>i</sup> Bonow RO, Masoudi FA, Rumsfeld JS, DeLong E, Estes NA 3rd, Goff DC Jr, Grady K, Green LA, Loth AR, Peterson ED, Piña IL, Radford MJ, Shahian DM; American College of Cardiology; American Heart Association Task Force on Performance Measures. ACC/AHA classification of care metrics: performance measures and quality metrics: a report of the American College of Cardiology/American Heart Association Task Force on Performance Measures. *J Am Coll Cardiol.* 2008 Dec 9;52(24):2113-7

<sup>ii</sup> Wernovsky G, Rome JJ, Tabbutt S, Rychik J, Cohen MS, Paridon SM, Webb G, Dodds KM, Gallagher MA, Fleck DA, Spray TL, Vetter VL, Gleason MM. Guidelines for the outpatient management of complex congenital heart disease. *Congenit Heart Dis.* 2006 Jan;1(1-2):10-26.

- 
- <sup>iii</sup> Khairy P, Clair M, Fernandes SM, Blume ED, Powell AJ, Newburger JW, Landzberg MJ, Mayer JE Jr. Cardiovascular outcomes after the arterial switch operation for D-transposition of the great arteries. *Circulation*. 2013 Jan 22;127(3):331-9.
- <sup>iv</sup> Rhodes LA, Wernovsky G, Keane JF, et al. Arrhythmias and intracardiac conduction after the arterial switch operation. *J Thorac Cardiovasc Surg*. 1995; 109(2):303-10.
- <sup>v</sup> Haas F, Wottke M, Poppert H, Meisner H. Long-term survival and functional follow-up in patients after the arterial switch operation. *Ann Thorac Surg*. 1999; 68(5):1692-7.
- <sup>vi</sup> Hayashi G, Kurosaki K, Echigo S, et al. Prevalence of arrhythmias and their risk factors mid- and long-term after the arterial switch operation. *Pediatr Cardiol*. 2006; 27(6):689-94.
- <sup>vii</sup> Khairy P, Clair M, Fernandes SM, et al. Cardiovascular outcomes after the arterial switch operation for D-transposition of the great arteries. *Circulation*. 2013; 127(3):331-9.
- <sup>viii</sup> Tobler D, Williams WG, Jegatheeswaran A, et al. Cardiac outcomes in young adult survivors of the arterial switch operation for transposition of the great arteries. *J Am Coll Cardiol*. 2010; 56(1):58-64.
- <sup>ix</sup> Silka MJ, Hardy BG, Menashe VD, Morris CD. A population-based prospective evaluation of risk of sudden cardiac death after operation for common congenital heart defects. *J Am Coll Cardiol*. 1998; 32(1):245-51.
- <sup>x</sup> Hutter PA, Kreb DL, Mantel SF, et al. Twenty-five years' experience with the arterial switch operation. *J Thorac Cardiovasc Surg*. 2002; 124(4):790-7.
- <sup>xi</sup> Mahle WT, McBride MG, Paridon SM. Exercise performance after the arterial switch operation for D-transposition of the great arteries. *Am J Cardiol*. 2011; 87(6):753-8.
- <sup>xii</sup> Pizzi MN, Franquet E, Aguadé-Bruix S, Manso B, Casaldàliga J, Cuberas-Borrós G, Romero-Farina G, Pinar J, Castell-Conesa J, García-Dorado D, Candell-Riera J. Long-term follow-up assessment after the arterial switch operation for correction of dextro-transposition of the great arteries by means of exercise myocardial perfusion-gated SPECT. *Pediatr Cardiol*. 2014 Feb;35(2):197-207.
- <sup>xiii</sup> Lalezari S, Bruggemans EF, Blom NA, Hazekamp MG. Thirty-year experience with the arterial switch operation. *Ann Thorac Surg*. 2011; 92(3):973-9.
- <sup>xiv</sup> Kempny A, Wustmann K, Borgia F, Dimopoulos K, Uebing A, Li W, Chen SS, Piorkowski A, Radley-Smith R, Yacoub MH, Gatzoulis MA, Shore DF, Swan L, Diller GP. Outcome in adult patients after arterial switch operation for transposition of the great arteries. *Int J Cardiol*. 2013 Sep 10;167(6):2588-93.
- <sup>xv</sup> Hutter PA, Thomeer BJ, Jansen P, et al. Fate of the aortic root after arterial switch operation. *Eur J Cardiothorac Surg*. 2001; 20(1):82-8.

- <sup>xvi</sup> Hourihan M, Colan SD, Wernovsky G, Maheswari U, Mayer JE Jr, Sanders SP. Growth of the aortic anastomosis, annulus, and root after the arterial switch procedure performed in infancy. *Circulation*. 1993; 88(2):615-20.
- <sup>xvii</sup> Choi BS, Kwon BS, Kim GB, et al. Long-term outcomes after an arterial switch operation for “simple” complete transposition of the great arteries. *Korean Circ J*. 2010; 40(1):23-30.
- <sup>xviii</sup> Bové T, De Meulder F, Vandenplas G, De Groote K, Panzer J, Suys B, DeWolf D, François K. Midterm assessment of the reconstructed arteries after the arterial switch operation. *Ann Thorac Surg*. 2008 Mar;85(3):823-30.
- <sup>xix</sup> Bellinger DC, Wypij D, Rivkin MJ, DeMaso DR, Robertson RL Jr, Dunbar-Masterson C, Rappaport LA, Wernovsky G, Jonas RA, Newburger JW. Adolescents with d-transposition of the great arteries corrected with the arterial switch procedure: neuropsychological assessment and structural brain imaging. *Circulation*. 2011 Sep 20;124(12):1361-9.
- <sup>xx</sup> Bellinger DC, Jonas RA, Rappaport LA, Wypij D, Wernovsky G, Kuban KC, Barnes PD, Holmes GL, Hickey PR, Strand RD, et al. Developmental and neurologic status of children after heart surgery with hypothermic circulatory arrest or low-flow cardiopulmonary bypass. *N Engl J Med*. 1995 Mar 2;332(9):549-5.
- <sup>xxi</sup> Hövels-Gürich HH, Seghaye MC, Schnitker R, Wiesner M, Huber W, Minkenber R, Kotlarek F, Messmer BJ, Von Bernuth G. Long-term neurodevelopmental outcomes in school-aged children after neonatal arterial switch operation. *J Thorac Cardiovasc Surg*. 2002 Sep;124(3):448-58.
- <sup>xxii</sup> Bellinger DC, Rappaport LA, Wypij D, Wernovsky G, Newburger JW. Patterns of developmental dysfunction after surgery during infancy to correct transposition of the great arteries. *J Dev Behav Pediatr*. 1997; 18(2):75-83.
- <sup>xxiii</sup> Calderon J, Jambaqué I, Bonnet D, Angeard N. Executive functions development in 5- to 7-year-old children with transposition of the great arteries: a longitudinal study. *Dev Neuropsychol*. 2014; 39(5):365-84.
- <sup>xxiv</sup> Calderon J, Bonnet D, Courtin C, Concordet S, Plumet MH, Angeard N. Executive function and theory of mind in school-aged children after neonatal corrective cardiac surgery for transposition of the great arteries. *Dev Med Child Neurol*. 2010; 52(12):1139-44.
- <sup>xxv</sup> Marino BS, Lipkin PH, Newburger JW, Peacock G, Gerdes M, Gaynor JW, Mussatto KA, Uzark K, Goldberg CS, Johnson WH Jr, Li J, Smith SE, Bellinger DC, Mahle WT; American Heart Association Congenital Heart Defects Committee, Council on Cardiovascular Disease in the Young, Council on Cardiovascular Nursing, and Stroke Council. Neurodevelopmental outcomes in children with congenital heart disease: evaluation and management: a scientific statement from the American Heart Association. *Circulation*. 2012 Aug 28;126(9):1143-72.
- <sup>xxvi</sup> Pasquali SK, Marino BS, Pudusseri A, Wernovsky G, Paridon SM, Walker SA, Cohen MS. Risk factors and comorbidities associated with obesity in children and adolescents after the arterial switch operation and Ross procedure. *Am Heart J*. 2009 Sep;158(3):473-9.



- <sup>xxvii</sup> Reid GJ, Irvine MJ, McCrindle BW, Sananes R, Ritvo PG, Siu SC, Webb GD. Prevalence and correlates of successful transfer from pediatric to adult health care among a cohort of young adults with complex congenital heart defects. *Pediatrics*. 2004 Mar;113(3 Pt 1):e197-205.
- <sup>xxviii</sup> Gurvitz MZ, Inkelas M, Lee M, Stout K, Escarce J, Chang RK. Changes in hospitalization patterns among patients with congenital heart disease during the transition from adolescence to adulthood. *J Am Coll Cardiol*. 2007 Feb 27;49(8):875-82.
- <sup>xxix</sup> Sable C, Foster E, Uzark K, Bjornsen K, Canobbio MM, Connolly HM, Graham TP, Gurvitz MZ, Kovacs A, Meadows AK, Reid GJ, Reiss JG, Rosenbaum KN, Sagerman PJ, Saidi A, Schonberg R, Shah S, Tong E, Williams RG; American Heart Association Congenital Heart Defects Committee of the Council on Cardiovascular Disease in the Young, Council on Cardiovascular Nursing, Council on Clinical Cardiology, and Council on Peripheral Vascular Disease. Best practices in managing transition to adulthood for adolescents with congenital heart disease: the transition process and medical and psychosocial issues: a scientific statement from the American Heart Association. *Circulation*. 2011 Apr 5;123(13):1454-85.
- <sup>xxx</sup> American Academy of Pediatrics; American Academy of Family Physicians; American College of Physicians; Transitions Clinical Report Authoring Group, Colley WC, Sagerman PJ. Supporting the health care transition from adolescence to adulthood in the medical home. *Pediatrics*. 2011; 128(1):182-200.
- <sup>xxxi</sup> Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents; National Heart, Lung, and Blood Institute. Expert panel on integrated guidelines for cardiovascular health and risk reduction in children and adolescents: summary report. *Pediatrics*. 2011; 128 Suppl 5:S213-56.
- <sup>xxxii</sup> Chowdhury D, Gurvitz M, Marelli A, Anderson J, Baker-Smith C, Diab KA, Edwards TC, Hougen T, Jedeikin R, Johnson JN, Karpawich P, Lai W, Lu JC, Mitchell S, Newburger JW, Penny DJ, Portman MA, Satou G, Teitel D, Villafane J, Williams R, Jenkins K; American College of Cardiology's Adult Congenital and Pediatric Cardiology Section's Ambulatory Pediatric Cardiology Quality Metrics Working Group. Development of Quality Metrics in Ambulatory Pediatric Cardiology. *J Am Coll Cardiol*. 2017 Feb 7;69(5):541-555.
- <sup>xxxiii</sup> Brook RH. The RAND/UCLA appropriateness method. In: McCormack KA, Moore SR, Siegel RA, editors. *Clinical Practice Guideline Development: Methodology Perspectives*. Rockville, MD: Agency for Healthcare Research and Policy, 1994:59–70.
- <sup>xxxiv</sup> Yeung E, Kay J, Roosevelt GE, Brandon M, Yetman AT. Lapse of care as a predictor for morbidity in adults with congenital heart disease. *Int J Cardiol*. 2008 Mar 28;125(1):62-5. Epub 2007 Apr 17. PubMed PMID: 1744243
- <sup>xxxv</sup> Warnes CA, Williams RG, Bashore TM, Child JS, Connolly HM, Dearani JA, del Nido P, Fasules JW, Graham TP Jr, Hijazi ZM, Hunt SA, King ME, Landzberg MJ, Miner PD, Radford MJ, Walsh EP, Webb GD, Smith SC Jr, Jacobs AK, Adams CD, Anderson JL, Antman EM, Buller CE, Creager MA, Ettinger SM, Halperin JL, Hunt SA, Krumholz HM, Kushner FG, Lytle BW, Nishimura RA, Page RL, Riegel B, Tarkington LG, Yancy CW; American College of Cardiology; American Heart Association Task Force on Practice Guidelines (Writing Committee to Develop Guidelines on the Management of Adults With Congenital Heart Disease); American

Society of Echocardiography; Heart Rhythm Society; International Society for Adult Congenital Heart Disease; Society for Cardiovascular Angiography and Interventions; Society of Thoracic Surgeons. ACC/AHA 2008 guidelines for the management of adults with congenital heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Develop Guidelines on the Management of Adults With Congenital Heart Disease). Developed in Collaboration With the American Society of Echocardiography, Heart Rhythm Society, International Society for Adult Congenital Heart Disease, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. 2008 Dec 2;52(23):e143-263.

<sup>xxxvi</sup> Angeli E, Formigari R, Pace Napoleone C, Oppido G, Ragni L, Picchio FM, Gargiulo G. Long-term coronary artery outcome after arterial switch operation for transposition of the great arteries. *Eur J Cardiothorac Surg*. 2010 Dec;38(6):714-20

<sup>xxxvii</sup> Prêtre R, Tamisier D, Bonhoeffer P, Mauriat P, Pouard P, Sidi D, Vouhé P. Results of the arterial switch operation in neonates with transposed great arteries. *Lancet*. 2001 Jun 9;357(9271):1826-30.

<sup>xxxviii</sup> Legendre A, Losay J, Touchot-Koné A, Serraf A, Belli E, Piot JD, Lambert V, Capderou A, Planche C. Coronary events after arterial switch operation for transposition of the great arteries. *Circulation*. 2003 Sep 9;108 Suppl 1:II186-90.

<sup>xxxix</sup> Massin M, Hövels-Gürich H, Däbritz S, Messmer B, von Bernuth G. Results of the Bruce treadmill test in children after arterial switch operation for simple transposition of the great arteries. *Am J Cardiol*. 1998 Jan 1;81(1):56-60.

<sup>xl</sup> Vandekerckhove KD, Blom NA, Lalezari S, Koolbergen DR, Rijlaarsdam ME, Hazekamp MG. Long-term follow-up of arterial switch operation with an emphasis on function and dimensions of left ventricle and aorta. *Eur J Cardiothorac Surg*. 2009 Apr;35(4):582-7; discussion 587-8.

<sup>xli</sup> Schwartz ML, Gauvreau K, del Nido P, Mayer JE, Colan SD. Long-term predictors of aortic root dilation and aortic regurgitation after arterial switch operation. *Circulation*. 2004 Sep 14;110(11 Suppl 1):II128-32.

<sup>xlii</sup> Hwang HY, Kim WH, Kwak JG, Lee JR, Kim YJ, Rho JR, Bae EJ, Noh CI. Mid-term follow-up of neoaortic regurgitation after the arterial switch operation for transposition of the great arteries. *Eur J Cardiothorac Surg*. 2006 Feb;29(2):162-7.

<sup>xliii</sup> Massin MM, Nitsch GB, Däbritz S, Seghaye MC, Messmer BJ, von Bernuth G. Growth of pulmonary artery after arterial switch operation for simple transposition of the great arteries. *Eur J Pediatr*. 1998 Feb;157(2):95-100.

<sup>xliv</sup> Ou P, Celermajer DS, Marini D, Agnoletti G, Vouhé P, Brunelle F, Le Quan Sang KH, Thalabard JC, Sidi D, Bonnet D. Safety and accuracy of 64-slice computed tomography coronary angiography in children after the arterial switch operation for transposition of the great arteries. *JACC Cardiovasc Imaging*. 2008 May;1(3):331-9

- <sup>xiv</sup> Hutter PA, Bennink GB, Ay L, Raes IB, Hitchcock JF, Meijboom EJ. Influence of coronary anatomy and reimplantation on the long-term outcome of the arterial switch. *Eur J Cardiothorac Surg*. 2000 Aug;18(2):207-13
- <sup>xv</sup> Rappaport LA, Wypij D, Bellinger DC, Helmers SL, Holmes GL, Barnes PD, Wernovsky G, Kuban KC, Jonas RA, Newburger JW. Relation of seizures after cardiac surgery in early infancy to neurodevelopmental outcome. Boston Circulatory Arrest Study Group. *Circulation*. 1998 Mar 3;97(8):773-9.
- <sup>xvii</sup> Pasquali SK, Marino BS, Powell DJ, McBride MG, Paridon SM, Meyers KE, Mohler ER, Walker SA, Kren S, Cohen MS. Following the arterial switch operation, obese children have risk factors for early cardiovascular disease. *Congenit Heart Dis*. 2010 Jan-Feb;5(1):16-24
- <sup>xviii</sup> Massin MM, Hövels-Gürich HH, Gérard P, Seghaye MC. Physical activity patterns of children after neonatal arterial switch operation. *Ann Thorac Surg*. 2006 Feb;81(2):665-70.
- <sup>xix</sup> Massin MM, Hövels-Gürich H, Seghaye MC. Atherosclerosis lifestyle risk factors in children with congenital heart disease. *Eur J Cardiovasc Prev Rehabil*. 2007 Apr;14(2):349-51
- <sup>i</sup> American Academy of Pediatrics; American Academy of Family Physicians; American College of Physicians; Transitions Clinical Report Authoring Group, Cooley WC, Sagerman PJ. Supporting the health care transition from adolescence to adulthood in the medical home. *Pediatrics*. 2011 Jul;128(1):182-200.
- <sup>ii</sup> Hilderson D, Saidi AS, Van Deyk K, Verstappen A, Kovacs AH, Fernandes SM, Canobbio MM, Fleck D, Meadows A, Linstead R, Moons P. Attitude toward and current practice of transfer and transition of adolescents with congenital heart disease in the United States of America and Europe. *Pediatr Cardiol*. 2009 Aug;30(6):786-93.
- <sup>iii</sup> Goossens E, Norekvål TM, Faerch J, Hody L, Olsen SS, Darmer MR, Jaarsma T, Moons P. Sexual counselling of cardiac patients in Europe: culture matters. *Int J Clin Pract*. 2011 Oct;65(10):1092-9.
- <sup>iiii</sup> Knauth A, Verstappen A, Reiss J, Webb GD. Transition and transfer from pediatric to adult care of the young adult with complex congenital heart disease. *Cardiol Clin*. 2006 Nov;24(4):619-29.
- <sup>liv</sup> <https://cvquality.acc.org/initiatives/ACPC-Quality-Network>
- <sup>lv</sup> Rudra HS, Mavroudis C, Backer CL, Kaushal S, Russell H, Stewart RD, Webb C, Sullivan C. The arterial switch operation: 25-year experience with 258 patients. *Ann Thorac Surg*. 2011 Nov;92(5):1742-6.
- <sup>lvi</sup> Daebritz SH, Nollert G, Sachweh JS, Engelhardt W, von Bernuth G, Messmer BJ. Anatomical risk factors for mortality and cardiac morbidity after arterial switch operation. *Ann Thorac Surg*. 2000; 69(6):1880-6.

---

<sup>lvi</sup> Losay J, Touchot A, Serraf A, Litvinova A, Lambert V, Piot JD, Lacour-Gayet F, Capderou A, Planche C. Late outcome after arterial switch operation for transposition of the great arteries. *Circulation*. 2001 Sep 18;104(12 Suppl 1):I121-6.

<sup>lviii</sup> Villafañe J, Lantin-Hermoso MR, Bhatt AB, Tweddell JS, Geva T, Nathan M, Elliott MJ, Vetter VL, Paridon SM, Kochilas L, Jenkins KJ, Beekman RH 3rd, Wernovsky G, Towbin JA; American College of Cardiology's Adult Congenital and Pediatric Cardiology Council.. D-transposition of the great arteries: the current era of the arterial switch operation. *J Am Coll Cardiol*. 2014 Aug 5;64(5):498-511.

<sup>lix</sup> Fricke TA, d'Udekem Y, Richardson M, Thuys C, Dronavalli M, Ramsay JM, Wheaton G, Grigg LE, Brizard CP, Konstantinov IE. Outcomes of the arterial switch operation for transposition of the great arteries: 25 years of experience. *Ann Thorac Surg*. 2012 Jul;94(1):139-45.

Accepted Article

Table 1. Candidate Quality Metrics for transposition of the great arteries (TGA) after arterial switch operation submitted to Steering Committee

Candidate Measure	Numerator	Denominator
<p>At least one echo in the first year of life, after ASO that reports on LV function, aortic root dimension, degree of aortic insufficiency (AI), patency of systemic and pulmonary outflows, branch pulmonary artery stenosis and coronary arteries.</p>	<p>Number of ASO patients reaching their first birthday for whom at least 1 echocardiogram has been performed and which documents left ventricular function, aortic root dimensions, degree of aortic regurgitation, nature of systemic and pulmonary outflow tracts and the branch pulmonary arteries</p>	<p>Number of ASO patients reaching their first birthday</p>

<p>Neurodevelopment assessment after ASO</p>	<p>Number of ASO patients age 24 months or more who have undergone formal, age-appropriate neurodevelopmental assessment within the previous 12 months and the number of patients ages 5 to 12 years who have undergone assessment within the previous 24 months</p>	<p>Number of ASO patients reaching the age of 24 months and the number of patients between 5 and 12 years</p>
<p>Assessment of lipid profile by age 11 years</p>	<p>Number of ASO patients age 11 years who have undergone assessment of lipid profile within previous 2 years</p>	<p>Number of ASO patients reaching their 11th birthday</p>
<p>Transition of care: patients with ASO (at age 18 or greater) with documented transition of care in the past 2 years</p>	<p>Patients following ASO, age 18 years with a transition plan documented in the medical record within the previous 2 years</p>	<p>Number of ASO patients age 18 years</p>