Erratum: Semi-automated pulmonary nodule interval segmentation using the NLST data

Yoganand Balagurunathana)

H.L.Moffitt Cancer Center (MCC), Tampa, FL, USA

Andrew Beers and Jayashree Kalpathy-Cramer

Massachusetts General Hospital (MGH), Boston, MA, USA

Michael McNitt-Gray

University of California Los Angeles (UCLA), Los Angeles, CA, USA

Lubomir Hadjiiski

University of Michigan (UMICH), Ann Arbor, MI, USA

Bensheng Zhao, Jianguo Zhu, and Hao Yang

Columbia University (CUMU), New York, NY, USA

Stephen S. F. Yip and Hugo J. W. L. Aerts

Radiation Oncology, Dana-Farber Cancer Institute (DFCC), Brigham and Women's Hospital (BWH), Harvard Medical School (HMC), Boston, MA, USA

Radiology, Dana-Farber Cancer Institute (DFCC), Brigham and Women's Hospital (BWH), Harvard Medical School (HMC), Boston, MA, USA

Sandy Napel

Stanford University (SU), CA, USA

Dmitrii Cherezov

H.L.Moffitt Cancer Center (MCC), Tampa, FL, USA University of South Florida (USF), Tampa, FL, USA

Kenny Cha and Heang-Ping Chan

University of Michigan (UMICH), Ann Arbor, MI, USA

Carlos Flores

University of California Los Angeles (UCLA), Los Angeles, CA, USA

Alberto Garcia and Robert Gillies

H.L.Moffitt Cancer Center (MCC), Tampa, FL, USA

Dmitry Goldgof^{a)}

H.L.Moffitt Cancer Center (MCC), Tampa, FL, USA University of South Florida (USF), Tampa, FL, USA

(Received 31 August 2018; accepted for publication 4 January 2018; published March 2018)

[https://doi.org/10.1002/mp.12766]

There is an error in the contributing author's name on the published article¹.

The correct spelling is Jianguo Zhu (misspelled as Jiangguo Zhu).

Page 12: Discussion Section:

Corrected to: The Food and Drug Administration (FDA)

Current: The Federal Drug Administration (FDA)

Corrected to:

Recently these nodules were segmented using the shape and location information (matched filter-based approach) to assess the sizes and volume estimation, a variability of 3.9 to 28% has been reported, they have shown a higher variability for smaller nodules.

Current:

Recently these nodules were given to a clinical radiologist to assess the sizes and volume estimation, a variability of 3.9 to 28% has been reported, they have shown a higher variability for smaller nodules.

REFERENCE

1. Balagurunathan Y, Beers A, Kalpathy-Cramer J, et al. Semi-automated pulmonary nodule interval segmentation using the NLST data. *Med Phys.* 2018;45:1093–1107.

^{a)}Authors to whom correspondence should be addressed. Eelectronic mails: goldgof@mail.usf.edu; yogab@moffitt.org.