

The Neurobiology of Central Sensitization

Steven E. Harte PhD, Richard E. Harris PhD, Daniel J. Clauw MD

Chronic Pain and Fatigue Research Center, Department of Anesthesiology, University of
Michigan, Ann Arbor, MI

Corresponding Author

Steven Harte, PhD

UM/CPFRC

24 Frank Lloyd Wright Dr.

PO Box 385, Lobby M

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/jabr.12137](https://doi.org/10.1111/jabr.12137)

This article is protected by copyright. All rights reserved

Ann Arbor, MI 48106

734-998-7053

seharte@med.umich.edu

Disclosures

Dr. Harte has received research funding from NIH, Aptinyx, Cerephex, Forest Laboratories, Eli Lilly and Merck; he has served as a consultant for Pfizer, Coy Labs, Analgesic Solutions, Aptinyx, and deCode Genetics. Dr. Harte is also member of Arbor Medical Innovations. Dr. Harris has received research funding from NIH; he also consulted with and received research funding from Pfizer. Dr. Clauw has received research funding from NIH, DoD, Cerephex, Forest, Merck, and Pfizer, and has served as a consultant for Tonix, Pfizer, Depomed, Sammumed, Aptinyx, and Zynerba.

DR. STEVEN E HARTE (Orcid ID : 0000-0002-2929-8694)

Article type : Special Issue

OVERVIEW

Central sensitization refers to the amplification of pain by central nervous system mechanisms. Classically described as a consequence of ongoing nociceptive input, it is increasingly recognized that central sensitization also occurs independent of peripheral injury or inflammation. Features of central sensitization have been identified in nearly all chronic pain conditions, and it is considered the primary underlying cause of pain in conditions such as fibromyalgia. Central sensitization is characterized in these conditions by widespread pain and multisite hyperalgesia/allodynia. Co-occurring symptoms include fatigue, mood and cognitive problems, sleep disturbances, and multisensory hypersensitivity. Individuals with central sensitization often report previous exposure to psychosocial or physical stressors, and a higher personal lifetime and family history of pain, with the latter findings supported by genetic studies. Neuroimaging studies of central sensitization show evidence of: changes in brain gray matter in pain processing regions; neurochemical imbalances; and altered resting brain-network connectivity between pro-nociceptive and antinociceptive brain areas. Immune system abnormalities have also been demonstrated in individuals with central sensitization. The recognition of central sensitization, and whether it is being driven by ongoing nociceptive input or it is occurring in the absence of a peripheral-driver, is critical for effective pain management.

INTRODUCTION

In this Special Issue of JABR, Woolf describes the first use of the term “central sensitization” as it relates to pain processing, as well as the early evolution of the meaning of this term. Described initially only in animal models, the term was first used by Woolf to refer: to a specific

spinal disorder mechanism that was responsible for augmenting ongoing peripheral nociceptive input (Woolf, 1983). Other preclinical pain researchers subsequently identified additional spinal mechanisms that amplify nociceptive signaling. The hallmarks of these processes were the presence of hyperalgesia and/or allodynia on quantitative sensory testing (QST) in animal models of nociception, often accompanied by enlargement of the nociceptive field and/or electrophysiological evidence of decreased firing threshold and increased discharge of spinal nociceptive neurons. Thus, the term central sensitization began to be used in a broader sense to convey the notion that an array of spinal mechanisms may be involved in amplifying or maintaining peripheral nociceptive input (Dougherty, Palecek, Paleckova, Sorkin, & Willis, 1992; Sluka & Westlund, 1992; Treede, Davis, Campbell, & Raja, 1992; Willis, 1988). Even though a substantial body of research had begun to demonstrate the important role for supraspinal mechanisms in pain processing (Gebhart, 1986; T. S. Jensen & Yaksh, 1992; Willis, 1985; Yaksh, 1988), in the 1980s and early 1990s the term central sensitization was originally applied: 1) only to situations where there was some initial ongoing peripheral nociceptive input; and 2) where the underlying mechanism(s) of this phenomenon was thought to lie in the spinal cord and dorsal root ganglion.

In the mid-1990's, many investigators began to leverage the term and concept of central sensitization as a way of explaining the pain hypersensitivity observed in a number of chronic pain states in humans, including neuropathic pain (Gracely, Lynch, & Bennett, 1992; Persson, Axelsson, Hallin, & Gustafsson, 1995), fibromyalgia (FM) (Anderberg, 2000; Gibson, Littlejohn, Gorman, Helme, & Granges, 1994; Granges & Littlejohn, 1993; Kosek, Ekholm, & Hansson, 1995; Sorensen, Bengtsson, Backman, Henriksson, & Bengtsson, 1995); headache (R. Jensen, Rasmussen, Pedersen, Lous, & Olesen, 1992; R. Jensen, Rasmussen, Pedersen, & Olesen, 1993); temporomandibular disorder (TMD) (Fillingim, Maixner, Kincaid, Sigurdsson, & Harris, 1996); and irritable bowel syndrome (IBS) (Mertz, Naliboff, Munakata, Niazi, & Mayer, 1995). As with the animal models, the original technique used to identify central sensitization in humans was QST. These early QST studies typically noted the same findings of secondary hyperalgesia or

allodynia seen in animal models of central sensitization, and this was used to infer that these same underlying processes may be playing a role in humans with these chronic pain conditions.

More recently, newer techniques, such as functional, chemical, and structural brain imaging, EEG, and other research techniques, have been used to substantiate and further identify the presence of, and underlying mechanisms driving, central sensitization. Central sensitization has now been suggested to play a role in hundreds of painful and other medical conditions. In fact, so much has been learned about the ubiquity of this problem in populations of chronic pain patients that there is now a broad consensus in the pain field that this might represent a third underlying core mechanism for pain, above and beyond nociceptive and neuropathic pain mechanisms that have been recognized for some time (C. Brummett et al., 2016; Kosek et al., 2016).

Chronic pain conditions that are now recognized mechanistically to fall within the spectrum of central sensitization include: FM; IBS (previously termed “spastic colitis” until the recognition that there was little “-itis” and that motility changes were not the major pathological feature); TMD (previously termed temporomandibular joint syndrome until it was recognized the problem was largely outside the joint) (Bair, Brownstein, et al., 2013; Bair, Ohrbach, et al., 2013; Fillingim et al., 2011; Greenspan et al., 2013); and urinary chronic pelvic pain syndromes (where, again, the condition previously called interstitial cystitis is now called bladder pain syndrome) (Bagarinao et al., 2014; Clemens et al., 2014; Farmer et al., 2015). This is not to say peripheral factors, or low-grade inflammation that is not identifiable clinically, do not play some role in these entities. But it is relevant that clinicians who care for individuals with these conditions, and who are quite adept at identifying (with blood tests, imaging, or endoscopy) peripheral damage or inflammation, have generally concluded that these are not inflammatory or peripheral-based disorders. A recent term, *Chronic Overlapping Pain Conditions (COPCs)*, has been coined by the NIH to indicate that FM, IBS, chronic fatigue syndrome (CFS), headache, interstitial cystitis/bladder pain syndrome, TMD, endometriosis, low back pain, and dry-eye disease may all represent conditions with overlapping clinical and pathophysiological features

(i.e., those related to central sensitization), where central factors may be playing a prominent or exclusive role in their pathogenesis (Levitt et al., 2017; Maixner, Fillingim, Williams, Smith, & Slade, 2016). Most of the initial studies examining the pathophysiology of central sensitization in humans have been in conditions such as FM and other *COPCs*, because these were amongst the first conditions where prominent central factors were identified.

However, these same features suggesting central sensitization are also commonly seen in individuals with chronic pain conditions and illnesses *where there is clearly ongoing nociceptive input*. A subset of individuals with any chronic pain condition will display features of central sensitization, including individuals with autoimmune disorders, any type of arthritis, cancer, sickle cell disease, and hypermobility syndromes (Clauw & Witter, 2009; Darbari, Ballas, & Clauw, 2014; Gwilym et al., 2009; Henry & Clauw, 2012; Phillips & Clauw, 2013). Because these individuals with central sensitization clinically resemble individuals with FM, they were previously labelled as having “secondary fibromyalgia” in the rheumatological literature. This form of central sensitization has not been nearly as well studied mechanistically in humans, in part because most studies of *COPCs* purposefully excluded individuals with demonstrable peripheral disease. However, what is known about this type of central sensitization is that it seems to be at least in part driven by ongoing nociceptive input because, when the input is removed, the features of central sensitization partially or entirely remit (Gwilym, Filippini, Douaud, Carr, & Tracey, 2010; Kosek & Ordeberg, 2000; Staud, Weyl, Bartley, Price, & Robinson, 2014).

This present review will provide a broad overview of these aggregate studies describing the current understanding of central sensitization as observed in humans, as well as identify unresolved issues in this field.

FEATURES OF CENTRAL SENSITIZATION GLEANED FROM EPIDEMIOLOGICAL OR OBSERVATIONAL STUDIES

Individuals who develop *COPCs* begin developing pain and other related central-sensitization symptoms (fatigue, sleep disturbance, sensory sensitivity) in early life. For example, individuals

who eventually go on to develop FM are more likely to experience headaches, dysmenorrhea, TMD, chronic fatigue, IBS and other functional GI disorders, interstitial cystitis/painful bladder syndrome, endometriosis, and other regional pain syndromes (especially back and neck pain) (Aaron & Buchwald, 2001; Hudson & Pope, 1994). What often appears clinically as a new episode of acute or subacute pain is just the newest region of the body experiencing pain (Warren et al., 2009). Consequently, many pain experts have suggested that *COPCs* are best understood as a single lifelong disease that merely tends to manifest in different bodily regions over time (Tracey & Bushnell, 2009; Williams & Clauw, 2009; Woolf, 2011).

In addition to *COPC*, patients having a high personal lifetime history of pain, often have a strong family history of chronic pain. The first-degree relatives of FM patients are eight-times more likely to have FM compared to family members of healthy controls, and they also report very high rates of other chronic pain conditions (Arnold et al., 2004). Furthermore, family members of individuals with FM are much more tender than the family members of controls, regardless of whether they have chronic pain or not (Buskila, Neumann, Hazanov, & Carmi, 1996; Hudson, Hudson, Pliner, Goldenberg, & Pope, 1985; Kato, Sullivan, Evengard, & Pedersen, 2006). This familial and personal co-aggregation of conditions has been called many other terms prior to *COPCs*, including *affective spectrum disorder* (Hudson, Goldenberg, Pope, Keck, & Schlesinger, 1992) and, more recently, *central sensitivity syndromes* (Yunus, 2008), and *chronic multi-symptom illnesses* (Yunus, 2008). The key symptoms besides pain that typically co-aggregate together are fatigue, memory difficulties, and mood disturbances (Fukuda et al., 1997; Fukuda et al., 1998). Twin studies suggest that approximately 50% of the risk of developing these conditions is genetic, and 50% environmental (Kato, Sullivan, Evengard, & Pedersen, 2009).

Environmental factors that are associated with the development of FM and other *COPCs* include various types of “stressors,” that typically involve acute pain for at least a few weeks. Psychosocial stress, including childhood trauma, is but one such stressor. FM and other *COPCs* are found at much higher than expected rates in individuals who have experienced certain types of infections (Buskila, Atzeni, & Sarzi-Puttini, 2008 [e.g., Epstein Barr Virus, Lyme disease,

Q fever, viral hepatitis]), physical trauma (e.g., motor vehicle collisions) (McLean et al., 2011), and deployment to war (J. D. Lewis et al., 2012). Of note, only 5-10% of individuals exposed to these stressors develop FM or CFS; the overwhelming majority of individuals who experience these same infections or other stressful events return to their baseline state of health. It is also likely that part of the reason these different stressors can seemingly lead to worsening of central sensitization symptoms is because of how these stressors affect activity level, sleep, or overall distress, any of which then can lead to worsening of pain and other symptoms (J. N. Ablin et al., 2013; Arnsen et al., 2007; Crook, Moldofsky, & Shannon, 1998; Moldofsky, 2010; Moldofsky, Saskin, & Lue, 1988). This may also be why treatments, such as exercise and improving sleep, seem to be so important and helpful in the overall treatment of patients with central sensitization.

As previously noted, the central sensitization or FM-phenotype is also commonly seen as a comorbidity in other chronic pain conditions such as osteoarthritis, rheumatoid arthritis, and systemic lupus erythematosus (SLE) (Buskila, Neumann, Vaisberg, Alkalay, & Wolfe, 1997; Buskila, Shnaider, et al., 1997; Clauw et al., 1997; McLean & Clauw, 2004). As many as 25% of patients correctly diagnosed with generalized inflammatory disorders such as SLE, rheumatoid arthritis, and ankylosing spondylitis will also meet the clinical criteria for FM (Clauw, 1995a, 1995b). This rate is even higher in individuals with conditions such as chronic low back pain, sickle cell disease, Ehler's Danlos and other hypermobility syndromes. However, in clinical practice, this co-expression is often unrecognized, especially when central sensitization develops after the nociceptive or neuropathic condition is diagnosed. In this setting, when comorbid central sensitization goes unrecognized, patients are often unnecessarily treated more aggressively with peripherally-direct interventions, such as surgery or immunosuppressive drugs, leading to increased morbidity without the desired therapeutic effects.

FM and other *COPCs* occur much more commonly in individuals with early life and current stress, and many if not most individuals will have a lifetime history of a psychiatric disorder such as depression or anxiety (Epstein et al., 1999). There is typically more psychiatric and

psychological comorbidity seen in tertiary-care settings or in individuals who are refractory to treatment. This bi-directional relationship between FM and psychiatric conditions is likely due in part to the fact that there are common triggers to both sets of conditions (e.g., early life stress or trauma), as well as shared pathophysiology (i.e., many of the neurotransmitters that affect pain transmission also affect mood, memory, fatigue, sleep). Other risk factors for developing FM that are potentially modifiable include poor sleep, obesity, physical inactivity, and poor job or life satisfaction. Similarly, cognitive factors, such as catastrophizing (the feeling that pain is very bad and associated with a poor prognosis for recovery) or movement-related fear, are poor prognostic factors in FM and other chronic pain states.

Animal Models As noted above, all of the early animal models of central sensitization required ongoing nociceptive input to drive central sensitization. In fact, this was one of the hallmark features of central sensitization in these models, in that the observed processes seemed to lessen or disappear when the ongoing peripheral nociceptive input was removed or blocked. This led to the early view that aggressively reducing ongoing nociceptive input to block subsequent central sensitization (i.e., pre-emptive analgesia [Woolf, 1991]) might be useful therapeutically. This general approach is still used, especially in perioperative medicine, where studies generally show that individuals with lower pain scores during the acute painful episode have less subsequent features of central sensitization following surgery (although it is often difficult to disentangle *cause vs. effect* in these studies, because individuals with less central sensitization would be expected to have less pain to an acute stimulus).

However, more recent animal models of both somatic and visceral pain emphasize that these same features of central sensitization can occur without any ongoing nociceptive input. For example, although few would purport that there is an animal model that mimics all of the key clinical features of FM, animal models can nonetheless be very helpful in understanding the pathogenesis of this condition (Sluka, 2009; Sluka & Clauw, 2016). Animals can clearly develop the critical features of central sensitization without any ongoing nociceptive input, including when exposed to sub-chronic swim stress (Suarez-Roca et al., 2006); neonatal separation from

their mothers (Pierce & Christianson, 2015); and many other non-painful stimuli (DeSantana, da Cruz, & Sluka, 2013; Sluka, Kalra, & Moore, 2001). Features of central sensitization and animal pain-like behaviors consistent with diffuse pain are also seen when central nervous system (CNS) neurotransmitters are purposefully altered in the directions found in FM. For example, chronic reserpine administration (which depletes bioamines) leads to features consistent with central sensitization (Taguchi et al., 2015), as does directly increasing glutamate levels in the insulae (Watson, 2016). This latter model of increasing insular glutamate was recently reverse-translated to show that the findings of “small fiber neuropathy” could even be induced simply by increasing CNS glutamate, as is known to occur in FM (Harte et al., 2017).

Genetic Factors The strong familial predisposition to FM and other chronic pain conditions has led many to study specific genetic polymorphisms that may be associated with a higher risk of developing FM. First, candidate-gene studies showed that genetic findings, such as the serotonin 5-HT_{2A} receptor polymorphism T/T phenotype; serotonin transporter; dopamine 4 receptor; and COMT (catecholamine *o*-methyl transferase) polymorphisms all were noted at higher-frequency in FM patients than controls. Some subsequent studies confirmed certain of these associations, whereas others did not (Buskila, Sarzi-Puttini, & Ablin, 2007; Diatchenko, Fillingim, Smith, & Maixner, 2013). Subsequent larger genome-wide linkage and candidate-gene studies identified other putative targets (Arnold et al., 2013; Smith et al., 2012). The linkage studies confirmed the strong genetic contribution to FM, and suggested linkage of FM to the chromosome 17p11.2-q11.2 region. The large candidate gene study identified three genes -- (GABRB3 [rs4906902, $p = 3.65 \times 10^{-6}$]; TAAR1 (rs8192619, $p = 1.11 \times 10^{-5}$); and GBP1 (rs7911, $p = 1.06 \times 10^{-4}$) with single nucleotide polymorphisms (SNPs) -- that differed in frequency between FM patients and healthy controls (Smith et al., 2012). TAAR1, along with several additional genes (RGS4, CNR1, and GRIA4) showed evidence of association with FM in a replication analysis using a second independent cohort of FM patients (Smith et al., 2012). In light of the fact that classic genetic studies have not yet identified strong, reproducible polymorphisms or haplotypes associated with FM, and because there is clear evidence of environmental factors (such as stress) playing a prominent role in the pathogenesis, other

groups have postulated that epigenetic findings might be important in FM (Ciampi de Andrade et al., 2017). This is a promising area that needs further research.

QST EVIDENCE OF CNS DISTURBANCES IN PAIN AND SENSORY PROCESSING

The physiological hallmark and defining feature of central sensitization is altered CNS pain-processing. This was originally identified clinically in conditions such as FM and TMD by identifying tenderness to palpation. Following these early clinical observations, many groups have used much more sophisticated QST methods to identify and characterize pain and sensory mechanisms in individuals suspected to have central sensitization (Arendt-Nielsen, 2015; Arendt-Nielsen & Yarnitsky, 2009; Backonja et al., 2013; Cruz-Almeida & Fillingim, 2014; Edwards, Sarlani, Wesselmann, & Fillingim, 2005; Greenspan et al., 2013; Pfau, Geber, Birklein, & Treede, 2012; Wylde et al., 2017). QST refers to procedures in which quantifiable stimuli (e.g., mechanical, thermal, or electric) are delivered in a systematic manner to the body. An individual's reaction to these stimuli, including non-verbal behavioral responses, such as AS withdrawal, ratings of perceived stimulus intensity and unpleasantness, and physiological responses, are used to measure sensory gain or loss. Data from QST studies suggest a wide, bell-shaped distribution in pain sensitivity across the general population. Most, but not all, individuals with *COPCs*, and a subset of individuals with any other chronic pain condition, will typically fall on the right side of the curve, with significant diffuse hypersensitivity outside the region of injury (i.e., secondary hyperalgesia and allodynia) (K. Ablin & Clauw, 2009; Diatchenko et al., 2006; Fillingim, 2005; Gwilym et al., 2009; Kleinbohl et al., 1999; Neddermeyer, Fluhr, & Lotsch, 2008; Nielsen, Staud, & Price, 2009; Tracey & Bushnell, 2009; Williams & Clauw, 2009). Importantly, this hypersensitivity can be present in both painful/symptomatic, as well as pain-free/non-symptomatic body sites, which strongly suggests a central pain mechanism. Some of the chronic pain conditions where widespread hyperalgesia/allodynia is consistently identified include FM, IBS, TMD, low back pain, tension headache, chronic pelvic pain, and vulvodynia (As-Sanie et al., 2013; Clauw & Chrousos, 1997; Gibson et al., 1994; Giesecke et al., 2004; Greenspan et al., 2011; Grinberg, Granot, Lowenstein, Abramov, & Weissman-Fogel, 2017; Leffler, Hansson, & Kosek, 2002; Petzke, Harris, Williams, Clauw, & Gracely, 2005; Whitehead et

al., 1990; O. H. G. Wilder-Smith, Tassonyi, & Arendt-Nielsen, 2002).

The efficiency of endogenous pain modulation networks is another marker of altered CNS pain processing measured by QST. Endogenous pain inhibitory networks can be activated in humans and animals during application of a noxious “conditioning” stimulus, resulting in attenuation of pain produced by a noxious “test” stimulus applied to a different body location -- an effect termed *conditioned pain modulation* (CPM). Although deficient CPM (i.e., reduced inhibition of test stimulus evoked pain) or even facilitation of test stimulus pain has been observed in a subset of healthy individuals (Potvin & Marchand, 2016; Schoen et al., 2016), it is more common in individuals with FM and other *COPCs* (Granot et al., 2008; Julien, Goffaux, Arsenault, & Marchand, 2005; Kosek & Ordeberg, 2000; Le Bars, Dickenson, & Besson, 1979; Mahomed, Downing, Jeal, & Coleman, 1976; Nir & Yarnitsky, 2015; C. H. Wilder-Smith & Robert-Yap, 2007; Yarnitsky et al., 2008). In fact, a 2012 meta-analysis confirmed that CPM is attenuated in the majority of chronic pain conditions (Lewis, Rice, & McNair, 2012). Decreased CPM is associated with a variety of clinical outcomes, including increased post-surgical pain and analgesia requirements, and the effectiveness of some centrally-acting analgesics (Edwards, 2005; Yarnitsky, 2015; Yarnitsky et al., 2008; Yarnitsky, Granot, Nahman-Averbuch, Khamaisi, & Granovsky, 2012). Functional neuroimaging studies have begun to reveal a network of cortical and brainstem regions involved in the normal CPM response in humans (Bogdanov et al., 2015; Piche, Arsenault, & Rainville, 2009; Sprenger, Bingel, & Buchel, 2011; Youssef, Macefield, & Henderson, 2016a, 2016b) that is consistent with earlier studies of endogenous analgesia in animals, and recently we identified an aberrant pattern of PAG-to-brainstem connectivity that may play a role in deficient CPM, or enhanced descending facilitation, in central sensitization (Harper, Ichesco, Schrepf, Hampson, et al., 2018).

Whereas CPM methods probe for the absence or impairment of inhibitory mechanisms, *temporal summation* of pain measures the amplification of nociceptive signaling related to central sensitization. Experimentally, *temporal summation* refers to an increase in the perceived intensity of pain in response to sequential stimuli of equal physical strength. This

response is thought to reflect the progressive increase in neuronal firing of dorsal horn neurons in response to repetitive C-fiber stimulation (i.e., windup) (Graven-Nielsen et al., 2000; Price, Mao, Frenk, & Mayer, 1994; Price et al., 2002; Staud, Vierck, Cannon, Mauderli, & Price, 2001; Weissman-Fogel et al., 2009); however, supraspinal mechanisms contribute to the regulation of this phenomenon (Cheng, Erpelding, Kucyi, DeSouza, & Davis, 2015). Temporal summation is a normal response that occurs in healthy humans, but is enhanced in patients with central sensitization, and has been shown to predict pain outcomes (Arendt-Nielsen et al., 2010; Maixner, Fillingim, Sigurdsson, Kincaid, & Silva, 1998; Sarlani, Grace, Reynolds, & Greenspan, 2004; Staud, Craggs, Perlstein, Robinson, & Price, 2008; Weissman-Fogel et al., 2009). For example, patients with osteoarthritis that report chronic pain after joint replacement surgery have significantly higher temporal summation pre-operatively and post-operatively, compared to patients that experience pain relief (Izumi, Petersen, Laursen, Arendt-Nielsen, & Graven-Nielsen, 2017; Petersen, Arendt-Nielsen, Simonsen, Wilder-Smith, & Laursen, 2015; Skou et al., 2013). This suggests that enhanced temporal summation may be a marker for vulnerable individuals with central sensitization that respond poorly to peripherally-directed pain interventions. CPM and temporal summation paradigms can also be used in combination to better characterize the balance of pro- and anti-nociceptive processing within individuals (Petersen, Graven-Nielsen, Simonsen, Laursen, & Arendt-Nielsen, 2016; Vaegter & Graven-Nielsen, 2016; Yarnitsky, Granot, & Granovsky, 2014). Additional QST measures that may show altered endogenous pain modulation in patients with central sensitization include assessments of pain after-sensations (pain sensations that linger after cessation of painful stimulation) (Staud, Koo, Robinson, & Price, 2007), and offset-analgesia (a disproportionate reduction in perceived pain following a subtle decrease in the intensity of painful stimulation) (Hermans et al., 2016; Oudejans, Smit, van Velzen, Dahan, & Niesters, 2015).

In addition to hypersensitivity to painful stimuli, there is accumulating evidence that many patients with central sensitization exhibit hypersensitivity to non-painful stimuli that bypass peripheral and spinal pathways. Patients with FM demonstrate perceptual amplification of auditory stimuli, and rate everyday sounds as more unpleasant compared to controls (Carrillo-

de-la-Pena, Vallet, Perez, & Gomez-Perretta, 2006; Dohrenbusch, Sodhi, Lamprecht, & Genth, 1997; Geisser, Glass, et al., 2008; McDermid, Rollman, & McCain, 1996; Wilbarger & Cook, 2011). Similar findings have been observed in TMD (Hollins et al., 2009), rheumatoid arthritis (McDermid et al., 1996), and IBS (Berman et al., 2002; Blomhoff, Jacobsen, Spetalen, Dahm, & Malt, 2000). FM patients have also been shown to have increased sensitivity to complex visual stress (Harte et al., 2016), light (Martenson et al., 2016), and unpleasant odors (Schweinhardt, Sauro, & Bushnell, 2008). These findings, coupled with evidence that auditory and visual sensitivities are often correlated with pain sensitivity in these patients (Geisser, Glass, et al., 2008; Harte et al., 2016), have led us and others to hypothesize that a generalized or global state of CNS sensory amplification may play a role in the pathogenesis of chronic pain disorders, and that these sensory measures may mark an important individual patient endophenotype of central sensitization (Geisser, Donnell, et al., 2008; Hollins et al., 2009; McDermid et al., 1996). *In fact, we propose that this might be one fundamental neurobiological difference between chronic pain patients with “top-down” central pain mechanisms, where increased sensitivity to a range of painful and non-painful sensory stimuli is present, and those with “bottom-up” central pain mechanisms, traditionally referred to as central sensitization, where only pain processing might be augmented.* This is supported by the fact that functional imaging studies find that the insula, a brain region that also plays a role in multisensory integration, is hyperactive in most individuals with central sensitization (Brooks & Tracey, 2007; Wager et al., 2013).

Brain Imaging Studies. The observation that QST outcomes in individuals with FM showed diffuse tenderness has led to subsequent brain neuroimaging studies that have been amongst the best “objective” evidence that the pain in FM and related-pain syndromes are “real,” and that the brain is playing a prominent role in pain pathophysiology (Harris & Clauw, 2006).

Early studies demonstrated increased neuronal activity in CNS sensory-processing regions in patients with central sensitization exposed to sensory stimuli that healthy individuals find innocuous (Cook et al., 2004a; Gracely et al., 2004; Gracely, Petzke, Wolf, & Clauw, 2002a; Hampson et al., 2013; Naliboff et al., 2001). More recent neuroimaging studies provide further

insight into the underlying CNS changes in central sensitization, and demonstrate significant structural, chemical, and functional alterations in pain-related brain areas. As a broad overview, changes in brain-gray matter are noted in pain-processing regions such as: the thalamus; periaqueductal gray (PAG); insula; cingulate and somatosensory cortices (Smallwood et al., 2013); neurochemical alterations, including increased levels of excitatory neurotransmitters (e.g., glutamate) and decreased levels of inhibitory neurotransmitters (e.g., γ -aminobutyric acid [GABA] are present in the insula (Harris & Clauw, 2012); and increased resting brain network connectivity to pro-nociceptive brain areas; and decreased connectivity to anti-nociceptive brain areas are also often present in central sensitization (Ichesco et al., 2014; K. B. Jensen et al., 2009; Loggia et al., 2013; Napadow, Kim, Clauw, & Harris, 2012). As such, multiple brain neuroimaging studies have found alterations in supraspinal pain processing in individuals with clinical features suggestive of central sensitization.

Functional magnetic resonance imaging (fMRI), in conjunction with QST, has been used to further demonstrate that patients with central sensitization exhibit a supraspinal-mediated multisensory hypersensitivity. We recently found that the insula was activated by an aversive visual stimulus, and that the degree of activation was strongly associated with patients' self-reported FM pain (Harte et al., 2016). Moreover, this activity within the insula was reduced following administration of pregabalin (but not placebo), and the analgesic effectiveness of this drug was associated with a concomitant reduction in the insula's response to the visual stimulus. In addition to sensitivity to visual stimuli, methods such as fMRI clearly demonstrate that individuals with FM display increased brain activity when given a mild pressure or heat stimulus that most individuals would feel as "touch" or "warmth," rather than "pain". During this experience of pain, similar brain activation patterns arise in the patients that are typically seen in pain-processing regions that normal pain free individuals experience during a noxious stimulus (Cook et al., 2004b; Gracely, Petzke, Wolf, & Clauw, 2002b). This indicates that individuals with central sensitization have a similar brain response to painful stimuli as controls, but at lower levels of stimulus intensity.

A more recent advance in fMRI research is to examine the extent that brain regions are “connected” to each other (i.e., simultaneously activated [or deactivated] (Ploner, Lee, Wiech, Bingel, & Tracey, 2010), simply at rest. The advantage of these types of resting-state analyses is that they provide a window into brain changes associated with chronic, ongoing spontaneous pain. Individuals with FM have increased connectivity between the insula and neural networks not normally thought to be involved with pain, such as the *Default Mode Network* (DMN), and the degree of this hyper-connectedness is related to the severity of ongoing pain (Napadow et al., 2012; Napadow et al., 2010). A different group has shown that, during a painful stimulus, connectivity is decreased between key anti-nociceptive regions (e.g., the brainstem – the origin of descending analgesic pathways) and a region they previously identified to be a potential source of dysfunctional pain inhibition in FM (K. B. Jensen et al., 2009; K. B. Jensen et al., 2012). Similar to the studies of visual stimulation described above, Lopez-Sola and colleagues demonstrated that individuals with central sensitization are more sensitive to a number of non-painful sensory stimuli, and that machine-learning paradigms can accurately distinguish FM from non-FM patients with over 90% accuracy using fMRI results (Lopez-Sola et al., 2014; Lopez-Sola et al., 2017).

Other neuroimaging techniques have been used to identify neurotransmitter and receptor abnormalities that may be driving the pain amplification seen in FM and other central sensitization pain disorders. Wood and colleagues used positron emission tomography (PET) to show that attenuated dopaminergic activity may be playing a role in pain transmission in FM, and we showed evidence of decreased *mu* opioid receptor availability (possibly due to an increased release of endogenous *mu* opioids) in FM (Harris et al., 2007; Wood et al., 2007). Recent work by Schrepf and colleagues used multi-modal fMRI-PET to demonstrate that this reduction in opioid receptor-binding is related to reduced antinociceptive brain responses in the anterior cingulate cortex – providing further support for a potential endogenous opioid mechanism in pain central sensitization (Schrepf et al., 2016). This latter finding, as well as previous studies showing increased levels of endogenous opioids in the cerebrospinal fluid (CSF) of FM patients (Baraniuk, Whalen, Cunningham, & Clauw, 2004), have been suggested as

evidence of why opioid analgesics appear to have poor efficacy in FM.

Our group and others have used proton spectroscopy ($^1\text{H-MRS}$) to probe other neurotransmitters, including glutamate and GABA. These studies have shown increases in brain concentrations of glutamate in pain-processing regions, such as the insula in FM (Harris, 2010; Harris et al., 2009; Harte, Clauw, Napadow, & Harris, 2013; Pyke, Osmotherly, & Baines, 2017). This finding has also been noted in the CFS of FM patients (Sarchielli, Di Filippo, Nardi, & Calabresi, 2007). Drugs that improve FM pain, such as pregabalin and gabapentin, are hypothesized to be working in part by reducing glutamatergic activity in the brain (Maneuf, Hughes, & McKnight, 2001). In support of this, we demonstrated that baseline glutamate levels and functional connectivity of the insula predicted which FM patients would successfully respond to pregabalin therapy (Harris et al., 2013). Patients with higher posterior insular glutamate and insular connectivity to the DMN had the most improvement in pain and hyperalgesia following pregabalin. Furthermore, improvement in pain was accompanied by normalization of the glutamate and functional connectivity findings, demonstrating that these abnormalities were at least partly responsible for pain pathogenesis (Harris et al., 2013). We have similarly shown that *decreased connectivity to antinociceptive* brain regions, such as the periaqueductal gray (PAG – a critical locus of descending analgesic pathways), predicts responsiveness to milnacipran, a serotonin norepinephrine reuptake inhibitor (Schmidt-Wilcke et al., 2014).

Conversely, $^1\text{H-MRS}$ has been used to demonstrate low levels of one of the body's major inhibitory neurotransmitters (GABA; Foerster et al., 2012; Harper, Ichesco, Schrepf, Halvorson, et al., 2018), and this likely accounts for the efficacy of medications such as gamma-hydroxybutyrate in FM (Russell et al., 2011). This finding may also suggest biological plausibility for the association between low-moderate alcohol consumption (compared to none or heavy drinking), and improved symptomatology and functionality in FM (C. H. Kim et al., 2013; Scott et al., 2018). Alcohol enhances inhibitory GABAergic neurotransmission (Chandler, Overton, Ruedi-Bettschen, & Platt, 2017; Chastain, 2006) and may exhibit a U-shaped curve for its

analgesic effects similar to its potentially beneficial cardiovascular effects. That imbalances between excitatory and inhibitory neurotransmitters are not found diffusely throughout the brain and, instead, appear to be localized to regions that contribute to multisensory processing (e.g., insula) (Brooks & Tracey, 2007; Craig, 2002), is consistent with the notion that a global sensory hyper-responsiveness is partly responsible for the pathophysiology of FM and related *COPCs* (K. Ablin & Clauw, 2009; Yunus, 2007). **Figure 1** illustrates the neurotransmitters that have been demonstrated to influence pain transmission in the CNS. This neurochemical profile helps illustrate why no single class of CNS analgesic is likely to work in every patient with pain of CNS origin.

INSERT FIGURE 1 ABOUT HERE

fMRI has also proven useful in determining how comorbid psychosocial factors influence pain processing in FM. For example, in FM patients with variable degrees of comorbid depression, anterior insula and amygdala activations were correlated with depressive symptoms, consistent with these “medial” and pre-frontal brain regions being involved with affective or motivational aspects of pain processing (and being more closely related to unpleasantness, rather than the sensory intensity of pain) (Berna et al., 2010). However, the degree of neuronal activation in more lateral structures generally thought to be associated with the sensory-processing of pain (i.e., location and intensity of the pain) were not associated with levels of depressive symptoms, or the presence or absence of major depression. This is consistent with substantial evidence that pain and depression are largely independent, but overlapping, physiological processes.

The role of neuroendocrine or autonomic abnormalities. Because of the link between *COPCs* and exposure to stress, and because both the neuroendocrine and autonomic nervous systems could cause many of the symptoms of these conditions, these factors have been fairly extensively studied, especially in FM and CFS (Clauw & Crofford, 2003; Crofford, 1998; Demitrack & Crofford, 1998). However, this research has generally yielded inconsistent

findings, and treatment studies targeting these systems have failed (Adler, Kinsley, Hurwitz, Mossey, & Goldenberg, 1999; Cohen et al., 2000; Crofford et al., 1994; Demitrack & Crofford, 1998; Martinez-Lavin, Hermosillo, Rosas, & Soto, 1998; Qiao, Vaeroy, & Morkrid, 1991). These studies note either hypo- or hyper-activity of both the hypothalamic pituitary adrenal axis (HPA) and sympathetic nervous system in individuals with FM and related conditions, and the precise abnormality varies from study to study. Moreover, these studies only find “abnormal” HPA or autonomic function in a very small percentage of patients, and there is tremendous overlap between cases and controls in any of these studies. Taken together, these factors are now generally thought to play some role in a subset of individuals, but not in all individuals with these conditions.

The inconsistency of these findings may be explained by the fact that nearly all of these studies were cross-sectional and assumed that, if HPA and/or autonomic dysfunction was found in these conditions, it must have *caused* the pain and other symptoms. Data now suggest the opposite. As noted above, more convincing data suggest that these factors (especially HPA abnormalities) might represent a *diathesis* or be *due to* pain or early life stress, rather than causing it. Notably, in two studies examining HPA function in FM, McLean and colleagues showed that salivary cortisol levels covaried with pain levels, and that CSF levels of corticotropin-releasing factor (CRH) were more closely related to an individual’s pain level or a history of early-life trauma than whether they were a FM patient or control (McLean et al., 2005; McLean et al., 2006). Because most previous studies of HPA and autonomic function in FM failed to control for pain levels, a previous history of trauma, or other comorbid disorders that could affect HPA or autonomic dysfunction, these inconsistencies are not surprising.

Evidence of abnormal cytokines and immune dysfunction in central sensitization. Although the prevailing view is that FM and related *COPCs* are not autoimmune disorders, and that classic anti-inflammatory agents are not of benefit in these conditions, there are some data suggesting that the immune system may be playing a role in their pathogenesis (Gur &

Oktayoglu, 2008). Perhaps the most consistent finding noted to date is a mild elevation in IL-8, which is a cytokine associated with sympathetic function (Bazzichi et al., 2007; Wallace, Bowman, Wormsley, & Peter, 1989). In other conditions closely linked to FM, such as interstitial cystitis, more sensitive assays of immune system function, that can be gleaned by stimulating peripheral immune cells, have been shown to be abnormal (Schrepf et al., 2015). We and others have speculated that diet or obesity could contribute to this low-grade inflammation in FM, and might be a potential target for therapy (Schrepf et al., 2017); while others have posited that this may provide evidence of microglia involvement in FM. These areas warrant further exploration, and are actively being investigated.

The role of “small fiber neuropathy” in central sensitization. Many groups have now shown evidence of decreased intra-epidermal nerve-fiber density (i.e., small fiber neuropathy) in FM (Caro & Winter, 2015; Doppler, Rittner, Deckart, & Sommer, 2015; S. H. Kim, Kim, Oh, & Clauw, 2008; Levine & Saperstein, 2015; Oaklander, Herzog, Downs, & Klein, 2013; Uceyler et al., 2013). However, debate remains regarding the meaning of these findings (Clauw, 2015). Reduced nerve-fiber density is a very non-specific finding that has now been noted in over 50 different pain and non-pain conditions (Clauw, 2015). Moreover, Harte and colleagues recently demonstrated that these findings could be induced in an animal model of central sensitization by increasing insular glutamate (Harte et al., 2017). We believe that this non-specific finding reflects an adaptive structural and functional reorganization of the peripheral nervous system (not unlike the changes in the CNS noted via voxel-based morphometry of the brain in chronic pain states (Smallwood et al., 2013), in the context of ongoing chronic pain and other neurological conditions. The hyper-excitability of C nociceptors noted in FM by Serra and colleagues could also be secondary to CNS disinhibition of peripheral systems, rather than indicating that there is primary pathology in the periphery (Serra et al., 2014). Regardless, the current evidence does not merit routine skin biopsies in suspected FM patients, and certainly does not yet suggest that this finding is of enough pathophysiological significance to support the (non-investigational) use of aggressive cytotoxic and anti-inflammatory regimens that some have proposed. When considering the available data regarding the pathophysiology of FM, including

the brain neuroimaging and multi-sensory hypersensitivity findings discussed above, the data strongly favor CNS factors rather than peripheral abnormalities, and these findings should continue to drive clinical care.

THE CONCEPT OF CENTRAL SENSITIZATION AS A CONTINUUM – RATHER THAN PRESENT OR ABSENT

Wolfe was the first to describe the concept of “*fibromyalgiansess*,” by showing the importance of “sub-syndromic” or “sub-threshold” FM (Wolfe, 2009). In a series of studies, he showed that, in individuals with conditions such as rheumatoid arthritis, low back pain or osteoarthritis, an individual’s FM score, derived with measures very similar to the 2010/11/16 American College of Rheumatology (ACR) FM criteria (Wolfe et al., 2016; Wolfe et al., 2011; Wolfe, Clauw, et al., 2010), was typically more predictive of pain and disability than more objective measures of activity of these illnesses, such as measures of inflammation or joint damage (Wolfe et al., 2014; Wolfe, Michaud, Li, & Katz, 2010). The survey version of these criteria, designed for clinical and epidemiological research, is entirely patient self-report, and can be administered on a single piece of paper (Wolfe et al., 2011). Our group uses the *Michigan Body Map* (C. M. Brummett et al., 2016) to assess the *Widespread Pain Index* (up to 19 body areas each counted as 1-point), as well as the *Symptom Severity Index* that queries the presence and severity of fatigue, sleep disturbances, memory difficulties (each scored 0-3 for presence and severity), as well as irritable bowel, headaches and mood problems (1-point each; total Symptom Severity Index Score = 0-12; **Figure 2**). The *Widespread Pain Index* and *Symptom Severity Index* are combined for a total FM score of 0-31. When employed as a dichotomous measure with a variety of “cut points” that can be used, this measure will roughly identify most of the same individuals as the original tender point-based 1990 FM criteria (except with many more males) (Wolfe et al., 2011; Wolfe, Clauw, et al., 2010). However, when FM is considered more as a physiological construct to determine where on the continuum of central sensitization an individual is, then this score can be used as a continuous measure (i.e., reflecting the degree of *fibromyalgiansess* or central sensitization) that can be useful in the diagnosis and treatment of virtually any patient with a rheumatic disorder that is experiencing pain (Wolfe, 2009).

INSERT FIGURE 2 ABOUT HERE

The FM score was recently shown to be associated with diffuse hyperalgesia on QST in females with knee osteoarthritis (Neville et al., In Press), and an index of widespread pain was used to identify aberrant brain structure and connectivity in pelvic pain patients with interstitial cystitis that was indistinguishable from women with FM (Kutch et al., 2017). Furthermore, a recent fMRI study by Basu and colleagues has shown that perhaps the most consistently-found feature of central sensitization in COPCs - increased connectivity between the DMN and the insula - was also found to correlate strongly with the degree of FM that is present in individuals with rheumatoid arthritis (Basu et al., 2018). Importantly, these data support the notion that central sensitization pathology exists in non-FM pain states, and that tools such as the FM survey criteria provide a useful self-report measure of this construct.

In several recent studies, this concept of *fibromyalginess* or sub-syndromal FM has been shown to be very clinically important. In these studies by Brummett and colleagues, individuals who were scheduled for either lower extremity joint replacement or hysterectomy completed a broad-battery of validated self-report measures prior to their surgery. It was hypothesized that higher FM scores on the 2011 FM survey criteria (Wolfe et al., 2011) would predict increased opioid requirements in the inpatient admission following surgery, as well as long-term postsurgical pain outcomes. Again, this measure is scored from 0-31, with a score of 13 typically used as the diagnostic cut-off point for FM. These studies demonstrated that, for each 1-point increase in this measure from 0 – 31, individuals needed an adjusted 7 – 9 mg more oral morphine equivalents to control their pain in the first 24-48 hours following surgery, and were 15-20% less likely to show pain improvement following surgery (Brummett et al., 2013; Brummett et al., 2015; Janda et al., 2015). These findings were independent of a number of preoperative characteristics, including age, sex, anxiety, depression, catastrophizing, and opioid use. More importantly, these findings were linear, and the same incremental increase in opioid

requirements and surgery non-responsiveness was observed both in individuals well below the threshold to diagnose FM, and in individuals exceeding this threshold. **Figure 3** shows two different individuals with osteoarthritis, neither of whom would meet criteria for FM, but who are at different points along the *fibromyalgiansess* continuum, and shows the marked difference in opioid responsiveness and improvement in pain following arthroplasty that these two individuals would experience. These data suggest that the degree of central sensitization reflected in this measure may help identify individuals in perioperative or other settings who are less likely to respond to peripherally-directed treatments such as surgery.

INSERT FIGURE 3 ABOUT HERE

SUMMARY AND CONCLUSION

As the use of the term central sensitization has expanded over the past several decades, so has our understanding of CNS processes that may lead to this phenomenon. Initially thought to be confined to the spinal cord, supraspinal mechanisms are now understood to play a significant role in many individuals who have this phenomenon present. One demarcation that is becoming increasingly clear, but has received little attention in the pain field, is how to differentiate whether an individual with this process has a problem that is being driven by ongoing nociceptive input (i.e., the original use of the term), or whether this process is occurring in the absence of a peripheral driver (as seems to be occurring in many individuals with *COPCs*). **Table 1** denotes these two sub-types of central sensitization as *bottom-up* (i.e., being driven by ongoing nociceptive input, and presumably found in individuals with autoimmune disorders, sickle cell disease, hypermobility syndromes), or *top-down* (i.e., where the primary problem may very well likely originate in supraspinal structures and does not require ongoing nociceptive input to maintain the process). The Table reviews what we now know about these two sub-types, but this is an area that requires much more investigation. The clinical reason for differentiating these two sub-types of altered CNS processing of pain should be obvious – the *bottom-up* form should be treated aggressively with interventions

aimed at reducing peripheral nociceptive drive, whereas CNS-directed therapies will be necessary for the *top-down* form. As our understanding of these two subtypes evolves, so should our semantic terms, and these authors suggest that the term central sensitization is only used to denote the *bottom-up* form, and a new term (e.g., *central hypersensitivity*) be used to denote the *top-down* form. We further propose that, taken together, both forms constitute a broader clinical construct of “centralized pain” that exists as a continuum across individuals with chronic pain. Many individuals with centralized pain are likely to have combinations of these two processes, and will need to be treated with both peripherally- and centrally-directed therapies.

INSERT TABLE 1 ABOUT HERE

REFERENCES

- Aaron, L. A., & Buchwald, D. (2001). A review of the evidence for overlap among unexplained clinical conditions. *Annals of Internal Medicine*, *134*(9), 868-881. doi:10.7326/0003-4819-134-9_Part_2-200105011-00011
- Ablin, J. N., Clauw, D. J., Lyden, A. K., Ambrose, K., Williams, D. A., Gracely, R. H., & Glass, J. M. (2013). Effects of sleep restriction and exercise deprivation on somatic symptoms and mood in healthy adults. *Clin Exp Rheumatol*, *31*(6 Suppl 79), S53-59.
- Ablin, K., & Clauw, D. J. (2009). From fibrositis to functional somatic syndromes to a bell-shaped curve of pain and sensory sensitivity: evolution of a clinical construct. *Rheum Dis Clin North Am*, *35*(2), 233-251. doi:10.1016/j.rdc.2009.06.006
- Adler, G. K., Kinsley, B. T., Hurwitz, S., Mossey, C. J., & Goldenberg, D. L. (1999). Reduced hypothalamic-pituitary and sympathoadrenal responses to hypoglycemia in women with fibromyalgia syndrome. *Am J Med*, *106*(5), 534-543.

- Anderberg, U. M. (2000). Comment on: Johns and Littlejohn, The role of sex hormones in pain response. *Pain*, *87*(1), 109-111.
- Arendt-Nielsen, L. (2015). Central sensitization in humans: assessment and pharmacology. *Handb Exp Pharmacol*, *227*, 79-102. doi:10.1007/978-3-662-46450-2_5
- Arendt-Nielsen, L., Nie, H., Laursen, M. B., Laursen, B. S., Madeleine, P., Simonsen, O. H., & Graven-Nielsen, T. (2010). Sensitization in patients with painful knee osteoarthritis. *Pain*, *149*(3), 573-581. doi:10.1016/j.pain.2010.04.003
- Arendt-Nielsen, L., & Yarnitsky, D. (2009). Experimental and clinical applications of quantitative sensory testing applied to skin, muscles and viscera. *J Pain*, *10*(6), 556-572. doi:10.1016/j.jpain.2009.02.002
- Arnold, L. M., Fan, J., Russell, I. J., Yunus, M. B., Khan, M. A., Kushner, I., . . . Iyengar, S. K. (2013). The fibromyalgia family study: a genome-wide linkage scan study. *Arthritis Rheum*, *65*(4), 1122-1128. doi:10.1002/art.37842
- Arnold, L. M., Hudson, J. I., Hess, E. V., Ware, A. E., Fritz, D. A., Auchenbach, M. B., . . . Keck, P. E., Jr. (2004). Family study of fibromyalgia. *Arthritis Rheum*, *50*(3), 944-952. doi:10.1002/art.20042
- Arnson, Y., Amital, D., Fostick, L., Silberman, A., Polliack, M. L., Zohar, J., . . . Amital, H. (2007). Physical activity protects male patients with post-traumatic stress disorder from developing severe fibromyalgia. *Clin Exp Rheumatol*, *25*(4), 529-533.
- As-Sanie, S., Harris, R. E., Harte, S. E., Tu, F. F., Neshewat, G., & Clauw, D. J. (2013). Increased pressure pain sensitivity in women with chronic pelvic pain. *Obstet Gynecol*, *122*(5), 1047-1055. doi:10.1097/AOG.0b013e3182a7e1f5
- Backonja, M. M., Attal, N., Baron, R., Bouhassira, D., Drangholt, M., Dyck, P. J., . . . Ziegler, D. (2013). Value of quantitative sensory testing in neurological and pain disorders: NeuPSIG consensus. *Pain*, *154*(9), 1807-1819. doi:10.1016/j.pain.2013.05.047

- Bagarinao, E., Johnson, K. A., Martucci, K. T., Ichesco, E., Farmer, M. A., Labus, J., . . . Mackey, S. (2014). Preliminary structural MRI based brain classification of chronic pelvic pain: A MAPP network study. *Pain, 155*(12), 2502-2509. doi:10.1016/j.pain.2014.09.002
- Bair, E., Brownstein, N. C., Ohrbach, R., Greenspan, J. D., Dubner, R., Fillingim, R. B., . . . Slade, G. D. (2013). Study protocol, sample characteristics, and loss to follow-up: the OPPERA prospective cohort study. *J Pain, 14*(12 Suppl), T2-19. doi:10.1016/j.jpain.2013.06.006
- Bair, E., Ohrbach, R., Fillingim, R. B., Greenspan, J. D., Dubner, R., Diatchenko, L., . . . Slade, G. D. (2013). Multivariable modeling of phenotypic risk factors for first-onset TMD: the OPPERA prospective cohort study. *J Pain, 14*(12 Suppl), T102-115. doi:10.1016/j.jpain.2013.09.003
- Baraniuk, J. N., Whalen, G., Cunningham, J., & Clauw, D. J. (2004). Cerebrospinal fluid levels of opioid peptides in fibromyalgia and chronic low back pain. *BMC Musculoskelet Disord, 5*, 48. doi:10.1186/1471-2474-5-48
- Basu, N., Kaplan, C. M., Ichesco, E., Larkin, T., Harris, R. E., Murray, A., . . . Clauw, D. J. (2018). Neurobiological features of fibromyalgia are also present among rheumatoid arthritis patients. *Arthritis Rheumatol.* doi:10.1002/art.40451
- Bazzichi, L., Rossi, A., Massimetti, G., Giannaccini, G., Giuliano, T., De Feo, F., . . . Bombardieri, S. (2007). Cytokine patterns in fibromyalgia and their correlation with clinical manifestations. *Clinical and Experimental Rheumatology, 25*(2), 225-230.
- Berman, S. M., Naliboff, B. D., Chang, L., Fitzgerald, L., Antolin, T., Camplone, A., & Mayer, E. A. (2002). Enhanced preattentive central nervous system reactivity in irritable bowel syndrome. *Am J Gastroenterol, 97*(11), 2791-2797. doi:10.1111/j.1572-0241.2002.07024.x
- Berna, C., Leknes, S., Holmes, E. A., Edwards, R. R., Goodwin, G. M., & Tracey, I. (2010). Induction of depressed mood disrupts emotion regulation neurocircuitry and enhances

pain unpleasantness. *Biological Psychiatry*, 67(11), 1083-1090.

doi:10.1016/j.biopsych.2010.01.014

Blomhoff, S., Jacobsen, M. B., Spetalen, S., Dahm, A., & Malt, U. F. (2000). Perceptual hyperreactivity to auditory stimuli in patients with irritable bowel syndrome. *Scand J Gastroenterol*, 35(6), 583-589.

Bogdanov, V. B., Vigano, A., Noirhomme, Q., Bogdanova, O. V., Guy, N., Laureys, S., . . . Schoenen, J. (2015). Cerebral responses and role of the prefrontal cortex in conditioned pain modulation: an fMRI study in healthy subjects. *Behavioural brain research*, 281, 187-198. doi:10.1016/j.bbr.2014.11.028

Brooks, J. C., & Tracey, I. (2007). The insula: a multidimensional integration site for pain. *Pain*, 128(1-2), 1-2. doi:10.1016/j.pain.2006.12.025

Brummett, C., Clauw, D., Harris, R., Harte, S., Hassett, A., & Williams, D. (2016). We agree with the need for a new term but disagree with the proposed terms. *Pain*, 157(12), 2876. doi:10.1097/j.pain.0000000000000717

Brummett, C. M., Bakshi, R. R., Goesling, J., Leung, D., Moser, S. E., Zollars, J. W., . . . Hassett, A. L. (2016). Preliminary validation of the Michigan Body Map. *Pain*, 157(6), 1205-1212. doi:10.1097/j.pain.0000000000000506

Brummett, C. M., Janda, A. M., Schueller, C. M., Tsodikov, A., Morris, M., Williams, D. A., & Clauw, D. J. (2013). Survey criteria for fibromyalgia independently predict increased postoperative opioid consumption after lower-extremity joint arthroplasty: a prospective, observational cohort study. *Anesthesiology*, 119(6), 1434-1443. doi:10.1097/ALN.0b013e3182a8eb1f

Brummett, C. M., Urquhart, A. G., Hassett, A. L., Tsodikov, A., Hallstrom, B. R., Wood, N. I., . . . Clauw, D. J. (2015). Characteristics of fibromyalgia independently predict poorer long-term analgesic outcomes following total knee and hip arthroplasty. *Arthritis Rheumatol*, 67(5), 1386-1394. doi:10.1002/art.39051

- Buskila, D., Atzeni, F., & Sarzi-Puttini, P. (2008). Etiology of fibromyalgia: the possible role of infection and vaccination. *Autoimmun Rev*, *8*(1), 41-43.
doi:10.1016/j.autrev.2008.07.023
- Buskila, D., Neumann, L., Hazanov, I., & Carmi, R. (1996). Familial aggregation in the fibromyalgia syndrome. *Semin Arthritis Rheum*, *26*(3), 605-611.
- Buskila, D., Neumann, L., Vaisberg, G., Alkalay, D., & Wolfe, F. (1997). Increased rates of fibromyalgia following cervical spine injury. A controlled study of 161 cases of traumatic injury [see comments]. *Arthritis & Rheumatism*, *40*(3), 446-452.
- Buskila, D., Sarzi-Puttini, P., & Ablin, J. N. (2007). The genetics of fibromyalgia syndrome. *Pharmacogenomics*, *8*(1), 67-74. doi:10.2217/14622416.8.1.67
- Buskila, D., Shnaider, A., Neumann, L., Zilberman, D., Hilzenrat, N., & Sikuler, E. (1997). Fibromyalgia in hepatitis C virus infection - Another infectious disease relationship. *Archives of Internal Medicine*, *157*(21), 2497-2500. doi:DOI 10.1001/archinte.157.21.2497
- Caro, X. J., & Winter, E. F. (2015). The Role and Importance of Small Fiber Neuropathy in Fibromyalgia Pain. *Curr Pain Headache Rep*, *19*(12), 55. doi:10.1007/s11916-015-0527-7
- Carrillo-de-la-Pena, M. T., Vallet, M., Perez, M. I., & Gomez-Perretta, C. (2006). Intensity dependence of auditory-evoked cortical potentials in fibromyalgia patients: a test of the generalized hypervigilance hypothesis. *J Pain*, *7*(7), 480-487.
doi:10.1016/j.jpain.2006.01.452
- Chandler, C. M., Overton, J. S., Ruedi-Bettschen, D., & Platt, D. M. (2017). GABAA Receptor Subtype Mechanisms and the Abuse-Related Effects of Ethanol: Genetic and Pharmacological Evidence. *Handb Exp Pharmacol*. doi:10.1007/164_2017_80
- Chastain, G. (2006). Alcohol, neurotransmitter systems, and behavior. *J Gen Psychol*, *133*(4), 329-335. doi:10.3200/GENP.133.4.329-335

- Cheng, J. C., Erpelding, N., Kucyi, A., DeSouza, D. D., & Davis, K. D. (2015). Individual Differences in Temporal Summation of Pain Reflect Pronociceptive and Antinociceptive Brain Structure and Function. *J Neurosci*, *35*(26), 9689-9700. doi:10.1523/JNEUROSCI.5039-14.2015
- Ciampi de Andrade, D., Maschietto, M., Galhardoni, R., Gouveia, G., Chile, T., Victorino Krepischi, A. C., . . . Brentani, H. P. (2017). Epigenetics insights into chronic pain: DNA hypomethylation in fibromyalgia-a controlled pilot-study. *Pain*, *158*(8), 1473-1480. doi:10.1097/j.pain.0000000000000932
- Clauw, D. J. (1995a). Fibromyalgia: more than just a musculoskeletal disease. *Am Fam Physician*, *52*(3), 843-851, 853-844.
- Clauw, D. J. (1995b). The pathogenesis of chronic pain and fatigue syndromes, with special reference to fibromyalgia. *Med Hypotheses*, *44*(5), 369-378.
- Clauw, D. J. (2015). What is the meaning of "small fiber neuropathy" in fibromyalgia? *Pain*. doi:10.1097/j.pain.0000000000000311
- Clauw, D. J., & Chrousos, G. P. (1997). Chronic pain and fatigue syndromes: overlapping clinical and neuroendocrine features and potential pathogenic mechanisms. *Neuroimmunomodulation*, *4*(3), 134-153. doi:10.1159/000097332
- Clauw, D. J., & Crofford, L. J. (2003). Chronic widespread pain and fibromyalgia: what we know, and what we need to know. *Best Practice & Research in Clinical Rheumatology*, *17*(4), 685-701. doi:10.1016/S1521-6942(03)00035-4
- Clauw, D. J., Schmidt, M., Radulovic, D., Singer, A., Katz, P., & Bresette, J. (1997). The relationship between fibromyalgia and interstitial cystitis. *Journal of Psychiatric Research*, *31*(1), 125-131. doi:Doi 10.1016/S0022-3956(96)00051-9
- Clauw, D. J., & Witter, J. (2009). Pain and rheumatology: thinking outside the joint. *Arthritis Rheum*, *60*(2), 321-324. doi:10.1002/art.24326

- Clemens, J. Q., Mullins, C., Kusek, J. W., Kirkali, Z., Mayer, E. A., Rodriguez, L. V., . . . Group, M. R. N. S. (2014). The MAPP research network: a novel study of urologic chronic pelvic pain syndromes. *BMC Urol*, *14*, 57. doi:10.1186/1471-2490-14-57
- Cohen, H., Neumann, L., Shore, M., Amir, M., Cassuto, Y., & Buskila, D. (2000). Autonomic dysfunction in patients with fibromyalgia: Application of power spectral analysis of heart rate variability [see comments]. *Seminars in Arthritis and Rheumatism*, *29*(4), 217-227.
- Cook, D. B., Lange, G., Ciccone, D. S., Liu, W. C., Steffener, J., & Natelson, B. H. (2004a). Functional imaging of pain in patients with primary fibromyalgia. *J Rheumatol*, *31*(2), 364-378. doi:0315162X-31-364 [pii]
- Cook, D. B., Lange, G., Ciccone, D. S., Liu, W. C., Steffener, J., & Natelson, B. H. (2004b). Functional imaging of pain in patients with primary fibromyalgia. *J Rheumatol.*, *31*(2), 364-378.
- Craig, A. D. (2002). How do you feel? Interoception: the sense of the physiological condition of the body. *Nat.Rev.Neurosci.*, *3*(8), 655-666.
- Crofford, L. J. (1998). The hypothalamic-pituitary-adrenal stress axis in fibromyalgia and chronic fatigue syndrome. *Z Rheumatol*, *57 Suppl 2*, 67-71.
- Crofford, L. J., Pillemer, S. R., Kalogeras, K. T., Cash, J. M., Michelson, D., Kling, M. A., . . . Wilder, R. L. (1994). Hypothalamic-pituitary-adrenal axis perturbations in patients with fibromyalgia. *Arthritis Rheum*, *37*(11), 1583-1592.
- Crook, J., Moldofsky, H., & Shannon, H. (1998). Determinants of disability after a work related musculetal injury [see comments]. *Journal of Rheumatology*, *25*(8), 1570-1577.
- Cruz-Almeida, Y., & Fillingim, R. B. (2014). Can quantitative sensory testing move us closer to mechanism-based pain management? *Pain Med*, *15*(1), 61-72. doi:10.1111/pme.12230

- Darbari, D. S., Ballas, S. K., & Clauw, D. J. (2014). Thinking beyond sickling to better understand pain in sickle cell disease. *Eur J Haematol*, *93*(2), 89-95. doi:10.1111/ejh.12340
- Demitrack, M. A., & Crofford, L. J. (1998). Evidence for and pathophysiologic implications of hypothalamic-pituitary-adrenal axis dysregulation in fibromyalgia and chronic fatigue syndrome. *Ann N Y Acad Sci*, *840*, 684-697.
- DeSantana, J. M., da Cruz, K. M., & Sluka, K. A. (2013). Animal models of fibromyalgia. *Arthritis Res Ther*, *15*(6), 222. doi:10.1186/ar4402
- Diatchenko, L., Fillingim, R. B., Smith, S. B., & Maixner, W. (2013). The phenotypic and genetic signatures of common musculoskeletal pain conditions. *Nat Rev Rheumatol*, *9*(6), 340-350. doi:10.1038/nrrheum.2013.43
- Diatchenko, L., Nackley, A. G., Slade, G. D., Bhalang, K., Belfer, I., Max, M. B., . . . Maixner, W. (2006). Catechol-O-methyltransferase gene polymorphisms are associated with multiple pain-evoking stimuli. *Pain*, *125*(3), 216-224. doi:10.1016/j.pain.2006.05.024
- Dohrenbusch, R., Sodhi, H., Lamprecht, J., & Genth, E. (1997). Fibromyalgia as a disorder of perceptual organization? An analysis of acoustic stimulus processing in patients with widespread pain. *Z Rheumatol*, *56*(6), 334-341.
- Doppler, K., Rittner, H. L., Deckart, M., & Sommer, C. (2015). Reduced dermal nerve fiber diameter in skin biopsies of patients with fibromyalgia. *Pain*, *156*(11), 2319-2325. doi:10.1097/j.pain.0000000000000285
- Dougherty, P. M., Palecek, J., Paleckova, V., Sorkin, L. S., & Willis, W. D. (1992). The role of NMDA and non-NMDA excitatory amino acid receptors in the excitation of primate spinothalamic tract neurons by mechanical, chemical, thermal, and electrical stimuli. *J Neurosci*, *12*(8), 3025-3041.
- Edwards, R. R. (2005). Individual differences in endogenous pain modulation as a risk factor for chronic pain. *Neurology*, *65*(3), 437-443. doi:10.1212/01.wnl.0000171862.17301.84

- Edwards, R. R., Sarlani, E., Wesselmann, U., & Fillingim, R. B. (2005). Quantitative assessment of experimental pain perception: multiple domains of clinical relevance. *Pain, 114*(3), 315-319. doi:10.1016/j.pain.2005.01.007
- Epstein, S. A., Kay, G., Clauw, D., Heaton, R., Klein, D., Krupp, L., . . . Zisook, S. (1999). Psychiatric disorders in patients with fibromyalgia. A multicenter investigation. *Psychosomatics, 40*(1), 57-63. doi:10.1016/S0033-3182(99)71272-7
- Farmer, M. A., Huang, L., Martucci, K., Yang, C. C., Maravilla, K. R., Harris, R. E., . . . Network, M. R. (2015). Brain White Matter Abnormalities in Female Interstitial Cystitis/Bladder Pain Syndrome: A MAPP Network Neuroimaging Study. *J Urol, 194*(1), 118-126. doi:10.1016/j.juro.2015.02.082
- Fillingim, R. B. (2005). Individual differences in pain responses. *Curr Rheumatol Rep, 7*(5), 342-347.
- Fillingim, R. B., Maixner, W., Kincaid, S., Sigurdsson, A., & Harris, M. B. (1996). Pain sensitivity in patients with temporomandibular disorders: relationship to clinical and psychosocial factors. *Clin J Pain, 12*(4), 260-269.
- Fillingim, R. B., Slade, G. D., Diatchenko, L., Dubner, R., Greenspan, J. D., Knott, C., . . . Maixner, W. (2011). Summary of findings from the OPPERA baseline case-control study: implications and future directions. *J Pain, 12*(11 Suppl), T102-107. doi:10.1016/j.jpain.2011.08.009
- Foerster, B. R., Petrou, M., Edden, R. A., Sundgren, P. C., Schmidt-Wilcke, T., Lowe, S. E., . . . Harris, R. E. (2012). Reduced insular gamma-aminobutyric acid in fibromyalgia. *Arthritis Rheum, 64*(2), 579-583. doi:10.1002/art.33339
- Fukuda, K., Dobbins, J. G., Wilson, L. J., Dunn, R. A., Wilcox, K., & Smallwood, D. (1997). An epidemiologic study of fatigue with relevance for the chronic fatigue syndrome. *J Psychiatr Res, 31*(1), 19-29.

- Fukuda, K., Nisenbaum, R., Stewart, G., Thompson, W. W., Robin, L., Washko, R. M., . . . Reeves, W. C. (1998). Chronic multisymptom illness affecting Air Force veterans of the Gulf War. *JAMA*, *280*(11), 981-988.
- Gebhart, G. F. (1986). Modulatory effects of descending systems on spinal dorsal horn neurons. In T. L. Yaksh (Ed.), *Spinal Afferent Processing* (pp. 391-416). New York: Plenum Press. (Reprinted from: NOT IN FILE).
- Geisser, M. E., Donnell, C. S., Petzke, F., Gracely, R. H., Clauw, D. J., & Williams, D. A. (2008). Comorbid somatic symptoms and functional status in patients with fibromyalgia and chronic fatigue syndrome: Sensory amplification as a common mechanism. *Psychosomatics*, *49*(3), 235-242. doi:DOI 10.1176/appi.psy.49.3.235
- Geisser, M. E., Glass, J. M., Rajcevska, L. D., Clauw, D. J., Williams, D. A., Kileny, P. R., & Gracely, R. H. (2008). A psychophysical study of auditory and pressure sensitivity in patients with fibromyalgia and healthy controls. *J Pain*, *9*(5), 417-422. doi:10.1016/j.jpain.2007.12.006
- Gibson, S. J., Littlejohn, G. O., Gorman, M. M., Helme, R. D., & Granges, G. (1994). Altered heat pain thresholds and cerebral event-related potentials following painful CO2 laser stimulation in subjects with fibromyalgia syndrome. *Pain*, *58*(2), 185-193.
- Giesecke, J., Reed, B. D., Haefner, H. K., Giesecke, T., Clauw, D. J., & Gracely, R. H. (2004). Quantitative sensory testing in vulvodynia patients and increased peripheral pressure pain sensitivity. *Obstet Gynecol*, *104*(1), 126-133. doi:10.1097/01.AOG.0000129238.49397.4e
- Gracely, R. H., Geisser, M. E., Giesecke, T., Grant, M. A., Petzke, F., Williams, D. A., & Clauw, D. J. (2004). Pain catastrophizing and neural responses to pain among persons with fibromyalgia. *Brain*, *127*(Pt 4), 835-843. doi:10.1093/brain/awh098
- Gracely, R. H., Lynch, S. A., & Bennett, G. J. (1992). Painful neuropathy: altered central processing maintained dynamically by peripheral input. *Pain*, *51*(2), 175-194.

- Gracely, R. H., Petzke, F., Wolf, J. M., & Clauw, D. J. (2002a). Functional magnetic resonance imaging evidence of augmented pain processing in fibromyalgia. *Arthritis Rheum*, *46*(5), 1333-1343. doi:10.1002/art.10225
- Gracely, R. H., Petzke, F., Wolf, J. M., & Clauw, D. J. (2002b). Functional magnetic resonance imaging evidence of augmented pain processing in fibromyalgia. *Arthritis Rheum.*, *46*(5), 1333-1343.
- Granges, G., & Littlejohn, G. (1993). Pressure pain threshold in pain-free subjects, in patients with chronic regional pain syndromes, and in patients with fibromyalgia syndrome. *Arthritis Rheum*, *36*(5), 642-646.
- Granot, M., Weissman-Fogel, I., Crispel, Y., Pud, D., Granovsky, Y., Sprecher, E., & Yarnitsky, D. (2008). Determinants of endogenous analgesia magnitude in a diffuse noxious inhibitory control (DNIC) paradigm: do conditioning stimulus painfulness, gender and personality variables matter? *Pain*, *136*(1-2), 142-149. doi:10.1016/j.pain.2007.06.029
- Graven-Nielsen, T., Aspegren Kendall, S., Henriksson, K. G., Bengtsson, M., Sorensen, J., Johnson, A., . . . Arendt-Nielsen, L. (2000). Ketamine reduces muscle pain, temporal summation, and referred pain in fibromyalgia patients. *Pain*, *85*(3), 483-491.
- Greenspan, J. D., Slade, G. D., Bair, E., Dubner, R., Fillingim, R. B., Ohrbach, R., . . . Maixner, W. (2013). Pain sensitivity and autonomic factors associated with development of TMD: the OPPERA prospective cohort study. *J Pain*, *14*(12 Suppl), T63-74 e61-66. doi:10.1016/j.jpain.2013.06.007
- Greenspan, J. D., Slade, G. D., Bair, E., Dubner, R., Fillingim, R. B., Ohrbach, R., . . . Maixner, W. (2011). Pain sensitivity risk factors for chronic TMD: descriptive data and empirically identified domains from the OPPERA case control study. *J Pain*, *12*(11 Suppl), T61-74. doi:10.1016/j.jpain.2011.08.006
- Grinberg, K., Granot, M., Lowenstein, L., Abramov, L., & Weissman-Fogel, I. (2017). A common pronociceptive pain modulation profile typifying subgroups of chronic pelvic pain

syndromes is interrelated with enhanced clinical pain. *Pain*, 158(6), 1021-1029.
doi:10.1097/j.pain.0000000000000869

Gur, A., & Oktayoglu, P. (2008). Status of immune mediators in fibromyalgia. *Curr Pain Headache Rep*, 12(3), 175-181.

Gwilym, S. E., Filippini, N., Douaud, G., Carr, A. J., & Tracey, I. (2010). Thalamic atrophy associated with painful osteoarthritis of the hip is reversible after arthroplasty: a longitudinal voxel-based morphometric study. *Arthritis Rheum*, 62(10), 2930-2940.
doi:10.1002/art.27585

Gwilym, S. E., Keltner, J. R., Warnaby, C. E., Carr, A. J., Chizh, B., Chessell, I., & Tracey, I. (2009). Psychophysical and functional imaging evidence supporting the presence of central sensitization in a cohort of osteoarthritis patients. *Arthritis Rheum*, 61(9), 1226-1234.
doi:10.1002/art.24837

Hampson, J. P., Reed, B. D., Clauw, D. J., Bhavsar, R., Gracely, R. H., Haefner, H. K., & Harris, R. E. (2013). Augmented central pain processing in vulvodynia. *J Pain*, 14(6), 579-589.
doi:10.1016/j.jpain.2013.01.767

Harper, D. E., Ichesco, E., Schrepf, A., Halvorson, M., Puiu, T., Clauw, D. J., . . . Network, M. R. (2018). Relationships between brain metabolite levels, functional connectivity, and negative mood in urologic chronic pelvic pain syndrome patients compared to controls: A MAPP research network study. *Neuroimage Clin*, 17, 570-578.
doi:10.1016/j.nicl.2017.11.014

Harper, D. E., Ichesco, E., Schrepf, A., Hampson, J. P., Clauw, D. J., Schmidt-Wilcke, T., . . . Harte, S. E. (2018). Resting Functional Connectivity of the Periaqueductal Gray Is Associated With Normal Inhibition and Pathological Facilitation in Conditioned Pain Modulation. *J Pain*. doi:10.1016/j.jpain.2018.01.001

Harris, R. E. (2010). Elevated excitatory neurotransmitter levels in the fibromyalgia brain. *Arthritis Res Ther*, 12(5), 141. doi:10.1186/ar3136

- Harris, R. E., & Clauw, D. J. (2006). How do we know that the pain in fibromyalgia is "real"? *Curr.Pain Headache Rep.*, *10*(6), 403-407.
- Harris, R. E., & Clauw, D. J. (2012). Imaging central neurochemical alterations in chronic pain with proton magnetic resonance spectroscopy. *Neurosci Lett*, *520*(2), 192-196. doi:10.1016/j.neulet.2012.03.042
- Harris, R. E., Clauw, D. J., Scott, D. J., McLean, S. A., Gracely, R. H., & Zubieta, J. K. (2007). Decreased central mu-opioid receptor availability in fibromyalgia. *J Neurosci*, *27*(37), 10000-10006. doi:10.1523/JNEUROSCI.2849-07.2007
- Harris, R. E., Napadow, V., Huggins, J. P., Pauer, L., Kim, J., Hampson, J., . . . Clauw, D. J. (2013). Pregabalin rectifies aberrant brain chemistry, connectivity, and functional response in chronic pain patients. *Anesthesiology*, *119*(6), 1453-1464. doi:10.1097/ALN.000000000000017
- Harris, R. E., Sundgren, P. C., Craig, A. D., Kirshenbaum, E., Sen, A., Napadow, V., & Clauw, D. J. (2009). Elevated insular glutamate in fibromyalgia is associated with experimental pain. *Arthritis Rheum*, *60*(10), 3146-3152. doi:10.1002/art.24849
- Harte, S. E., Clauw, D. J., Hayes, J. M., Feldman, E. L., St Charles, I. C., & Watson, C. J. (2017). Reduced intraepidermal nerve fiber density after a sustained increase in insular glutamate: a proof-of-concept study examining the pathogenesis of small fiber pathology in fibromyalgia. *Pain Rep*, *2*(3), e590. doi:10.1097/PR9.0000000000000590
- Harte, S. E., Clauw, D. J., Napadow, V., & Harris, R. E. (2013). Pressure Pain Sensitivity and Insular Combined Glutamate and Glutamine (Glx) Are Associated with Subsequent Clinical Response to Sham But Not Traditional Acupuncture in Patients Who Have Chronic Pain. *Med Acupunct*, *25*(2), 154-160. doi:10.1089/acu.2013.0965
- Harte, S. E., Ichesco, E., Hampson, J. P., Peltier, S. J., Schmidt-Wilcke, T., Clauw, D. J., & Harris, R. E. (2016). Pharmacologic attenuation of cross-modal sensory augmentation within the chronic pain insula. *Pain*, *157*(9), 1933-1945. doi:10.1097/j.pain.0000000000000593

- Henry, N. L., & Clauw, D. J. (2012). Thinking beyond the tumor to better understand chronic symptoms in breast cancer survivors. *Breast Cancer Res Treat*, *133*(2), 413-416. doi:10.1007/s10549-011-1804-8
- Hermans, L., Calders, P., Van Oosterwijck, J., Verschelde, E., Bertel, E., & Meeus, M. (2016). An Overview of Offset Analgesia and the Comparison with Conditioned Pain Modulation: A Systematic Literature Review. *Pain Physician*, *19*(6), 307-326.
- Hollins, M., Harper, D., Gallagher, S., Owings, E. W., Lim, P. F., Miller, V., . . . Maixner, W. (2009). Perceived intensity and unpleasantness of cutaneous and auditory stimuli: an evaluation of the generalized hypervigilance hypothesis. *Pain*, *141*(3), 215-221. doi:10.1016/j.pain.2008.10.003
- Hudson, J. I., Goldenberg, D. L., Pope, H. G., Jr., Keck, P. E., Jr., & Schlesinger, L. (1992). Comorbidity of fibromyalgia with medical and psychiatric disorders. *Am J Med*, *92*(4), 363-367.
- Hudson, J. I., Hudson, M. S., Pliner, L. F., Goldenberg, D. L., & Pope, H. G., Jr. (1985). Fibromyalgia and major affective disorder: a controlled phenomenology and family history study. *Am J Psychiatry*, *142*(4), 441-446. doi:10.1176/ajp.142.4.441
- Hudson, J. I., & Pope, H. G. (1994). The concept of affective spectrum disorder: relationship to fibromyalgia and other syndromes of chronic fatigue and chronic muscle pain. *Baillieres Clin Rheumatol*, *8*(4), 839-856.
- Ichesco, E., Schmidt-Wilcke, T., Bhavsar, R., Clauw, D. J., Peltier, S. J., Kim, J., . . . Harris, R. E. (2014). Altered resting state connectivity of the insular cortex in individuals with fibromyalgia. *J Pain*, *15*(8), 815-826 e811. doi:10.1016/j.jpain.2014.04.007
- Izumi, M., Petersen, K. K., Laursen, M. B., Arendt-Nielsen, L., & Graven-Nielsen, T. (2017). Facilitated temporal summation of pain correlates with clinical pain intensity after hip arthroplasty. *Pain*, *158*(2), 323-332. doi:10.1097/j.pain.0000000000000764

- Janda, A. M., As-Sanie, S., Rajala, B., Tsodikov, A., Moser, S. E., Clauw, D. J., & Brummett, C. M. (2015). Fibromyalgia survey criteria are associated with increased postoperative opioid consumption in women undergoing hysterectomy. *Anesthesiology*, *122*(5), 1103-1111. doi:10.1097/ALN.0000000000000637
- Jensen, K. B., Kosek, E., Petzke, F., Carville, S., Fransson, P., Marcus, H., . . . Ingvar, M. (2009). Evidence of dysfunctional pain inhibition in Fibromyalgia reflected in rACC during provoked pain. *Pain*, *144*(1-2), 95-100. doi:10.1016/j.pain.2009.03.018
- Jensen, K. B., Loitole, R., Kosek, E., Petzke, F., Carville, S., Fransson, P., . . . Kong, J. (2012). Patients with fibromyalgia display less functional connectivity in the brain's pain inhibitory network. *Mol Pain*, *8*, 32. doi:10.1186/1744-8069-8-32
- Jensen, R., Rasmussen, B. K., Pedersen, B., Lous, I., & Olesen, J. (1992). Cephalic muscle tenderness and pressure pain threshold in a general population. *Pain*, *48*(2), 197-203.
- Jensen, R., Rasmussen, B. K., Pedersen, B., & Olesen, J. (1993). Muscle tenderness and pressure pain thresholds in headache. A population study. *Pain*, *52*(2), 193-199.
- Jensen, T. S., & Yaksh, T. L. (1992). Brainstem excitatory amino acid receptors in nociception: microinjection mapping and pharmacological characterization of glutamate-sensitive sites in the brainstem associated with algogenic behavior. *Neuroscience*, *46*(3), 535-547.
- Julien, N., Goffaux, P., Arsenault, P., & Marchand, S. (2005). Widespread pain in fibromyalgia is related to a deficit of endogenous pain inhibition. *Pain*, *114*(1-2), 295-302. doi:10.1016/j.pain.2004.12.032
- Kato, K., Sullivan, P. F., Evengard, B., & Pedersen, N. L. (2006). Importance of genetic influences on chronic widespread pain. *Arthritis Rheum*, *54*(5), 1682-1686. doi:10.1002/art.21798
- Kato, K., Sullivan, P. F., Evengard, B., & Pedersen, N. L. (2009). A population-based twin study of functional somatic syndromes. *Psychol Med*, *39*(3), 497-505. doi:10.1017/S0033291708003784

- Kim, C. H., Vincent, A., Clauw, D. J., Luedtke, C. A., Thompson, J. M., Schneekloth, T. D., & Oh, T. H. (2013). Association between alcohol consumption and symptom severity and quality of life in patients with fibromyalgia. *Arthritis Res Ther*, *15*(2), R42. doi:10.1186/ar4200
- Kim, S. H., Kim, D. H., Oh, D. H., & Clauw, D. J. (2008). Characteristic electron microscopic findings in the skin of patients with fibromyalgia: preliminary study. *Clin Rheumatol*, *27*(2), 219-223. doi:10.1007/s10067-007-0739-2
- Kleinbohl, D., Holzl, R., Moltner, A., Rommel, C., Weber, C., & Osswald, P. M. (1999). Psychophysical measures of sensitization to tonic heat discriminate chronic pain patients. *Pain*, *81*(1-2), 35-43. doi:[http://dx.doi.org/10.1016/S0304-3959\(98\)00266-8](http://dx.doi.org/10.1016/S0304-3959(98)00266-8)
- Kosek, E., Cohen, M., Baron, R., Gebhart, G. F., Mico, J. A., Rice, A. S., . . . Sluka, A. K. (2016). Do we need a third mechanistic descriptor for chronic pain states? *Pain*, *157*(7), 1382-1386. doi:10.1097/j.pain.0000000000000507
- Kosek, E., Ekholm, J., & Hansson, P. (1995). Increased pressure pain sensibility in fibromyalgia patients is located deep to the skin but not restricted to muscle tissue. *Pain*, *63*(3), 335-339.
- Kosek, E., & Ordeberg, G. (2000). Lack of pressure pain modulation by heterotopic noxious conditioning stimulation in patients with painful osteoarthritis before, but not following, surgical pain relief. *Pain*, *88*(1), 69-78.
- Kutch, J. J., Ichesco, E., Hampson, J. P., Labus, J. S., Farmer, M. A., Martucci, K. T., . . . Network, M. R. (2017). Brain signature and functional impact of centralized pain: a multidisciplinary approach to the study of chronic pelvic pain (MAPP) network study. *Pain*, *158*(10), 1979-1991. doi:10.1097/j.pain.0000000000001001
- Le Bars, D., Dickenson, A. H., & Besson, J. M. (1979). Diffuse noxious inhibitory controls (DNIC). I. Effects on dorsal horn convergent neurones in the rat. *Pain*, *6*(3), 283-304.

- Leffler, A. S., Hansson, P., & Kosek, E. (2002). Somatosensory perception in a remote pain-free area and function of diffuse noxious inhibitory controls (DNIC) in patients suffering from long-term trapezius myalgia. *Eur J Pain*, *6*(2), 149-159. doi:10.1053/eujp.2001.0312
- Levine, T. D., & Saperstein, D. S. (2015). Routine use of punch biopsy to diagnose small fiber neuropathy in fibromyalgia patients. *Clin Rheumatol*, *34*(3), 413-417. doi:10.1007/s10067-014-2850-5
- Levitt, A. E., Galor, A., Chowdhury, A. R., Felix, E. R., Sarantopoulos, C. D., Zhuang, G. Y., . . . Levitt, R. C. (2017). Evidence that Dry Eye Represents a Chronic Overlapping Pain Condition. *Mol Pain*, *13*, 1744806917729306. doi:10.1177/1744806917729306
- Lewis, G. N., Rice, D. A., & McNair, P. J. (2012). Conditioned pain modulation in populations with chronic pain: a systematic review and meta-analysis. *J Pain*, *13*(10), 936-944. doi:10.1016/j.jpain.2012.07.005
- Lewis, J. D., Wassermann, E. M., Chao, W., Ramage, A. E., Robin, D. A., & Clauw, D. J. (2012). Central sensitization as a component of post-deployment syndrome. *NeuroRehabilitation*, *31*(4), 367-372. doi:10.3233/NRE-2012-00805
- Loggia, M. L., Kim, J., Gollub, R. L., Vangel, M. G., Kirsch, I., Kong, J., . . . Napadow, V. (2013). Default mode network connectivity encodes clinical pain: an arterial spin labeling study. *Pain*, *154*(1), 24-33. doi:10.1016/j.pain.2012.07.029
- Lopez-Sola, M., Pujol, J., Wager, T. D., Garcia-Fontanals, A., Blanco-Hinojo, L., Garcia-Blanco, S., . . . Deus, J. (2014). Altered functional magnetic resonance imaging responses to nonpainful sensory stimulation in fibromyalgia patients. *Arthritis Rheumatol*, *66*(11), 3200-3209. doi:10.1002/art.38781
- Lopez-Sola, M., Woo, C. W., Pujol, J., Deus, J., Harrison, B. J., Monfort, J., & Wager, T. D. (2017). Towards a neurophysiological signature for fibromyalgia. *Pain*, *158*(1), 34-47. doi:10.1097/j.pain.0000000000000707

- Mahomedy, M. C., Downing, J. W., Jeal, D. E., & Coleman, A. J. (1976). Anaesthetic induction for Caesarean section with propanidid. *Anaesthesia*, *31*(2), 205-211.
- Maixner, W., Fillingim, R., Sigurdsson, A., Kincaid, S., & Silva, S. (1998). Sensitivity of patients with painful temporomandibular disorders to experimentally evoked pain: evidence for altered temporal summation of pain. *Pain*, *76*(1-2), 71-81.
- Maixner, W., Fillingim, R. B., Williams, D. A., Smith, S. B., & Slade, G. D. (2016). Overlapping Chronic Pain Conditions: Implications for Diagnosis and Classification. *J Pain*, *17*(9 Suppl), T93-T107. doi:10.1016/j.jpain.2016.06.002
- Maneuf, Y. P., Hughes, J., & McKnight, A. T. (2001). Gabapentin inhibits the substance P-facilitated K(+)-evoked release of [(3)H]glutamate from rat caudial trigeminal nucleus slices. *Pain*, *93*(2), 191-196.
- Martenson, M. E., Halawa, O. I., Tonsfeldt, K. J., Maxwell, C. A., Hammack, N., Mist, S. D., . . . Heinricher, M. M. (2016). A possible neural mechanism for photosensitivity in chronic pain. *Pain*, *157*(4), 868-878. doi:10.1097/j.pain.0000000000000450
- Martinez-Lavin, M., Hermosillo, A. G., Rosas, M., & Soto, M. E. (1998). Circadian studies of autonomic nervous balance in patients with fibromyalgia: a heart rate variability analysis. *Arthritis Rheum*, *41*(11), 1966-1971. doi:10.1002/1529-0131(199811)41:11<1966::AID-ART11>3.0.CO;2-O
- McDermid, A. J., Rollman, G. B., & McCain, G. A. (1996). Generalized hypervigilance in fibromyalgia: evidence of perceptual amplification. *Pain*, *66*(2-3), 133-144.
- McLean, S. A., & Clauw, D. J. (2004). Predicting chronic symptoms after an acute "stressor"-- lessons learned from 3 medical conditions. *Med Hypotheses*, *63*(4), 653-658. doi:10.1016/j.mehy.2004.03.022
- McLean, S. A., Diatchenko, L., Lee, Y. M., Swor, R. A., Domeier, R. M., Jones, J. S., . . . Liberzon, I. (2011). Catechol O-methyltransferase haplotype predicts immediate musculoskeletal

- neck pain and psychological symptoms after motor vehicle collision. *J Pain*, 12(1), 101-107. doi:10.1016/j.jpain.2010.05.008
- McLean, S. A., Williams, D. A., Harris, R. E., Kop, W. J., Groner, K. H., Ambrose, K., . . . Clauw, D. J. (2005). Momentary relationship between cortisol secretion and symptoms in patients with fibromyalgia. *Arthritis Rheum*, 52(11), 3660-3669. doi:10.1002/art.21372
- McLean, S. A., Williams, D. A., Stein, P. K., Harris, R. E., Lyden, A. K., Whalen, G., . . . Clauw, D. J. (2006). Cerebrospinal fluid corticotropin-releasing factor concentration is associated with pain but not fatigue symptoms in patients with fibromyalgia. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*, 31(12), 2776-2782. doi:10.1038/sj.npp.1301200
- Mertz, H., Naliboff, B., Munakata, J., Niazi, N., & Mayer, E. A. (1995). Altered Rectal Perception Is a Biological Marker of Patients with Irritable-Bowel-Syndrome. *Gastroenterology*, 109(1), 40-52. doi:Doi 10.1016/0016-5085(95)90267-8
- Moldofsky, H. (2010). Rheumatic manifestations of sleep disorders. *Curr Opin Rheumatol*, 22(1), 59-63. doi:10.1097/BOR.0b013e328333b9cc
- Moldofsky, H., Saskin, P., & Lue, F. A. (1988). Sleep and symptoms in fibrositis syndrome after a febrile illness. *J Rheumatol*, 15(11), 1701-1704.
- Naliboff, B. D., Derbyshire, S. W., Munakata, J., Berman, S., Mandelkern, M., Chang, L., & Mayer, E. A. (2001). Cerebral activation in patients with irritable bowel syndrome and control subjects during rectosigmoid stimulation. *Psychosom Med*, 63(3), 365-375.
- Napadow, V., Kim, J., Clauw, D. J., & Harris, R. E. (2012). Decreased intrinsic brain connectivity is associated with reduced clinical pain in fibromyalgia. *Arthritis Rheum*, 64(7), 2398-2403. doi:10.1002/art.34412

- Napadow, V., LaCount, L., Park, K., As-Sanie, S., Clauw, D. J., & Harris, R. E. (2010). Intrinsic brain connectivity in fibromyalgia is associated with chronic pain intensity. *Arthritis Rheum*, *62*(8), 2545-2555. doi:10.1002/art.27497
- Neddermeyer, T. J., Fluhr, K., & Lotsch, J. (2008). Principle components analysis of pain thresholds to thermal, electrical, and mechanical stimuli suggests a predominant common source of variance. *Pain*, *138*(2), 286-291. doi:10.1016/j.pain.2007.12.015
- Neville, S. J., Clauw, A., Moser, S. E., Urquhart, A. G., Clauw, D. J., Brummett, C. M., & Harte, S. E. (In Press). Association between the 2011 fibromyalgia survey criteria and multisite pain sensitivity. *Clin J Pain*.
- Nielsen, C. S., Staud, R., & Price, D. D. (2009). Individual differences in pain sensitivity: measurement, causation, and consequences. *J Pain*, *10*(3), 231-237. doi:10.1016/j.jpain.2008.09.010
- Nir, R. R., & Yarnitsky, D. (2015). Conditioned pain modulation. *Curr Opin Support Palliat Care*, *9*(2), 131-137. doi:10.1097/SPC.0000000000000126
- Oaklander, A. L., Herzog, Z. D., Downs, H. M., & Klein, M. M. (2013). Objective evidence that small-fiber polyneuropathy underlies some illnesses currently labeled as fibromyalgia. *Pain*, *154*(11), 2310-2316. doi:10.1016/j.pain.2013.06.001
- Oudejans, L. C., Smit, J. M., van Velzen, M., Dahan, A., & Niesters, M. (2015). The influence of offset analgesia on the onset and offset of pain in patients with fibromyalgia. *Pain*, *156*(12), 2521-2527. doi:10.1097/j.pain.0000000000000321
- Persson, J., Axelsson, G., Hallin, R. G., & Gustafsson, L. L. (1995). Beneficial effects of ketamine in a chronic pain state with allodynia, possibly due to central sensitization. *Pain*, *60*(2), 217-222.
- Petersen, K. K., Arendt-Nielsen, L., Simonsen, O., Wilder-Smith, O., & Laursen, M. B. (2015). Presurgical assessment of temporal summation of pain predicts the development of

chronic postoperative pain 12 months after total knee replacement. *Pain*, 156(1), 55-61.
doi:10.1016/j.pain.0000000000000022

Petersen, K. K., Graven-Nielsen, T., Simonsen, O., Laursen, M. B., & Arendt-Nielsen, L. (2016). Preoperative pain mechanisms assessed by cuff algometry are associated with chronic postoperative pain relief after total knee replacement. *Pain*, 157(7), 1400-1406.
doi:10.1097/j.pain.0000000000000531

Petzke, F., Harris, R. E., Williams, D. A., Clauw, D. J., & Gracely, R. H. (2005). Differences in unpleasantness induced by experimental pressure pain between patients with fibromyalgia and healthy controls. *Eur J Pain*, 9(3), 325-335.
doi:10.1016/j.ejpain.2004.09.001

Pfau, D. B., Geber, C., Birklein, F., & Treede, R. D. (2012). Quantitative sensory testing of neuropathic pain patients: potential mechanistic and therapeutic implications. *Curr Pain Headache Rep*, 16(3), 199-206. doi:10.1007/s11916-012-0261-3

Phillips, K., & Clauw, D. J. (2013). Central pain mechanisms in the rheumatic diseases: future directions. *Arthritis Rheum*, 65(2), 291-302. doi:10.1002/art.37739

Piche, M., Arsenault, M., & Rainville, P. (2009). Cerebral and cerebrospinal processes underlying counterirritation analgesia. *J Neurosci*, 29(45), 14236-14246.
doi:10.1523/JNEUROSCI.2341-09.2009

Pierce, A. N., & Christianson, J. A. (2015). Stress and chronic pelvic pain. *Prog Mol Biol Transl Sci*, 131, 509-535. doi:10.1016/bs.pmbts.2014.11.009

Ploner, M., Lee, M. C., Wiech, K., Bingel, U., & Tracey, I. (2010). Prestimulus functional connectivity determines pain perception in humans. *Proceedings of the National Academy of Sciences of the United States of America*, 107(1), 355-360.
doi:10.1073/pnas.0906186106

- Potvin, S., & Marchand, S. (2016). Pain facilitation and pain inhibition during conditioned pain modulation in fibromyalgia and in healthy controls. *Pain, 157*(8), 1704-1710.
doi:10.1097/j.pain.0000000000000573
- Price, D. D., Mao, J., Frenk, H., & Mayer, D. J. (1994). The N-methyl-D-aspartate receptor antagonist dextromethorphan selectively reduces temporal summation of second pain in man. *Pain, 59*(2), 165-174.
- Price, D. D., Staud, R., Robinson, M. E., Mauderli, A. P., Cannon, R., & Vierck, C. J. (2002). Enhanced temporal summation of second pain and its central modulation in fibromyalgia patients. *Pain, 99*(1-2), 49-59.
- Pyke, T. L., Osmotherly, P. G., & Baines, S. (2017). Measuring Glutamate Levels in the Brains of Fibromyalgia Patients and a Potential Role for Glutamate in the Pathophysiology of Fibromyalgia Symptoms: A Systematic Review. *Clin J Pain, 33*(10), 944-954.
doi:10.1097/AJP.0000000000000474
- Qiao, Z. G., Vaeroy, H., & Morkrid, L. (1991). Electrodermal and microcirculatory activity in patients with fibromyalgia during baseline, acoustic stimulation and cold pressor tests. *J Rheumatol, 18*(9), 1383-1389.
- Russell, I. J., Holman, A. J., Swick, T. J., Alvarez-Horine, S., Wang, Y. G., Guinta, D., & Sodium Oxybate, F. M. S. G. (2011). Sodium oxybate reduces pain, fatigue, and sleep disturbance and improves functionality in fibromyalgia: results from a 14-week, randomized, double-blind, placebo-controlled study. *Pain, 152*(5), 1007-1017.
doi:10.1016/j.pain.2010.12.022
- Sarchielli, P., Di Filippo, M., Nardi, K., & Calabresi, P. (2007). Sensitization, glutamate, and the link between migraine and fibromyalgia. *Curr.Pain Headache Rep., 11*(5), 343-351.
- Sarlani, E., Grace, E. G., Reynolds, M. A., & Greenspan, J. D. (2004). Evidence for up-regulated central nociceptive processing in patients with masticatory myofascial pain. *J Orofac Pain, 18*(1), 41-55.

- Schmidt-Wilcke, T., Ichesco, E., Hampson, J. P., Kairys, A., Peltier, S., Harte, S., . . . Harris, R. E. (2014). Resting state connectivity correlates with drug and placebo response in fibromyalgia patients. *Neuroimage Clin*, *6*, 252-261. doi:10.1016/j.nicl.2014.09.007
- Schoen, C. J., Ablin, J. N., Ichesco, E., Bhavsar, R. J., Kochlefl, L., Harris, R. E., . . . Harte, S. E. (2016). A novel paradigm to evaluate conditioned pain modulation in fibromyalgia. *J Pain Res*, *9*, 711-719. doi:10.2147/JPR.S115193
- Schrepf, A., Bradley, C. S., O'Donnell, M., Luo, Y., Harte, S. E., Kreder, K., . . . Multidisciplinary Approach to the Study of Chronic Pelvic Pain Research, N. (2015). Toll-like receptor 4 and comorbid pain in Interstitial Cystitis/Bladder Pain Syndrome: a multidisciplinary approach to the study of chronic pelvic pain research network study. *Brain Behav Immun*, *49*, 66-74. doi:10.1016/j.bbi.2015.03.003
- Schrepf, A., Harper, D. E., Harte, S. E., Wang, H., Ichesco, E., Hampson, J. P., . . . Harris, R. E. (2016). Endogenous opioidergic dysregulation of pain in fibromyalgia: a PET and fMRI study. *Pain*, *157*(10), 2217-2225. doi:10.1097/j.pain.0000000000000633
- Schrepf, A., Harte, S. E., Miller, N., Fowler, C., Nay, C., Williams, D. A., . . . Rothberg, A. (2017). Improvement in the Spatial Distribution of Pain, Somatic Symptoms, and Depression After a Weight Loss Intervention. *J Pain*, *18*(12), 1542-1550. doi:10.1016/j.jpain.2017.08.004
- Schweinhardt, P., Sauro, K. M., & Bushnell, M. C. (2008). Fibromyalgia: a disorder of the brain? *Neuroscientist*, *14*(5), 415-421. doi:10.1177/1073858407312521
- Scott, J. R., Hassett, A. L., Schrepf, A. D., Brummett, C. M., Harris, R. E., Clauw, D. J., & Harte, S. E. (2018). Moderate Alcohol Consumption Is Associated with Reduced Pain and Fibromyalgia Symptoms in Chronic Pain Patients. *Pain Med*. doi:10.1093/pm/pny032
- Serra, J., Collado, A., Sola, R., Antonelli, F., Torres, X., Salgueiro, M., . . . Bostock, H. (2014). Hyperexcitable C nociceptors in fibromyalgia. *Ann Neurol*, *75*(2), 196-208. doi:10.1002/ana.24065

- Skou, S. T., Graven-Nielsen, T., Rasmussen, S., Simonsen, O. H., Laursen, M. B., & Arendt-Nielsen, L. (2013). Widespread sensitization in patients with chronic pain after revision total knee arthroplasty. *Pain, 154*(9), 1588-1594. doi:10.1016/j.pain.2013.04.033
- Sluka, K. A. (2009). Is it possible to develop an animal model of fibromyalgia? *Pain, 146*(1-2), 3-4. doi:10.1016/j.pain.2009.07.032
- Sluka, K. A., & Clauw, D. J. (2016). Neurobiology of fibromyalgia and chronic widespread pain. *Neuroscience, 338*, 114-129. doi:10.1016/j.neuroscience.2016.06.006
- Sluka, K. A., Kalra, A., & Moore, S. A. (2001). Unilateral intramuscular injections of acidic saline produce a bilateral, long-lasting hyperalgesia. *Muscle Nerve, 24*(1), 37-46.
- Sluka, K. A., & Westlund, K. N. (1992). An experimental arthritis in rats: dorsal horn aspartate and glutamate increases. *Neurosci Lett, 145*(2), 141-144. doi:Doi 10.1016/0304-3940(92)90006-S
- Smallwood, R. F., Laird, A. R., Ramage, A. E., Parkinson, A. L., Lewis, J., Clauw, D. J., . . . Robin, D. A. (2013). Structural brain anomalies and chronic pain: a quantitative meta-analysis of gray matter volume. *J Pain, 14*(7), 663-675. doi:10.1016/j.jpain.2013.03.001
- Smith, S. B., Maixner, D. W., Fillingim, R. B., Slade, G., Gracely, R. H., Ambrose, K., . . . Diatchenko, L. (2012). Large candidate gene association study reveals genetic risk factors and therapeutic targets for fibromyalgia. *Arthritis Rheum, 64*(2), 584-593. doi:10.1002/art.33338
- Sorensen, J., Bengtsson, A., Backman, E., Henriksson, K. G., & Bengtsson, M. (1995). Pain analysis in patients with fibromyalgia. Effects of intravenous morphine, lidocaine, and ketamine. *Scand J Rheumatol, 24*(6), 360-365.
- Sprenger, C., Bingel, U., & Buchel, C. (2011). Treating pain with pain: supraspinal mechanisms of endogenous analgesia elicited by heterotopic noxious conditioning stimulation. *Pain, 152*(2), 428-439. doi:10.1016/j.pain.2010.11.018

- Staud, R., Craggs, J. G., Perlstein, W. M., Robinson, M. E., & Price, D. D. (2008). Brain activity associated with slow temporal summation of C-fiber evoked pain in fibromyalgia patients and healthy controls. *Eur J Pain*, *12*(8), 1078-1089. doi:10.1016/j.ejpain.2008.02.002
- Staud, R., Koo, E., Robinson, M. E., & Price, D. D. (2007). Spatial summation of mechanically evoked muscle pain and painful aftersensations in normal subjects and fibromyalgia patients. *Pain*, *130*(1-2), 177-187. doi:10.1016/j.pain.2007.03.015
- Staud, R., Vierck, C. J., Cannon, R. L., Mauderli, A. P., & Price, D. D. (2001). Abnormal sensitization and temporal summation of second pain (wind-up) in patients with fibromyalgia syndrome. *Pain*, *91*(1-2), 165-175.
- Staud, R., Weyl, E. E., Bartley, E., Price, D. D., & Robinson, M. E. (2014). Analgesic and anti-hyperalgesic effects of muscle injections with lidocaine or saline in patients with fibromyalgia syndrome. *Eur J Pain*, *18*(6), 803-812. doi:10.1002/j.1532-2149.2013.00422.x
- Suarez-Roca, H., Silva, J. A., Arcaya, J. L., Quintero, L., Maixner, W., & Pinerua-Shuhaibar, L. (2006). Role of mu-opioid and NMDA receptors in the development and maintenance of repeated swim stress-induced thermal hyperalgesia. *Behavioural brain research*, *167*(2), 205-211. doi:10.1016/j.bbr.2005.09.006
- Taguchi, T., Katanosaka, K., Yasui, M., Hayashi, K., Yamashita, M., Wakatsuki, K., . . . Mizumura, K. (2015). Peripheral and spinal mechanisms of nociception in a rat reserpine-induced pain model. *Pain*, *156*(3), 415-427. doi:10.1097/01.j.pain.0000460334.49525.5e
- Tracey, I., & Bushnell, M. C. (2009). How neuroimaging studies have challenged us to rethink: is chronic pain a disease? *J Pain*, *10*(11), 1113-1120.
- Treede, R. D., Davis, K. D., Campbell, J. N., & Raja, S. N. (1992). The plasticity of cutaneous hyperalgesia during sympathetic ganglion blockade in patients with neuropathic pain. *Brain*, *115* (Pt 2), 607-621.

- Uceyler, N., Zeller, D., Kahn, A. K., Kewenig, S., Kittel-Schneider, S., Schmid, A., . . . Sommer, C. (2013). Small fibre pathology in patients with fibromyalgia syndrome. *Brain*, *136*(Pt 6), 1857-1867. doi:10.1093/brain/awt053
- Vaegter, H. B., & Graven-Nielsen, T. (2016). Pain modulatory phenotypes differentiate subgroups with different clinical and experimental pain sensitivity. *Pain*, *157*(7), 1480-1488. doi:10.1097/j.pain.0000000000000543
- Wager, T. D., Atlas, L. Y., Lindquist, M. A., Roy, M., Woo, C. W., & Kross, E. (2013). An fMRI-based neurologic signature of physical pain. *N Engl J Med*, *368*(15), 1388-1397. doi:10.1056/NEJMoa1204471
- Wallace, D., Bowman, R. L., Wormsley, S. B., & Peter, J. B. (1989). Cytokines and immune regulation in patients with fibrositis [letter] [published erratum appears in *Arthritis Rheum* 1989 Dec;32(12):1607]. *Arthritis Rheum.*, *32*(10), 1334-1335.
- Warren, J. W., Howard, F. M., Cross, R. K., Good, J. L., Weissman, M. M., Wesselmann, U., . . . Clauw, D. J. (2009). Antecedent nonbladder syndromes in case-control study of interstitial cystitis/painful bladder syndrome. *Urology*, *73*(1), 52-57. doi:10.1016/j.urology.2008.06.031
- Watson, C. J. (2016). Insular balance of glutamatergic and GABAergic signaling modulates pain processing. *Pain*, *157*(10), 2194-2207. doi:10.1097/j.pain.0000000000000615
- Weissman-Fogel, I., Granovsky, Y., Crispel, Y., Ben-Nun, A., Best, L. A., Yarnitsky, D., & Granot, M. (2009). Enhanced presurgical pain temporal summation response predicts post-thoracotomy pain intensity during the acute postoperative phase. *J Pain*, *10*(6), 628-636. doi:10.1016/j.jpain.2008.12.009
- Whitehead, W. E., Holtkotter, B., Enck, P., Hoelzl, R., Holmes, K. D., Anthony, J., . . . Schuster, M. M. (1990). Tolerance for rectosigmoid distention in irritable bowel syndrome. *Gastroenterology*, *98*(5 Pt 1), 1187-1192.

- Wilbarger, J. L., & Cook, D. B. (2011). Multisensory hypersensitivity in women with fibromyalgia: implications for well being and intervention. *Arch Phys Med Rehabil*, *92*(4), 653-656. doi:10.1016/j.apmr.2010.10.029
- Wilder-Smith, C. H., & Robert-Yap, J. (2007). Abnormal endogenous pain modulation and somatic and visceral hypersensitivity in female patients with irritable bowel syndrome. *World J Gastroenterol*, *13*(27), 3699-3704.
- Wilder-Smith, O. H. G., Tassonyi, E., & Arendt-Nielsen, L. (2002). Preoperative back pain is associated with diverse manifestations of central neuroplasticity. *Pain*, *97*(3), 189-194. doi:Pii S0304-3959(01)00430-4
Doi 10.1016/S0304-3959(01)00430-4
- Williams, D. A., & Clauw, D. J. (2009). Understanding fibromyalgia: lessons from the broader pain research community. *J Pain*, *10*(8), 777-791. doi:10.1016/j.jpain.2009.06.001
- Willis, W. D., Jr. (1985). Central nervous system mechanisms for pain modulation. *Appl Neurophysiol*, *48*(1-6), 153-165.
- Willis, W. D., Jr. (1988). Dorsal horn neurophysiology of pain. *Ann N Y Acad Sci*, *531*, 76-89.
- Wolfe, F. (2009). Fibromyalgiansess. *Arthritis Rheum*, *61*(6), 715-716. doi:10.1002/art.24553
- Wolfe, F., Clauw, D. J., Fitzcharles, M. A., Goldenberg, D. L., Hauser, W., Katz, R. L., . . . Walitt, B. (2016). 2016 Revisions to the 2010/2011 fibromyalgia diagnostic criteria. *Semin Arthritis Rheum*, *46*(3), 319-329. doi:10.1016/j.semarthrit.2016.08.012
- Wolfe, F., Clauw, D. J., Fitzcharles, M. A., Goldenberg, D. L., Hauser, W., Katz, R. S., . . . Winfield, J. B. (2011). Fibromyalgia criteria and severity scales for clinical and epidemiological studies: a modification of the ACR Preliminary Diagnostic Criteria for Fibromyalgia. *J Rheumatol*, *38*(6), 1113-1122. doi:10.3899/jrheum.100594

- Wolfe, F., Clauw, D. J., Fitzcharles, M. A., Goldenberg, D. L., Katz, R. S., Mease, P., . . . Yunus, M. B. (2010). The American College of Rheumatology preliminary diagnostic criteria for fibromyalgia and measurement of symptom severity. *Arthritis Care Res (Hoboken)*, *62*(5), 600-610. doi:10.1002/acr.20140
- Wolfe, F., Michaud, K., Busch, R. E., Katz, R. S., Rasker, J. J., Shahouri, S. H., . . . Hauser, W. (2014). Polysymptomatic distress in patients with rheumatoid arthritis: understanding disproportionate response and its spectrum. *Arthritis Care Res (Hoboken)*, *66*(10), 1465-1471. doi:10.1002/acr.22300
- Wolfe, F., Michaud, K., Li, T., & Katz, R. S. (2010). Chronic conditions and health problems in rheumatic diseases: comparisons with rheumatoid arthritis, noninflammatory rheumatic disorders, systemic lupus erythematosus, and fibromyalgia. *J Rheumatol*, *37*(2), 305-315. doi:10.3899/jrheum.090781
- Wood, P. B., Schweinhardt, P., Jaeger, E., Dagher, A., Hakyemez, H., Rabiner, E. A., . . . Chizh, B. A. (2007). Fibromyalgia patients show an abnormal dopamine response to pain. *Eur J Neurosci*, *25*(12), 3576-3582. doi:10.1111/j.1460-9568.2007.05623.x
- Woolf, C. J. (1983). Evidence for a central component of post-injury pain hypersensitivity. *Nature*, *306*(5944), 686-688.
- Woolf, C. J. (1991). Generation of acute pain: central mechanisms. *Br Med Bull*, *47*(3), 523-533.
- Woolf, C. J. (2011). Central sensitization: implications for the diagnosis and treatment of pain. *Pain*, *152*(3 Suppl), S2-15. doi:10.1016/j.pain.2010.09.030
- Wylde, V., Sayers, A., Odutola, A., Gooberman-Hill, R., Dieppe, P., & Blom, A. W. (2017). Central sensitization as a determinant of patients' benefit from total hip and knee replacement. *Eur J Pain*, *21*(2), 357-365. doi:10.1002/ejp.929
- Yaksh, T. L. (1988). CNS mechanisms of pain and analgesia. *Cancer Surv*, *7*(1), 5-28.

- Yarnitsky, D. (2015). Role of endogenous pain modulation in chronic pain mechanisms and treatment. *Pain, 156 Suppl 1*, S24-31. doi:10.1097/01.j.pain.0000460343.46847.58
- Yarnitsky, D., Crispel, Y., Eisenberg, E., Granovsky, Y., Ben-Nun, A., Sprecher, E., . . . Granot, M. (2008). Prediction of chronic post-operative pain: pre-operative DNIC testing identifies patients at risk. *Pain, 138*(1), 22-28. doi:10.1016/j.pain.2007.10.033
- Yarnitsky, D., Granot, M., & Granovsky, Y. (2014). Pain modulation profile and pain therapy: between pro- and antinociception. *Pain, 155*(4), 663-665. doi:10.1016/j.pain.2013.11.005
- Yarnitsky, D., Granot, M., Nahman-Averbuch, H., Khamaisi, M., & Granovsky, Y. (2012). Conditioned pain modulation predicts duloxetine efficacy in painful diabetic neuropathy. *Pain, 153*(6), 1193-1198. doi:10.1016/j.pain.2012.02.021
- Youssef, A. M., Macefield, V. G., & Henderson, L. A. (2016a). Cortical influences on brainstem circuitry responsible for conditioned pain modulation in humans. *Hum Brain Mapp, 37*(7), 2630-2644. doi:10.1002/hbm.23199
- Youssef, A. M., Macefield, V. G., & Henderson, L. A. (2016b). Pain inhibits pain; human brainstem mechanisms. *Neuroimage, 124*(Pt A), 54-62. doi:10.1016/j.neuroimage.2015.08.060
- Yunus, M. B. (2007). Fibromyalgia and overlapping disorders: the unifying concept of central sensitivity syndromes. *Semin Arthritis Rheum, 36*(6), 339-356. doi:10.1016/j.semarthrit.2006.12.009
- Yunus, M. B. (2008). Central sensitivity syndromes: a new paradigm and group nosology for fibromyalgia and overlapping conditions, and the related issue of disease versus illness. *Semin Arthritis Rheum, 37*(6), 339-352. doi:10.1016/j.semarthrit.2007.09.003

Figure 1. Neurotransmitter systems that generally facilitate (left side) or inhibit (right side) central nervous system (CNS) pain transmission. The arrows indicate the levels of these neurotransmitters in the CNS of individuals with fibromyalgia (FM), and the boxes indicate drugs that have been shown to be effective in FM that are likely working in part via those neurotransmitters.





Figure 2. The 2011 Survey Criteria for Fibromyalgia (Wolfe et al., 2011) using the Michigan Body Map (C. M. Brummett et al., 2016).

Fibromyalgia Symptoms (Modified ACR 2010 Fibromyalgia Diagnostic Criteria)

1. Please indicate below if you have had pain or tenderness over the past 7 days in each of the areas listed below. Check the boxes in the diagram below for each area in which you have had pain or tenderness. Be sure to mark right and left sides separately.



2. Using the following scale, indicate for each item your severity over the past week by checking the appropriate box.

No problem
Slight or mild problems: generally mild or intermittent
Moderate: considerable problems; often present and/or at a moderate level
Severe: continuous, life-disturbing problems

	No problem	Slight or mild	Moderate	Severe
a. Fatigue	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b. Trouble thinking or remembering	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
c. Waking up tired (unrefreshed)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

3. During the past 6 months have you had any of the following symptoms?

	No	Yes
a. Pain or cramps in lower abdomen	<input type="checkbox"/>	<input type="checkbox"/>
b. Depression	<input type="checkbox"/>	<input type="checkbox"/>
c. Headache	<input type="checkbox"/>	<input type="checkbox"/>

4. Have the symptoms in questions 2-3 and pain been present at a similar level for at least 3 months?

	No	Yes
	<input type="checkbox"/>	<input type="checkbox"/>

5. Do you have a disorder that would otherwise explain the pain?

	No	Yes
	<input type="checkbox"/>	<input type="checkbox"/>

Figure 3. Fibromyalginess (FMness) scores from individuals undergoing lower extremity arthroplasty for osteoarthritis.^{227,228} The red line indicates the score that is used to classify an individual as being FM positive. Two different hypothetical participants, who do not meet criteria for FM, are compared with respect to the amount of opioids (in oral morphine equivalents) required for pain control during a 24-48 hour inpatient period and the likelihood of achieving 50% improvement in pain at six months following surgery.





Table 1. Differences between “*top down*” and “*bottom up*” forms of central sensitization.



Clauw DJ. Refresher Courses, 16th World Congress on Pain. 2016.

Author Manuscript