



Laser therapy for treatment of peri-implant mucositis and peri-implantitis: An American Academy of Periodontology best evidence review

Guo-Hao Lin^{1,2} | Fernando Suárez López del Amo^{1,3} | Hom-Lay Wang¹

¹Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry, Ann Arbor, MI

²Department of Orofacial Sciences, University of California San Francisco School of Dentistry, San Francisco, CA

³Department of Periodontics, University of Oklahoma College of Dentistry, Norman, OK

Correspondence

Hom-Lay Wang, Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry, 1011 North University Avenue, Ann Arbor, MI 48109-1078.
Email: homlay@umich.edu

Abstract

Background: Peri-implant diseases are prevalent, with numerous therapies studied in an attempt to combat this condition. The present review aims to systematically evaluate the effectiveness of laser therapy with non-surgical or surgical therapy in managing peri-implant mucositis and peri-implantitis.

Methods: An electronic search of three databases and a hand search of peer-reviewed journals for relevant articles published (in English) from January 1980 to June 2016 were performed. Human clinical trials of ≥ 10 patients with peri-implant diseases, treated with surgical or non-surgical approaches and laser therapy, and a follow-up period of ≥ 6 months, were included. Random-effects meta-analyses were performed to analyze weighted mean difference (WMD) and confidence interval for the recorded variables according to PRISMA guidelines. Risk of bias assessment was also performed for randomized controlled trials included.

Results: From 22 articles selected, 11 were included in the meta-analyses. The outcomes of using lasers as a monotherapy could not be evaluated since no controlled studies were identified. Therefore, all reported results were the outcomes of applying lasers as an adjunct to surgical/non-surgical treatment. For the non-surgical approach, WMD of probing depth (PD), clinical attachment level (CAL), bleeding on probing (BOP), plaque index (PI), marginal bone level (MBL) and recession (REC) was 0.15 mm ($P = 0.50$), -0.10 mm ($P = 0.32$), 21.08% ($P = 0.02$), -0.07 ($P = 0.002$), -0.22 mm ($P = 0.04$) and -0.11 mm ($P = 0.34$), respectively. For the surgical approach with a long-term follow up, WMD of PD, CAL, BOP, and PI was 0.45 mm ($P = 0.11$), 0.22 mm ($P = 0.56$), 7.26% ($P = 0.76$) and -0.09 ($P = 0.84$), respectively.

Conclusions: Current evidence shows laser therapy in combination with surgical/non-surgical therapy provided minimal benefit in PD reduction, CAL gain, amount of REC improvement, and PI reduction in the treatment of peri-implant diseases. Lasers when used as an adjunct to non-surgical therapy might result in more BOP reduction in the short term. However, current evidence allowed for analysis of only Er:YAG, CO₂, and diode lasers. Studies on others failed to have controlled evidence supporting their evaluation.

KEYWORDS

Decontamination, dental implants, lasers, meta-analyses, peri-implantitis, systematic review



Laser, an acronym for light amplification by stimulated emission of radiation, was introduced in periodontics in the 1990s¹⁻³ as a tool in diagnostics, surgeries, and physiologic studies.⁴ It is a device that emits light coherently into a small, intense, and nearly non-divergent beam with sufficient energy to cut through hard and soft tissues. The effect of a laser depends on the energy emitted and absorption by the target tissue. Basically, this energy represents a monochromatic light that is collimated into a focused beam interacting with the targeted tissue by being scattered, transmitted, absorbed, or reflected (see supplementary Figure 1 in online *Journal of Periodontology*). The power of the energy could result in effects such as warming, coagulation, or vaporization of the tissues, based on different energy levels.⁴ Therefore, lasers have myriad applications in periodontics and implant dentistry, which includes non-surgical and surgical periodontal therapy,^{5,6} gingivectomy and crown-lengthening procedures,^{7,8} as well as decontamination of implants with peri-implantitis.⁹

Given the increased use of dental implants in oral reconstructions, it is no surprise that recent epidemiologic research demonstrated mean prevalence rates of 43% and 22% for peri-implant mucositis and peri-implantitis, respectively.¹⁰ Likewise, there is an increase in the number of studies conducted seeking an effective and predictable therapy for peri-implant diseases. It was shown that peri-implant mucositis could be effectively treated by mechanical non-surgical therapy,¹¹ and peri-implantitis generally requires surgical interventions.^{12,13} Consequently, a variety of surgical approaches, such as access flap debridement, resective and regenerative procedures, implantoplasty, and lasers for implant surface decontamination, have been proposed.¹⁴

With only limited evidence in specific scenarios, the employment of lasers around implants has been proven effective in different clinical therapies, such as non-surgical treatment¹⁵ and the combination of lasers and surgical treatment;^{16,17} however, conflicting results have been reported when lasers were used as a method for implant surface detoxification.¹⁶⁻¹⁸ A recent review concluded that laser therapy provided identical outcomes as other surface detoxification methods with regard to probing depth (PD) reduction, clinical attachment level (CAL) gain, and radiographic bone fill.¹⁹ The tremendous heterogeneity in treatment outcomes after the application of lasers for peri-implant diseases could be attributed to three main factors: 1) the multifactorial etiology of peri-implant infections, including host- and implant-related factors, could play a major role in treatment outcomes;²⁰⁻²³ 2) the wide range of lasers investigated had different properties and settings (i.e., wavelength, power, waveform, pulse duration, energy/pulse, density of the energy, duration of the exposure, angulation of the energy toward the targeted tissue, peak power of the pulse, and the properties of tissue),⁴ all of which could strongly influence the treatment

outcomes; and 3) the frequency of use might have an impact on treatment outcomes as several studies demonstrated positive results with repeated application of lasers to peri-implant defects.¹⁵

Despite the increasing number of investigations conducted using lasers for implant surface detoxification, significant heterogeneity and controversy still exist. Hence, the aim of this systematic review and meta-analyses is to evaluate the potential of lasers in the detoxification and treatment of peri-implant mucositis and peri-implantitis.

1 | MATERIALS AND METHODS

1.1 | Population, Intervention, Comparison, Outcome (PICO) question²⁴

The focused question of this systematic review was: “Do lasers used alone or as adjuncts provide better treatment and patient outcomes in the management of peri-implant mucositis or peri-implantitis”? The population selected comprised individuals with peri-implant mucositis or peri-implantitis. The intervention investigated was the use of lasers alone or as adjuncts in surgical/non-surgical therapies. The selected outcomes to be compared between individuals treated with laser and those treated without were changes in: 1) PD, 2) CAL, 3) percentage of bleeding on probing (BOP), 4) plaque index (PI), 5) recession (REC), and 6) marginal bone level (MBL).

1.2 | Selection criteria

Prospective and retrospective human case series (CS), non-randomized controlled clinical trials (CCTs), or randomized controlled clinical trials (RCTs) published (in English) from January 1980 to June 2016 were screened. Inclusion criteria were: 1) had ≥ 10 patients diagnosed with peri-implant mucositis or peri-implantitis and treated with lasers surgically or non-surgically; 2) had a follow-up period of ≥ 6 months; and 3) reported outcomes of one of the clinical parameters (PD, CAL, BOP, PI, REC, or MBL) after the therapy. Exclusion criteria were studies published as: 1) editorials; 2) letters or comments and non-English citations; 3) animal/in vitro studies; 4) review articles; and 5) case reports/series with < 10 patients.

1.3 | Screening process

Two independent examiners (G-HL and FSLA) conducted the literature search using three databases (Ovid MEDLINE, EMBASE, and Dentistry and Oral Sciences Source). The search terms used, where mh represented the MeSH terms, tiab represented title and/or abstract, ti represented title, ab represented abstract, la represented language, pt represented publication type, exp represented explosion,



lim represented limit, it represented item types, and DE represented heading or keyword, in MEDLINE/PubMed were: (“peri-implantitis”[mh] OR “peri-implant mucositis”[tiab] OR “peri-implant”[tiab] OR “peri-implants”[tiab] OR “peri-implantitis”[tiab] OR peri-implant[tiab] OR peri-implants[tiab] OR periimplantitis[tiab]) AND (“laser therapy”[mh] OR “lasers, solid-state”[mh] OR laser[tiab] OR lasers[tiab]) AND English[la] NOT (letter[pt] OR comment[pt] OR editorial[pt]) The search terms used in EMBASE were: ‘periimplantitis’/exp OR ‘peri—implant’:ab,ti OR ‘peri—implants’:ab,ti OR ‘peri-implantitis’:ab,ti OR peri-implant*:ab,ti AND (‘laser’/exp OR ‘solid state laser’/exp OR laser*:ab,ti) AND [english]/lim NOT ([animals]/lim NOT [humans]/lim) NOT (‘letter’/exp OR ‘editorial’/exp OR note:it OR erratum:it), and the search terms used in Dentistry and Oral Sciences Source were: (DE “PERI—implantitis” OR TI “peri—implant*” OR AB “peri—implant*”) AND (DE “Lasers” OR TI “Laser*” OR AB “Laser*”)

A hand search was also performed for all print only journals (January 1980 to June 2016). Furthermore, a search in the references of included papers as well as the related systematic reviews was conducted for publications not electronically identified. The two reviewers (G-HL and FSLA) examined the pre-identified articles in full text, and their eligibility for this review was confirmed after discussion. The level of agreement between reviewers regarding study inclusion was calculated with κ statistics.

1.4 | Statistical analyses

Two reviewers (G-HL and FSLA) independently extracted data from papers that met inclusion criteria. Any disagreements were reconciled after discussion with a third reviewer (H-LW). Demographics, such as study design, sample size, numbers of implants, follow-up period, treatment outcome measurements, and study conclusion, were extracted and recorded for each selected study. Authors of the selected studies were contacted if additional information regarding the study was needed for the review and meta-analyses.

The primary outcome was PD reduction, and the secondary outcome was changes in recorded peri-implant parameters. The pooled weighted mean difference (WMD) and the 95% confidence interval (CI) of each variable were estimated using a computer program.* Random effects meta-analyses of the selected studies were applied to minimize bias caused by methodologic differences among studies. Forest plots were generated to represent WMD and 95% CI in primary and secondary outcomes for all included studies using number of dental implants as the unit of analysis. Heterogeneity was assessed with χ^2 test and I^2 test, which ranged between 0% and 100% with the lower values representing less hetero-

geneity. If a study presented more than one test/control arm, the outcomes of each test/control group were combined. The reporting of these meta-analyses adhered to the PRISMA (Preferred Reporting Items for Systematic Review and Meta-analyses) statement.²⁵

1.5 | Risk of bias assessment

Criteria used to assess the quality of the selected RCTs were modified from the RCTs checklist of the Cochrane Center²⁶ and the CONSORT statement,²⁷ which provided guidelines for the following parameters: 1) sequence generation; 2) allocation concealment method; 3) masking of examiner; 4) adequate handling of incomplete outcome data; and 5) absence of selective outcome reporting. The degree of bias was categorized as low risk (if all criteria were met), moderate risk (when only one criterion was missing), or high risk (if two or more criteria were missing). Two reviewers (G-HL and FSLA) assessed all included articles independently (see supplementary Table 1 in online *Journal of Periodontology*).

1.6 | Level of evidence assessment

The level of currently available evidence for lasers in treatment of peri-implant diseases was examined. The authors categorized the evidence level based on the following criteria: 1) “no evidence” (represents no RCTs or CCTs were identified to warrant the treatment benefit); 2) “limited evidence” (represents ≤ 3 RCTs or CCTs were identified); and 3) “some evidence” (represents > 3 RCTs or CCTs were identified) (see supplementary Table 2 in online *Journal of Periodontology*).

2 | RESULTS

The screening process is shown in Figure 1. Electronic and hand searches yielded 237 articles, of which 27 were selected for full-text evaluation after screening of titles and abstracts. Five articles^{28–32} were then further excluded due to the following: 1) being an in vitro study;³¹ 2) use of a light-emitting diode source instead of laser;^{30,32} and 3) no clinical data reported.^{28,29} Twenty-two total articles^{9,15–18,33–49} were included in this systematic review. The main features and conclusions of the included studies are summarized in Table 1 (non-surgical) and Table 2 (surgical). Eleven papers (nine RCTs^{9,16,18,34,37,39,41,43,45} and two CCTs^{48,49}) that met the inclusion criteria were included for meta-analyses. It is worth noting that currently no controlled studies are identified offering evidence of lasers used as a monotherapy in the treatment of peri-implantitis. The κ value for inter-reviewer agreement for potentially relevant articles was 0.93 (titles and abstracts) and 0.97 (full-text articles), indicating an “almost perfect” agreement between the two reviewers.⁵⁰

* RevMan v.5.0, The Cochrane Collaboration, London, U.K.




FIGURE 1 Flowchart illustrating the publication selection process

2.1 | Features of the included studies

Among the 22 human clinical trials, lasers were used as an adjunct to non-surgical interventions in 13 studies (Table 1).^{15,34–37,40–43,45,47–49} In the other nine studies,^{9,16–18,33,38,39,44,46} lasers were used with a surgical approach (Table 2). Although the current review aims to investigate the effect of adjunctive laser treatment on clinical outcomes of peri-implant diseases, most of the included articles focused on peri-implantitis, and only three studies^{47–49} used lasers as an adjunct in the treatment of peri-implant mucositis. Seven studies^{9,16–18,33,38,44} introduced guided bone regeneration (GBR) procedures as part of surgical treatment with either non-resorbable^{16,33} or absorbable^{9,17,18,38,44} membranes. Most of the treated implants in the selected studies were rough-surfaced implants,^{9,15–18,33–35,38–49} whereas smooth-surfaced implants were more commonly found in the non-surgical laser-treated approach.^{36,37} Among the selected studies that examined a surgical treatment modality, the diode laser was used in three studies,^{33,39,46} the carbon dioxide (CO₂) laser in two studies,^{16,17} and the erbium:yttrium-aluminum-garnet (Er:YAG) laser in four studies.^{9,18,38,44} For studies that used a non-surgical treatment approach, the diode laser was used in eight studies,^{15,40–43,45,48,49} and

the Er:YAG laser was introduced in five studies.^{34–37,47} The selected articles generally included mechanical hand curettage with plastic, titanium, or carbon fiber curets. However, air abrasives,^{16,36,37,40,41,43} chlorhexidine rinse,^{9,34} or locally delivered antibiotics^{41,43} were applied in some studies. The follow-up period ranged from 6^{18,39,44,46} to 60¹⁶ months in studies that used the surgical approach and 6 to 12 months in those with the non-surgical approach.^{15,34–37,40–43,45,47–49}

Meta-analyses conducted in the current study only include CCTs and RCTs with data comparing the clinical parameters between groups with and without adjunctive laser treatment. For non-surgical treatment, all included studies had a follow-up period of 6 to 12 months; therefore, only short-term clinical outcomes could be analyzed. For surgical groups, two studies^{9,16} reported data after a 48-month follow-up period; therefore, short-term and long-term outcomes were analyzed separately. Two articles by Schwarz et al.^{9,38} were follow-up studies of an early study.¹⁸ Similarly, Renvert et al.³⁷ and Persson et al.³⁶ retrieved data from the same population. Therefore, repeated data were not included in meta-analyses. Most studies^{9,15–18,34–49} reported a PD and BOP reduction with CAL gain when defects were treated surgically or non-surgically. Higher mean PD reduction was



TABLE 1 Features of the Included Articles Using a Non-Surgical Approach

Author/ year	Study design	Groups	Type	Laser settings	Patients (n)	Implants (n)	Follow-up (months)	Diagnosis	Treatment outcomes (difference; mean \pm SD)	Conclusions					
								Mucositis/peri- implantitis	PD (mm)	CAL (mm)	MBL (mm)	BOP (%)	PI	REC (mm)	
Schwarz et al. 2005 ^{34a}	RCT	T: Laser C: MI (plastic curets + CHX)	Er:YAG	2,940 nm, 12.7 J/cm ²	T: 10 C: 10	T: 16 C: 14	6	Peri-implantitis	T: 0.8 \pm 0.83 C: 0.7 \pm 1.05	T: 0.7 \pm 0.65 C: 0.6 \pm 1.05		T: 52.13 \pm 10.19 C: 21.57 \pm 14.01	T: 0 \pm 0.33 C: 0 \pm 0.36	T: -0.1 \pm 0.43 C: -0.1 \pm 0.58 (increase)	Laser group resulted in a statistically significant higher reduction of BOP than control.
Schwarz et al. 2006 ³⁵	CS	Laser	Er:YAG	2,940 nm, 12.7 J/cm ²	n = 12 (data of only eight patients extracted)	n = 12 (data of only eight implants extracted)	24 (only 6-month data retrieved)	Peri-implantitis	1.86 \pm 0.61	0.87 \pm 0.71		68.63 \pm 12.85	0.13 \pm 0.16	0.24 \pm 0.39 (increase)	Single course of non-surgical treatment of peri-implantitis using Er:YAG laser was not sufficient for the maintenance of failing implants.
Persson et al. 2011 ³⁶	RCT	T: Laser C: Air-abrasive	Er:YAG	100 ml/pulse and 10 Hz (12.7 J/cm ²)	T: 21 C: 21	T: 55 C: 45	6	Peri-implantitis	T: 0.9 \pm 0.8 C: 0.8 \pm 0.5					Both methods failed to reduce bacterial counts. Clinical improvements were limited.	
Renvert et al. 2011 ^{37a}	RCT	T: Laser C: Air-abrasive	Er:YAG	100 ml/pulse and 10 Hz (12.7 J/cm ²)	T: 21 C: 21	T: 55 C: 45	6	Peri-implantitis	T: 0.8 \pm 0.5 C: 0.9 \pm 0.8	T: -0.3 \pm 0.9 C: -0.1 \pm 0.8				Clinical improvement was similar between treatments using an Er:YAG laser or air-abrasive for debridement.	
Deppe et al. 2013 ⁴⁰	Pilot	1: Air-abrasive + laser 2: severe bone loss	Diode	660 nm, 100 mW with 60 mW/cm ²	T: 16 C: 16	T: 18 C: 18	6	Peri-implantitis	1: 0.4 \pm 0.63 2: -0.7 \pm 0.91 (increased PD)	1: 1.4 \pm 0.97 2: -0.2 \pm 0.75 (CAL loss)	1: -0.3 \pm 0.75 2: 1.9 \pm 0.81		1: 0.2 \pm 0.43 2: 0.7 \pm 1.29	Non-surgical diode laser treatment could stop bone resorption in moderate peri-implant defects but not in severe defects.	
Schär et al. 2013 ^{41a}	RCT	T: MI + air polishing + H ₂ O ₂ + laser C: MI (titanium curets) + air polishing + minocycline	Diode	660 nm, 100 mW	T: 20 C: 20	T: 20 C: 20	6	Peri-implantitis	T: 0.36 \pm 0.36 C: 0.49 \pm 0.50	T: 0.16 \pm 0.48 C: 0.19 \pm 0.44		BOP- positive sites T: 4.03 \pm 1.66 to 1.51 \pm 1.41 C: 4.41 \pm 1.47 to 2.10 \pm 1.55	T: 0.13 \pm 0.10 C: 0.18 \pm 0.14	T: 0.20 \pm 0.58 C: 0.30 \pm 0.66	Both modalities had comparable reductions in inflammation and PD. Complete resolution of inflammation was not achieved with either therapy.

(Continues)



TABLE 1 (Continued)

Author/ Study year	design	Groups	Type	Laser settings	Patients (n)	Implants (n)	Follow-up (months)	Diagnosis	Treatment outcomes (difference; mean ± SD)	Conclusions					
								Mucositis/peri-implantitis	PD (mm)	CAL (mm)	MBL (mm)	BOP (%)	PI	REC (mm)	
Thierbach and Eger 2013 ⁴²	CS	MI (plastic curets) + laser W: with pus WO: without pus	Diode	660 nm × 10 seconds	28	W: 17 WO: 33	7	Peri-implantitis	W: 0.43 ± 0.1 WO: 1.13 ± 0.17	W: 1.79 ± 0.36 WO: 1.3 ± 0.93		W: 85.91 ± 11.53 WO: 54.98 ± 28.19			Non-surgical treatment was effective in managing peri-implantitis without pus formation. Peri-implantitis with pus formation required a combination of surgical and regenerative procedures.
Bassetti et al. 2014 ^{43a}	RCT	T: MI + air-polishing + laser C: MI (titanium curets) + air-polishing + minocycline	Diode	660 nm, 100 mW × 10 seconds	T: 20 C: 20	T: 20 C: 20	12	Peri-implantitis	T: 0.11 ± 0.44 C: 0.56 ± 0.52	T: 0.08 ± 0.54 C: 0.31 ± 0.45		T: 2.29 ± 0.97 C: 2.86 ± 0.88	T: 0.12 ± 0.10 C: 0.21 ± 0.09	T: 0.03 ± 0.57 C: 0.27 ± 0.71 (decrease)	Non-surgical mechanical debridement with adjunctive diode laser was equally effective in the reduction of inflammation as adjunctive delivery of minocycline microspheres.
Ansan et al. 2015 ^{45a}	RCT	T: MI + laser C: MI (plastic curets)	Diode	810 nm, 1W pulsed mode with 3 J/cm ² × 1 minute	10	T: 24 C: 24	6	Peri-implantitis	T: 0.17 ± 0.41 C: 0.21 ± 0.24	T: -0.66 ± 0.28 C: -0.28 ± 0.32		T: 100% to 95.8% C: 100% to 100%	T: 91.7% to 54.2% C: 91.7% to 41.7%		Diode laser did not yield any additional positive influence on the peri-implant healing compared with conventional scaling alone.
Mettraux et al. 2016 ¹⁵	CS	MI (carbon fiber curets) + laser	Diode	810 nm, 2.5 W, 50 Hz, 30s × 3	15	23	24	Peri-implantitis	Deepest PD buccal: 7.5 ± 2.6-3.6 ± 0.7 Deepest PD oral: 7.7 ± 2.1-3.8 ± 0.9			% of implants with ≥1 site with BOP: 100% to 43%			Non-surgical treatment of peri-implantitis with the use of diode laser resulted in significant clinical improvements for at least 2 years.
Schwarz et al. 2015 ⁴⁷	CS	M: peri-implant mucositis, MI (carbon curets)+CHX P: Peri-implantitis, laser	Er:YAG	2.94 × 1 minute, 12.7 J/cm ²	M: 17 P: 17	M: 24 P: 21	6	Both	P: 1.0 ± 0.9 M: 0.1 ± 0.6			P: 27.9 ± 18.5 M: 38.0 ± 23.1			Non-surgical treatment of peri-implant mucositis using MI + CHX or peri-implantitis using laser was associated with significant short-term clinical improvements.

(Continues)



TABLE 1 (Continued)

Author/ year	Study design	Groups	Type	Laser settings	Patients (n)	Implants (n)	Follow-up (months)	Diagnosis Mucositis/peri- implantitis	PD (mm)	CAL (mm)	MBL (mm)	BOP (%)	PI	REC (mm)	Conclusions
Al Amri et al. 2016 ^{48a}	CCT	T: MI + laser C: MI (ultrasonic scaler)	Diode	660 nm, 100 mW	T: 34 C: 33	T: 34 C: 33	12	Both	T: 15.8 ± 1.27 C: 15.2 ± 1.05		T: -0.1 ± 0.1 C: 0.0 ± 0.21	T: 33.9 ± 4.87 C: 21.2 ± 3.3			In patients with type 2 diabetes, MI with adjunctive diode laser treatment was more effective compared with MI alone.
Lerario et al. 2016 ^{49a}	CCT	T: MI + laser C: MI (ultrasonic scaler and titanium curets)	Diode	810 nm, 1 W x 30 seconds	T: 21 C: 6	T: 101 C: 24	12	Both	T: 2.66 ± 1.07 C: 0.94 ± 1.13			T: 85.14 C: 27.78			Diode laser seemed to be a valuable tool in the treatment of mucositis and peri-implantitis. Significant PD and BOP reduction was observed.

T = test; C = control; MI = mechanical instrumentation; CHX = chlorhexidine rinse; Er:YAG = erbium:yttrium-aluminum-garnet.

^a Article included in meta-analyses.



TABLE 2 Features of the Included Articles Using a Surgical Approach

Author/ year	Study design	Groups	Type	Laser settings	Patients (n)	Implants (n)	Follow-up (months)	Diagnosis Mucositis/peri- implantitis	Treatment outcomes (difference; mean ± SD)					Conclusions		
									PD (mm)	CAL (mm)	MBL (mm)	BOP (%)	PI		REC (mm)	
Haas et al. 2000 ³³	CS	GBR + laser	Diode	906 nm	17	24	mean 9.5	Peri-implantitis	2.0 ± 1.9							Photosensitizing treatment and GBR resulted in significant reduction of the peri-implant bone defect.
Deppe et al. 2007 ^{16a}	CCT	T1: air abrasive + resective + laser T2: air abrasive + GBR + laser C1: air abrasive + resective/GBR C2: air abrasive + GBR	CO ₂	2.5W, 175 J/cm ² , 12 × 5 seconds	T1: 10 T2: 9 C1: 6 C2: 7	T1: 22 T2: 17 C1: 19 C2: 15	60	Peri-implantitis	T1: 2.3 ± 0.9 T2: 3.2 ± 1.1 C1: 1.9 ± 1.1 C2: 2.6 ± 1.2	T1: 0.6 ± 0.8 T2: 3.2 ± 0.9 C1: 0.3 ± 0.8 C2: 3.0 ± 0.8	T1: 0.6 ± 1.3 T2: 3.1 ± 0.9 C1: -0.1 ± 1.0 C2: 2.7 ± 0.9	T1: 0.4 ± 0.7 T2: 1.4 ± 0.7 C1: 1.0 ± 0.7 C2: 0.3 ± 0.9				CO ₂ laser decontamination might be more efficacious than conventional decontamination in deep and narrow bony defects.
Romanos and Nentwig 2008 ¹⁷	CS	MI (titanium curets) + GBR + laser	CO ₂	2.84W, 1 minute	15	19	45	Peri-implantitis	3.52 ± 0.49							CO ₂ laser decontamination in combination with GBR could be effective in treating peri-implantitis.
Schwarz et al. 2011 ^{18a}	RCT	T: MI + GBR + laser C: MI (plastic curets) + GBR	Er:YAG	11.4 J/cm ² , 10 pulses per second	T: 15 C: 15	T: 19 C: 16	6	Peri-implantitis	T: 1.7 ± 1.4 C: 2.4 ± 1.5	T: 1.5 ± 1.4 C: 2.2 ± 1.4	T: 47.8 ± 35.5 C: 55.0 ± 31.1	T: 0.4 ± 0.5 C: 0.5 ± 0.6	T: -0.2 ± 0.2 C: -0.2 ± 0.3			Laser treatment did not have significant benefit on surface decontamination.
Schwarz et al. 2012 ³⁸	RCT	T: MI + GBR + laser C: MI (plastic curets) + GBR	Er:YAG	11.4 J/cm ² , 10 pulses per second	T: 10 C: 14	T: 10 C: 14	24	Peri-implantitis	T: 1.1 ± 2.2 C: 1.5 ± 2.0	T: 1.0 ± 2.2 C: 1.2 ± 2.2	T: 75.0 ± 32.6 C: 54.9 ± 30.3	T: 0.2 ± 0.6 C: 0.0 ± 0.8	T: -0.1 ± 0.4 C: -0.3 ± 0.6			Treatment outcomes following surgical therapy of peri-implantitis might be influenced by factors other than the method of surface decontamination.

(Continues)



TABLE 2 (Continued)

Author/ year	Study design	Groups	Type	Laser settings	Patients (n)	Implants (n)	Follow-up (months)	Diagnosis Mucositis/peri- implantitis	Treatment outcomes (difference; mean \pm SD)					Conclusions	
									PD (mm)	CAL (mm)	MBL (mm)	BOP (%)	PI		REC (mm)
Schwarz et al. 2013 ^{9a}	RCT	T: MI + GBR + laser C: MI (plastic cure) + GBR	Er:YAG	11.4 J/cm ² , 10 pulses per second	T: 7 C: 10	T: 9 C: 12	48	Peri-implantitis	T: 1.3 \pm 1.8 C: 1.2 \pm 1.9	T: 1.2 \pm 2.0 C: 1.5 \pm 2.0	MBL (mm)	T: 71.6 \pm 24.9 C: 85.2 \pm 16.4	T: 0.4 \pm 0.7 C: 0.0 \pm 1.1	T: -0.1 \pm 0.3 C: -0.3 \pm 0.9	Treatment outcomes following surgical therapy were not influenced by factors other than the method of surface decontamination.
Schwarz et al. 2014 ^{4a}	CS	Laser + MI (plastic curets) + implanto- plasty + GBR + CTG	Er:YAG	11.4 J/cm ²	10	13	6	Peri-implantitis	2.53 \pm 1.80	2.07 \pm 1.93		74.4 \pm 28.5	0.23 \pm 0.59	0.46 \pm 0.77	Regenerative and resective surgical therapy with soft tissue grafts might control peri-implantitis lesions without compromising the esthetic outcome.
Bombeckari et al. 2013 ^{39a}	RCT	T: MI + laser C: MI (plastic scaler) + CHX	Diode	810 nm, 1W, 20 seconds \times 5	T: 20 C: 20	T: 20 C: 20	6	Peri-implantitis	T: 1.0 \pm 0.31 C: 0.30 \pm 0.52	T: 0.54 \pm 0.09 C: 0.10 \pm 0.01		T: 60 \pm 50 C: 30 \pm 48		A significantly lower proinflammatory index of peri-implantitis was observed in the laser group at 6 months of follow-up.	
Papadopoulos et al. 2015 ⁴⁶	RCT	T: MI + laser C: MI (plastic curets)	Diode	980 nm, 0.8W, 3 times	T: 8 C: 8	T: 8 C: 8	6	Peri-implantitis	T: 1.48 C: 1.21	T: 0.65 C: 0.17		T: 57.4 C: 62.5	T: 9.6 C: 16.7	The adjunctive use of the diode laser was not beneficial in the surgical management of peri-implantitis.	

GBR = guided bone regeneration; T = test; C = control; MI = mechanical instrumentation; CO₂ = carbon dioxide; Er:YAG = erbium:yttrium-aluminum-garnet; CTG = connective tissue graft; CHX = chlorhexidine.
^aArticle included in meta-analyses.

generally achieved in the group augmented with bone grafts and membranes. Slight MBL loss was reported in some non-surgical treatment groups using lasers.^{37,45,49} When compared with mechanical debridement and antiseptics in combination with a surgical approach, the addition of laser treatment showed slight-to-no benefit in PD/BOP reduction and CAL gain. In the non-surgical treatment group, the addition of lasers showed significant reduction of BOP compared with the non-laser treatment group. However, a slight but significant MBL loss was also detected in the laser treatment group.

2.2 | Short-term (≤ 12 months) outcomes of non-surgical treatment in combination with lasers

Of the included studies, six articles^{34,37,41,43,45,49} were selected and pooled in the meta-analyses. These six studies presented a follow-up period of 6 to 12 months, of which four studies^{41,43,45,49} used the diode laser and another two studies^{34,37} used the Er:YAG laser.

A total of six articles^{34,37,41,43,45,49} were pooled to evaluate PD reduction. The results presented WMD of 0.24 mm (95% CI = -0.38 to 0.85 mm, *P* = 0.45, four studies were included^{41,43,45,49}), -0.07 mm (95% CI = -0.32 to 0.18 mm, *P* = 0.57, two studies were included^{34,37}), and 0.15 mm (95% CI = -0.28 to 0.57 mm, *P* = 0.50) for diode laser, Er:YAG laser, and overall comparison, respectively. No statistical significance was found (Figure 2A) for any of the comparison between groups. The comparisons presented a low heterogeneity for Er:YAG laser (*P* = 0.59) and high heterogeneity for diode laser (*P* < 0.001) and overall comparisons (*P* < 0.001).

In terms of CAL gain, three articles^{34,41,43} were analyzed. The results presented WMD of -0.12 mm (95% CI = -0.33 to 0.09 mm, *P* = 0.25, two studies were included^{41,43}), 0.10 mm (95% CI = -0.54 to 0.74 mm, *P* = 0.76, only one study was included³⁴), and -0.10 mm (95% CI = -0.30 to 0.10 mm, *P* = 0.32) for diode laser, Er:YAG laser, and overall comparison, respectively. No statistical significance was found (Figure 2B) for any comparisons between groups. The comparisons presented low heterogeneity for diode laser (*P* = 0.35) and overall comparisons (*P* = 0.52).

Pertaining to BOP reduction, two articles were analyzed.^{34,48} The results presented WMD of 12.70% (95% CI = 10.71% to 14.69%, *P* < 0.001, one study⁴⁸), 30.56% (95% CI = 21.68% to 39.44%, *P* < 0.001, one study³⁴), and 21.08% (95% CI = 3.61% to 38.55%, *P* = 0.02) for diode laser, Er:YAG laser, and overall comparison, respectively. Statistical significance was found, favoring laser treatment group (Figure 2C). However, the comparison presented a high heterogeneity for overall comparison (*P* = 0.001).

Regarding PI reduction, three^{34,41,43} articles were analyzed. The results presented WMD of -0.07 (95% CI = -0.12




FIGURE 2 Meta-analyses of clinical parameters in non-surgical approach between groups with adjunctive laser therapy (test) and without laser therapy (control) among selected studies. A) Comparison of PD reduction; B) comparison of CAL gain; C) comparison of BOP reduction; D) comparison of PI; E) comparison of MBL loss; F) comparison of REC increase. IV = independent variable

to -0.03 , $P = 0.002$, two studies included^{41,43}), 0.00 (95% CI = -0.25 to 0.25 , $P > 0.99$, one study³⁴), and -0.07 (95% CI = -0.12 to -0.03 , $P = 0.002$) for diode laser, Er:YAG laser, and overall comparisons, respectively. Statistical significance was found for diode laser and overall comparisons, favoring non-laser treatment group (Figure 2D). The comparisons presented a low heterogeneity for diode laser ($P = 0.41$) and overall comparison ($P = 0.60$).

The outcome of MBL was reported in three articles.^{37,45,48} The results presented WMD of -0.23 mm (95% CI = -0.50 to 0.04 mm, $P = 0.10$, two studies were included^{45,48}), -0.20 mm (95% CI = -0.53 to 0.13 mm, $P = 0.24$, one study³⁷), and -0.22 mm (95% CI = -0.43 to -0.01 , $P = 0.04$) for diode laser, Er:YAG laser, and overall comparisons, respectively. Statistical significance was found for diode laser and overall comparisons, favoring control group (Figure 2E). The comparisons presented a high heterogeneity for diode laser ($P = 0.003$) and overall comparison ($P = 0.01$).

In terms of REC, three articles were analyzed.^{34,41,43} The results presented WMD of -0.17 mm (95% CI = -0.44 to 0.11 mm, $P = 0.24$, two studies were included^{41,43}), 0.00 mm (95% CI = -0.37 to 0.37 mm, $P > 0.99$, one study³⁴), and -0.11 mm (95% CI = -0.33 to 0.11 mm, $P = 0.34$) for diode laser, Er:YAG laser, and overall comparison, respectively. No statistical significance was found (Figure 2F) for any of the comparison between groups. The comparisons presented a low heterogeneity for diode laser ($P = 0.62$) and overall comparisons ($P = 0.69$).

2.3 | Short-term (≤ 12 months) outcomes of surgical treatment in combination with lasers

Of the included studies, two articles^{18,39} were selected and pooled in meta-analyses. Both articles reported treatment outcomes after a 6-month follow-up.

For PD reduction, an overall WMD of 0.08 mm (95% CI = -1.28 to 1.44 mm) between laser treatment group and conventional treatment group with no statistical significance ($P = 0.91$) was found (Figure 3A). The comparisons presented a high heterogeneity among selected studies (P value for χ^2 test = 0.006 and I^2 test = 87%).

In terms of CAL gain, an overall WMD of -0.03 mm (95% CI = -1.13 to 1.07 mm), with no statistical significance ($P = 0.96$) was found (Figure 3B). The comparisons presented a high heterogeneity among selected studies (P value for χ^2 test = 0.02 and I^2 test = 83%).

An overall WMD of 9.88% (95% CI = -26.46% to 46.21%), with no statistical significance ($P = 0.59$) with regard to BOP reduction was found (Figure 3C). The comparisons presented a moderate-to-high heterogeneity among selected studies (P value for χ^2 test = 0.05 and I^2 test = 73%).

Since there was only one comparative study¹⁸ reporting differences in PI and REC between test and control groups, the meta-analyses could not be performed. Based on this article, mean difference was -0.10 (95% CI = -0.47 to 0.27) for PI and 0.0 mm (95% CI = -0.17 to 0.17 mm) for REC, with no detectable statistical significance ($P = 0.60$ for PI; $P > 0.99$ for REC) between groups. In addition, no short-term studies reported outcomes of MBL change; therefore, short-term MBL change between groups could not be evaluated.

2.4 | Long-term (≥ 48 months) outcomes of surgical treatment in combination with lasers

Among the included studies, two articles^{9,16} were selected and pooled in the meta-analyses to evaluate the long-term treatment outcomes. One study⁹ was a 4-year follow-up of a previous study.¹⁸ Another study, by Deppe et al.,¹⁶ reported treatment outcomes after a 60-month follow-up.

An overall WMD of 0.45 mm (95% CI = -0.10 to 1.00 mm) between laser treatment group and conventional treatment group, with no statistical significance ($P = 0.11$) was found for PD reduction (Figure 4A). The comparisons presented a low heterogeneity among selected studies (P value for χ^2 test = 0.64 and I^2 test = 0%).

In terms of CAL gain, an overall WMD of 0.22 mm (95% CI = -0.52 to 0.95 mm), with no statistical significance ($P = 0.56$) was found (Figure 4B). The comparisons presented a low heterogeneity among selected studies (P value for χ^2 test = 0.52 and I^2 test = 0%).

Pertaining to BOP reduction, an overall WMD of 7.26% (95% CI = -38.77% to 53.29%), with no statistical significance ($P = 0.76$) was found (Figure 4C). The comparisons presented a high heterogeneity among selected studies (P value for χ^2 test = 0.03 and I^2 test = 80%).

An overall WMD of -0.09 (95% CI = -0.95 to 0.77), with no statistical significance ($P = 0.84$) was found for PI reduction (Figure 4D). The comparisons presented a moderate heterogeneity among selected studies (P value for χ^2 test = 0.06 and I^2 test = 71%).

Only one comparative study reported MBL¹⁶ differences between test and control groups, likewise for differences in REC.⁹ As a result, the meta-analysis could not be performed. Based on these studies, mean difference of MBL and REC was 0.64 mm (95% CI = -0.24 to 1.52 mm) and 0.20 mm (95% CI = -0.35 to 0.75 mm), respectively, with no statistical significance ($P = 0.15$ for MBL and $P = 0.47$ for REC) detected between groups.

2.5 | Risk of bias assessment

The results of the risk of bias assessment for included RCTs are summarized in supplementary Table 1 in the online *Journal of Periodontology*. Four studies^{34,36,37,41}




FIGURE 3 Meta-analyses for the comparison of clinical parameters in surgical approach between groups with adjunctive laser therapy (test) and without laser therapy (control) among selected studies after a short follow-up period of 6 to 12 months. A) comparison of PD reduction; B) comparison of CAL gain; C) comparison of BOP reduction. IV = independent variable

had low risk of bias, three studies^{18,39,46} had moderate risk of bias, and three studies^{9,43,45} had high risk of bias.

2.6 | Level of evidence assessment

The results of level of evidence assessment are reported in supplementary Table 2 in the online *Journal of Periodon-*

tology. No evidence is currently available to support the use of lasers in treatment of peri-implant mucositis. As for non-surgical treatment of peri-implantitis, some evidence presented controversial clinical benefits of adjunct laser treatment in the short term, but no evidence was found to support the long-term benefits. As for surgical treatment of peri-implantitis, limited evidence presented controversial short-term clinical benefits and no long-term benefits.




FIGURE 4 Meta-analyses for the comparison of clinical parameters in surgical approach between groups with adjunctive laser therapy (test) and without laser therapy (control) among selected studies after a long-term follow-up period of 48 to 60 months. A) comparison of PD reduction; B) comparison of CAL gain; C) comparison of BOP reduction; D) comparison of PI. IV = independent variable

3 | DISCUSSION

While previous studies^{51,52} suggested that peri-implant mucositis might be successfully treated if detected early and

when combined with non-surgical treatment, a complete resolution of peri-implant inflammation was not commonly obtained.⁵³ Therefore, the current review aims to identify the potential benefit of adjunctive laser therapy in the

treatment of peri-implant mucositis and peri-implantitis. However, among the selected 22 human clinical trials, only three studies^{47–49} included patients with peri-implant mucositis; other studies only selected patients with peri-implantitis. In a study by Schwarz et al.,⁴⁷ carbon curets and chlorhexidine rinse for surface detoxification were used to treat peri-implant mucositis, while Er:YAG laser was used to treat peri-implantitis. Although both groups achieved short-term significant clinical improvements, a complete disease resolution was not achieved in the majority of the study patients. Another two studies^{48,49} compared the effect of mechanical debridement with or without adjunctive diode laser treatment on clinical outcomes of peri-implant diseases and reported significant reduction of PD and BOP if the laser was used. However, since the disease status (peri-implant mucositis or peri-implantitis) of the participants was not clearly demarcated in the analysis of the treatment outcomes, the evidence supporting the use of lasers to treat peri-implant mucositis was limited. Currently the efficacy of non-surgical treatment of peri-implant mucositis with or without using lasers could not be warranted, and more RCTs should be conducted in the future to investigate this topic.

Pertaining to treatment of peri-implantitis, the current review and meta-analyses failed to detect significant PD reduction and CAL gain when lasers were used together with non-surgical and surgical therapies. This result was consistent with previously published systematic reviews.^{19,54} However, the current review identified two recent articles^{39,46} that introduced the diode laser in the surgical management of peri-implantitis. While both studies demonstrated promising outcomes in PD reduction and CAL gain when the laser was used versus mechanical debridement only, the WMD presented less than 1 mm for both parameters. Therefore, the adjunctive use of diode laser in surgical treatment of peri-implantitis appeared to be effective; however, its efficacy still required further investigations.

When considering the benefit of GBR procedures in treating peri-implant defects, most of the defects that had GBR achieved better outcomes in terms of PD reduction, CAL gain, and MBL gain.^{9,16–18,33,38,44} However, an addition of lasers to GBR procedures did not present further improvement to the peri-implant parameters.^{9,16,18,38} Therefore, GBR procedures should be considered as a standard treatment modality irrespective of the use of lasers when dealing with intrabony peri-implant defects.⁴⁴

Peri-implant tissue recession after surgical therapy with and without adjunctive laser treatment was only reported from a series of publications by Schwarz et al.^{9,18,38} In their studies, the same population was followed up to 48 months, and slight recession, ranging from 0.1 to 0.3 mm, was observed. For the intervention, no significant change in soft tissue level (ranging from 0.03 to 0.3 mm) was observed in studies^{34,41,43} that evaluated non-surgical therapy with and without laser. Therefore,

the current evidence revealed that adjunctive laser treatment combined with surgical/non-surgical therapies did not interfere with peri-implant soft tissue levels.

Conflicting data on BOP reduction was reported in selected studies.^{9,18,34,38,39,46,48} Some studies reported significant BOP reduction when lasers were introduced.^{34,39,48} On the contrary, other studies^{9,18,38,46} did not find significant BOP reduction as compared with conventional treatment. It appeared that the adjunctive use of lasers with non-surgical therapy might decrease BOP after a short follow-up period of 6 to 12 months. This could be explained by the coagulation or vaporization of the tissues after laser treatment, and reflects the truth that a long-term control of tissue inflammation might be related to maintenance protocol instead of active treatment. However, when adjunctive lasers were used in surgical treatment, there was no significant BOP reduction detected.^{9,16} Interestingly, previous studies have reported a higher percentage of BOP around dental implants despite lower plaque scores and fewer signs of inflammation.^{21,55} Also, higher sensitivity of probing around dental implants compared with probing around teeth has been reported;⁵⁶ therefore, clinicians should be aware that the presence of BOP might not accurately represent the inflammatory status of the peri-implant tissues. Also, absence of BOP has been reported as a good indicator of healthy peri-implant mucosa, but presence of BOP might have limited diagnostic value.⁵⁷ As such, clinicians should evaluate this parameter cautiously and not apply the absence of BOP as the only criterion of treatment outcomes.

The results of this meta-analysis showed no additional benefit in PI reduction when lasers were used as an adjunct to conventional intervention. Interestingly most data reported a minimal difference in PI (a change of < 1) between baseline and the final follow-up appointment in groups with and without laser treatment. Therefore, it could be assumed that the outcome of PI reduction might not be significantly related to treatment modalities. Other factors, such as maintenance protocol⁵⁸ or the use of locally delivered antibiotics,^{41,43} might have a larger impact on PI reduction.

Most studies that evaluated surgical interventions used GBR in the treatment of both test and control groups.^{9,16–18,33,38,44} Therefore, MBL could not be accurately evaluated since this difference might not be related to the adjunctive use of laser therapy. When performing non-surgical treatment, a statistically significant MBL loss was detected (-0.22 mm with 95% CI = -0.43 to -0.01 , $P = 0.04$) in the laser-treated group compared with the control group. This might be a result of uncontrolled temperature increase in the laser-treated area, thus further jeopardizing the healing outcomes.^{45,59} However, when interpreting this result, clinicians should be aware that only three studies^{37,45,48} were pooled in the meta-analyses and this difference was minimal in terms of clinical significance. Nonetheless, after repeated application of a diode laser, Mettraux et al.¹⁵ reported sound



improvement in clinical parameters for implants affected by peri-implantitis. Results from this study opened a new area of future research to investigate the need for repeated laser applications to decontaminate an implant surface.

Comparisons of bacterial profile between treatment with and without adjunctive laser therapy were recorded in six studies.^{28,29,36,39,43,45} Among these six studies, three studies^{28,29,39} introduced surgical approaches, while the remaining three studies had non-surgical treatment.^{36,43,45} Bach et al.,²⁸ in an early study, reported that the elimination of certain pathogenic bacteria with diode laser could be maintained at the 60-month follow-up in 10 of 15 patients. Moreover, significant reduction of black-pigmented, Gram-negative anaerobic bacteria was noted throughout the study period. Similarly, Dörtbudak et al.²⁹ used a diode laser in combination with surgical treatment to evaluate reduction of bacteria counts in patients with peri-implantitis. The results of their study showed that the combined treatment significantly reduced the bacterial counts of *Aggregatibacter actinomycetemcomitans*, *Porphyromonas gingivalis*, and *Prevotella intermedia*. However, this study lacked a control group, and complete elimination of bacteria was never achieved. Recently, Bombeccari et al.³⁹ concluded that a diode laser, when used as an adjunct to surgical treatment, was able to reduce the bacterial biofilm by 95.2% colony-forming units (CFUs) per milliliter compared to 80.85% CFU reduction in conventional treatment group. As for non-surgical intervention, all three studies^{37,43,45} that introduced laser therapy as an adjunct reported no additional benefit on the peri-implant microbiota compared with conventional mechanical debridement after a 6- to 12-month follow-up period. Interestingly, it has been reported that results of microbiologic testing are often inconsistent due to different laboratory processing,^{60,61} thus this information should always be interpreted cautiously. To date, there are very few clinical trials that have assessed the efficacy of various laser treatments on microbiologic outcomes; therefore, a firm conclusion cannot be drawn.

There were a number of limitations in the current systematic review and meta-analyses: 1) only 11 papers with comparable data were included in the meta-analyses; 2) a few analyses presented with high heterogeneity (this heterogeneity was related to the presence of confounding factors within and among the selected studies, for example, different study designs, follow-up periods, various lasers settings, etc.); 3) owing to the limited available data, patient-centered outcome measures, cost-effectiveness of lasers, and microbiologic data were not analyzed in the current review; 4) the systematic review only included studies written in English, which could result in a selection bias; 5) because peri-implant diseases are multifactorial and affected by numerous local and systemic factors, further investigations should evaluate the influence of both implant- and host-related factors as they both play a major role in treatment outcomes; and 6) the various

definitions of peri-implant diseases as well as different measurement methods of clinical parameters might also influence the presented results. Moreover, the main parameter used for evaluation of disease status (PD) is subject to multiple variables that might affect its accuracy.

4 | CONCLUSIONS

Current evidence allowed for analysis of only Er:YAG, CO₂, and diode lasers. Studies on others failed to have controlled evidence to support their evaluation. Since the types of lasers analyzed have different modes of action, the limited number of included studies and patients/implants evaluated makes it difficult to warrant their therapeutic values. Data on adjunctive laser treatment for peri-implant mucositis are scarce. Therefore, future clinical trials are needed to evaluate the potential benefit of this approach. Based on the results of meta-analyses, when treating peri-implantitis surgically, no differences in PD reduction, CAL gain, amount of REC, and PI reduction were found between groups with and without adjunctive laser treatment. However, controversial results have been reported in the literature. Also based on the results of meta-analyses, when treating peri-implantitis non-surgically, adjunctive laser treatment might result in more BOP reduction in the short term. However, no long-term data were available to warrant this benefit. Limited evidence showed that non-surgical treatment with adjunctive laser therapy might result in slightly more MBL loss compared with conventional treatment and also showed a potential reduction in dark pigmented, gram-negative anaerobic bacteria when applying adjunctive diode laser therapy to surgical treatment of peri-implantitis. However, this benefit was not detected in non-surgical treatment with adjunctive laser therapy.

ACKNOWLEDGMENTS AND DISCLOSURES

The American Academy of Periodontology Best Evidence Consensus meeting on lasers was sponsored by J. Morita USA (Irvine, California). The authors thank Dr. Jia-Hui Fu (Department of Periodontics, National University of Singapore, Singapore) for her assistance in proofreading this manuscript. The authors report no conflicts of interest related to this review.

REFERENCES

1. Midda M, Renton-Harper P. Lasers in dentistry. *Br Dent J.* 1991;170:343–346.
2. Midda M. The use of lasers in periodontology. *Curr Opin Dent.* 1992;2:104–108.
3. Midda M. Lasers in periodontics. *Periodontal Clin Investig.* 1992;14:14–20.
4. Cobb CM. Lasers in periodontics: a review of the literature. *J Periodontol.* 2006;77:545–564.



5. Behdin S, Monje A, Lin GH, Edwards B, Othman A, Wang HL. Effectiveness of laser application for periodontal surgical therapy: systematic review and meta-analysis. *J Periodontol*. 2015;86:1352–1363.
6. Nguyen NT, Byarlay MR, Reinhardt RA, Marx DB, Meinberg TA, Kaldahl WB. Adjunctive non-surgical therapy of inflamed periodontal pockets during maintenance therapy using diode laser: a randomized clinical trial. *J Periodontol*. 2015;86:1133–1140.
7. Aral CA, Dilber E, Aral K, Sarica Y, Sivrikoz ON. Management of cyclosporine and nifedipine-induced gingival hyperplasia. *J Clin Diagn Res*. 2015;9:ZD12–ZD15.
8. Kumar P, Rattan V, Rai S. Comparative evaluation of healing after gingivectomy with electrocautery and laser. *J Oral Biol Craniofac Res*. 2015;5:69–74.
9. Schwarz F, Hegewald A, John G, Sahm N, Becker J. Four-year follow-up of combined surgical therapy of advanced peri-implantitis evaluating two methods of surface decontamination. *J Clin Periodontol*. 2013;40:962–967.
10. Derks J, Tomasi C. Peri-implant health and disease. A systematic review of current epidemiology. *J Clin Periodontol*. 2015;42(Suppl. 16):S158–S171.
11. Renvert S, Roos-Jansåker AM, Claffey N. Non-surgical treatment of peri-implant mucositis and peri-implantitis: a literature review. *J Clin Periodontol*. 2008;35(Suppl.8):305–315.
12. Heitz-Mayfield LJ, Mombelli A. The therapy of peri-implantitis: a systematic review. *Int J Oral Maxillofac Implants*. 2014;29(Suppl):325–345.
13. Renvert S, Polyzois IN. Clinical approaches to treat peri-implant mucositis and peri-implantitis. *Periodontol 2000*. 2015;68:369–404.
14. Chan HL, Lin GH, Suarez F, MacEachern M, Wang HL. Surgical management of peri-implantitis: a systematic review and meta-analysis of treatment outcomes. *J Periodontol*. 2014;85:1027–1041.
15. Mettraux GR, Sculean A, Bürgin WB, Salvi GE. Two-year clinical outcomes following non-surgical mechanical therapy of peri-implantitis with adjunctive diode laser application. *Clin Oral Implants Res*. 2016;27:845–849.
16. Deppe H, Horch HH, Neff A. Conventional versus CO₂ laser-assisted treatment of peri-implant defects with the concomitant use of pure-phase beta-tricalcium phosphate: a 5-year clinical report. *Int J Oral Maxillofac Implants*. 2007;22:79–86.
17. Romanos GE, Nentwig GH. Regenerative therapy of deep peri-implant infrabony defects after CO₂ laser implant surface decontamination. *Int J Periodontics Restorative Dent*. 2008;28:245–255.
18. Schwarz F, Sahm N, Iglhaut G, Becker J. Impact of the method of surface debridement and decontamination on the clinical outcome following combined surgical therapy of peri-implantitis: a randomized controlled clinical study. *J Clin Periodontol*. 2011;38:276–284.
19. Mailoa J, Lin GH, Chan HL, MacEachern M, Wang HL. Clinical outcomes of using lasers for peri-implantitis surface detoxification: a systematic review and meta-analysis. *J Periodontol*. 2014;85:1194–1202.
20. Heitz-Mayfield LJ, Schmid B, Weigel C, et al. Does excessive occlusal load affect osseointegration? An experimental study in the dog. *Clin Oral Implants Res*. 2004;15:259–268.
21. Karoussis IK, Müller S, Salvi GE, Heitz-Mayfield LJ, Brägger U, Lang NP. Association between periodontal and peri-implant conditions: a 10-year prospective study. *Clin Oral Implants Res*. 2004;15:1–7.
22. Klokkevold PR, Han TJ. How do smoking, diabetes, and periodontitis affect outcomes of implant treatment. *Int J Oral Maxillofac Implants*. 2007;22(Suppl):173–202.
23. Salvi GE, Carollo-Bittel B, Lang NP. Effects of diabetes mellitus on periodontal and peri-implant conditions: update on associations and risks. *J Clin Periodontol*. 2008;35(Suppl.8):398–409.
24. Schardt C, Adams MB, Owens T, Keitz S, Fontelo P. Utilization of the PICO framework to improve searching PubMed for clinical questions. *BMC Med Inform Decis Mak*. 2007;7:16.
25. Moher D, Liberati A, Tetzlaff J, Altman DG, Group PRISMA. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6:e1000097.
26. Higgins JP, Green S. *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available at: <http://www.cochrane-handbook.org>. Accessed June 27th, 2016.
27. Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: updated guidelines for reporting parallel group randomized trials. *Ann Intern Med*. 2010;152:726–732.
28. Bach G, Neckel C, Mall C, Krekeler G. Conventional versus laser-assisted therapy of periimplantitis: a five-year comparative study. *Implant Dent*. 2000;9:247–251.
29. Dörtbudak O, Haas R, Bernhart T, Mailath-Pokorny G. Lethal photosensitization for decontamination of implant surfaces in the treatment of peri-implantitis. *Clin Oral Implants Res*. 2001;12:104–108.
30. De Angelis N Felice P, Grusovin MG, Camurati A, Esposito M. The effectiveness of adjunctive light-activated disinfection (LAD) in the treatment of peri-implantitis: 4-month results from a multicentre pragmatic randomised controlled trial. *Eur J Oral Implantology*. 2012;5:321–331.
31. Eick S, Markauskaite G, Nietzsche S, Laugisch O, Salvi GE, Sculean A. Effect of photoactivated disinfection with a light-emitting diode on bacterial species and biofilms associated with periodontitis and peri-implantitis. *Photodiagn Photodyn Ther*. 2013;10:156–167.
32. Esposito M, Grusovin MG, De Angelis N Camurati A, Campailla M, Felice P. The adjunctive use of light-activated disinfection (LAD) with FotoSan is ineffective in the treatment of peri-implantitis: 1-year results from a multicentre pragmatic randomised controlled trial. *Eur J Oral Implantology*. 2013;6:109–119.
33. Haas R, Baron M, Dörtbudak O, Watzek G. Lethal photosensitization, autogenous bone, and e-PTFE membrane for the treatment of peri-implantitis: preliminary results. *Int J Oral Maxillofac Implants*. 2000;15:374–382.
34. Schwarz F, Sculean A, Rothamel D, Schwenzer K, Georg T, Becker J. Clinical evaluation of an Er:yAG laser for nonsurgical treatment of peri-implantitis: a pilot study. *Clin Oral Implants Res*. 2005;16:44–52.
35. Schwarz F, Bieling K, Bonsmann M, Latz T, Becker J. Non-surgical treatment of moderate and advanced periimplantitis lesions: a controlled clinical study. *Clin Oral Investig*. 2006;10:279–288.



36. Persson GR, Roos-Jansåker AM, Lindahl C, Renvert S. Microbiologic results after non-surgical erbium-doped:yttrium, aluminum, and garnet laser or air-abrasive treatment of peri-implantitis: a randomized clinical trial. *J Periodontol*. 2011;82:1267–1278.
37. Renvert S, Lindahl C, Roos Jansåker AM, Persson GR. Treatment of peri-implantitis using an Er:yAG laser or an air-abrasive device: a randomized clinical trial. *J Clin Periodontol*. 2011;38:65–73.
38. Schwarz F, John G, Mainusch S, Sahm N, Becker J. Combined surgical therapy of peri-implantitis evaluating two methods of surface debridement and decontamination. A two-year clinical follow up report. *J Clin Periodontol*. 2012;39:789–797.
39. Bombeccari GP, Guzzi G, Gualini F, Gualini S, Santoro F, Spadari F. Photodynamic therapy to treat periimplantitis. *Implant Dent*. 2013;22:631–638.
40. Deppe H, Mücke T, Wagenpfeil S, Kesting M, Sculean A. Non-surgical antimicrobial photodynamic therapy in moderate vs severe peri-implant defects: a clinical pilot study. *Quintessence Int*. 2013;44:609–618.
41. Schär D, Ramseier CA, Eick S, Arweiler NB, Sculean A, Salvi GE. Anti-infective therapy of peri-implantitis with adjunctive local drug delivery or photodynamic therapy: six-month outcomes of a prospective randomized clinical trial. *Clin Oral Implants Res*. 2013;24:104–110.
42. Thierbach R, Eger T. Clinical outcome of a nonsurgical and surgical treatment protocol in different types of peri-implantitis: a case series. *Quintessence Int*. 2013;44:137–148.
43. Bassetti M, Schär D, Wicki B, et al. Anti-infective therapy of peri-implantitis with adjunctive local drug delivery or photodynamic therapy: 12-month outcomes of a randomized controlled clinical trial. *Clin Oral Implants Res*. 2014;25:279–287.
44. Schwarz F, Sahm N, Becker J. Combined surgical therapy of advanced peri-implantitis lesions with concomitant soft tissue volume augmentation. A case series. *Clin Oral Implants Res*. 2014;25:132–136.
45. Arısan V, Karabuda ZC, Arıcı SV, Topçuoğlu N, Külekçi G. A randomized clinical trial of an adjunct diode laser application for the nonsurgical treatment of peri-implantitis. *Photomed Laser Surg*. 2015;33:547–554.
46. Papadopoulos CA, Vouros I, Menexes G, Konstantinidis A. The utilization of a diode laser in the surgical treatment of peri-implantitis. A randomized clinical trial. *Clin Oral Investig*. 2015;19:1851–1860.
47. Schwarz F, John G, Hegewald A, Becker J. Non-surgical treatment of peri-implant mucositis and peri-implantitis at zirconia implants: a prospective case series. *J Clin Periodontol*. 2015;8:783–788.
48. Al Amri MD, Kellesarian SV, Ahmed A, Al-Kheraif AA, Romanos GE, Javed F. Efficacy of periimplant mechanical debridement with and without adjunct antimicrobial photodynamic therapy in patients with type 2 diabetes mellitus. *Photodiagn Photodyn Ther*. 2016;14:166–169.
49. Lerario F, Roncati M, Gariffo A, et al. Non-surgical periodontal treatment of peri-implant diseases with the adjunctive use of diode laser: preliminary clinical study. *Lasers Med Sci*. 2016;31:1–6.
50. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33:159–174.
51. Pontoriero R, Tonelli MP, Carnevale G, Mombelli A, Nyman SR, Lang NP. Experimentally induced peri-implant mucositis. A clinical study in humans. *Clin Oral Implants Res*. 1994;5:254–259.
52. Salvi GE, Aglietta M, Eick S, Sculean A, Lang NP, Ramseier CA. Reversibility of experimental peri-implant mucositis compared with experimental gingivitis in humans. *Clin Oral Implants Res*. 2012;23:182–190.
53. Schwarz F, Becker K, Renvert S. Efficacy of air polishing for the non-surgical treatment of peri-implant diseases: a systematic review. *J Clin Periodontol*. 2015;42:951–959.
54. Kotsakis GA, Konstantinidis I, Karoussis IK, Ma X, Chu H. Systematic review and meta-analysis of the effect of various laser wavelengths in the treatment of peri-implantitis. *J Periodontol*. 2014;85:1203–1213.
55. Brägger U, Bürgin WB, Hämmerle CH, Lang NP. Associations between clinical parameters assessed around implants and teeth. *Clin Oral Implants Res*. 1997;8:412–421.
56. Gerber JA, Tan WC, Balmer TE, Salvi GE, Lang NP. Bleeding on probing and pocket probing depth in relation to probing pressure and mucosal health around oral implants. *Clin Oral Implants Res*. 2009;20:75–78.
57. Lang NP, Wilson TG, Corbet EF. Biological complications with dental implants: their prevention, diagnosis and treatment. *Clin Oral Implants Res*. 2000;11(Suppl. 1):146–155.
58. Monje A, Aranda L, Diaz KT, et al. Impact of maintenance therapy for the prevention of peri-implant diseases: a systematic review and meta-analysis. *J Dent Res*. 2016;95:372–379.
59. Fontana CR, Kurachi C, Mendonça CR, Bagnato VS. Temperature variation at soft periodontal and rat bone tissues during a medium-power diode laser exposure. *Photomed Laser Surg*. 2004;22:519–522.
60. Mellado JR, Freedman AL, Salkin LM, Stein MD, Schneider DB, Cutler RH. The clinical relevance of microbiologic testing: a comparative analysis of microbiologic samples secured from the same sites and cultured in two independent laboratories. *Int J Periodontics Restorative Dent*. 2001;21:232–239.
61. Salkin LM, Freedman AL, Mellado JR, Stein MD, Schneider DB, Butler L. The clinical relevance of microbiologic testing. Part 2: a comparative analysis of microbiologic samples secured simultaneously from the same sites and cultured in the same laboratory. *Int J Periodontics Restorative Dent*. 2003;23:121–127.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Lin G-H, López del Amo FS, Wang H-L. Laser therapy for treatment of peri-implant mucositis and peri-implantitis: An American Academy of Periodontology best evidence review. *J Periodontol*. 2018;89:766–782. <https://doi.org/10.1902/jop.2017.160483>