

Zhang Jianfeng (Orcid ID: 0000-0002-3591-3219)
Qin Pengmin (Orcid ID: 0000-0002-3569-6365)
Huang Zirui (Orcid ID: 0000-0002-5949-0206)

Title:

Disrupted neural variability during propofol-induced sedation and unconsciousness

Short title: Propofol disrupts neural variability

Author names and affiliations

Zirui Huang^{b,1,*}, Jun Zhang^{a,1}, Jinsong Wu^{c,*}, Xiaoge Liu^a, Jianghui Xu^a, Jianfeng Zhang^d, Pengmin Qin^e, Rui Dai^f, Zhong Yang^g, Ying Mao^c, Anthony G. Hudetz^b and Georg Northoff^{h,i,j}

^a Department of Anesthesiology, Huashan Hospital, Fudan University, Shanghai, 200040, PR China

^b Department of Anesthesiology and Center for Consciousness Science, University of Michigan, Ann Arbor, 48109, USA

^c Department of Neurosurgery, Huashan Hospital, Fudan University, Shanghai, 200040, PR China

^d College of Biomedical Engineering and Instrument Science, Zhejiang University, Hangzhou, 310000, PR China

^e School of Psychology, South China Normal University, Guangzhou, 510631, PR China

^f State Key Laboratory of Brain and Cognitive Science, Institute of Biophysics, Chinese Academy of Sciences, Beijing, 100101, PR China

^g Department of Radiology, Huashan Hospital, Fudan University, Shanghai, 200040, PR China

^h Institute of Mental Health Research, University of Ottawa, Ottawa, ON, K1Z 7K4, Canada

ⁱ Center for Cognition and Brain Disorders, Hangzhou Normal University, Hangzhou, 311121, PR China

^j Mental Health Centre, Zhejiang University School of Medicine, Hangzhou, 310000, PR China

¹*Authors contributed equally to this work.*

**Corresponding to:*

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/hbm.24304](https://doi.org/10.1002/hbm.24304)

Dr. Zirui Huang, Department of Anesthesiology and Center for Consciousness Science, University of Michigan Medical School. Domino Farms Lby M Ste 3100, 24 Frank Lloyd Wright Dr, Ann Arbor, MI 48105. Email: huangzu@umich.edu.

Dr. Jinsong Wu, Department of Anesthesiology, Huashan Hospital, Fudan University. 12#, Wulumuqi Zhong Road, Shanghai 200040, China. Email: wjsongc@126.com.

Abstract

Variability quenching is a widespread neural phenomenon in which trial-to-trial variability (TTV) of neural activity is reduced by repeated presentations of a sensory stimulus. However, its neural mechanism and functional significance remain poorly understood. Recurrent network dynamics are suggested as a candidate mechanism of TTV, and they play a key role in consciousness. We thus asked whether the variability-quenching phenomenon is related to the level of consciousness. We hypothesized that TTV reduction would be compromised during reduced level of consciousness by propofol anesthetics. We recorded fMRI signals of resting-state and stimulus-induced activities in three conditions: wakefulness, sedation and unconsciousness (i.e. deep anesthesia). We measured the average (trial-to-trial mean, TTM) and variability (TTV) of auditory stimulus-induced activity under the three conditions. We also examined another form of neural variability (temporal variability, TV), which quantifies the overall dynamic range of ongoing neural activity across time, during both the resting-state and the task. We found that 1) TTM decreased gradually from wakefulness through sedation to anesthesia, 2) stimulus-induced TTV reduction normally seen during wakefulness was abolished during both sedation and anesthesia, 3) TV increased in the task state as compared to resting-state during both wakefulness and sedation, but not anesthesia. Together, our results reveal distinct effects of propofol on the two forms of neural variability (TTV and TV). They imply that the anesthetic disrupts recurrent network dynamics, thus prevents the stabilization of cortical activity states. These findings shed new light on the temporal dynamics of neuronal variability and its alteration

during anesthetic-induced unconsciousness.

Keywords: neural variability; trial-to-trial variability; temporal variability; recurrent network; fMRI; sedation; anesthesia; propofol; consciousness

Introduction

A close relationship between spontaneous and stimulus-induced neural activities, based on the assumption of a linear superposition, is a commonly held tenet of the neuroscientific literature [Arieli et al., 1996; Azouz and Gray, 1999; Becker et al., 2011; Fox et al., 2006]. Given this assumption, the variance of the spontaneous and stimulus-induced activities across trials should be additive as a direct consequence of the Law of Total Variance [He, 2013]. Then, one would expect to find an increase of total variability after stimulus onset.

A number of experiments have challenged this view by showing a decrease in trial-to-trial variability, commonly called TTV reduction, after stimulus-onset. TTV reduction has been consistently observed in cellular recordings [Chang et al., 2012; Churchland et al., 2010; Finn et al., 2007; Monier et al., 2003; Scaglione et al., 2011; White et al., 2012], EEG/ECoG [He and Zempel, 2013; Schurger et al., 2015], and fMRI [Ferri et al., 2015; He, 2013; Huang et al., 2017]. These data were collected across a variety of species and brain areas and under a broad range of stimulus conditions suggesting that TTV reduction driven by stimulation or task was a widespread neural phenomenon. These findings also demonstrated that the spontaneous and stimulus-induced activities were not independent but interacting with each. This interaction was thought to be mediated by active noise suppression [White et al., 2012],

characterized as negative interaction [He, 2013] or nonlinear phase-dependence [Huang et al., 2017].

Behaviorally, stronger TTV reduction is associated with faster reaction time [He, 2013], superior perceptual abilities [Arazi et al., 2017; Baria et al., 2017; Schurger et al., 2015] and superior memory recall [Xue et al., 2010]. Therefore, it is plausible to assume that neuronal mechanisms involved in TTV reduction contribute to the disambiguation of environmental signals and enhance the efficiency of cortical information processing [Churchland et al., 2010; Finn et al., 2007; Monier et al., 2003; White et al., 2012].

The mechanism underlying TTV reduction has been explored by a series of neuronal network modeling studies. For instance, the TTV reduction by stimuli can arise from recurrent processing during which the states of cortical activity are stabilized by inputs [Deco and Hugues, 2012; Litwin-Kumar and Doiron, 2012; Mazzucato et al., 2015; Rajan et al., 2010; Sussillo and Abbott, 2009]. Recurrent network dynamics, consisting of distinct feedforward and feedback information flow, have been suggested as a mechanism of consciousness [Boly et al., 2011; Lamme and Roelfsema, 2000; Mashour, 2014; Moutard et al., 2015; Tononi et al., 2016; Tononi and Edelman, 1998]. This led us to ask whether the stimulus-related TTV reduction also depends upon the level of consciousness. We hypothesized that TTV reduction would be reduced or absent when the level of consciousness is reduced. This hypothesis was supported by previous studies demonstrating that various anesthetics diminished recurrent processing through the suppression of anterior-posterior cortical feedback signaling [Alkire et al., 2008; Imas et al., 2005; Lee et al., 2013; Mashour and Hudetz, 2018].

On the first sight, neural variability may seem detrimental for accurate cognitive processing, perception, and potentially consciousness. However, several studies suggest that temporal variability (TV), defined as a fluctuation of neural activity over time, is beneficial for the neural system's adaptability, efficiency and cognitive performance [Dai et al., 2016; Garrett et al., 2010; Garrett et al., 2011a; Garrett et al., 2013a; Garrett et al., 2013b; Garrett et al., 2014; McIntosh et al., 2008; McIntosh et al., 2010; Vakorin et al., 2011a]. Moreover, higher TV may reflect greater network complexity that is more capable for modulating its activity to explore a larger repertoire of brain states [Faisal et al., 2008; Garrett et al., 2011b; Hudetz et al., 2015; Mišić et al., 2011; Shew et al., 2009; Vakorin et al., 2011b]. Conversely, when TV is low, such as during reduced level of consciousness, the capacity of the brain to explore different configurations/states would be reduced [Baria et al., 2018; Huang et al., 2014a; Huang et al., 2014b; Huang et al., 2016; Hudetz et al., 2015; Tagliazucchi et al., 2016; Zhang et al., 2018].

In summary, on one hand, higher neural variability appears important for cognitive functioning and consciousness; on the other hand, a reduction of variability seems essential for stimulus processing or task response. The crucial point to dissolve the apparent contradiction is the distinction between the two forms of neural variability – the quenching of stimulus-induced variability (i.e., TTV reduction), and the variability of ongoing neural activity across time (i.e., TV during resting-state or entire task state). As alluded to above, the high TV of ongoing activity can be considered as the brain's exploration of possible states within a region of neuronal activity space [Churchland et al., 2010; Deco and Hugues, 2012; He, 2013; Nogueira et al., 2018; Ponce-Alvarez et al., 2015; Tononi et al., 2016], whereas a

superimposed stimulation would shrink the available neuronal activity space to a subset with high stimulus-relevance; thus reducing the neural states from “exploratory” to “certainty”.

How the two forms of neural variability (TV and TTV) may be affected in reduced states of consciousness has not been elicited in the same experimental paradigm. Given its significance, we set out to investigate this question by an approach using anesthetic modulation of the level of consciousness. Specifically, we tested the hypothesis, the lack of TTV reduction during anesthetic administration as determined from fMRI-BOLD signals of spontaneous (resting-state) and stimulus-induced activities in three conditions: wakefulness, propofol-induced sedation and unconsciousness (i.e. deep anesthesia). We measured the average (trial-to-trial mean, TTM) and variability (TTV) of the stimulus-induced activity in response to auditory stimuli. We also measured the TV of ongoing activity during both resting-state and task, and calculated the TV difference between resting and task states (i.e. task modulation of TV), under the three conditions.

Materials and Methods

Subjects

The study was approved by the Institutional Review Board (IRB) of Huashan Hospital, Fudan University. Informed consent was obtained by all the subjects to participant in the study. Twenty right-handed subjects were recruited (male/female: 8/12; age: 32-64 years), who were undergoing an elective trans-sphenoidal approach for resection of a pituitary microadenoma. The pituitary microadenomas were diagnosed by their size (<10 mm in diameter without growing out of the sella)

Author Manuscript

based on radiological examinations and plasma endocrinal parameters. These subjects were ASA (American Society of Anesthesiologists) physical status I or II grade, with no history of craniotomy, cerebral neuropathy, vital organ dysfunction or administration of neuropsychiatric drugs. The subjects had no contraindication for an MRI examination, such as vascular clips or metallic implants. Among them, five subjects had to be excluded from the study and further data analysis because of excessive movements. Also, see below for more details on motion correction and subject exclusion.

Anesthesia Protocol

Subjects fasted for at least 8 hours from solids and 2 hours from liquids before participating the study. During the fMRI study, vital signs including blood pressure, heart rate, pulse oximetry (SpO_2), and partial pressure of carbon dioxide ($PaCO_2$) were continuously monitored (Table 1). The subjects received propofol sedation and deep anesthesia, during which intravenous anesthetic propofol was infused through an intravenous catheter placed into a vein of the right hand or forearm. Propofol acts mainly on $GABA_A$ receptors, potentiating GABAergic interneurons in the cortex and enhancing neuronal inhibition [Brown et al., 2011; Franks, 2008]. Propofol was administered using a target controlled infusion (TCI) pump to obtain constant effect-site concentration, as estimated by the pharmacokinetic model [Marsh et al., 1991]. Remifentanyl (1.0 $\mu\text{g}/\text{kg}$) and succinylcholine (1.5 mg/kg) were administered to facilitate endotracheal intubation at deep anesthesia. TCI concentrations were increased in 0.1 $\mu\text{g}/\text{ml}$ steps beginning at 1.0 $\mu\text{g}/\text{ml}$ until reaching the appropriate effect-site concentration. A 5-min equilibration period was allowed to insure equilibration of propofol repartition

between compartments.

The TCI propofol were maintained at a stable effect-site concentration of 1.3 $\mu\text{g/ml}$ for sedation, and 4.0 $\mu\text{g/ml}$ for deep anesthesia of which the dose reliably induces an unconscious state [Xu et al., 2009]. The level of consciousness was evaluated clinically throughout the study with the Ramsay scale [Ramsay et al., 1974]. The subjects were asked to strongly squeeze the hand of the investigator. The subject was considered fully awake if the response to verbal command (“squeeze strongly my hand!”) is clear and strong (Ramsay 1-2), in mild sedation if the response to verbal command is clear but slow (Ramsay 3-4), and in deep sedation or general anesthesia if there is no response to verbal command (Ramsay 5–6). For each assessment, Ramsay scale verbal commands were repeated twice (before remifentanyl and succinylcholine were administered at deep anesthesia) (Table 1). As anticipated, the corresponding Ramsay scores ($n=11$; mean \pm SD) for 1.3 $\mu\text{g/ml}$ and 4.0 $\mu\text{g/ml}$ of propofol concentration were 3.8 ± 1.7 and 6.0 ± 0 .

The subjects breathed spontaneously during wakefulness and sedation. During deep anesthesia, the subjects were ventilated with intermittent positive pressure ventilation, setting tidal volume at 8–10 ml/kg, respiratory rate 10-12 beats per minute, and maintaining PetCO₂ (partial pressure of end tidal CO₂) at 35-45 mmHg. Two certified anesthesiologists were present throughout the study, and complete resuscitation equipment were always available. Subjects wore earplugs and headphones during the fMRI scanning.

Data acquisition

Functional magnetic resonance imaging (fMRI) acquisition consisted of 8-min eyes closed resting-state and 18-min task-state (auditory stimuli) scans for each level of consciousness (wakefulness, sedation and anesthesia). The subject's head was fixed in the scan frame and padded with spongy cushions to minimize head movement. The subjects were informed to take a comfortable supine position, to relax with their eyes closed during scanning (with an eye patch). The subjects were instructed not to concentrate on anything in particular during resting-state scan, and listen to the sound during task scan. A Siemens 3T scanner (Siemens MAGNETOM, Germany) with a standard head coil was used to acquire gradient-echo EPI images of the whole brain (33 slices, repetition time/echo time [TR/TE] = 2,000/30 ms, slice thickness 5 = mm, field of view = 210 mm, flip angle = 90°, image matrix: 64 x 64); 240 scans (8 min) were acquired for the resting-states, and 540 scans (18 min) were acquired for the task. In addition, high-resolution anatomical images were acquired.

Task-fMRI design

A sparse event-related design was adopted with 60 trials. This included 30 own and 30 other (an unknown person's name) names that were delivered in a pseudo-random order. The names were presented by a familiar voice (recorded from one of the patients' family members) with an audio clip lasting for 0.5 s (see [Lipsman et al., 2014; Qin et al., 2012] for details of this well-established paradigm). The unknown names were matched to the own name by gender and number of syllables. In the presented study, own and other names were collapsed into a single condition (auditory stimuli) in order to increase the total number of trials which is beneficial for TTV calculation (see below). Each

audio clip was followed by inter-trial-intervals (ITIs) ranging unpredictably from 15.5 to 25.5s (2s-step). The benefit of long ITIs was avoiding potential nonlinearities associated with overlapping hemodynamic responses between preceding and subsequent trials, especially for trial-to-trial variability calculation [Huang et al., 2017]. All stimuli were programmed using E-Prime 2.0 (Psychology Software Tools, Pittsburgh, PA) and delivered via an audiovisual stimulus presentation system designed for an MRI environment. The volume of the headphones was adjusted to the comfort level. The subjects were required to pay attention and passively listen to the names without behavioral response or judgment.

Data preprocessing

Preprocessing steps were implemented in AFNI (Cox 1996; <http://afni.nimh.nih.gov/afni>) including: (1) discarding the first four frames of each fMRI run; (2) slice timing correction; (3) rigid body correction/realignment within and across runs; (4) co-registration with high-resolution anatomical images; (5) spatial normalization into Talaraich stereotactic space; (6) resampling to $3\times 3\times 3$ mm³ voxels; (7) regressing out linear and non-linear drift (equivalent to a high-pass filtering of 0.0067 Hz), head motion and its temporal derivative, and mean time series from the white matter (WM) and cerebrospinal fluid (CSF) to control for non-neural noise [Fox et al., 2005]; (8) spatial smoothing with a 8 mm full-width at half-maximum isotropic Gaussian kernel; (9) the time-course per voxel was normalized into z-value by subtracting the mean and dividing by the standard deviation of BOLD signals across all scans, accounting for differences in variance of non-neural origin (e.g., distance from

head coil) [He, 2011; Stephens et al., 2013].

The issue of motion artifacts was addressed rigorously as minor group differences in motion have been shown to artificially create between-groups differences [van Dijk et al., 2012; Power et al., 2012; Power et al., 2014]. We first calculated the indices of the amount of motion (shift and rotation) for each subject [Zang et al., 2007]. Subjects whose head motion's shift > 3 mm or rotation $> 3^\circ$ were excluded. For the remaining subjects, frame-wise displacement (FD) motion censoring was calculated using frame-wise Euclidean Norm (square root of the sum squares) of the six-dimension motion derivatives. A frame (TR) and its each previous frame were censored out from the data (using AFNI function 3dTproject) if the given frame's derivative value has a Euclidean Norm of $FD > 0.4$ mm [Huang et al., 2018]. Furthermore, to further exclude the potential confound of head motion, the head motion indices was included as covariates during group-level analysis.

It has been shown that including or excluding global signal regression (GSR) during preprocessing may affect the interpretations of the results [Chai et al., 2012; Murphy et al., 2009; Murphy and Fox, 2017; Saad et al., 2012]. For this reason, we presented and compared the results both with and without GSR for all the following analyses.

The average of stimulus-induced activity

The average of stimulus-induced BOLD signal changes at each time point across trials was calculated within the time window after stimulus presentation (0s-15s). The averaged time-course of each voxel of each subject was normalized by subtracting the mean value at stimulus onset (0s-1s). The

peak value of evoked activity (4-5s) was defined as TTM (trial-to-trial mean) index. Unlike conventional general linear model (GLM) analysis, we adopted TTM to quantify the stimulus-evoked activity and thus to make the results comparable with the TTV index (see below). Also note that we compared the results between TTM and estimated activity (regression coefficients, β value) using GLM; the results of the two were approximately the same. Whole-brain voxel-wise one sample t-tests against zero were performed at the group level for each condition, i.e. wakefulness, sedation and deep anesthesia. Unless otherwise stated, all resulting z-maps were thresholded at $\alpha < 0.05$ at the cluster level, where the corresponding p value at the voxel level was 0.005 with a cluster size of 85 voxels. This was achieved using AFNI's upgraded function 3dClustSim that simulates noise volume assuming the spatial auto-correlation function is given by a mixed-model rather than a Gaussian-shaped function [Cox et al., 2017].

The variability of stimulus-induced activity

The standard deviation (SD) of BOLD signal changes at each time point across trials was calculated within the time window after stimulus presentation (0s-15s). The SD time-course of each voxel of each subject was normalized by subtracting and dividing by the SD at stimulus onset (0-1s) [He, 2013; Huang et al., 2017]. The normalized SD value during the peak of evoked activity (4-5s) was defined as TTV index [Huang et al., 2017]. Whole-brain voxel-wise one sample t-tests against zero were performed at the group level.

Analysis of temporal variability – task vs. rest

The standard deviation (SD) of BOLD signal changes across time for the entire task and rest periods, respectively, was calculated for each condition (wakefulness, sedation and deep anesthesia), yielding six SD maps per subject. The task-modulation of TV was defined as the difference map between task and rest (task minus rest) for each condition. The significance of task-modulation effect of TV was examined for each network (see below for the network definition) using one sample t-tests against zero at the group level with FDR correction.

Definition of functional networks

We adopted a well-established network template derived from CONN's ICA analyses of HCP dataset in 497 subjects (www.conn-toolbox.org). The network template included the default mode network (DMN), frontoparietal (FP), salience/cingulo-opercular (Sal), language (Lan), dorsal attention (DA), sensorimotor (SM), and visual (Vis). In addition to these networks, we also included two brain areas (defined by AFNI atlas): bilateral primary auditory cortex (Audi) and thalamus (Tha). They were chosen due to the nature of our task design (auditory stimuli) such that we expected to see robust stimulus-induced activity in these regions.

Results

The average and variability of stimulus-induced activity (TTM vs. TTV)

We found graded diminished TTM in both amplitude and spatial extension from wakefulness over sedation to deep anesthesia, where the TTM was limited to primary auditory cortices during anesthesia ([Figure 1A](#)). After GRS, it revealed “deactivation” in several brain areas including frontal, parietal and

visual cortices during wakefulness, while there was minimal impact of GSR on the results of sedation and deep anesthesia (Figure 1B).

We observed significant TTV reduction after stimulus onset in especially the posterior part of brain including bilateral inferior parietal lobe and posterior cingulate cortex (PCC) during wakefulness (Figure 2A). Note that these significant TTV values were negative (namely, reduction), because of the normalization performed against the stimulus onset (see Methods); it thus indicated how the TTV changes relative to the baseline. Furthermore, the GSR helped identify a few more regions including angular gyrus, auditory, medial frontal and anterior cingulate cortices (Figure 2B). In contrast, no significant TTV change was found during either sedation or deep anesthesia.

We further examined the TTM and TTV in various functional networks (see Figure 3 for Audi and DMN as example regions; Figure 4A for network template illustration). During wakefulness, significantly increased TTM was seen in Audi, Sal, Lan, SM and Tha without GSR. After GSR, the statistical significance was improved with a few more regions detected including positive activation in SM and deactivation in DMN, FP and Vis (Figure 4B). During sedation, a significant increase of TTM was only seen in Audi, while significant increases of TTM were also found in Lan and SM after GSR (Figure 4B). During deep anesthesia, Audi was the only network showing significant TTM increase both with and without GSR (Figure 4B). For TTV during wakefulness, significant reduction was seen in Audi, DMN and FP before GSR; Sal and Lan were further detected using GSR. No significant result of TTV was seen in either sedation or deep anesthesia (Figure 4C).

Task-modulation of temporal variability (task TV vs. rest TV)

We first examined the TV differences between different conditions (e.g. wakefulness vs. sedation) during the resting-state. We found significantly decreased TV in FP and Sal during sedation, and DMN, FP and Vis during deep anesthesia, when compared to wakefulness; no-GSR and GSR showed similar results (Figure 5A and 5C). We next tested the TV differences during the task between different conditions. No TV difference was found between wakefulness and sedation, while many networks including DMN, FP, Lan, DA and Vis, showed significant differences between wakefulness and anesthesia. The standard deviations of BOLD signals across trials during stimulus onset (0s-1s) and peak (4s-5s), originally used for TTV calculation, were also presented in Figure 5 as an additional support. Most importantly, we sought to examine how the task context (repeated auditory stimuli with unpredictable inter-trial-intervals) modulates the TV compared with its resting-state in different conditions. We observed significant task-modulation of TV (task > rest) in Lan during wakefulness, and in Audi, FP, Sal, Lan during sedation before GSR (Figure 5B). Interestingly, the GSR improved the significance such that all significant networks seen during sedation came out to be significant during wakefulness (Figure 5D). Lastly, no significant task-modulation of TV was found during deep anesthesia either with or without GSR (Figure 5B and 5D).

Discussion

We demonstrated distinct effects of propofol on the average (TTM), variability (TV) of stimulus-induced activity, and the task modulation of temporal variability (TV) of ongoing activity (see

Table 2 for a summary). There was a graded attenuation of TTM from wakefulness through sedation to anesthesia. Importantly, the stimulus-induced TTV reduction normally seen during wakefulness was abolished during both sedation and anesthesia. In addition, there was an increase of TV in the task state as compared to resting-state during both wakefulness and sedation, but not anesthesia.

Graded attenuation of TTM from wakefulness through sedation to anesthesia

We found a gradual decrease of TTM in both amplitude and spatial distribution from wakefulness (widespread brain areas) through sedation (auditory cortex, language and sensorimotor networks) to deep anesthesia (auditory cortex only). Our results are in line with earlier observations that propofol attenuates stimulus-induced activity during auditory and noxious stimulation [Adapa et al., 2014; Davis et al., 2007; DiFrancesco et al., 2013; Dueck et al., 2005; Frölich et al., 2017; Liu et al., 2012; Mhuirheartaigh et al., 2010; Mhuirheartaigh et al., 2013]. These effects may be due to an overall impairment of hierarchy cortical processing and information integration [Alkire et al., 2008; Davis et al., 2007].

Disrupted TTV reduction during sedation and anesthesia

A series of neurophysiological and neuroimaging experiments have demonstrated substantial TTV reduction after stimulus onset in wakeful animals and humans [Churchland et al., 2010; He, 2013; Huang et al., 2017]. Likewise, we found significant stimulus-induced TTV reduction in various brain networks during wakefulness.

Extending the former studies, we also demonstrated that the TTV reduction was abolished by the

anesthetic propofol. Previous studies showed that TTV reduction derives from a nonlinear interaction between spontaneous and stimulus-induced activities [He, 2013; Huang et al., 2017]. We thereby tentatively suggest that anesthetics may disrupt the nonlinearity of neural dynamics in response to external stimuli. Propofol is known to potentiate postsynaptic inhibition by GABAergic interneurons in the cortex, which may contribute to the breakdown of anterior-posterior cortical feedback signaling [Alkire et al., 2008; Imas et al., 2005; Lee et al., 2013]. Thus, these effects may corrupt the recurrent neural circuits necessary for TTV reduction. To delineate the precise neural mechanisms underlying the anesthetic modulation of TTV reduction will require further investigation.

The variability of spontaneous neural activity can be understood to arise from a dynamic system that engages in an ongoing exploration of metastable network configurations selected from a “dynamic repertoire” [Deco et al., 2014; Deco and Hugues, 2012; Hudetz et al., 2015; Kelso, 2012; Litwin-Kumar and Doiron, 2012; Tagliazucchi et al., 2016]. Specifically, it is assumed that the network has access to a different number of stable attractors during spontaneous vs. stimulus-induced activity. During spontaneous activity, the network explores metastable configurations within a multi-attractor regime by means of its inherent chaotic behavior. The presentation of stimulus reduces the set of available attractors by eliminating the transition to other attractors, resulting in a net decrease in neural variability across trials [Deco and Hugues, 2012]. The lack of TTV reduction during sedation and anesthesia may reflect the reduced capacity of the dynamical system to access or stabilize network states and form specific attractors in response to external stimuli.

Reduced resting-state TV but preserved task-modulation of TV during sedation

We observed a decrease of resting-state TV in frontoparietal networks during sedation, with a further decrease of TV in both frontoparietal and DMN regions during deep anesthesia. This observation is in line with our prior finding that resting-state TV was closely related to the level of consciousness [Huang et al., 2014a; Huang et al., 2014b; Huang et al., 2016; Zhang et al., 2018]; further supported by a recent fMRI study in rats [Baria et al., 2018]. The reduction of resting-state TV is compatible with former studies suggesting a decrease of the brain's dynamic repertoire during anesthesia, sleep and unresponsive wakefulness [Barttfeld et al., 2015; Chennu et al., 2014; Hudetz et al., 2015; King and Dehaene, 2014; Sarasso et al., 2015; Tagliazucchi et al., 2016]. However, the exact linkage between the reduced temporal variability of fMRI-BOLD signals and the reduced temporal variability of dynamic functional connectivity patterns (the dynamic repertoire) during altered states of consciousness remains to be determined.

Interestingly, we found significant task-modulation of TV (task > rest) in auditory cortex and higher-order cortical regions (frontoparietal, salience and language networks) during both wakefulness and sedation, whereas no such modulation was found during deep anesthesia. The increase of TV during task may indicate that the neural system increased its dynamic range upon cognitive demand. One could envision that the wider dynamic range would provide a larger region of neuronal activity space to accommodate greater stimulus uncertainty as compared to the resting-state in which cognitive demand is lower due to less stimulus uncertainty. Thus, the task-related TV increase may be the sign of

an adaptive behavior of the neural system engaged in enhancing its efficiency and cognitive performance [Garrett et al., 2013a; Garrett et al., 2013b].

In summary, the higher TV in task-state vs. resting-state during sedation suggests that the relevant brain regions preserved their responsiveness and adaptability in terms of their capacity to augment the dynamic range of ongoing activity, presumably to accommodate the external stimulus' demand. In contrast, during anesthesia, brain activity appeared isolated from the external environment such that the cortical activity space was dramatically shrunk, almost unresponsive to stimuli, with TTV reduction and task-modulation of TV both abolished.

Conclusions

Our results reveal distinct effects of propofol on the two forms of neural variability – trial-to-trial variability (TTV) and temporal variability (TV). They imply that the anesthetic disrupts recurrent network dynamics, preventing the stabilization of cortical activity states. These findings shed new light on the temporal dynamics of neuronal variability and its alteration during anesthetic-induced unconsciousness.

Comments on global signal regression

We presented and compared the results both with and without global signal regression to ensure our results are not artefactually driven by the fMRI global signal itself [Murphy and Fox, 2017]. Overall, global signal regression improved the significance for all measurements while did not affect our general conclusion. Specifically, more regions were detected with significant TTV reduction after

GSR during wakefulness, for both whole brain analysis (Figure 2B) and network analysis (the right panel in Figure 4C). In addition, more networks showed significant task modulation of TV after GSR during wakefulness (Figure 5D) comparing to the results without GSR (Figure 5B). These improvements may be due to the benefit of GSR by further removing non-neuronal sources of global variance such as respiration and movement [Power et al., 2014; Yan et al., 2013]. Note that the above improvements did not change the sign (positive or negative) of TTV and TV values. However, we did find that GSR changes the sign for TTM values that several brain regions/networks became “deactivated” (Figure 1B and the right panel in Figure 4B). The question on whether these regions were truly deactivated in response to stimuli may be analogous to asking if there exist true negative correlations (anti-correlations) during resting-state in the context of GSR. This, however, is still under debate [Murphy and Fox, 2017]. We would tentatively suggest that the deactivations of TTM may be relative values against the global signal; it would be otherwise considered to be true deactivations if we could observe them for both GSR and no-GSR. Taking a step back, if we assume that the apparent deactivations have some functional significance (albeit unknown at this point), then the fact that no such deactivation was seen in either sedation or anesthesia may point to another dysfunction induced by propofol anesthetics that warrants future investigations.

Limitations

Several limitations of our study need to be considered: 1) We here used a passive name listening paradigm instead of goal-directed tasks (visual stimulus detection or semantic judgments) requiring

button responses as previous fMRI reported [He, 2013; Huang et al., 2017]. This was due to the consideration that the subjects were unable to make explicit motor response during anesthesia; it would thus be otherwise incomparable between different conditions (e.g. with motor response during wakefulness vs. without motor response during anesthesia). It is noteworthy that the evidence of TTV reduction from previous studies was not solely drawn from goal-directed tasks. For instance, neuronal TTV reduction was seen in wakeful rats during passive visual stimulation (e.g. white-black flashing) [White et al., 2012], as well as in monkeys and cats during presenting visual sine-wave gratings [Churchland et al., 2010]. Therefore, the TTV reduction through a presumed active noise suppression mechanism [White et al., 2012] may not be necessarily associated with active behavioral response (e.g. goal-directed).

2) We did not observe widespread TTV reduction across the entire cortex as previously shown [He, 2013; Huang et al., 2017]. This may be due to the simplicity of the task with lower cognitive load and task demands, as well as a relatively lower number of trials in a sparse-event related design within limited clinical fMRI scanning time.

3) Churchland et al. [2010] reported a TTV reduction of the stimulus-induced neuronal activity in many cortical areas regardless of whether the animal is anesthetized or awake. However, both our data in humans and a previous study in rats [White et al., 2012] showed that the TTV reduction was not maintained under anesthetized states. It remains an open question about which of the observations are closer to the physiological reality, and/or to what extent the TTV reduction depends upon tasks,

modalities (visual, auditory, tactile, etc.), attentional and conscious states.

4) It remains to be determined if our results can be generalized to other classes of general anesthetics, including halogenated ethers such as sevoflurane or non-GABAergic drugs such as ketamine. Therefore, future investigation of both forms of neural variability in those drugs where consciousness is lost is necessary to flesh out the full scientific and clinical implications of our findings.

Acknowledgments

This study was supported by Medical Guidance Supporting Project from Shanghai Municipal Science and Technology Committee (No. 17411961400 to Jun Zhang), grants of the National Institute of General Medical Sciences of the National Institutes of Health (R01-GM103894 to Anthony G. Hudetz), EJLB-Michael Smith Foundation, Canada Institute of Health Research (CIHR), Hope of Depression Foundation (HDRF), and National Natural Science Foundation of China (No. 31271195 to Georg Northoff). We declare no conflict of interest for all authors.

References

- Adapa RM, Davis MH, Stamatakis EA, Absalom AR, Menon DK (2014): Neural correlates of successful semantic processing during propofol sedation. *Hum Brain Mapp* 35:2935–2949. <http://doi.wiley.com/10.1002/hbm.22375>.
- Alkire MT, Hudetz AG, Tononi G (2008): Consciousness and anesthesia. *Science* 322:876–80. <http://www.sciencemag.org/cgi/doi/10.1126/science.1149213>.
- Arazi A, Gonen-Yaacovi G, Dinstein I (2017): The Magnitude of Trial-By-Trial Neural Variability Is Reproducible over Time and across Tasks in Humans. *eneuro* 4:ENEURO.0292-17.2017. <http://www.ncbi.nlm.nih.gov/pubmed/29279861>.
- Arieli A, Sterkin A, Grinvald A, Aertsen A (1996): Dynamics of ongoing activity: explanation of the large variability in evoked cortical responses. *Science* 273:1868–71. <http://www.ncbi.nlm.nih.gov/pubmed/8791593>.
- Azouz R, Gray CM (1999): Cellular mechanisms contributing to response variability of cortical neurons in vivo. *J Neurosci* 19:2209–23. <http://www.ncbi.nlm.nih.gov/pubmed/10066274>.
- Baria AT, Centeno M V., Ghantous ME, Chang PC, Procissi D, Apkarian AV (2018): BOLD temporal variability differentiates wakefulness from anesthesia-induced unconsciousness. *J Neurophysiol* 119:834–848. <http://www.ncbi.nlm.nih.gov/pubmed/29212921>.
- Baria AT, Maniscalco B, He BJ (2017): Initial-state-dependent, robust, transient neural dynamics

encode conscious visual perception. Ed. Ole Jensen. PLOS Comput Biol 13:e1005806.

<http://www.ncbi.nlm.nih.gov/pubmed/29176808>.

Barttfeld P, Uhrig L, Sitt JD, Sigman M, Jarraya B (2015): Signature of consciousness in the dynamics of resting-state brain activity. Proc Natl Acad Sci 112:E5219–E5220.

<http://www.pnas.org/lookup/doi/10.1073/pnas.1515029112>.

Becker R, Reinacher M, Freyer F, Villringer A, Ritter P (2011): How Ongoing Neuronal Oscillations Account for Evoked fMRI Variability. J Neurosci 31:11016–11027.

<http://www.ncbi.nlm.nih.gov/pubmed/21795550>.

Boly M, Garrido MI, Gosseries O, Bruno M-A, Boveroux P, Schnakers C, Massimini M, Litvak V, Laureys S, Friston K (2011): Preserved feedforward but impaired top-down processes in the vegetative state. Science (80-) 332:858–862. <http://www.ncbi.nlm.nih.gov/pubmed/21566197>.

Brown EN, Purdon PL, Van Dort CJ (2011): General anesthesia and altered states of arousal: a systems neuroscience analysis. Annu Rev Neurosci 34:601–28.

<http://www.ncbi.nlm.nih.gov/pubmed/21513454><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3390788>.

Chai XJ, Castañán AN, Öngür D, Whitfield-Gabrieli S (2012): Anticorrelations in resting state networks without global signal regression. Neuroimage 59:1420–1428.

<http://www.ncbi.nlm.nih.gov/pubmed/21889994>.

Chang MH, Armstrong KM, Moore T (2012): Dissociation of response variability from firing rate effects in frontal eye field neurons during visual stimulation, working memory, and attention. *J Neurosci* 32:2204–16. <http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.2967-11.2012>.

Chennu S, Finoia P, Kamau E, Allanson J, Williams GB, Monti MM, Noreika V, Arnatkeviciute A, Canales-Johnson A, Olivares F, Cabezas-Soto D, Menon DK, Pickard JD, Owen AM, Bekinschtein TA (2014): Spectral Signatures of Reorganised Brain Networks in Disorders of Consciousness. Ed. Bard Ermentrout. *PLoS Comput Biol* 10:e1003887. <http://www.ncbi.nlm.nih.gov/pubmed/25329398>.

Churchland MM, Yu BM, Cunningham JP, Sugrue LP, Cohen MR, Corrado GS, Newsome WT, Clark AM, Hosseini P, Scott BB, Bradley DC, Smith MA, Kohn A, Movshon JA, Armstrong KM, Moore T, Chang SW, Snyder LH, Lisberger SG, Priebe NJ, Finn IM, Ferster D, Ryu SI, Santhanam G, Sahani M, Shenoy K V (2010): Stimulus onset quenches neural variability: a widespread cortical phenomenon. *Nat Neurosci* 13:369–378. <http://www.ncbi.nlm.nih.gov/pubmed/20173745>.

Cox RW, Chen G, Glen DR, Reynolds RC, Taylor PA (2017): FMRI Clustering in AFNI: False-Positive Rates Redux. *Brain Connect* 7:152–171. <http://www.ncbi.nlm.nih.gov/pubmed/28398812>.

Dai R, Huang Z, Tu H, Wang L, Tanabe S, Weng X, He S, Li D (2016): Interplay between Heightened Temporal Variability of Spontaneous Brain Activity and Task-Evoked Hyperactivation in the Blind. *Front Hum Neurosci* 10:632.

<http://journal.frontiersin.org/article/10.3389/fnhum.2016.00632/full>.

Davis MH, Coleman MR, Absalom AR, Rodd JM, Johnsrude IS, Matta BF, Owen AM, Menon DK

(2007): Dissociating speech perception and comprehension at reduced levels of awareness. *Proc*

Natl Acad Sci U S A 104:16032–7. <http://www.pnas.org.libproxy.ucl.ac.uk/content/104/41/16032>.

Deco G, Hugues E (2012): Neural Network Mechanisms Underlying Stimulus Driven Variability

Reduction. Ed. Tim Behrens. *PLoS Comput Biol* 8:e1002395.

<http://www.ncbi.nlm.nih.gov/pubmed/22479168>.

Deco G, McIntosh AR, Shen K, Hutchison RM, Menon RS, Everling S, Hagmann P, Jirsa VK (2014):

Identification of optimal structural connectivity using functional connectivity and neural

modeling. *J Neurosci* 34:7910–6.

<http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.4423-13.2014>.

DiFrancesco MW, Robertson SA, Karunanayaka P, Holland SK (2013): BOLD fMRI in infants under

sedation: Comparing the impact of pentobarbital and propofol on auditory and language

activation. *J Magn Reson Imaging* 38:1184–95. <http://doi.wiley.com/10.1002/jmri.24082>.

van Dijk KRA, Sabuncu MR, Buckner RL (2012): The influence of head motion on intrinsic functional

connectivity MRI. *Neuroimage* 59:431–438. <http://www.ncbi.nlm.nih.gov/pubmed/21810475>.

Dueck MH, Petzke F, Gerbershagen HJ, Paul M, Hesselmann V, Girmus R, Krug B, Sorger B, Goebel R,

Lehrke R, Sturm V, Boerner U (2005): Propofol attenuates responses of the auditory cortex to

acoustic stimulation in a dose-dependent manner: a fMRI study. *Acta Anaesthesiol Scand* 49:784–91. <http://doi.wiley.com/10.1111/j.1399-6576.2005.00703.x>.

Faisal AA, Selen LPJ, Wolpert DM (2008): Noise in the nervous system. *Nat Rev Neurosci* 9:292–303. <http://www.nature.com/doi/10.1038/nrn2258>.

Ferri F, Costantini M, Huang Z, Perrucci MG, Ferretti A, Romani GL, Northoff G (2015): Intertrial Variability in the Premotor Cortex Accounts for Individual Differences in Peripersonal Space. *J Neurosci* 35:16328–39. <http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.1696-15.2015>.

Finn IM, Priebe NJ, Ferster D (2007): The Emergence of Contrast-Invariant Orientation Tuning in Simple Cells of Cat Visual Cortex. *Neuron* 54:137–152. <http://www.ncbi.nlm.nih.gov/pubmed/17408583>.

Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME (2005): The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl Acad Sci U S A* 102:9673–8. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1157105&tool=pmcentrez&rendertype=abstract>.

Fox MD, Snyder AZ, Zacks JM, Raichle ME (2006): Coherent spontaneous activity accounts for trial-to-trial variability in human evoked brain responses. *Nat Neurosci* 9:23–25. <http://www.ncbi.nlm.nih.gov/pubmed/16341210>.

Franks NP (2008): General anaesthesia: from molecular targets to neuronal pathways of sleep and arousal. *Nat Rev Neurosci* 9:370–386. <http://www.ncbi.nlm.nih.gov/pubmed/18425091>.

Frölich MA, Banks C, Ness TJ (2017): The Effect of Sedation on Cortical Activation: A Randomized Study Comparing the Effects of Sedation With Midazolam, Propofol, and Dexmedetomidine on Auditory Processing. *Anesth Analg* 124:1603–1610. <http://insights.ovid.com/crossref?an=00000539-201705000-00037>.

Garrett DD, Kovacevic N, McIntosh AR, Grady CL (2010): Blood Oxygen Level-Dependent Signal Variability Is More than Just Noise. *J Neurosci* 30:4914–4921. <http://www.ncbi.nlm.nih.gov/pubmed/20371811>.

Garrett DD, Kovacevic N, McIntosh AR, Grady CL (2011a): The Importance of Being Variable. *J Neurosci* 31:4496–4503. <http://www.ncbi.nlm.nih.gov/pubmed/21430150>.

Garrett DD, Samanez-Larkin GR, MacDonald SWS, Lindenberger U, McIntosh AR, Grady CL (2013a): Moment-to-moment brain signal variability: A next frontier in human brain mapping? *Neurosci Biobehav Rev* 37:610–624. <http://www.ncbi.nlm.nih.gov/pubmed/23458776>.

Garrett DD, Kovacevic N, McIntosh AR, Grady CL (2013b): The modulation of BOLD variability between cognitive states varies by age and processing speed. *Cereb Cortex* 23:684–93. <https://academic.oup.com/cercor/article-lookup/doi/10.1093/cercor/bhs055>.

Garrett DD, McIntosh AR, Grady CL (2011b): Moment-to-moment signal variability in the human

brain can inform models of stochastic facilitation now. *Nat Rev Neurosci* 12:612; author reply 612. <http://www.nature.com/articles/nrn3061-c1>.

Garrett DD, McIntosh AR, Grady CL (2014): Brain signal variability is parametrically modifiable. *Cereb Cortex* 24:2931–40. <https://academic.oup.com/cercor/article-lookup/doi/10.1093/cercor/bht150>.

He BJ (2013): Spontaneous and Task-Evoked Brain Activity Negatively Interact. *J Neurosci* 33:4672–4682. <http://www.ncbi.nlm.nih.gov/pubmed/23486941>.

He BJ (2011): Scale-free properties of the functional magnetic resonance imaging signal during rest and task. *J Neurosci* 31:13786–95. <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3197021&tool=pmcentrez&rendertype=abstract>.

He BJ, Zempel JM (2013): Average is optimal: an inverted-U relationship between trial-to-trial brain activity and behavioral performance. Ed. Olaf Sporns. *PLoS Comput Biol* 9:e1003348. <http://dx.plos.org/10.1371/journal.pcbi.1003348>.

Huang Z, Dai R, Wu X, Yang Z, Liu D, Hu J, Gao L, Tang W, Mao Y, Jin Y, Wu X, Liu B, Zhang Y, Lu L, Laureys S, Weng X, Northoff G (2014a): The self and its resting state in consciousness: An investigation of the vegetative state. *Hum Brain Mapp* 35:1997–2008. <http://doi.wiley.com/10.1002/hbm.22308>.

- Huang Z, Liu X, Mashour GA, Hudetz AG (2018): Timescales of Intrinsic BOLD Signal Dynamics and Functional Connectivity in Pharmacologic and Neuropathologic States of Unconsciousness. *J Neurosci* 38:2304–2317. <http://www.ncbi.nlm.nih.gov/pubmed/29386261>.
- Huang Z, Wang Z, Zhang J, Dai R, Wu J, Li Y, Liang W, Mao Y, Yang Z, Holland G, Zhang J, Northoff G (2014b): Altered temporal variance and neural synchronization of spontaneous brain activity in anesthesia. *Hum Brain Mapp* 35:5368–5378. <http://doi.wiley.com/10.1002/hbm.22556>.
- Huang Z, Zhang J, Longtin A, Dumont G, Duncan NW, Pokorny J, Qin P, Dai R, Ferri F, Weng X, Northoff G (2017): Is there a nonadditive interaction between spontaneous and evoked activity? Phase-dependence and its relation to the temporal structure of scale-free brain activity. *Cereb Cortex* 27:1037–1059. <http://cercor.oxfordjournals.org/cgi/doi/10.1093/cercor/bhv288>.
- Huang Z, Zhang J, Wu J, Qin P, Wu X, Wang Z, Dai R, Li Y, Liang W, Mao Y, Yang Z, Zhang J, Wolff A, Northoff G (2016): Decoupled temporal variability and signal synchronization of spontaneous brain activity in loss of consciousness: An fMRI study in anesthesia. *Neuroimage* 124:693–703. <http://linkinghub.elsevier.com/retrieve/pii/S1053811915007831>.
- Hudetz AG, Liu X, Pillay S (2015): Dynamic repertoire of intrinsic brain states is reduced in propofol-induced unconsciousness. *Brain Connect* 5:10–22. <http://online.liebertpub.com/doi/abs/10.1089/brain.2014.0230>.
- Imas OA, Ropella KM, Ward BD, Wood JD, Hudetz AG (2005): Volatile anesthetics disrupt frontal-posterior recurrent information transfer at gamma frequencies in rat. *Neurosci Lett*

387:145–50. <http://linkinghub.elsevier.com/retrieve/pii/S0304394005006890>.

Kelso JAS (2012): Multistability and metastability: understanding dynamic coordination in the brain.

Philos Trans R Soc Lond B Biol Sci 367:906–18.

<http://rstb.royalsocietypublishing.org/cgi/doi/10.1098/rstb.2011.0351>.

King J-R, Dehaene S (2014): Characterizing the dynamics of mental representations: the temporal

generalization method. Trends Cogn Sci 18:203–210.

<http://www.ncbi.nlm.nih.gov/pubmed/24593982>.

Lamme VA, Roelfsema PR (2000): The distinct modes of vision offered by feedforward and recurrent

processing. Trends Neurosci 23:571–9. <http://www.ncbi.nlm.nih.gov/pubmed/11074267>.

Lee U, Ku S, Noh G, Baek S, Choi B, Mashour G a (2013): Disruption of frontal-parietal

communication by ketamine, propofol, and sevoflurane. Anesthesiology 118:1264–1275.

<http://www.ncbi.nlm.nih.gov/pubmed/23695090>.

Lipsman N, Nakao T, Kanayama N, Krauss JK, Anderson A, Giacobbe P, Hamani C, Hutchison WD,

Dostrovsky JO, Womelsdorf T, Lozano AM, Northoff G (2014): Neural overlap between resting

state and self-relevant activity in human subcallosal cingulate cortex – Single unit recording in an

intracranial study. Cortex 60:139–144. <http://www.ncbi.nlm.nih.gov/pubmed/25438747>.

Litwin-Kumar A, Doiron B (2012): Slow dynamics and high variability in balanced cortical networks

with clustered connections. Nat Neurosci 15:1498–1505.

<http://www.ncbi.nlm.nih.gov/pubmed/23001062>.

Liu X, Lauer KK, Ward BD, Rao SM, Li SJ, Hudetz AG (2012): Propofol disrupts functional interactions between sensory and high-order processing of auditory verbal memory. *Hum Brain Mapp* 33:2487–2498. <http://doi.wiley.com/10.1002/hbm.21385>.

Marsh B, Morton N, Kenny GNC (1991): Pharmacokinetic model driven infusion of propofol in children. *Br J Anaesth* 67:41–48. <http://www.ncbi.nlm.nih.gov/pubmed/1859758>.

Mashour GA (2014): Top-down mechanisms of anesthetic-induced unconsciousness. *Front Syst Neurosci* 8:115. <http://www.ncbi.nlm.nih.gov/pubmed/25002838>.

Mashour GA, Hudetz AG (2018): Neural Correlates of Unconsciousness in Large-Scale Brain Networks. *Trends Neurosci* 41:150–160. <http://www.ncbi.nlm.nih.gov/pubmed/29409683>.

Mazzucato L, Fontanini A, La Camera G (2015): Dynamics of Multistable States during Ongoing and Evoked Cortical Activity. *J Neurosci* 35:8214–8231. <http://www.ncbi.nlm.nih.gov/pubmed/26019337>.

McIntosh AR, Kovacevic N, Lippe S, Garrett D, Grady C, Jirsa V (2010): The development of a noisy brain. *Arch Ital Biol* 148:323–37. <http://www.ncbi.nlm.nih.gov/pubmed/21175017>.

McIntosh AR, Kovacevic N, Itier RJ (2008): Increased Brain Signal Variability Accompanies Lower Behavioral Variability in Development. Ed. Karl J. Friston. *PLoS Comput Biol* 4:e1000106. <http://www.ncbi.nlm.nih.gov/pubmed/18604265>.

- Mhuirheartaigh RN, Rosenorn-Lannig D, Wise R, Jbabdi S, Rogers R, Tracey I (2010): Cortical and subcortical connectivity changes during decreasing levels of consciousness in humans: a functional magnetic resonance imaging study using propofol. *J Neurosci* 30:9095–102. <http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.5516-09.2010>.
- Mišić B, Vakorin VA, Paus T, McIntosh AR (2011): Functional embedding predicts the variability of neural activity. *Front Syst Neurosci* 5:90. <http://www.ncbi.nlm.nih.gov/pubmed/22164135>.
- Monier C, Chavane F, Baudot P, Graham LJ, Frégnac Y (2003): Orientation and direction selectivity of synaptic inputs in visual cortical neurons: a diversity of combinations produces spike tuning. *Neuron* 37:663–80. <http://www.ncbi.nlm.nih.gov/pubmed/12597863>.
- Moutard C, Dehaene S, Malach R (2015): Spontaneous Fluctuations and Non-linear Ignitions: Two Dynamic Faces of Cortical Recurrent Loops. *Neuron* 88:194–206. <http://www.ncbi.nlm.nih.gov/pubmed/26447581>.
- Murphy K, Birn RM, Handwerker DA, Jones TB, Bandettini PA (2009): The impact of global signal regression on resting state correlations: Are anti-correlated networks introduced? *Neuroimage* 44:893–905. <http://www.ncbi.nlm.nih.gov/pubmed/18976716>.
- Murphy K, Fox MD (2017): Towards a consensus regarding global signal regression for resting state functional connectivity MRI. *Neuroimage* 154:169–173. <http://linkinghub.elsevier.com/retrieve/pii/S1053811916306711>.

- Ní Mhuircheartaigh R, Warnaby C, Rogers R, Jbabdi S, Tracey I (2013): Slow-wave activity saturation and thalamocortical isolation during propofol anesthesia in humans. *Sci Transl Med* 5:208ra148. <http://www.ncbi.nlm.nih.gov/pubmed/24154602>.
- Nogueira R, Lawrie S, Moreno-Bote R (2018): Neuronal Variability as a Proxy for Network State. *Trends Neurosci* 41:170–173. <http://www.ncbi.nlm.nih.gov/pubmed/29602335>.
- Ponce-Alvarez A, He BJ, Hagmann P, Deco G (2015): Task-Driven Activity Reduces the Cortical Activity Space of the Brain: Experiment and Whole-Brain Modeling. Ed. Lyle J. Graham. *PLOS Comput Biol* 11:e1004445. <http://www.ncbi.nlm.nih.gov/pubmed/26317432>.
- Power JD, Barnes KA, Snyder AZ, Schlaggar BL, Petersen SE (2012): Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage* 59:2142–2154. <http://linkinghub.elsevier.com/retrieve/pii/S1053811911011815>.
- Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE (2014): Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage* 84:320–41. <http://linkinghub.elsevier.com/retrieve/pii/S1053811913009117>.
- Qin P, Liu Y, Shi J, Wang Y, Duncan N, Gong Q, Weng X, Northoff G (2012): Dissociation between anterior and posterior cortical regions during self-specificity and familiarity: a combined fMRI-meta-analytic study. *Hum Brain Mapp* 33:154–64. <http://doi.wiley.com/10.1002/hbm.21201>.

- Rajan K, Abbott LF, Sompolinsky H (2010): Stimulus-dependent suppression of chaos in recurrent neural networks. *Phys Rev E* 82:11903. <http://www.ncbi.nlm.nih.gov/pubmed/20866644>.
- Ramsay MA, Savege TM, Simpson BR, Goodwin R (1974): Controlled sedation with alphaxalone-alphadolone. *Br Med J* 2:656–9. <http://www.ncbi.nlm.nih.gov/pubmed/4835444>.
- Saad ZS, Gotts SJ, Murphy K, Chen G, Jo HJ, Martin A, Cox RW (2012): Trouble at Rest: How Correlation Patterns and Group Differences Become Distorted After Global Signal Regression. *Brain Connect* 2:25–32. <http://www.ncbi.nlm.nih.gov/pubmed/22432927>.
- Sarasso S, Boly M, Napolitani M, Gosseries O, Charland-Verville V, Casarotto S, Rosanova M, Casali AG, Brichant J-F, Boveroux P, Rex S, Tononi G, Laureys S, Massimini M (2015): Consciousness and Complexity during Unresponsiveness Induced by Propofol, Xenon, and Ketamine. *Curr Biol* 25:3099–3105. <http://www.ncbi.nlm.nih.gov/pubmed/26752078>.
- Scaglione A, Moxon KA, Aguilar J, Foffani G (2011): Trial-to-trial variability in the responses of neurons carries information about stimulus location in the rat whisker thalamus. *Proc Natl Acad Sci U S A* 108:14956–61. <http://www.pnas.org/cgi/doi/10.1073/pnas.1103168108>.
- Schurger A, Sarigiannidis I, Naccache L, Sitt JD, Dehaene S (2015): Cortical activity is more stable when sensory stimuli are consciously perceived. *Proc Natl Acad Sci* 112:E2083–E2092. <http://www.ncbi.nlm.nih.gov/pubmed/25847997>.
- Shew WL, Yang H, Petermann T, Roy R, Plenz D (2009): Neuronal avalanches imply maximum

dynamic range in cortical networks at criticality. *J Neurosci* 29:15595–600.

<http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.3864-09.2009>.

Stephens GJ, Honey CJ, Hasson U (2013): A place for time: the spatiotemporal structure of neural

dynamics during natural audition. *J Neurophysiol* 110:2019–26.

<http://www.ncbi.nlm.nih.gov/pubmed/23926041><http://www.pubmedcentral.nih.gov/articler>

[ender.fcgi?artid=PMC3841928](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3841928/).

Sussillo D, Abbott LF (2009): Generating Coherent Patterns of Activity from Chaotic Neural Networks.

Neuron 63:544–557. <http://www.ncbi.nlm.nih.gov/pubmed/19709635>.

Tagliazucchi E, Chialvo DR, Siniatchkin M, Brichant J, Laureys S (2016): Large-scale signatures of

unconsciousness are consistent with a departure from critical dynamics. *J R Soc Interface*

13:1–34.

<http://rsif.royalsocietypublishing.org/content/13/114/20151027><http://rsif.royalsocietypublis>

<http://rsif.royalsocietypublishing.org/content/13/114/20151027?rss=1><http://rsif.royalsocietypublishing.org/content/royi>

[nterface/13/114/20151027.full.pdf](http://rsif.royalsocietypublishing.org/content/royinterface/13/114/20151027.full.pdf).

Tononi G, Edelman GM (1998): Consciousness and complexity. *Science* 282:1846–51.

<http://www.ncbi.nlm.nih.gov/pubmed/9836628>.

Tononi G, Boly M, Massimini M, Koch C (2016): Integrated information theory: from consciousness to

its physical substrate. *Nat Rev Neurosci* 17:450–61.

<http://dx.doi.org/10.1038/nrn.2016.44><http://www.ncbi.nlm.nih.gov/pubmed/27225071>.

Vakorin VA, Lippe S, McIntosh AR (2011a): Variability of Brain Signals Processed Locally Transforms into Higher Connectivity with Brain Development. *J Neurosci* 31:6405–6413. <http://www.ncbi.nlm.nih.gov/pubmed/21525281>.

Vakorin VA, Mišić B, Krakovska O, McIntosh AR (2011b): Empirical and theoretical aspects of generation and transfer of information in a neuromagnetic source network. *Front Syst Neurosci* 5:96. <http://www.ncbi.nlm.nih.gov/pubmed/22131968>.

White B, Abbott LF, Fiser J (2012): Suppression of cortical neural variability is stimulus- and state-dependent. *J Neurophysiol* 108:2383–92. <http://jn.physiology.org/cgi/doi/10.1152/jn.00723.2011>.

Xu Z, Liu F, Yue Y, Ye T, Zhang B, Zuo M, Xu M, Hao R, Xu Y, Yang N, Che X (2009): C50 for propofol-remifentanyl target-controlled infusion and bispectral index at loss of consciousness and response to painful stimulus in Chinese patients: a multicenter clinical trial. *Anesth Analg* 108:478–83. <http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00000539-200902000-00015>.

Xue G, Dong Q, Chen C, Lu Z, Mumford JA, Poldrack RA (2010): Greater Neural Pattern Similarity Across Repetitions Is Associated with Better Memory. *Science* (80-) 330:97–101. <http://www.ncbi.nlm.nih.gov/pubmed/20829453>.

Yan C-G, Craddock RC, Zuo X-N, Zang Y-F, Milham MP (2013): Standardizing the intrinsic brain:

towards robust measurement of inter-individual variation in 1000 functional connectomes.

Neuroimage 80:246–62. <http://linkinghub.elsevier.com/retrieve/pii/S105381191300428X>.

Zang YF, He Y, Zhu CZ, Cao QJ, Sui MQ, Liang M, Tian LX, Jiang TZ, Wang YF (2007): Altered baseline brain activity in children with ADHD revealed by resting-state functional MRI. *Brain Dev* 29:83–91. <http://www.ncbi.nlm.nih.gov/pubmed/16919409>.

Zhang J, Huang Z, Chen Y, Zhang J, Ghinda D, Nikolova Y, Wu J, Xu J, Bai W, Mao Y, Yang Z, Duncan N, Qin P, Wang H, Chen B, Weng X, Northoff G (2018): Breakdown in the temporal and spatial organization of spontaneous brain activity during general anesthesia. *Hum Brain Mapp* 39:2035–2046. <http://www.ncbi.nlm.nih.gov/pubmed/29377435>.

Figure Legends

Figure 1. The average of stimulus-induced activity across trials (trial-to-trial mean, TTM) during the peak period (4-5s) in wakefulness, propofol-induced sedation and deep anesthesia. Group-level z-maps (one sample t-tests against zero) were corrected at $\alpha < 0.05$ at the cluster level. **A.** Results without global signal regression (No-GSR). **B.** Results with GSR.

Figure 2. The variability of stimulus-induced activity across trials (trial-to-trial variability, TTV) during the peak period (4-5s) in wakefulness, propofol-induced sedation and deep anesthesia. Group-level z-maps (one sample t-tests against zero) were corrected at $\alpha < 0.05$ at the cluster level. **A.** Results without global signal regression (No-GSR). **B.** Results with GSR. Note that these significant TTV changes are negative values (namely reduction), because of the normalization against the stimulus onset (see Methods); it thus indicates how the TTV changes relative to the baseline.

Figure 3. Time courses of average and variability of stimulus-induced activity across trials within the time window after stimulus presentation (0s-15s). The results are shown for the auditory cortex (Audi) and default-mode network (DMN). Shaded areas indicate the peak period (4-5s) of stimulus-induced activity, where trial-to-trial mean (TTM) and trial-to-trial variability (TTV) were calculated. Results without and with global signal regression (No-GSR and GSR) are shown. Error bars indicate \pm SEM.

Figure 4. Trial-to-trial mean (TTM) and trial-to-trial variability (TTV) in various functional networks. **A.** network template includes auditory cortex (Audi), default mode network (DMN), frontoparietal (FP), salience/cingulo-opercular (Sal), language (Lan), dorsal attention (DA), sensorimotor (SM), visual (Vis), and thalamus (Tha). **B.** TTM values and statistical significances of different networks. **C.** TTV values and statistical significances of different networks. Results without and with global signal regression (No-GSR and GSR) are shown. *Significant differences against zero by one-sample t-tests. #Significant differences against wakefulness by paired sample t-tests (i.e. sedation vs. wakefulness, and anesthesia vs. wakefulness). §Significant differences against sedation by paired sample t-tests (i.e. anesthesia vs. sedation). Error bars indicate \pm SEM. All results were FDR corrected. The insets in B and C show the effect size (Cohen's d) for the mean differences against zero, where $d=0.2$ is considered a 'small' effect size, 0.5 represents a 'medium' effect size and 0.8 a 'large' effect size. The range limit of y-axis is from 0-0.8 for illustrative purpose. Overall, all reported significant results are close to or above $d=0.8$.

Figure 5. Temporal variability (TV) during resting-state (Rest), task-state (Task), as well as the standard deviations of BOLD signals across trials during stimulus onset (Onset) and peak period (Peak) originally used for TTV calculation. **A.** TV without global signal regression (No-GSR). #Significant differences against wakefulness by paired sample t-tests (i.e. sedation vs. wakefulness, and anesthesia vs. wakefulness). §Significant differences against sedation by paired sample t-tests (i.e. anesthesia vs.

sedation). Error bars indicate \pm SEM. **B.** A visualization of the TV (No-GSR) in a variability “space” across brain networks to highlight the comparisons of Task vs. Rest (task-modulation of TV). Red frames indicate significant task-modulation effects of TV. Results with GSR are shown in **C** and **D**. All results were FDR corrected. The right panels in **B** and **D** show the effect size (Cohen’s d) for task vs. rest, where $d=0.2$ is considered a 'small' effect size, 0.5 represents a 'medium' effect size and 0.8 a 'large' effect size. The range limit of y-axis is from 0-0.8 for illustrative purpose. Overall, all reported significant results are close to or above $d=0.8$.