



Guidelines for reporting of statistics for clinical research in urology

Melissa Assel¹, Daniel Sjoberg¹, Andrew Elders² , Xuemei Wang³, Dezheng Huo⁴, Albert Botchway⁵, Kristin Delfino⁵, Yunhua Fan⁶, Zhiguo Zhao⁸, Tatsuki Koyama⁸, Brent Hollenbeck⁹, Rui Qin¹⁰, Whitney Zahnd¹¹, Emily C. Zabor¹, Michael W. Kattan⁷ and Andrew J. Vickers¹ 

¹Memorial Sloan Kettering Cancer Center, New York, NY, USA, ²Glasgow Caledonian University, Glasgow, UK, ³The University of Texas, MD Anderson Cancer Center, Houston, TX, ⁴The University of Chicago, Chicago, ⁵Southern Illinois University School of Medicine, Springfield, IL, ⁶University of Minnesota, Minneapolis, MN, ⁷Cleveland Clinic, Cleveland, OH, ⁸Vanderbilt University Medical Center, Nashville, TN, USA, ⁹University of Michigan, Nanjing, China, ¹⁰Janssen Research & Development, Raritan, NJ, and ¹¹University of South Carolina, Charleston, SC, USA

In an effort to improve the quality of statistics in the clinical urology literature, statisticians at *European Urology*, *The Journal of Urology*, *Urology* and *BJU International* came together to develop a set of guidelines to address common errors of statistical analysis, reporting, and interpretation. Authors should 'break any of the guidelines if it makes scientific sense to do so', but would need to

provide a clear justification. Adoption of the guidelines will in our view not only increase the quality of published papers in our journals but improve statistical knowledge in our field in general.

Keywords

statistics, guidelines, analysis

Introduction

It is widely acknowledged that the quality of statistics in the clinical research literature is poor. This is true for urology just as it is for other medical specialties. In 2005, Scales et al. [1] published a systematic evaluation of the statistics in papers appearing in a single month in one of the four leading urology medical journals: *European Urology*, *The Journal of Urology*, *Urology* and *BJU International*. They reported widespread errors, including 71% of papers with comparative statistics having at least one statistical flaw. These findings mirror many others in the literature, see, for instance, the review given by Lang and Altman [2]. The quality of statistical reporting in urology journals has no doubt improved since 2005, but remains unsatisfactory.

The four urology journals in the Scales et al. [1] review have come together to publish a shared set of statistical guidelines, adapted from those in use at one of the journals, *European Urology*, since 2014 [3]. The guidelines will also be adopted by *European Urology Focus* and *European Urology Oncology*. Statistical reviewers at the four journals will systematically assess submitted manuscripts using the guidelines to improve statistical analysis, reporting, and interpretation. Adoption of the guidelines will, in our view, not only increase the quality of published papers in our journals but improve statistical knowledge in our field in general. Asking an author to follow

a guideline about, say, the fallacy of accepting the null hypothesis would no doubt result in a better paper, but we hope that it would also enhance the author's understanding of hypothesis tests.

The guidelines are didactic, based on the consensus of the statistical consultants to the journals. We avoided, where possible, making specific analytical recommendations and focused instead on analyses or methods of reporting statistics that should be avoided. We intend to update the guidelines over time and hence encourage readers who question the value or rationale of a guideline to write to the authors.

- 1 The golden rule: Break any of the guidelines if it makes scientific sense to do so.** Science varies too much to allow methodological or reporting guidelines to apply universally.
- 2 Reporting of design and statistical analysis**
 - 2.1 Follow existing reporting guidelines for the type of study you are reporting.** These include Consolidated Standards of Reporting Trials (CONSORT) for randomised trials, REporting Recommendations for Tumor MARKer Prognostic Studies (REMARK) for marker studies, Transparent Reporting of a multivariable prediction model for Individual Prognosis Or Diagnosis (TRIPOD) for prediction

models, STrengthening the Reporting of OBServational studies in Epidemiology (STROBE) for observational studies, or Assessing the Methodological quality of Systematic Reviews (AMSTAR) for systematic reviews. Statisticians and methodologists have contributed extensively to a large number of reporting guidelines. The first is widely recognised to be the CONSORT statement on the reporting of randomised trials, but there are now many other guidelines, covering a wide range of different types of study. Reporting guidelines can be downloaded from the Equator Web site (<http://www.equator-network.org>).

- 2.2 *Describe cohort selection fully. It is insufficient to state, for instance, 'the study cohort consisted of 1144 patients treated for BPH at our institution'.* The cohort needs to be defined in terms of dates (e.g., 'presenting March 2013 to December 2017'), inclusion criteria (e.g., 'IPSS >12'), and whether patients were selected to be included (e.g., for a research study) vs being a consecutive series. Exclusions should be described one by one, with the number of patients omitted for each exclusion criterion to give the final cohort size [e.g., 'patients with prior surgery ($n = 43$), allergies to 5 α -reductase inhibitors ($n = 12$), and missing data on baseline prostate volume ($n = 86$) were excluded to give a final cohort for analysis of 1003 patients']. Note that inclusion criteria can be omitted if obvious from context (e.g., no need to state 'undergoing radical prostatectomy for histologically confirmed prostate cancer'); on the other hand, dates may need to be explained if their rationale could be questioned (e.g., 'March 2013, when our specialist voiding clinic was established, to December 2017').
- 2.3 *Describe the practical steps of randomisation in randomised trials.* Although this reporting guideline is part of the CONSORT statement, it is so critical and so widely misunderstood that it bears repeating. The purpose of randomisation is to prevent selection bias. This can be achieved only if those consenting patients cannot guess a patient's treatment allocation before registration in the trial or change it afterward. This safeguard is known as 'allocation concealment'. Stating merely that 'a randomisation list was created by a statistician' or that 'envelope randomisation was used' does not ensure allocation concealment: a list could have been posted in the nurse's station for all to see; envelopes can be opened and resealed. Investigators need to specify the exact logistic steps taken to ensure allocation concealment. The best method is to use a password-protected computer database.
- 2.4 *The statistical methods should describe the study questions and the statistical approaches used to address each question.* Many statistical methods sections state

only something like 'Mann–Whitney was used for comparisons of continuous variables and Fisher's exact for comparisons of binary variables'. This says little more than 'the inference tests used were not grossly erroneous for the type of data'. Instead, statistical methods sections should lay out each primary study question separately: carefully detail the analysis associated with each and describe the rationale for the analytical approach, where this is not obvious or if there are reasonable alternatives. Special attention and description should be provided for rarely used statistical techniques.

- 2.5 *The statistical methods should be described in sufficient detail to allow replication by an independent statistician given the same data set.* Vague reference to 'adjusting for confounders' or 'non-linear approaches' is insufficiently specific to allow replication, a cornerstone of the scientific method. All statistical analyses should be specified in the Methods section, including details such as the covariates included in a multivariable model. All variables should be clearly defined where there is room for ambiguity. For instance, avoid saying that 'Gleason grade was included in the model'; state instead 'Gleason grade group was included in four categories 1, 2, 3 and 4 or 5'.

3 Inference and P values (see also 'Use and interpretation of P values' below)

- 3.1 *Do not accept the null hypothesis. In a court case, defendants are declared guilty or not guilty; there is no verdict of 'innocent'.* Similarly, in a statistical test, the null hypothesis is rejected or not rejected. If the P value is ≥ 0.05 , investigators should avoid conclusions such as 'the drug was ineffective', 'there was no difference between groups', or 'response rates were unaffected'. Instead, authors should use phrases such as 'we did not see evidence of a drug effect', 'we were unable to demonstrate a difference between groups', or simply 'there was no statistically significant difference in response rates'.
- 3.2 *P values just above 5% are not a trend, and they are not moving.* Avoid saying that a P value such as 0.07 shows a 'trend' (which is meaningless) or 'approaches statistical significance' (because the P value is not moving). Alternative language might be: 'although we saw some evidence of improved response rates in patients receiving the novel procedure, differences between groups did not meet conventional levels of statistical significance'.
- 3.3 *P values and 95% CIs do not quantify the probability of a hypothesis.* A P value of, say, 0.03 does not mean that there is 3% probability that the findings are due to chance. Additionally, a 95% CI should not be interpreted as a 95% certainty the true parameter value

is in the range of the 95% CI. The correct interpretation of a P value is the probability of finding the observed or more extreme results when the null hypothesis is true; the 95% CI will contain the true parameter value 95% of the time were a study to be repeated many times using different samples.

- 3.4 *Do not use CIs to test hypotheses.* Investigators often interpret CIs in terms of hypotheses. For instance, investigators might claim that there is a statistically significant difference between groups because the 95% CI for the odds ratio (OR) excludes 1. Such claims are problematic because CIs are concerned with estimation, not inference. Moreover, the mathematical method to calculate CIs may be different from those used to calculate P values. It is perfectly possible to have a 95% CI that includes no difference between groups even though the P value is <0.05 or vice versa. For instance, in a study of 100 patients in two equal groups, with event rates of 70% and 50%, the P value from Fisher's exact test is 0.066 but the 95% CI for the OR is 1.03–5.26. The 95% CI for the risk difference and risk ratio also exclude no difference between groups.
- 3.5 *Take care interpreting results when reporting multiple P values.* The more questions you ask, the more likely you are to get a spurious answer to at least one of them. For example, if you report P values for five independent true null hypotheses, the probability that you will falsely reject at least one is not 5%, but $>20\%$. Although formal adjustment of P values is appropriate in some specific cases, such as genomic studies, a more common approach is simply to interpret P values in the context of multiple testing. For instance, if an investigator examines the association of 10 variables with three different endpoints, thereby testing 30 separate hypotheses, a P value of 0.04 should not be interpreted in the same way as if the study tested only a single hypothesis with a P value of 0.04.
- 3.6 *Do not report separate P values for each of two different groups in order to address the question of whether there is a difference between groups.* One scientific question means one statistical hypothesis tested by one P value. To illustrate the error of using two P values to address one question, take the case of a randomised trial of drug vs placebo to reduce voiding symptoms, with 30 patients in each group. The authors might report that symptom scores improved by 6 (standard deviation 14) points in the drug group ($P = 0.03$ by one-sample t -test) and 5 (standard deviation 15) points in the placebo group ($P = 0.08$). However, the study hypothesis concerns the difference between drug and placebo. To test a single hypothesis, a single P value is needed. A two-sample t -test for these data gives a P value of 0.8; unsurprising, given that the scores in each group were virtually the same, confirming that it would be unsound to conclude that the drug was effective based on the finding that change was significant in the drug group but not in the placebo controls.
- 3.7 *Use interaction terms in place of subgroup analyses.* A similar error to the use of separate tests for a single hypothesis is when an intervention is shown to have a statistically significant effect in one group of patients but not another. One approach that is more appropriate is to use what is known as an 'interaction term' in a statistical model. For instance, to determine whether a drug reduced pain scores more in women than men, the model might be as follows: $\{\text{Final Pain Score}\} = \beta_0 + \beta_1\{\text{Baseline Pain Score}\} + \beta_2\{\text{Drug}\} + \beta_3\{\text{Sex}\} + \beta_4\{\text{Drug}\} \times \{\text{Sex}\}$. It is sometimes appropriate to report estimates and CIs within subgroups of interest, but P values should be avoided.
- 3.8 *Tests for change over time are generally uninteresting.* A common analysis is to conduct a paired t -test comparing, say, erectile function in older men at baseline with erectile function after 5 years of follow-up. The null hypothesis here is that 'erectile function does not change over time', which is known to be false. Investigators are encouraged to focus on estimation rather than inference, reporting, for example, the mean change over time along with a 95% CI.
- 3.9 *Avoid using statistical tests to determine the type of analysis to be conducted.* Numerous statistical tests are available that can be used to determine how a hypothesis test should be conducted. For instance, investigators might conduct a Shapiro–Wilk test for normality to determine whether to use a t -test or Mann–Whitney, Cochran's Q to decide whether to use a fixed- or random-effects approach in a meta-analysis, or use a t -test for between-group differences in a covariate to determine whether that covariate should be included in a multivariable model. The problem with these sorts of approaches is that they are often testing a null hypothesis that is known to be false. For instance, no data set perfectly follows a normal distribution. Moreover, it is often questionable that changing the statistical approach in the light of the test is actually of benefit. Statisticians are far from unanimous as to whether Mann–Whitney is always superior to t -test when data are non-normal, or that fixed effects are invalid under study heterogeneity, or that the criterion of adjusting for a variable should be whether it is significantly different between groups. Investigators should generally follow a pre-specified analytical plan, only altering the analysis if the data unambiguously point to a better alternative.

3.10 *When reporting P values, be clear about the hypothesis tested and ensure that the hypothesis is a sensible one. P values test very specific hypotheses. When reporting a P value in the results section, state the hypothesis being tested unless this is completely clear. Take, for instance, the statement 'Pain scores were higher in group 1 and similar in groups 2 and 3 ($P = 0.02$)'. It is ambiguous whether the P value of 0.02 is testing group 1 vs groups 2 and 3 combined or the hypothesis that pain score is the same in all three groups. Clarity about the hypotheses being tested can help avoid the testing of inappropriate hypotheses. For instance, P values for differences between groups at baseline in a randomised trial is testing a null hypothesis that is known to be true (informally, that any observed differences between groups are due to chance).*

4 Reporting of study estimates

4.1 *Use appropriate levels of precision. Reporting a P value of 0.7345 suggests that there is an appreciable difference between P values of 0.7344 and 0.7346. Reporting that 16.9% of 83 patients responded entails a precision (to the nearest 0.1%) that is nearly 200-times greater than the width of the CI (10% to 27%). Reporting in a clinical study that the mean calorie consumption was 2069.9 suggests that calorie consumption can be measured extremely precisely by a food questionnaire. Some might argue that being overly precise is irrelevant, because the extra numbers can always be ignored. The counter-argument is that investigators should think very hard about every number they report, rather than just carelessly cutting and pasting numbers from the statistical software printout. The specific guidelines for precision are as follows:*

- Report P values to a single significant figure unless the P is close to 0.05 (say, 0.01 to 0.2), in which case, report two significant figures. Do not report 'not significant' ('NS') for P values of ≥ 0.05 . Very low P values can be reported as $P < 0.001$ or similar. A P value can indeed be 1, although some investigators prefer to report this as > 0.9 . For instance, the following P values are reported to appropriate precision: < 0.001 , 0.004, 0.045, 0.13, 0.3, 1.
- Report percentages, rates and probabilities to two significant figures, e.g., 75%, 3.4%, 0.13%.
- Do not report P values of zero, as any experimental result has a non-zero probability.
- Do not give decimal places if a probability or proportion is 1 (e.g., a P value of 1.00 or a percentage of 100.00%). The decimal places suggest it is possible to have, say, a P value of 1.05. There is a similar consideration for data that can only take integer values.

It makes sense to state that, for instance, the mean number of pregnancies was 2.4, but not that 29% of women reported 1.0 pregnancies.

- There is generally no need to report estimates to more than three significant figures.
 - Hazard ratios (HRs) and ORs are normally reported to two decimal places, although this can be avoided for high ORs (e.g., 18.2 rather than 18.17).
- 4.2 *Avoid redundant statistics in cohort descriptions. Authors should be selective about the descriptive statistics reported and ensure that each and every number provides unique information. Authors should avoid reporting descriptive statistics that can be readily derived from data that have already been provided. For instance, there is no need to state that 40% of a cohort were men and 60% were women, choose one or the other. Another common error is to include a column of descriptive statistics for two groups separately and then the whole cohort combined. If, say, the median age is 60 years in group 1 and 62 years in group 2, we do not need to be told that the median age in the cohort as a whole is close to 61 years.*
- 4.3 *For descriptive statistics, median and quartiles are preferred over means and standard deviations (or standard errors); range should be avoided. The median and quartiles provide all sorts of useful information, for instance, that 50% of patients had values above the median or between the quartiles. The range gives the values of just two patients and so is generally uninformative of the data distribution.*
- 4.4 *Report estimates for the main study questions. A clinical study typically focuses on a limited number of scientific questions. Authors should generally provide an estimate for each of these questions. In a study comparing two groups, for instance, authors should give an estimate of the difference between groups, and avoid giving only data on each group separately or, simply saying that the difference was or was not significant. In a study of a prognostic factor, authors should give an estimate of the strength of the prognostic factor, such as an OR or HR, as well as reporting a P value testing the null hypothesis of no association between the prognostic factor and outcome.*
- 4.5 *Report CIs for the main estimates of interest. Authors should generally report a 95% CI around the estimates relating to the key research questions, but not other estimates given in a paper. For instance, in a study comparing two surgical techniques, the authors might report adverse event rates of 10% and 15%; however, the key estimate in this case is the difference between groups, so this estimate, 5%, should be reported along*

with a 95% CI (e.g., 1–9%). CIs should not be reported for the estimates within each group (e.g., adverse event rate in group A of 10%, 95% CI 7–13%). Similarly, CIs should not be given for statistics such as mean age or gender ratio.

- 4.6 *Do not treat categorical variables as continuous.* A variable such as Gleason grade groups are scored 1–5, but it is not true that the difference between group 3 and 4 is half as great as the difference between group 2 and 4. Variables such as Gleason grade group should be reported as categories (e.g., 40% grade group 1, 20% group 2, 20% group 3, 20% group 4 and 5) rather than as a continuous variable (e.g., mean Gleason score of 2.4). Similarly, categorical variables such as Gleason grade should be entered into regression models not as a single variable (e.g., a HR of 1.5 per 1-point increase in Gleason grade group) but as multiple categories (e.g., a HR of 1.6 comparing Gleason grade group 2 to group 1 and HR of 3.9 comparing group 3 to group 1).
- 4.7 *Avoid categorisation of continuous variables unless there is a convincing rationale.* A common approach to a variable such as age is to define patients as either old (≥ 60 years) or young (< 60 years) and then enter age into analyses as a categorical variable, reporting, for example, that ‘patients aged ≥ 60 years had twice the risk of an operative complication than patients aged < 60 years’. In epidemiological and marker studies, a common approach is to divide a variable into quartiles and report a statistic such as a HR for each quartile compared to the lowest (‘reference’) quartile. This is problematic because it assumes that all values of a variable within a category are the same. For instance, it is likely not the case that a patient aged 61 years has the same risk as a patient aged 90 years, but a very different risk to a patient aged 59 years. It is generally preferable to leave variables in a continuous form, reporting, for instance, how risk changes with a 10-year increase in age. Non-linear terms can also be used, to avoid the assumption that the association between age and risk follows a straight line.
- 4.8 *Do not use statistical methods to obtain thresholds for clinical practice.* There are various statistical methods available to dichotomise a continuous variable. For instance, outcomes can be compared either side of several different thresholds, and the optimal threshold chosen as the one associated with the smallest P value. Alternatively, investigators might choose a threshold that leads to the highest value of sensitivity and specificity, that is, the point closest to the top left-hand corner of a receiver operating characteristic (ROC) curve. Such methods are inappropriate for determining clinical thresholds because they do not consider clinical consequences. The ROC curve approach, for instance, assumes that sensitivity and specificity are of equal value, whereas it is generally worse to miss disease than to treat unnecessarily. The smallest P value approach tests strength of evidence against the null hypothesis, which has little to do with the relative benefits and harms of a treatment or further diagnostic evaluation.
- 4.9 *The association between a continuous predictor and outcome can be demonstrated graphically, particularly by using non-linear modelling.* In high-school mathematics, we often thought about the relationship between y and x by plotting a line on a graph, with a scatterplot added in some cases. This also holds true for many scientific studies. In the case of a study of age and complication rates, for instance, an investigator could plot age on the x -axis against risk of a complication on the y -axis and show a regression line, perhaps with a 95% CI. Non-linear modelling is often useful because it avoids assuming a linear relationship and allows the investigator to determine questions such as whether risk starts to increase disproportionately beyond a given age.
- 4.10 *Do not ignore significant heterogeneity in meta-analyses.* Informally speaking, heterogeneity statistics test whether variations between the results of different studies in a meta-analysis are consistent with chance, or whether such variation reflects, at least in part, true differences between studies. If heterogeneity is present, authors need to do more than merely report the P value and focus on the random-effects estimate. Authors should investigate the sources of heterogeneity and try to determine the factors that lead to differences in study results, e.g., by identifying common features of studies with similar findings or idiosyncratic aspects of studies with outlying results.
- 4.11 *For time-to-event variables, report the number of events but not the proportion.* Take the case of a study that reported: ‘of 60 patients accrued, 10 (17%) died’. While it is important to report the number of events, patients entered the study at different times and were followed for different periods, so the reported proportion of 17% is meaningless. The standard statistical approach to time-to-event variables is to calculate probabilities, such as the risk of death being 60% by 5 years or the median survival, the time at which the probability of survival first drops below 50%, being 52 months.
- 4.12 *For time-to-event analyses, report median follow-up for patients without the event or the number followed without an event at a given follow-up time.* It is often useful to describe how long a cohort has been followed. To illustrate the appropriate methods of doing so, take the case of a cohort of 1000 paediatric

patients with cancer treated in 1970 and followed to 2010. If the cure rate was only 40%, the median follow-up for all patients might only be a few years, whilst the median follow-up for patients who survived was 40 years. This latter statistic gives a much better impression of how long the cohort had been followed. Now assume that in 2009, a second cohort of 2000 patients was added to the study. The median follow-up for survivors will now be around a year, which is again misleading. An alternative would be to report a statistic such as ‘312 patients have been followed without an event for at least 35 years’.

- 4.13 *For time-to-event analyses, describe when follow-up starts and when and how patients are censored. A common error is that investigators use a censoring date, which leads to an overestimate of survival. For example, when assessing metastasis-free survival a patient without a record of metastasis should be censored on the date of the last time the patient was known to be free of metastasis (e.g., negative bone scan, undetectable PSA), not at the date of last patient contact (which may not have involved assessment of metastasis). For overall survival, date of last patient contact would be an acceptable censoring date because the patient was indeed known to be event-free at that time. When assessing cause-specific endpoints, special consideration should be given to the cause of death. The endpoints ‘disease-specific survival’ and ‘disease-free survival’ have specific definitions and require careful attention to methods. With disease-specific survival, authors need to consider carefully how to handle death due to other causes. One approach is to censor patients at the time of death, but this can lead to bias in certain circumstances, such as when the predictor of interest is associated with other cause of death and the probability of other cause of death is moderate or high. Competing risk analysis is appropriate in these situations. With disease-free survival, both evidence of disease (e.g., disease recurrence) and death from any cause are counted as events, and so censoring at the time of other cause of death is inappropriate. If investigators are specifically interested only in the former, and wish to censor deaths from other causes, they should define their endpoint as ‘freedom from progression’.*
- 4.14 *For time-to-event analyses, avoid reporting mean follow-up or survival time, or estimates of survival in those who had the event. All three estimates are problematic in the context of censored data.*
- 4.15 *For time-to-event analyses, make sure that all predictors are known at time zero or consider alternative approaches such as a landmark analysis or time-dependent covariates. In many cases, variables of*

interest vary over time. As a simple example, imagine we were interested in whether PSA velocity predicted time to progression in patients with prostate cancer on active surveillance. The problem is that PSA is measured at various times after diagnosis. Unless they were being careful, investigators might use time from diagnosis in a Kaplan–Meier or Cox regression but use PSA velocity calculated on PSAs measured at 1- and 2-year follow-up. As another example, investigators might determine whether response to chemotherapy predicts cancer survival, but measure survival from the time of the first dose, before response is known. It is obviously invalid to use information only known ‘after the clock starts’. There are two main approaches to this problem. A ‘landmark analysis’ is often used when the variable of interest is generally known within a short and well-defined period of time, such as adjuvant therapy or chemotherapy response. In brief, the investigators start the clock at a fixed ‘landmark’ (e.g., 6 months after surgery). Patients are only eligible if they are still at risk at the landmark (e.g., patients who recur before 6 months are excluded) and the status of the variable is fixed at that time (e.g., a patient who gets chemotherapy at 7 months is defined as being in the no adjuvant group). Alternatively, investigators can use a time-dependent variable approach. In brief, this ‘resets the clock’ each time new information is available about a variable. This would be the approach most typically used for the PSA velocity and progression example.

- 4.16 *When presenting Kaplan–Meier figures, present the number at risk and truncate follow-up when numbers are low. Giving the number at risk is useful for helping to understand when patients were censored. When presenting Kaplan–Meier figures, a good general rule is to truncate follow-up when the number at risk in any group falls below five (or even 10) as the tail of a Kaplan–Meier distribution is very unstable.*

5 Multivariable models and diagnostic tests

- 5.1 *Multivariable, propensity and instrumental variable analyses are not a magic wand. Some investigators assume that multivariable adjustment ‘removes confounding’, ‘makes groups similar’ or ‘mimics a randomised trial’. There are two problems with such claims. First, the value of a variable recorded in a data set is often approximate and so may mask differences between groups. For instance, clinical stage might be used as a covariate in a study comparing treatments for localised prostate cancer. But stage T2c might constitute a small nodule on each prostate lobe or, alternatively, most of the prostate consisting of a large,*

hard mass. The key point is that if one group has more T2c disease than the other, it is also likely that the T2c's in that group will fall towards the more aggressive end of the spectrum. Multivariable adjustment has the effect of making the rates of T2c in each group the same, but does not ensure that the type of T2c is identical. Second, a model only adjusts for a small number of measured covariates. That does not exclude the possibility of important differences in unmeasured (or even unmeasurable) covariates. A common assumption is that propensity methods somehow provide better adjustment for confounding than traditional multivariable methods. Except in certain rare circumstances, such as when the number of covariates is large relative to the number of events, propensity methods give extremely similar results to multivariable regression. Similarly, instrumental variables analyses depend on the availability of a good instrument, which is less common than is often assumed. In many cases, the instrument is not strongly associated with the intervention, leading to a large increase in the 95% CI or, in some cases, an underestimate of treatment effects.

- 5.2 *Avoid stepwise selection. Investigators commonly choose which variables to include in a multivariable model by first determining which variables are statistically significant on univariable analysis; alternatively, they may include all variables in a single model but then remove any that are not significant.* This type of data-dependent variable selection in regression models has several undesirable properties, increasing the risk of overfit and making many statistics, such as the 95% CI, highly questionable. Use of stepwise selection should be restricted to a limited number of circumstances, such as during the initial stages of developing a model, if there is poor knowledge of what variables might be predictive.
- 5.3 *Avoid reporting estimates such as ORs or HRs for covariates when examining the effects of interventions.* In a typical observational study, an investigator might explore the effects of two different approaches to radical prostatectomy on recurrence while adjusting for covariates such as stage, grade and PSA. It is rarely worth reporting estimates such as ORs or HRs for the covariates. For instance, it is well known that a high Gleason score is strongly associated with recurrence: reporting a HR of say, 4.23, is not helpful and a distraction from the key finding, the HR between the two types of surgery.
- 5.4 *Rescale predictors to obtain interpretable estimates.* Predictors sometimes have a moderate association with outcome and can take a large range of values. This can lead to uninterpretable estimates. For instance, the OR for cancer per year of age might be given as 1.02 (95% CI 1.01–1.02; $P < 0.001$). It is not helpful to have the upper bound of a CI be equivalent to the central estimate; a better alternative would be to report an OR per 10 years of age. This is simply achieved by creating a new variable equal to age divided by 10 to obtain an OR of 1.16 (95% CI 1.10–1.22; $P < 0.001$) per 10-year difference in age.
- 5.5 *Avoid reporting both univariate and multivariable analyses unless there is a good reason.* Comparison of univariate and multivariable models can be of interest when trying to understand mechanisms. For instance, if race is a predictor of outcome on univariate analysis, but not after adjustment for income and access to care, one might conclude that poor outcome in African-Americans is explained by socioeconomic factors. However, the routine reporting of estimates from both univariate and multivariable analysis is discouraged.
- 5.6 *Avoid ranking predictors in terms of strength.* It is tempting for authors to rank predictors in a model, claiming, for instance, 'the novel marker was the strongest predictor of recurrence'. Most commonly, this type of claim is based on comparisons of ORs or HRs. Such rankings are not meaningful as, amongst other reasons, it depends on how variables are coded. For instance, the OR for human kallikrein 2 (hK2), and hence whether or not it is an apparently 'stronger' predictor than PSA, will depend on whether it is entered in nanogrammes or picogrammes per millilitre. Further, it is unclear how one should compare model coefficients when both categorical and continuous variables are included. Finally, the prevalence of a categorical predictor also matters: a predictor with an OR is 3.5 but a prevalence of 0.1% is less important than one with a 50% prevalence and an OR of 2.0.
- 5.7 *Discrimination is a property not of a multivariable model but rather of the predictors and the data set.* Although model building is generally seen as a process of fitting coefficients, discrimination is largely a property of what predictors are available. For instance, we have excellent models for prostate cancer outcome primarily because Gleason score is very strongly associated with malignant potential. In addition, discrimination is highly dependent on how much a predictor varies in the data set. As an example, a model to predict erectile dysfunction that includes age will have much higher discrimination for a population sample of adult men than for a group of older men presenting at a urology clinic, because there is a greater variation in age in the population sample. Authors need to consider these points when drawing conclusions about the discrimination of models. This

is also why authors should be cautious about comparing the discrimination of different multivariable models where these were assessed in different datasets.

- 5.8 *Correction for overfit is strongly recommended for internal validation.* In the same way that it is easy to predict last week's weather, a prediction model generally has very good properties when evaluated on the same data set used to create the model. This problem is generally described as overfit. Various methods are available to correct for overfit, including cross-validation and bootstrap resampling. Note that such methods should include all steps of model building. For instance, if an investigator uses stepwise methods to choose which predictors should go into the model and then fits the coefficients, a typical cross-validation approach would be to: (i) split the data into 10 groups, (ii) use stepwise methods to select predictors using the first nine groups, (iii) fit coefficients using the first nine groups, (iv) apply the model to the 10th group to obtain predicted probabilities, and (v) repeat steps ii–iv until all patients in the data set have a predicted probability derived from a model fitted to a data set that did not include that patient's data. Statistics such as the area under the curve (AUC) are then calculated using the predicted probabilities directly.
- 5.9 *Calibration should be reported and interpreted correctly.* Calibration is a critical component of a statistical model: the main concern for any patient is whether the risk given by a model is close to his or her true risk. It is rarely worth reporting calibration for a model created and tested on the same data set, even if techniques such as cross-validation are used. This is because calibration is nearly always excellent on internal validation. Where a pre-specified model is tested on an independent data set, calibration should be displayed graphically in a calibration plot. The Hosmer–Lemeshow test addresses an inappropriate null hypothesis and should be avoided. Note also that calibration depends upon both the model coefficients and the dataset being examined. A model cannot be inherently 'well calibrated'. All that can be said is that predicted and observed risk are close in a specific data set, representative of a given population.
- 5.10 *Avoid reporting sensitivity and specificity for continuous predictors or a model.* Investigators often report sensitivity and specificity at a given threshold for a continuous predictor (e.g., a PSA level of 10 ng/mL), or report specificity at a given sensitivity (e.g., 90%). Reporting sensitivity and specificity is not of value because it is unclear how high sensitivity or specificity would have to be so as to be high enough to justify clinical use. Similarly, it is very difficult to determine which of two tests, one with a higher sensitivity and

the other with a higher specificity, is preferable because clinical value depends on the prevalence of disease and the relative harms of a false-positive compared with a false-negative result. In the case of reporting specificities at fixed sensitivity, or vice versa, it is all but impossible to choose the sensitivity rationally. For instance, a team of investigators may state that they want to know specificity at 80% sensitivity, because they want to ensure they catch 80% of cases. But 80% might be too low if prevalence is high or too high if prevalence is low.

- 5.11 *Report the clinical consequences of using a test or a model.* In place of statistical abstractions, such as sensitivity and specificity, or a ROC curve, authors are encouraged to choose illustrative thresholds and then report results in terms of clinical consequences. As an example, consider a study in which a marker is measured in a group of patients undergoing biopsy. Authors could report that if a given level of the marker had been used to determine biopsy, then a certain number of biopsies would have been conducted and a certain number of cancers found and missed.
- 5.12 *Interpret decision curves with careful reference to threshold probabilities.* It is insufficient merely to report that, for instance, 'the marker model had highest net benefit for threshold probabilities of 35–65%'. Authors need to consider whether those threshold probabilities are rational. If the study reporting benefit between 35% and 65% concerned detection of high-grade prostate cancer, few if any urologists would demand that a patient have at least a one-in-three chance of high-grade disease before recommending biopsy. The authors would therefore need to conclude that the model was not of benefit.

6 Conclusions and interpretation

- 6.1 *Draw a conclusion, do not just repeat the results.* Conclusion sections are often simply a restatement of the results. For instance, 'a statistically significant relationship was found between body mass index (BMI) and disease outcome' is not a conclusion. Authors instead need to state implications for research and/or clinical practice. For instance, a conclusion section might call for research to determine whether the association between BMI is causal or make a recommendation for more aggressive treatment of patients with higher BMI.
- 6.2 *Avoid using words such as 'may' or 'might'.* A conclusion such as that a novel treatment 'may' be of benefit would only be untrue if it had been proven that the treatment was ineffective. Indeed, that the treatment may help would have been the rationale for the study in the first place. Using words such as 'may'

in the conclusion is equivalent to stating, 'we know no more at the end of this study than we knew at the beginning', reason enough to reject a paper for publication.

- 6.3 *A statistically significant P value does not imply clinical significance.* A small *P* value means only that the null hypothesis has been rejected. This may or may not have implications for clinical practice. For instance, that a marker is a statistically significant predictor of outcome does not imply that treatment decisions should be made on the basis of that marker. Similarly, a statistically significant difference between two treatments does not necessarily mean that the former should be preferred to the latter. Authors need to justify any clinical recommendations by carefully analysing the clinical implications of their findings.
- 6.4 *Avoid pseudo-limitations such as 'small sample size' and 'retrospective analysis', consider instead sources of potential bias and the mechanism for their effect on findings.* Authors commonly describe study limitations in a rather superficial way, such as, 'small sample size and retrospective analysis are limitations'. But a small sample size may be immaterial if the results of the study are clear. For instance, if a treatment or predictor is associated with a very large OR, a large sample size might be unnecessary. Similarly, a retrospective design might be entirely appropriate, as in the case of a marker study with very long-term follow-up, and have no discernible disadvantages compared to a prospective study. Discussion of limitations should include both the likelihood and effect size of possible bias.
- 6.5 *Consider the impact of missing data and patient selection.* It is rare that complete data are obtained from all patients in a study. A typical paper might report, for instance, that of 200 patients, eight had data missing on important baseline variables and 34 did not complete the end of study questionnaire, leading to a final data set of 158. Similarly, many studies include a relatively narrow subset of patients, such as 50 patients referred for imaging before surgery, out of the 500 treated surgically during that timeframe. In both cases, it is worth considering analyses to investigate whether patients with missing data or who were not selected for treatment were different in some way from those who were included in the analyses. Although statistical adjustment for missing data is complex and is warranted only in a limited set of circumstances, basic analyses to understand the characteristics of patients with missing data are relatively straightforward and are often helpful.
- 6.6 *Consider the possibility and impact of ascertainment bias.* Ascertainment bias occurs when an outcome

depends on a test, and the propensity for a patient to be tested is associated with the predictor. PSA screening provides a classic example: prostate cancer is found by biopsy, but the main reason why men are biopsied is because of an elevated PSA. A study in a population subject to PSA screening will therefore overestimate the association between PSA and prostate cancer. Ascertainment bias can also be caused by the timing of assessments. For instance, frequency of biopsy in prostate cancer active surveillance will depend on prior biopsy results and PSA levels, and this induces an association between those predictors and time to progression.

- 6.7 *Do not confuse outcome with response. Amongst subgroups of patients undergoing the same treatment, patients with poorer outcomes may still be good candidates for that treatment.* Investigators often compare outcomes in different subgroups of patients all receiving the same treatment. A common error is to conclude that patients with poor outcome are not good candidates for that treatment and should receive an alternative approach. This is to confuse differences between patients for differences between treatments. As a simple example, patients with large tumours are more likely to recur after surgery than patients with small tumours, but that cannot be taken to suggest that resection is not indicated for patients with tumours larger than a certain size. Indeed, surgery is generally more strongly indicated for patients with aggressive (but localised) disease and such patients are unlikely to do well on surveillance.
- 6.8 *Be cautious about causal attribution: correlation does not imply causation.* It is well-known that 'correlation does not imply causation' but authors often slip into this error in making conclusions. The introduction and methods section might insist that the purpose of the study is merely to determine whether there is an association between, say, treatment frequency and treatment response, but the conclusions may imply that, for instance, more frequent treatment would improve response rates.

Use and Interpretation of *P* values

That *P* values are widely misused and misunderstood is apparent from even the most cursory reading of the medical literature. One of the most common errors is accepting the null hypothesis, for instance, concluding from a *P* value of 0.07 that a drug is ineffective or that two surgical techniques are equivalent. This particular error is described in detail in guideline 3.1. The more general problem, which we address here, is that *P* values are often given excessive weight in the interpretation of a study. Indeed, studies are often classed by investigators into 'positive' or 'negative' based on statistical

significance. Gross misuse of P values has led some to advocate banning the use of P values completely [4].

We follow the American Statistical Association statement on P values and encourage all researchers to read either the full statement [5] or the summary [6]. In particular, we emphasise that the P value is just one statistic that helps interpret a study; it does not determine our interpretations. Drawing conclusions for research or clinical practice from a clinical research study requires evaluation of the strengths and weakness of study methodology, the results of other pertinent data published in the literature, biological plausibility, and effect size. Sound and nuanced scientific judgment cannot be replaced by just checking whether one of the many statistics in a paper is or is not $P < 0.05$.

Concluding Remarks

These guidelines are not intended to cover all medical statistics but rather the statistical approaches most commonly used in clinical research papers in urology. It is quite possible for a paper to follow all of the guidelines yet be statistically flawed or to break numerous guidelines and still be statistically sound. On balance, however, the analysis, reporting and interpretation of clinical urological research will be improved by adherence to these guidelines.

Funding

Supported in part by the Sidney Kimmel Center for Prostate and Urologic Cancers, P50-CA92629 Specialized Program of Research Excellence (SPORE) grant from the National Cancer Institute (NCI) to Dr H. Scher, and the P30-

CA008748 National Institutes of Health (NIH)/NCI Cancer Center Support Grant to Memorial Sloan-Kettering Cancer Center.

Conflict of Interest

None declared.

References

- 1 Scales CD Jr, Norris RD, Peterson BL, Preminger GM, Dahm P. Clinical research and statistical methods in the urology literature. *J Urol* 2005; 174: 1374–9
- 2 Lang TA, Altman DG. Basic statistical reporting for articles published in biomedical journals: the “Statistical Analyses and Methods in the Published Literature” or the SAMPL Guidelines. *Int J Nurs Stud* 2015; 52: 5–9
- 3 Vickers AJ, Sjoberg DD. Guidelines for reporting of statistics in *European Urology*. *Eur Urol* 2015; 67: 181–7
- 4 Woolston C. Psychology journal bans P values. *Nature* 2015; 519: 9
- 5 Wasserstein RL, Lazar NA. The ASA’s statement on p -values: context, process, and purpose. *Am Stat* 2016; 70: 129–33
- 6 American Statistical Association. American Statistical Association releases statement on statistical significance and P -values: provides principles to improve the conduct and interpretation of quantitative science. Available at: <https://www.amstat.org/asa/files/pdfs/P-ValueStatement.pdf>. Accessed December 2018

Correspondence: Andrew J. Vickers, Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY, USA.

e-mail: vickersa@mskcc.org

Abbreviations: BMI, body mass index; CONSORT, Consolidated Standards of Reporting Trials; HR, hazard ratio; OR, odds ratio; ROC, receiver operating characteristic.