

DR ELIZABETH C VERNA (Orcid ID : 0000-0002-9658-3751)

DR AVIN AGGARWAL (Orcid ID : 0000-0001-8239-7563)

Article type : D - Comprehensive Review

Liver Transplantation for Hepatocellular Carcinoma: Management After the Transplant

Elizabeth C. Verna, MD, MS*¹, Yuval A. Patel, MD, MHS*², Avin Aggarwal, MD³, Archita P. Desai, MD⁴, Catherine Frenette, MD⁵, Anjana A. Pillai, MD⁶, Reena Salgia, MD⁷, Anil Seetharam, MD⁸, Pratima Sharma, MD, MS⁹, Courtney Sherman, MD¹⁰, Georgios Tsoulfas, MD¹¹, Francis Y. Yao, MD¹⁰ *This manuscript is a work-product of the American Society of Transplantation Liver Intestine Community of Practice*

*Co-First Authors

¹Center for Liver Disease and Transplantation, Columbia University, New York, NY, USA

²Division of Gastroenterology, Department of Medicine, Duke University, Durham, NC, USA

³Department of Medicine, Division of Gastroenterology and Hepatology, University of Arizona College of Medicine, Tucson, AZ, USA

⁴Division of Gastroenterology, Department of Medicine, Indiana University, Indianapolis, IN, USA

⁵Scripps Center for Organ Transplantation, Scripps Green Hospital, La Jolla, CA, USA

⁶Center for Liver Diseases, University of Chicago Medicine, Chicago, IL, USA

⁷ Department of Gastroenterology/Hepatology, Henry Ford Hospital, Detroit, MI, USA

⁸ Transplant Hepatology, University of Arizona College of Medicine, Phoenix, AZ, USA

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/AJT.15697](https://doi.org/10.1111/AJT.15697)

This article is protected by copyright. All rights reserved

⁹Michigan Medicine, University of Michigan, Ann Arbor, MI, USA

¹⁰ Division of Gastroenterology, Department of Medicine, University of California, San Francisco, San Francisco, CA, USA

¹¹Department of Surgery, Aristotle University School of Medicine, Thessaloniki Greece

Corresponding Author:

Elizabeth C. Verna

email: ev77@cumc.columbia.edu

Abbreviations:

alpha-fetoprotein (AFP); calcineurin inhibitor (CNI); cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4); death protein 1/programmed death ligand 1 (PD-1/PDL-1); des-gamma-carboxyprothrombin (DCP); ¹⁸F-fluorodeoxyglucose positron emission tomography/CT (¹⁸F-FDG PET/CT); hepatocellular carcinoma (HCC); liver transplantation (LT); local regional therapy (LRT); mammalian target of rapamycin (mTOR); microvascular invasion (MVI); neutrophil/lymphocyte ratio (NLR); transcatheter arterial chemoembolization (TACE); University of California San Francisco (UCSF); vascular endothelial growth factor (VEGF); Yttrium-90 microspheres (Y-90);

Abstract:

Hepatocellular carcinoma (HCC) is an increasingly common indication for liver transplantation (LT) in the United States and in many parts of the world. In the last decade, significant work has been done to better understand how to risk stratify LT candidates for recurrence of HCC following transplant using a combination of biomarker and imaging findings. However, despite the high frequency of HCC in the LT population, guidance regarding post-transplant management is lacking. In particular, there is no current evidence to support specific post-LT surveillance strategies, leading to

significant heterogeneity in practices. In addition, there are no current recommendations regarding recurrence prevention, including immunosuppression regimen or secondary prevention with adjuvant chemotherapy. Finally, guidance on treatment of disease recurrence is also lacking and there is significant controversy about the use of immunotherapy in transplant recipients due to the risk of rejection. Thus, outcomes for patients with recurrence are poor. This paper therefore provides a comprehensive review of the current literature on post-LT management of patients with HCC, and identifies gaps in our current knowledge that are in urgent need of further investigation.

Introduction:

In this era of rising hepatocellular carcinoma (HCC) incidence, HCC is an increasingly common indication for liver transplantation (LT). In 2015, HCC was the indication for 24% of liver transplant registrants and 27% of liver transplants, rendering it the most common reason for LT and waitlist additions, regardless of underlying etiology.¹

LT began to evolve as a therapy for HCC when the incidental finding of small HCC in explanted livers was not found to alter outcomes as compared to explants without HCC². The landmark study by Mazzaferro in 1996 then established LT as an effective treatment for early HCC defined by the Milan Criteria (one lesion ≤ 5 cm or 3 lesions all ≤ 3 cm without evidence of vascular invasion or extrahepatic spread³. Survival following LT for HCC has improved over time with advances in care and is similar to that of nonmalignant indications⁴⁻⁸.

Though LT for HCC is a highly effective cure for early stage disease, guidance regarding tailored post-transplant management of this unique population to optimize outcomes is lacking. Given the heterogeneity of HCC burden and tumor biology seen in patients that present for transplantation, it is essential for transplant providers to consider the unique features of a given patient's HCC when devising their post-transplant management plan. This paper aims to comprehensively review the current literature on post-transplant management of patients with HCC, as well as identify gaps in our current knowledge that are in urgent need of further investigation.

HCC Recurrence: Magnitude of the Problem

While LT provides an excellent treatment option for long-term survival in selected patients with HCC, post-transplant HCC recurrence is an important negative predictor of post-transplant survival. In the initial study by Mazzaferro and colleagues evaluating the role of LT in cirrhotic patients with small unresectable HCC, 8% of patients experienced HCC recurrence by four years after LT³. Subsequently, studies focused on patients within Milan criteria using pre-transplant data have described post-transplant recurrence in approximately 10-16% of patients⁹⁻¹¹. A 2015 systematic review including a heterogeneous group of 61 studies demonstrated a mean rate of HCC recurrence of 16% and mean time from transplant to HCC recurrence of 13 months (range 2-132 months).¹² Of note, nearly 51% of LT recipients in this review were ultimately classified as beyond Milan based upon examination of patient pathology. Notably, as many centers worldwide transplant patients with HCC that are beyond Milan criteria, the magnitude of HCC recurrence may be larger than proposed by these estimates.

Prediction of HCC Recurrence

Multiple tumor related factors have been identified to predict the risk of HCC recurrence after LT, and these factors have been incorporated in a number of proposed pre-LT and post-LT prognostic models (Tables 1 and 2).

Well-Established Pre-Transplant Factors Predicting HCC Recurrence

Radiographic tumor burden Traditionally, tumor diameter and number of tumor nodules have been used as criteria determining eligibility for LT, based on the observation that these morphologic parameters are associated with microvascular invasion (MVI) and HCC recurrence after LT. The Milan criteria³ have been incorporated into the UNOS/OPTN policy for priority listing with MELD-exception for LT since 2002 (modified as UNOS T2 criteria). Modest expansion of the upper limits of tumor burden, including the University of California San Francisco (UCSF) criteria^{13,14} and the “up-to-7” criteria¹⁵, have resulted in survival only slightly below that using Milan criteria. Further expansion of tumor criteria would result in significantly reduced post-transplant survival, as illustrated in the “Metro-ticket” paradigm of “the further the distance, the higher the price”.¹⁵ The decision regarding expansion of transplant tumor

criteria must be governed by the ability to achieve a minimal survival threshold to justify expansion and not cause harm to non-HCC patients on the waiting list.^{3,16}

A strategy combining tumor burden with assessment of response to local regional therapy (LRT) over time as a marker of favorable tumor biology has gained broader acceptance. Patients who exhibit tumor progression despite LRT have significantly worse post-LT outcomes when compared to those who demonstrate treatment response or stable disease following LRT¹⁷⁻¹⁹. Down-staging represents a structured approach that aims at merging tumor morphologic parameters with objective and sustained response to LRT²⁰ as an additional risk stratification tool. This approach is supported by the observation that post-LT survival outcomes in those who have been successfully down-staged to conventional Milan criteria and are not significantly different than those who meet Milan criteria at presentation.²⁰ In an effort to standardize criteria for down-staging of HCC prior to LT, OPTN/UNOS recently adopted the UCSF / Region 5 down-staging protocol with specific inclusion criteria^{20,21} as a new national policy for granting automatic MELD exception for LT. Initial tumor burden beyond these criteria have been suggested to have an adverse impact on both the probability of successful down-staging²² and worse post-LT survival.²³ It has been suggested that there is a strong correlation between the initial tumor burden assessed by the sum of the largest tumor diameter and the probability of successful down-staging to within Milan criteria.²² Furthermore, extending the initial tumor burden beyond the current UNOS down-staging inclusion criteria is associated with a significantly worse 3-year post-LT survival in an analysis of UNOS data.²³

Despite advances in cross-sectional imaging over the last few decades, radiographic under-staging of HCC still occurs in up to 25-30% of patients.²⁴⁻²⁷ Furthermore, misdiagnosis has been reported in 11-25% (no HCC found in explant).^{28,29} To reduce the misdiagnosis rate, a revised UNOS/OPTN imaging policy for HCC was implemented, requiring structured reporting, standardized imaging protocols for LT centers, and equipment specifications for dynamic contrast-enhanced multiphase CT or MRI.^{30,31}

Alpha-fetoprotein. Growing evidence suggests alpha-fetoprotein (AFP) to be a powerful prognostic biomarker in LT³². AFP has been included in virtually every

prognostic model in LT for HCC (Tables 1 and 2), but there is no consensus on an AFP cut off value to be applied as an exclusion criterion for LT. AFP levels as low as 16 to 20 ng/mL have been associated with poor post-LT outcomes^{3,32}. A much higher AFP level of >1000 ng/mL before LT is associated with particularly poor post-LT survival, regardless of the tumor burden^{33,34}. A new UNOS policy has recently been implemented, requiring those with an AFP >1000 ng/mL to show a decrease in AFP to <500 ng/mL with LRT before LT can be undertaken. This policy is supported by a recent analysis of UNOS data showing that a decrease in AFP from >1000 to 101-499 ng/mL before LT, mostly as a result of LRT, would result in a greater than two-fold reduction in post-LT mortality and an almost threefold reduction in HCC recurrence.³⁵

Many other approaches in using AFP to refine LT selection criteria have been explored. A multi-center French study incorporated a number of AFP thresholds within a prognostic score that included AFP, largest tumor diameter, and number of tumor nodules³⁴. The Metroticket version 2.0 presented a continuum of AFP in combination with the sum of the largest tumor diameter and number of tumor nodules to predict post-LT survival³⁶. A positive AFP slope resulting from rising AFP values over time also appears to correlate with a higher risk of post-LT HCC recurrence^{19,37-40}, but the AFP slope threshold that predicts worse post-LT outcomes has varied from one study to another. It has also been shown in several large series that pre-LT AFP response to LRT significantly impacts recurrence.^{41,42} It is important to note that AFP is not entirely specific to HCC-related production and can be elevated in the setting of hepatic inflammation, including in the setting of viral hepatitis.

Well-Established Explant Features Predicting HCC Recurrence

Microvascular Invasion. The presence of microvascular invasion (MVI) on explant is strongly associated with HCC recurrence and reduced survival after LT^{43,44} (Table 2). The incidence of MVI is almost twice as high in tumors larger than 5 cm compared to smaller tumors^{27,45-48}. Patients with multiple tumors were 2 times more likely to have MVI as a solitary tumor⁴⁹. AFP >1000 ng/ml⁴⁸ and positive uptake on PET⁵⁰ have also been associated with a greater likelihood of MVI. Nevertheless, the lack of a standardized histological definition of MVI creates heterogeneity across

studies, and the presence or absence of MVI cannot be reliably determined prior to LT (45). Biopsy of the tumor has a very low sensitivity for MVI detection⁵¹. A noninvasive approach involving a contrast-enhanced computed tomography based biomarker derived from a HCC “venous invasion gene” gene expression signature, referred to as “radiogenomics”⁵¹, has shown promise, but the usefulness of radiogenomics in LT candidates with early stage HCC requires further study. Similarly, the creation of a radiographic-radiomic model using clinical and contrast-enhanced computed tomography factors to predict MVI for HCC may have an application for early stage HCC patients considering LT, though further research is needed.⁵²

Histological Grade of Tumor Differentiation. Poorly differentiated tumor grade has been identified in many studies as an important risk factor for HCC recurrence after LT^{46,53,54} (Table 2). Several groups have proposed using pre-LT biopsies of the largest tumor to exclude patients with poorly differentiated HCC from LT, while placing no restrictions in the upper limits in tumor burden as long as there was no radiographic evidence of macro-vascular invasion. Based on this approach, the University of Padova group reported a 5-year actuarial survival rate of 75% and a recurrence-free probability of 92%⁵⁵. Using the same approach but also excluding those with cancer-related symptoms and poor performance status, the University of Toronto reported a 5-year post-LT patient survival of 69% (versus 78% with Milan) and HCC recurrence probability of 30% (versus 13% with Milan)⁵⁶. Poorly differentiated tumor grade and vascular invasion were found in the explant in 8% and 40% respectively. A major concern about liberalizing tumor burden and relying on biopsy assessment of tumor histologic grade is that the overall agreement of pre-operative needle core biopsy with explant histopathology was poor according to one study, in which a significantly lower percentage of cases were identified as poorly differentiated tumor grade by biopsy compared to explant (15 versus 28%)⁴⁸. In addition, in a more recent series, the correlation with explant pathology was poor, and had no utility over Milan criteria alone in predicting post-LT recurrence.⁵⁷ At present, given the limitations of histologic assessment noted in current available evidence, no guidelines support routine use of biopsy to help guide eligibility of transplant for those with HCC.

Other Potential Predictors of HCC Recurrence

Other Biomarkers. Aside from AFP, a number of serum biomarkers have been reported to be associated with the risk of HCC recurrence, including des-gamma-carboxyprothrombin (DCP)⁵⁸⁻⁶⁰, AFP-L3%, and absolute AFP-L3⁵⁸. Some studies have also suggested that inflammatory biomarkers, including neutrophil/lymphocyte ratio (NLR) and platelet/lymphocyte ratio, can predict tumor recurrence after LT^{61,62}. Nevertheless, other studies did not confirm the association between these inflammatory biomarkers and outcome after LT for HCC mostly within Milan criteria^{63,64}. The role of these biomarkers in clinical practice, especially with respect to candidate selection for LT, has not yet been clearly defined.

Genomic biomarkers have the potential to discriminate HCC recurrence risk. In one study, allelic imbalance in chromosome 9/18 microsatellites was found to strongly correlate with recurrence in HCC beyond Milan Criteria⁶⁵. In another study on patients with tumor beyond Milan criteria, the presence of progenitor cell markers (either CK19 or S2 signatures) had significantly greater HCC recurrence rate and lower survival than those without these gene signatures⁶⁶.

Positron emission tomography scan. Several studies have suggested that ¹⁸F-fluorodeoxyglucose positron emission tomography/CT (¹⁸F-FDG PET/CT) was a prognostic marker in LT for HCC, showing good correlation between ¹⁸F-FDG avid tumors and unfavorable histo-pathological tumor characteristics including microvascular invasion and poorly differentiated tumor grade^{50,67,68}. Non-uptake of tumor on ¹⁸F-FDG PET/CT may allow the use of very liberal tumor size criteria in living donor LT according to studies from Korea and Japan^{69,70}.

Hepatitis C virus infection. Hepatitis C virus (HCV)-associated cirrhosis is a well-established risk factor for HCC, and antiviral treatment with viral eradication is associated with a significant decline in HCC risk overall. Recent reports however have raised concern that antiviral therapy with directly-acting antivirals may lead to an unexpected increased risk of HCC recurrence after potentially curative therapy, including resection and ablation.⁷¹⁻⁷³ This concern has not been supported by subsequent studies and pooled or meta-analyses.⁷⁴⁻⁷⁷ In addition, while data on post-transplant recurrence specifically are limited, DAA therapy has not been associated with

increased post-LT recurrence risk.⁷⁸ Thus antiviral treatment decisions should be made in the context of organ access and clinical characteristics of the patient⁷⁹, but should not be avoided specifically to prevent HCC recurrence.

Donor characteristics. Several donor characteristics, including older donor age^{80,81} and non-local livers⁸¹, have been implicated as risk factors for HCC recurrence following LT. These findings require confirmation. The effects of graft type (living donor versus deceased donor) on HCC recurrence after LT have yielded conflicting results. When comparing living donor LT and deceased donor LT in the post-MELD prioritization era, a meta-analysis showed no significant difference in HCC recurrence rates between these grafts⁸². However, it should be emphasized that having a potential live donor at listing compared to not has been associated with increased survival for patients with HCC, likely related to a lower dropout rate and a shorter waiting period for LT.⁸³ In one study, those receiving allografts donated after cardiac death did not experience a higher rate of recurrence compared to donation after brain death organs⁸⁴.

Prognostic Models

Proposed models based on variables that are available before LT generally follow the principle of combining tumor morphologic parameters (diameter, number, or volume) with measures or surrogates of tumor “biology” metrics (Table 1). They include AFP and other biomarkers (DCP, NLR). Pre-LT biopsy of the tumors has also been used to determine histologic grade of differentiation and to exclude poorly differentiated tumors for LT. While these prognostic models were intended to improve selection of candidates with good post-LT outcomes, the patient populations were highly heterogeneous and most of these prognostic models were based on retrospective data collection. Consequently, these models still require vigorous testing and prospective validation using well-defined criteria for the upper limits in tumor burden and specific thresholds for these biomarkers.

The post-LT prognostic models are based on explant tumor histopathologic characteristics, and they most commonly include measures of tumor burden, tumor grade of differentiation, and the presence or absence of vascular invasion (Table 2).

The majority of these prognostic models also include AFP and/or other biomarkers. The post-LT prognostic models help identify candidates at high risk for HCC recurrence, and provide a reference for the expected incidence of HCC recurrence. These models are also helpful in the development of standardized post-LT surveillance strategies, and the identification of the appropriate subgroups at sufficiently high risk for HCC recurrence to be considered for clinical trials using neoadjuvant therapy to reduce the risk of HCC recurrence.

Knowledge Gaps

- AFP is currently the most widely utilized serum biomarker but there is no consensus on standardized cut off values and/or dynamic changes in AFP levels to guide clinical decision making.
- Standardization of the definition of MVI is needed for both reporting and research purposes.
- Most other biomarkers remain investigational and have not yet been adopted in routine practice, and better biomarkers are still needed.
- Further exploration of the impact of specific liver-directed therapies and the correlation with post-LT risk of HCC recurrence are needed.
- Many pre- and post-LT prognostic models have been proposed, but still require prospective independent validation. An ideal pre-transplant model would lead to improved post-LT outcomes by incorporating surrogates of tumor “biology” metrics with conventional morphologic parameters. Post-LT models would be useful in developing standardized post-LT surveillance strategies, and identification of subgroups to be considered for future neoadjuvant therapy to reduce HCC recurrence.

Post-Transplant HCC Surveillance

Surveillance may improve survival through access to earlier and perhaps curative treatment of recurrence⁸⁵⁻⁸⁸. To be most effective, post-LT HCC surveillance should be done in the window of time that covers most recurrences, includes imaging of locations where recurrences are known to occur and with modalities that have good sensitivity

and specificity. Unfortunately, while studies have looked at timing, risk factors, and characteristics of HCC recurrence, there remains a lack of data to guide serum and imaging tests and their frequency. Specifically, there are no trials to date studying surveillance protocols and their impact on post-LT outcomes.

HCC Recurrence Patterns

While the timing of post-transplant tumor recurrence is variable, peak HCC recurrence occurs around 2-3 years after transplant^{84,89} (Table 3). Early HCC recurrence defined as within the first year of LT portends the worst prognosis.^{82, 83, 87} This could occur due to non-detectable extrahepatic metastases that may be present before LT and as a consequence of circulating HCC clones engrafting and growing in a target organ after LT. The plausible explanation for late HCC recurrence (within 2-5 years of LT) could be a second unknown hit that may lead to late engrafting of HCC cells that are less in number and remained latent for a long time during the post-LT period. HCC recurrence after >5 years of LT, though infrequent, has been described⁸⁰. These data suggest that while surveillance should be most intense during the first 2 years, it should be maintained until at least 5 years post-LT for high risk patients as longer time to recurrence is a predictor for improved outcomes after treatment of recurrent HCC^{85,90-92}.

The pattern of HCC recurrence extends from hepatic to extrahepatic and from single site recurrence to multiple site recurrence^{80,85,87,88,93-95} (Table 3). The most common extrahepatic sites are lungs and bones. However, HCC recurrence can also occur in adrenal glands, soft tissue (e.g. local recurrence of biopsy tracks), peritoneum, and brain.

Recommendations for Post-LT Surveillance

There are currently no clinical trial data to guide a recommendation for a specific surveillance protocol though expert groups have published consensus statements based upon the data available.¹² With recent advances in systemic treatment, the implementation of surveillance programs may result in earlier detection that may improve survival in patients with recurrence who are able to be treated with surgical,

local, or systemic treatment. Therefore, all liver transplant recipients with HCC should be enrolled in a surveillance program.

As there are no specific evidence-based risk stratification criteria that can be formally recommended, there is significant variation in surveillance practices between programs. In general, we recommend cross sectional imaging with either multiphase CT or multiphase MRI should be performed at least every 6 months for at least 2-3 years post-transplant as the majority of patients recur in this time. Many programs continue surveillance until 5 years post-transplant based upon the low likelihood of recurrence beyond this point, however data on the best or most cost-effective length of surveillance are also not available to guide recommendations. Given the small amount of data for AFP in patients post-transplant, and the relatively low cost of this test, AFP level should be drawn every 6 months for five years^{96,97}. There are not enough data to validate use of other biomarkers such as AFP-L3% or DCP in a routine surveillance program after LT for HCC.

Imaging for lung metastases with non-contrast CT chest should also be performed every 6 months in this period as well. Bone metastases may also be a site of recurrence, but given the low sensitivity of bone scan we do not recommend routine imaging with bone scan. Patients may present either with an elevation of alkaline phosphatase or with pain at the site of bone metastasis, and in case of these signs or symptoms further imaging may be performed at that time.

Risk-based screening has been proposed by several groups, including by Mehta and colleagues recommending screening based on RETREAT score (no screening if RETREAT score = 0 due to estimated recurrence risk < 3%, every 6 months for 2 years if score = 1-3, up to 5 years for those if score = 4, and every 3 to 4 months for 2 years followed by every 6 months until year 5 if score \geq 5)^{98,99}. While the RETREAT criteria could be considered for clinical use, these data must be prospectively validated and there are no published prospective studies of specific of risk stratification-based surveillance strategies. Ultimately risk-based screening would be desirable if it were shown to be safe and perhaps most cost-effective.

Knowledge Gaps

- Prospective validation of imaging every 6 months as a surveillance strategy is needed in the liver transplant recipient population. In addition, data regarding the relative benefits of different lengths of surveillance are needed.
- Risk-stratified surveillance protocols must also be rigorously studied, perhaps including whether there are patients at sufficiently low risk that there is no benefit to surveillance.
- Whether post-LT surveillance improves HCC treatment remains to be proven

Strategies for Prevention of HCC Recurrence

Immunosuppression

The immunosuppression regimen of the patient is a possible factor in preventing HCC recurrence, as the immune system is a major defense against cancer, either by attacking dysplastic cells themselves or by controlling viruses linked to cancer ¹⁰⁰. Additionally, as the immunosuppression regimen can change over time, so can the significance of HCC recurrence post-LT, as early recurrence may have a different biological behavior and identity compared to late HCC recurrence ¹⁰¹.

Table 4 lists the different types of immunosuppression medications and the possible manner in which they can affect HCC recurrence post-LT, including the potential mechanisms involved. Though medication profiles may be strategically used to address HCC recurrence, in practice multiple factors are weighted in formulating the optimal immunosuppression regimen for a specific patient such as consideration of side effects and comorbidities including kidney dysfunction.

A class of immunosuppression medications that appears to play a central role in most efforts to control HCC recurrence are the mammalian target of rapamycin (mTOR) inhibitors. Reducing calcineurin inhibitors (CNIs) and adding a mTOR inhibitor may reduce HCC recurrence, given antiproliferative properties against HCC ¹⁰². Specifically, mTOR involves two signaling pathways: mTOR complex1, which is responsible for cell proliferation based on regulatory signals from the immune system, and mTOR complex 2, which affects cellular metabolism ¹⁰³. The two main existing mTOR inhibitors are sirolimus (affecting complex 1 and 2) and everolimus (selective for complex 1).

Sirolimus has demonstrated antiangiogenic activities linked to a decrease in production of vascular endothelial growth factor (VEGF) and inhibited response of vascular endothelial cells to stimulation by VEGF^{104,105}. Data from small single center retrospective studies, as well as two meta-analyses, suggest that compared to CNIs, use of sirolimus reduced the risk of HCC recurrence after LT¹⁰⁶⁻¹⁰⁹^{106,110-114}. Efficacy of this strategy may vary depending on the stage of the initial disease or the presence of hepatitis C virus¹¹⁵. The SILVER study is a multicenter randomized controlled trial seeking to obtain definitive answers.¹¹⁶ In this study, though there appeared to be an advantage in the sirolimus group regarding improvement in recurrence-free survival in the first 3-5 years; this benefit is subsequently lost with further follow-up. Everolimus is approved for immunosuppression in LT recipients with low dose CNIs and likely has similar effects as sirolimus. However, available evidence is from small, single center studies that lack adequate power to make clear recommendations.¹¹⁷ Factors that result in reluctance to use mTOR inhibitors immediately post-LT include concern for increased risk of hepatic artery thrombosis and impaired wound healing^{118,119}. Therefore, the main impediment to widespread use of mTOR inhibitors for HCC recurrence prevention has been the lack of high-quality studies that demonstrate a benefit. Though mTOR inhibitors have shown some promise to reduce or prevent HCC recurrence in smaller studies, the highest quality data through the SILVER study does not show a benefit. At this time, there is a need for further well-designed multicenter studies to better delineate their use in the post-LT population.

There are certain combination strategies of immunosuppression medications with a greater promise of success at minimizing risk of HCC recurrence. Medications such as steroids and CNIs possibly contribute to HCC recurrence, whereas others such as the mTOR inhibitors, and, to a lesser extent MMF, may have a beneficial effect. Though the optimal combination has yet to be found, an overall goal of reducing CNIs with possible eventual complete withdrawal and avoidance of azathioprine and anti-thymocyte globulin is consistent with a strategy to minimize risk of HCC recurrence. As a counterbalance, the addition of mTOR inhibitors with or without MMF, can help maintain a safe immunosuppression level to avoid rejection, while at the same time not endangering renal function and possibly preventing HCC recurrence. In the context of

acute cellular rejection (ACR), the benefit of increased immunosuppression to attenuate alloreactivity must be balanced against the risk of promoting oncogenesis. At present, no randomized controlled trials powered to detect differences in de novo tumor formation or HCC recurrence between ACR treatment strategies are available.

Though we seek an optimal medication strategy, the existence of multiple medication-, patient- and disease-related variables make this challenging. Furthermore, a generalizable medication strategy may not be optimal. Rather, the biological behavior of HCC may prove to be better targeted by a patient-targeted approach, or otherwise, a “personalized” strategy towards immunosuppression. The identification of non-invasive “biomarkers of tolerance” may help identify those patients where “safe” immunosuppression withdrawal may reduce the risk of HCC recurrence.

Adjuvant systemic therapy

There is no evidence to support the use of adjuvant systemic chemotherapy to prevent post-LT HCC recurrence and this approach is not currently recommended. There is no evidence that adjuvant systemic therapy using doxorubicin provides benefit in preventing HCC recurrence after LT ¹²⁰. In an RCT of 58 patients, oxaliplatin based adjuvant chemotherapy for patients with HCC beyond Milan criteria showed improved 1- and 3- year overall survival in the adjuvant oxaliplatin group. However, there was no difference in the 3-year disease-free survival in both the groups ¹²¹.

Licartin is a promising antineoplastic agent for adjuvant chemotherapy for patients with advanced HCC¹²². It is an immune-radioconjugate containing metuximab, an antibody fragment targeting the HCC-associated antigen HAb18G/CD147, a member of CD147 family that is overexpressed in HCC and fibroblasts. Its expression is associated with cancer cell progression and increased adhesion, invasion, and metastasis. Metuximab is conjugated to the radioisotope Iodine I¹²². In a randomized controlled trial of 66 HCC patients beyond Milan Criteria the Licartin group received three doses every 28 days three weeks after LT and had significantly lower HCC recurrence rate compared to the control group (27% vs 57%; p=0.017)¹²³.

The role of pre-emptive adjuvant therapy with Sorafenib was initially considered promising but newer data demonstrates there is likely minimal added benefit. Most of

these studies are limited to a small number of subjects, single center studies, and case series¹²⁴⁻¹²⁶. One of the largest retrospective studies examined the role of adjuvant Sorafenib (25 patients) versus standard of care (20 patients) in patients with advanced HCC on explant pathology. The authors concluded that the use of pre-emptive Sorafenib did not improve recurrence free survival or overall survival¹²⁷. Similarly, data on the use of neoadjuvant sorafenib among patients with potentially curative resection or ablation therapy did not demonstrate a benefit in terms of recurrence.¹²⁸

Immunotherapies have not been studied for prevention of post-transplant recurrence. Currently, a clinical trial examining the effects of nivolumab in patients with high risk of recurrence after curative resection or ablation is underway but does not include LT recipients.¹²⁹

Knowledge Gaps

- The optimal combination of immunosuppressive regimens to decrease HCC recurrence is currently unknown.
- Through their antiproliferative properties, mTOR inhibitors are believed to have the most promising role in preventing recurrence although data are limited and not widely replicated.
- Data on the use of adjuvant systemic chemotherapy in preventing HCC recurrence are needed.
- With the changing landscape of HCC and approval of new systemic chemotherapeutic agents, future studies are warranted to characterize the efficacy and safety of these agents as adjuvant therapy in LT recipients.

Management of HCC Recurrence

HCC recurrences are usually asymptomatic and diagnosed as a result of surveillance imaging. Management is mainly focused at treatment of biopsy-proven HCC recurrences once they occur given strategies for prevention are limited.

Surgical Resection

Surgical resection for recurrent HCC has been shown to be an independent predictor of long-term survival if isolated to a single organ^{85,87,130-132}. In the largest single center experience, of 106 total recurrences, 25 underwent surgical resection with overall better survival with this approach. In this series, a multivariable risk model was also constructed to predict post-recurrence survival.¹³² In a retrospective analysis, Valdivieso and colleagues identified 11/23 patients with HCC recurrence after LT who underwent surgical resection due to intrahepatic and extrahepatic disease; of these, 8 patients obtained a R-0 resection¹¹. Second recurrence was seen in 50% of the R-0 cases and all 3 of the patients with R-1 resection. Despite this, overall recurrence-free survival was significantly higher in the R-0 cohort (32.3 months vs 6.9 months; $p=0.006$) with a five-year survival rate of 27% (vs 0% at 3 years in the other cohort). Similarly, Regalia and colleagues performed surgical resection of 7/21 patients with post-LT HCC recurrence with both intra- and extrahepatic disease¹³¹. Patients who underwent surgical resection had a 57% survival rate at 4 years vs 14% survival rate in non-surgical candidates ($p<0.02$). Finally, in a recent French study, 22/70 patients with recurrent HCC after LT who underwent surgical resection (2 intrahepatic and 20 extrahepatic) had a median survival of 35 vs 15 months for non-resected patients ($p<0.001$)⁸⁷. In carefully selected patients with good functional status and limited disease, resection of isolated extrahepatic recurrent disease in regional lymph nodes, adrenals, and lungs have shown favorable survival¹³³⁻¹³⁵.

Locoregional Therapies

Data on liver directed therapy for the treatment of recurrent HCC after LT are lacking and limited to small case series. Zhai and colleagues examined the efficacy of microwave ablation (MWA) in the treatment of intrahepatic recurrence of HCC in 11 LT recipients¹³⁶. Patients tolerated the procedure well with only 3 cases having tumor progression within 1-7 months after MWA. Survival rates at 1 and 2 years were 51.5% and 15.3% respectively with an average survival time of 17.3 months. Ko and colleagues evaluated the efficacy of transcatheter arterial chemoembolization (TACE) in a cohort of 28 patients with recurrent HCC after living donor LT¹³⁷. Although tumor size

was reduced by $\geq 25\%$ in 19/28 patients, intra- or extrahepatic recurrence occurred in 93% of patients during the six-month follow-up period after the intervention. The 1-, 3- and 5- year survival rates were 48%, 6%, and 0%, respectively. A single case report described the use of Yttrium-90 microspheres (Y-90) in a LT recipient with intrahepatic HCC recurrence 22 months after transplantation who failed both initial resection and adjuvant chemotherapy¹³⁸. Follow-up imaging 2 months post-treatment showed tumor necrosis and excellent treatment response but no long-term data were reported.

Sorafenib and other RAF kinase inhibitors

The use of systemic chemotherapy in post-LT HCC recurrence is limited to small studies exploring the use of sorafenib in this setting. Treatment with sorafenib shows a median survival of 12 months (range 1.45-20.1)^{104,139-143}. In approximately 80% of these cases, an mTOR inhibitor was combined for immunosuppression. In a recent meta-analysis by Mancuso and colleagues, patients who received sorafenib for recurrent HCC after LT had an overall 1-year survival ranging from 18-90% with a pooled estimate of 1 year survival at 63%¹⁴⁴. In a recent analysis of the US cohort of the GIDEON registry, the safety and tolerability of sorafenib was studied in patients with HCC recurrence who had undergone LT or resection¹⁴⁵. Most adverse events occurred in the first 4 weeks of treatment and the incidence of adverse events resulting in discontinuation of drug were similar in both groups. There is no evidence of increased toxicity from sorafenib in the post-transplant setting compared to patients with primary HCC; however, the use of concomitant mTOR inhibitors can increase the rate of dose reduction or discontinuation due to severe side effects. There is also one multicenter retrospective series demonstrating the safety of the use of regorafenib in LT recipients who failed to response to sorafenib.¹⁴⁶

Data on the use of other systemic chemotherapy agents in this setting are also limited. The use of metronomic capecitabine in 38 patients with post LT HCC recurrence showed a median survival of 22 months compared to 7 months ($p < 0.01$) in patients who received best supportive care with an acceptable safety profile compared to sorafenib.¹⁴⁷

Immunotherapy

Despite the promise of checkpoint inhibitors in the treatment of several malignancies with excellent outcomes, the use of immunotherapy in transplant recipients is challenging due to the potentially increased risk of allograft rejection and graft loss.¹⁴⁸ Immune checkpoint inhibitors target cell death protein 1/programmed death ligand 1 (PD-1/PDL-1) or cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4), all of which are important negative regulators of T-cell immune function.¹⁴⁹ Immune checkpoint inhibitors work by restarting an effective anti-tumor immune response, allowing immune system destruction of tumor cells. However, through this same mechanism, the immune system can also lose the ability to recognize self from non-self and thereby cause immune system mediated adverse events similar to autoimmune diseases. Of importance in solid organ transplant recipients, the PD-1:PDL pathway plays an important role in regulating alloimmunity and transplant tolerance.¹⁵⁰ Therefore, transplant recipients are at particular risk for allograft rejection when immune checkpoint inhibitors are used as cancer therapies. Theoretically, usage of such immunomodulatory agents in transplant recipients may promote the development of allograft injury, severe rejection, and even death.¹⁵¹

Nevertheless, HCC represents a promising target for anticancer immunotherapies such as immune checkpoint inhibitors targeting CTLA-4 and PD-1/PDL-1.¹⁵¹ Recently, two PD-1 checkpoint inhibitors, nivolumab and pembrolizumab, have been approved for advanced HCC as second line therapies. Both medications received conditional FDA approval based on limited Phase II data, and unfortunately, recent phase III data showed pembrolizumab failed to improve progression-free or overall survival in patients with advanced HCC.

There have been multiple case reports in the literature of the use of CTLA-4 antibodies and PD-1/PDL-1 antibodies in patients after solid organ transplant. These reports have been in liver, kidney, heart, and corneal transplants, for several tumor types including HCC, melanoma, and non-small cell lung cancer.¹⁵¹⁻¹⁵⁷ There has only been one small case series from the Mayo Clinic on their single center experience of PD-1 inhibitor therapy in 7 liver transplant recipients, 6 with HCC and one with metastatic melanoma.¹⁵⁵ Overall, the reported graft rejection rates in the literature have

ranged from 25-54%, with median time to graft rejection 8-19 days after initiation of therapy.^{153,157} Several reports have detailed successful treatment of graft rejection with steroids, mycophenolate mofetil, and anti-thymocyte globulin, but there have also been several reports of rapid graft failure and death from severe allograft rejection. While there are reports of tumor response in transplant recipients with immune checkpoint inhibitors, it seems that the response rates are much lower than that seen in non-transplant patients.¹⁵³

Overall, there appears to be no clear association with time from transplant to treatment or particular immune suppression strategies to predict or mitigate risk of rejection. Interestingly, there are emerging data that staining of liver allograft for PD-1 lymphocyte expression may be predictive, where lack of PD-1 staining lymphocytes in the allograft may be associated with a lower risk of rejection¹⁵⁵. However, until there are more robust clinical trials that allow prediction and clarification of the risk of graft rejection and loss in transplant recipients, the immune checkpoint inhibitors must be used with extreme caution in these patients.

Knowledge Gaps

- Despite availability of various treatment strategies, including the use of chemotherapeutic regimens, surgical resection, and liver directed therapy, long term recurrence free survival is currently rarely achieved and is an important area of further research.
- Data are limited regarding the use of immune checkpoint inhibitors in this population, particularly given the potential increased risk of rejection. Given the lack of robust data or guidelines, further studies should employ a multidisciplinary and multimodal approach tailored to each patient.

Conclusions:

As the incidence of HCC continues to rise, patients with HCC will be an increasingly large proportion of our transplant population. While we have learned a significant amount about how to cure patients of HCC with transplant and risk stratify for

recurrence, there remain many opportunities for improvement in our post-transplant management. We therefore have provided a roadmap of knowledge gaps aimed at guiding the development of studies that will significantly advance post-transplant care of these patients, including in the areas of post-transplant surveillance, prophylactic measures including alteration of the standard immunosuppression regimen, and treatment of established HCC recurrence.

Disclosure

The authors of this manuscript have conflicts to disclose as described by the American Journal of Transplantation. Dr. Francis Yao receives research support from Fuji film Wako Diagnostics; Anjana Pillai is on the Speaker's Bureau for Simply Speaking Hepatitis, Eisai Inc, BTG and Medical Advisory Board for Wako Diagnostics and Exelixis; Catherine Frenette is on the speakers bureau for Bayer, BMS, Eisai and Exelixis and is on the advisory board for Eisai, Wako and Exelixis and has reserach support from Bayer.

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

Table 1. Summary of Proposed Pre-Transplant Selection Models (selected)*

Pre-LT Selection Model	Tumor Burden	Biomarker(s)	Additional Criteria	5-year Post-LT Overall Survival	AUROC
US National Policy¹⁵⁸	Milan or down-staged to Milan	AFP >1000 ng/ml reduced to <500		80%	
French AFP Model³⁴	Size and number (lowest risk: largest tumor \leq 3 cm and \leq 3 tumors)	AFP (lowest risk: \leq 100 ng/ml)		68% if risk score \leq 2 vs. 47% if risk score >2	0.7
Metro-Ticket 2.0³⁶	Tumor number + size of largest	AFP			0.72

	tumor				
TTV-AFP Model ¹⁵⁹	Total tumor volume $\leq 115 \text{ cm}^3$	AFP ≤ 400 ng/ml		75% (at 4 years) for those >Milan but within TTV-AFP	0.8
Extended Toronto Criteria (ETC) ⁵⁶	No upper limits		1) Biopsy of largest tumor: poorly differentiated tumors excluded 2) No cancer-related symptoms	68% for those >Milan but within ETC	
Pre-MORAL ⁶¹	Largest tumor size (lowest risk: ≤ 3 cm)	AFP (lowest risk: < 200 ng/ml)	NLR (lower risk < 5)	5-yr RFS: 99% low risk 70% medium-risk 56% high-risk	0.82
HALT-HCC ⁶⁴	Hypotenuse between tumor number & largest tumor size**	InAFP	MELD-Na		0.61
NYCA Score ⁴²	Tumor number + size of largest tumor	AFP response to treatment		Low risk 75%, acceptable risk 62%, high risk 40%	0.73

*Proposed criteria based on living donor liver transplant are not included

**By Pythagorean Theorem ($A^2 + B^2 = C^2$); e.g. a patient with 3 lesions with largest 4 cm would receive tumor burden score of 5

Table 2. Summary of Proposed Post-Transplant Prognostic Models (selected)*

Post-LT Prognostic Model	Tumor Burden	Biomarker(s)	Other Histologic features	5-year Recurrence risk/ Survival	AUROC
RETREAT⁹⁸	Sum of largest viable tumor diameter and number of viable tumor	AFP	Vascular invasion	5-yr recurrence risk 2.9% for RETREAT score 0; 75% for RETREAT score ≥ 5	0.75
US HCC Consortium¹⁶⁰	Tumor diameter Tumor number	AFP NLR	Vascular invasion Tumor differentiation	Based on Normogram	0.76
Post-MORAL⁶¹	Tumor diameter Tumor number		Vascular invasion Tumor differentiation	5-yr recurrence-free survival ranging from 97% in low risk group to 22% in very high risk group	0.87
Decaens et al⁹²	Tumor diameter Tumor number		Vascular invasion Tumor differentiation	5-yr recurrence risk 14.5% for risk score < 4 and 51.5% for risk score ≥ 5	

*Proposed criteria based on living donor liver transplant are not included

Table 3. HCC recurrence time and location

Study and time period	Patients with Recurrence	Time to recurrence (months)	Hepatic Recurrence	Extrahepatic Recurrence	Multi-site Recurrence
Roayaie et al 1998-2002	57/311 (18%)	12.3 (1.5-60.3)	9 (16%)	30 (53%)	18 (32%)
Cescon et al 1997-2009	34/283 (12%)	12 (1-118)	3 (9%)	7 (21%) lung=3 bone=2 peritoneum=2	24 (71%)
Escartin et al 1988-2005	28/184 (15%)	Early<12 months=5.7, late≥12 months=33.5	7 (25%)	21 (75%) lung=7 bone=5, adrenal=2 peritoneum=2 skin=2 lymph node=2 CNS=1	11 (39%)
Valdivieso et al 1996-2008	23/182 (9%)	23.4	2 (9%)	16 (70%) lung=9 bone=3 adrenal=2 lymph node=2	5 (22%)
Kornberg et al 1994-2007	16/60 (27%)	23 (4-58)	4 (25%)	12 (75%) lung=5 bone=4 adrenal=1 peritoneum=1 CNS=1	

Sharma et al 2002-2008	17/94 (18%)	25.2	6 (4%)	6 (4%) lung=1 adrenal=3 abdominal soft tissue=2	11 (65%)
Mehta et al 2002-2012	84/721 (12%)	13	22 (26%)	84 (100%) lung=37, bone=25, peritoneum=22	21 (25%)
Fernandez- Sevilla et al 1991-2013	70/493 (14.2%)	17	2 (2.8%)	51 (72.9%)	17 (24.3%)
Sapisochin et al 2000-2012	121/780 (15.5%)	14	16 (13.2%)	63 (52.1%)	42 (34.7%)

Table 4. Immunosuppression and HCC recurrence

Type of Immunosuppression Medication	Role in HCC recurrence after LT	Proposed Mechanism
Steroids ¹⁶¹⁻¹⁶⁴	Contributing possibly	Reduce strength of immune inflammatory response, inhibit apoptosis and promote migration of malignant cells
Calcineurin inhibitors (CNIs) ¹⁶⁵⁻¹⁶⁸	Contributing possibly	High levels of CNIs oncologically detrimental; decreased HCC recurrence with lower trough levels
Antimetabolites ¹⁶⁹⁻¹⁷¹	Azathioprine possibly contributing; Mycophenolate mofetil (MMF) role unclear	Role of azathioprine as carcinogen; Although MMF has an anti-angiogenesis and anti-proliferative action it has not been shown to decrease HCC recurrence; however it is used in order to decrease CNI levels
Induction agents ^{139,172}	Antithymocyte globulin	Helpful in reducing CNI levels, but because of

	(ATG) and Basiliximab role unclear by themselves	their disappearing effect on CD25+ cells, including tumor specific cytotoxic T cells, their overall effect is not clear
mTOR inhibitors (sirolimus and everolimus) ^{106,110-114,116}	Possibly decreasing HCC recurrence, although data not clear	Anticancer properties through mTOR inhibition, although when treatment is initiated may be critical regarding effectiveness; everolimus may stabilize HCC progression

References:

1. Yang JD, Larson JJ, Watt KD, et al. Hepatocellular Carcinoma Is the Most Common Indication for Liver Transplantation and Placement on the Waitlist in the United States. *Clinical gastroenterology and hepatology : the official clinical practice journal of the American Gastroenterological Association*. 2017;15(5):767-775.e763.
2. Iwatsuki S, Starzl TE, Sheahan DG, et al. Hepatic resection versus transplantation for hepatocellular carcinoma. *Annals of surgery*. 1991;214(3):221-228; discussion 228-229.
3. Mazzaferro V, Regalia E, Doci R, et al. Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med*. 1996;334(11):693-699.

4. Figueras J, Jaurrieta E, Valls C, et al. Survival after liver transplantation in cirrhotic patients with and without hepatocellular carcinoma: a comparative study. *Hepatology (Baltimore, Md)*. 1997;25(6):1485-1489.
5. Min AD, Saxena R, Thung SN, et al. Outcome of hepatitis C patients with and without hepatocellular carcinoma undergoing liver transplant. *The American journal of gastroenterology*. 1998;93(11):2148-2153.
6. Molmenti EP, Klintmalm GB. Liver transplantation in association with hepatocellular carcinoma: an update of the International Tumor Registry. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2002;8(9):736-748.
7. Yoo HY, Patt CH, Geschwind JF, Thuluvath PJ. The outcome of liver transplantation in patients with hepatocellular carcinoma in the United States between 1988 and 2001: 5-year survival has improved significantly with time. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2003;21(23):4329-4335.
8. Wong SN, Reddy KR, Keeffe EB, et al. Comparison of clinical outcomes in chronic hepatitis B liver transplant candidates with and without hepatocellular carcinoma. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2007;13(3):334-342.
9. Escartin A, Sapisochin G, Bilbao I, et al. Recurrence of hepatocellular carcinoma after liver transplantation. *Transplantation proceedings*. 2007;39(7):2308-2310.
10. Plessier A, Codes L, Consigny Y, et al. Underestimation of the influence of satellite nodules as a risk factor for post-transplantation recurrence in patients with small hepatocellular carcinoma. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2004;10(2 Suppl 1):S86-90.
11. Valdivieso A, Bustamante J, Gastaca M, et al. Management of hepatocellular carcinoma recurrence after liver transplantation. *Transplantation proceedings*. 2010;42(2):660-662.
12. de'Angelis N, Landi F, Carra MC, Azoulay D. Managements of recurrent hepatocellular carcinoma after liver transplantation: A systematic review. *World journal of gastroenterology*. 2015;21(39):11185-11198.
13. Yao FY, Ferrell L, Bass NM, Bacchetti P, Ascher NL, Roberts JP. Liver transplantation for hepatocellular carcinoma: comparison of the proposed UCSF criteria with the Milan criteria and the Pittsburgh modified TNM criteria. *Liver Transpl*. 2002;8(9):765-774.

14. Yao FY, Ferrell L, Bass NM, et al. Liver transplantation for hepatocellular carcinoma: expansion of the tumor size limits does not adversely impact survival. *Hepatology*. 2001;33(6):1394-1403.
15. Mazzaferro V, Llovet JM, Miceli R, et al. Predicting survival after liver transplantation in patients with hepatocellular carcinoma beyond the Milan criteria: a retrospective, exploratory analysis. *The Lancet Oncology*. 2009;10(1):35-43.
16. Volk ML, Vijan S, Marrero JA. A novel model measuring the harm of transplanting hepatocellular carcinoma exceeding Milan criteria. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2008;8(4):839-846.
17. Otto G, Herber S, Heise M, et al. Response to transarterial chemoembolization as a biological selection criterion for liver transplantation in hepatocellular carcinoma. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2006;12(8):1260-1267.
18. Millonig G, Graziadei IW, Freund MC, et al. Response to preoperative chemoembolization correlates with outcome after liver transplantation in patients with hepatocellular carcinoma. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2007;13(2):272-279.
19. Lai Q, Avolio AW, Graziadei I, et al. Alpha-fetoprotein and modified response evaluation criteria in solid tumors progression after locoregional therapy as predictors of hepatocellular cancer recurrence and death after transplantation. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2013;19(10):1108-1118.
20. Yao FY, Mehta N, Flemming J, et al. Downstaging of hepatocellular cancer before liver transplant: long-term outcome compared to tumors within Milan criteria. *Hepatology (Baltimore, Md)*. 2015;61(6):1968-1977.
21. Mehta N, Guy J, Frenette CT, et al. Excellent Outcomes of Liver Transplantation Following Down-Staging of Hepatocellular Carcinoma to Within Milan Criteria: A Multicenter Study. *Clinical gastroenterology and hepatology : the official clinical practice journal of the American Gastroenterological Association*. 2018;16(6):955-964.
22. Sinha J, Mehta N, Dodge JL, Poltavskiy E, Roberts J, Yao F. Are There Upper Limits in Tumor Burden for Down-Staging of Hepatocellular Carcinoma to Liver Transplant? Analysis of the All-Comers Protocol. *Hepatology*. 2019.

23. Mehta N, Dodge JL, Grab JD, Yao FY. National Experience on Down-staging of Hepatocellular Carcinoma before Liver Transplant: Influence of Tumor Burden, AFP, and Wait Time. *Hepatology*. 2019.
24. Harper AM, Edwards E, Washburn WK, Heimbach J. An early look at the Organ Procurement and Transplantation Network explant pathology form data. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2016;22(6):757-764.
25. Llovet JM, Schwartz M, Mazzaferro V. Resection and liver transplantation for hepatocellular carcinoma. *Seminars in liver disease*. 2005;25(2):181-200.
26. Schwartz ME. Liver transplantation for hepatocellular carcinoma: the best treatment, but for which patient? *Hepatology (Baltimore, Md)*. 1996;24(6):1539-1541.
27. Shetty K, Timmins K, Brensinger C, et al. Liver transplantation for hepatocellular carcinoma validation of present selection criteria in predicting outcome. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2004;10(7):911-918.
28. Freeman RB, Mithoefer A, Ruthazer R, et al. Optimizing staging for hepatocellular carcinoma before liver transplantation: A retrospective analysis of the UNOS/OPTN database. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2006;12(10):1504-1511.
29. Mehta N, Dodge JL, Roberts JP, Hirose R, Yao FY. Misdiagnosis of hepatocellular carcinoma in patients receiving no local-regional therapy prior to liver transplant: An analysis of the Organ Procurement and Transplantation Network explant pathology form. *Clinical transplantation*. 2017;31(11).
30. Pomfret EA, Washburn K, Wald C, et al. Report of a national conference on liver allocation in patients with hepatocellular carcinoma in the United States. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2010;16(3):262-278.
31. Wald C, Russo MW, Heimbach JK, Hussain HK, Pomfret EA, Bruix J. New OPTN/UNOS policy for liver transplant allocation: standardization of liver imaging, diagnosis, classification, and reporting of hepatocellular carcinoma. *Radiology*. 2013;266(2):376-382.

32. Hakeem AR, Young RS, Marangoni G, Lodge JP, Prasad KR. Systematic review: the prognostic role of alpha-fetoprotein following liver transplantation for hepatocellular carcinoma. *Alimentary pharmacology & therapeutics*. 2012;35(9):987-999.
33. Hameed B, Mehta N, Sapisochin G, Roberts JP, Yao FY. Alpha-fetoprotein level > 1000 ng/mL as an exclusion criterion for liver transplantation in patients with hepatocellular carcinoma meeting the Milan criteria. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2014;20(8):945-951.
34. Duvoux C, Roudot-Thoraval F, Decaens T, et al. Liver transplantation for hepatocellular carcinoma: a model including alpha-fetoprotein improves the performance of Milan criteria. *Gastroenterology*. 2012;143(4):986-994.e983; quiz e914-985.
35. Mehta N, Dodge JL, Roberts JP, Hirose R, Yao FY. Alpha-Fetoprotein Decrease from > 1,000 to < 500 ng/mL in Patients with Hepatocellular Carcinoma Leads to Improved Posttransplant Outcomes. *Hepatology*. 2019;69(3):1193-1205.
36. Mazzaferro V, Sposito C, Zhou J, et al. Metroticket 2.0 Model for Analysis of Competing Risks of Death After Liver Transplantation for Hepatocellular Carcinoma. *Gastroenterology*. 2018;154(1):128-139.
37. Han K, Tzimas GN, Barkun JS, et al. Preoperative alpha-fetoprotein slope is predictive of hepatocellular carcinoma recurrence after liver transplantation. *Canadian journal of gastroenterology = Journal canadien de gastroenterologie*. 2007;21(1):39-45.
38. Giard JM, Mehta N, Dodge JL, Roberts JP, Yao FY. Alpha-Fetoprotein Slope >7.5 ng/mL per Month Predicts Microvascular Invasion and Tumor Recurrence After Liver Transplantation for Hepatocellular Carcinoma. *Transplantation*. 2018;102(5):816-822.
39. Dumitra TC, Dumitra S, Metrakos PP, et al. Pretransplantation alpha-fetoprotein slope and milan criteria: strong predictors of hepatocellular carcinoma recurrence after transplantation. *Transplantation*. 2013;95(1):228-233.
40. Vibert E, Azoulay D, Hoti E, et al. Progression of alphafetoprotein before liver transplantation for hepatocellular carcinoma in cirrhotic patients: a critical factor. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2010;10(1):129-137.
41. Agopian VG, Harlander-Locke MP, Ruiz RM, et al. Impact of Pretransplant Bridging Locoregional Therapy for Patients With Hepatocellular Carcinoma Within Milan Criteria Undergoing Liver

- Transplantation: Analysis of 3601 Patients From the US Multicenter HCC Transplant Consortium. *Ann Surg.* 2017;266(3):525-535.
42. Halazun KJ, Tabrizian P, Najjar M, et al. Is it Time to Abandon the Milan Criteria?: Results of a Bicoastal US Collaboration to Redefine Hepatocellular Carcinoma Liver Transplantation Selection Policies. *Ann Surg.* 2018;268(4):690-699.
 43. Parfitt JR, Marotta P, Alghamdi M, et al. Recurrent hepatocellular carcinoma after transplantation: use of a pathological score on explanted livers to predict recurrence. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society.* 2007;13(4):543-551.
 44. Welling TH, Eddinger K, Carrier K, et al. Multicenter Study of Staging and Therapeutic Predictors of Hepatocellular Carcinoma Recurrence Following Transplantation. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society.* 2018;24(9):1233-1242.
 45. Bhattacharjya S, Bhattacharjya T, Quaglia A, et al. Liver transplantation in cirrhotic patients with small hepatocellular carcinoma: an analysis of pre-operative imaging, explant histology and prognostic histologic indicators. *Digestive surgery.* 2004;21(2):152-159; discussion 159-160.
 46. Jonas S, Bechstein WO, Steinmuller T, et al. Vascular invasion and histopathologic grading determine outcome after liver transplantation for hepatocellular carcinoma in cirrhosis. *Hepatology.* 2001;33(5):1080-1086.
 47. Tsai TJ, Chau GY, Lui WY, et al. Clinical significance of microscopic tumor venous invasion in patients with resectable hepatocellular carcinoma. *Surgery.* 2000;127(6):603-608.
 48. Pawlik TM, Delman KA, Vauthey JN, et al. Tumor size predicts vascular invasion and histologic grade: Implications for selection of surgical treatment for hepatocellular carcinoma. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society.* 2005;11(9):1086-1092.
 49. Gouw AS, Balabaud C, Kusano H, Todo S, Ichida T, Kojiro M. Markers for microvascular invasion in hepatocellular carcinoma: where do we stand? *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society.* 2011;17 Suppl 2:S72-80.
 50. Kornberg A, Freesmeyer M, Barthel E, et al. 18F-FDG-uptake of hepatocellular carcinoma on PET predicts microvascular tumor invasion in liver transplant patients. *American journal of*

- transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2009;9(3):592-600.
51. Banerjee S, Wang DS, Kim HJ, et al. A computed tomography radiogenomic biomarker predicts microvascular invasion and clinical outcomes in hepatocellular carcinoma. *Hepatology (Baltimore, Md)*. 2015;62(3):792-800.
 52. Xu X, Zhang HL, Liu QP, et al. Radiomic analysis of contrast-enhanced CT predicts microvascular invasion and outcome in hepatocellular carcinoma. *J Hepatol*. 2019;70(6):1133-1144.
 53. Klintmalm GB. Liver transplantation for hepatocellular carcinoma: a registry report of the impact of tumor characteristics on outcome. *Annals of surgery*. 1998;228(4):479-490.
 54. Tamura S, Kato T, Berho M, et al. Impact of histological grade of hepatocellular carcinoma on the outcome of liver transplantation. *Archives of surgery (Chicago, Ill : 1960)*. 2001;136(1):25-30; discussion 31.
 55. Cillo U, Vitale A, Grigoletto F, et al. Intention-to-treat analysis of liver transplantation in selected, aggressively treated HCC patients exceeding the Milan criteria. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2007;7(4):972-981.
 56. Sapisochin G, Goldaracena N, Laurence JM, et al. The extended Toronto criteria for liver transplantation in patients with hepatocellular carcinoma: A prospective validation study. *Hepatology (Baltimore, Md)*. 2016;64(6):2077-2088.
 57. Court CM, Harlander-Locke MP, Markovic D, et al. Determination of hepatocellular carcinoma grade by needle biopsy is unreliable for liver transplant candidate selection. *Liver Transpl*. 2017;23(9):1123-1132.
 58. Chaiteerakij R, Zhang X, Addissie BD, et al. Combinations of biomarkers and Milan criteria for predicting hepatocellular carcinoma recurrence after liver transplantation. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2015;21(5):599-606.
 59. Lee JH, Cho Y, Kim HY, et al. Serum Tumor Markers Provide Refined Prognostication in Selecting Liver Transplantation Candidate for Hepatocellular Carcinoma Patients Beyond the Milan Criteria. *Annals of surgery*. 2016;263(5):842-850.
 60. Fujiki M, Takada Y, Ogura Y, et al. Significance of des-gamma-carboxy prothrombin in selection criteria for living donor liver transplantation for hepatocellular carcinoma. *American journal of*

- transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2009;9(10):2362-2371.
61. Halazun KJ, Najjar M, Abdelmessih RM, et al. Recurrence After Liver Transplantation for Hepatocellular Carcinoma: A New MORAL to the Story. *Annals of surgery*. 2017;265(3):557-564.
 62. Lai Q, Castro Santa E, Rico Juri JM, Pinheiro RS, Lerut J. Neutrophil and platelet-to-lymphocyte ratio as new predictors of dropout and recurrence after liver transplantation for hepatocellular cancer. *Transplant international : official journal of the European Society for Organ Transplantation*. 2014;27(1):32-41.
 63. Parisi I, Tsochatzis E, Wijewantha H, et al. Inflammation-based scores do not predict post-transplant recurrence of hepatocellular carcinoma in patients within Milan criteria. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2014;20(11):1327-1335.
 64. Sasaki K, Firl DJ, Hashimoto K, et al. Development and validation of the HALT-HCC score to predict mortality in liver transplant recipients with hepatocellular carcinoma: a retrospective cohort analysis. *The lancet Gastroenterology & hepatology*. 2017;2(8):595-603.
 65. Schwartz M, Dvorchik I, Roayaie S, et al. Liver transplantation for hepatocellular carcinoma: extension of indications based on molecular markers. *Journal of hepatology*. 2008;49(4):581-588.
 66. Miltiadiou O, Sia D, Hoshida Y, et al. Progenitor cell markers predict outcome of patients with hepatocellular carcinoma beyond Milan criteria undergoing liver transplantation. *Journal of hepatology*. 2015;63(6):1368-1377.
 67. Lee SD, Kim SH, Kim YK, et al. (18)F-FDG-PET/CT predicts early tumor recurrence in living donor liver transplantation for hepatocellular carcinoma. *Transplant international : official journal of the European Society for Organ Transplantation*. 2013;26(1):50-60.
 68. Kornberg A, Kupper B, Tannapfel A, et al. Patients with non-[18 F]fludeoxyglucose-avid advanced hepatocellular carcinoma on clinical staging may achieve long-term recurrence-free survival after liver transplantation. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2012;18(1):53-61.
 69. Lee SD, Lee B, Kim SH, et al. Proposal of new expanded selection criteria using total tumor size and (18)F-fluorodeoxyglucose - positron emission tomography/computed tomography for living

- donor liver transplantation in patients with hepatocellular carcinoma: The National Cancer Center Korea criteria. *World journal of transplantation*. 2016;6(2):411-422.
70. Kaido T, Ogawa K, Mori A, et al. Usefulness of the Kyoto criteria as expanded selection criteria for liver transplantation for hepatocellular carcinoma. *Surgery*. 2013;154(5):1053-1060.
71. Reig M, Marino Z, Perello C, et al. Unexpected high rate of early tumor recurrence in patients with HCV-related HCC undergoing interferon-free therapy. *J Hepatol*. 2016;65(4):719-726.
72. Kozbial K, Moser S, Schwarzer R, et al. Unexpected high incidence of hepatocellular carcinoma in cirrhotic patients with sustained virologic response following interferon-free direct-acting antiviral treatment. *J Hepatol*. 2016;65(4):856-858.
73. Conti F, Buonfiglioli F, Scuteri A, et al. Early occurrence and recurrence of hepatocellular carcinoma in HCV-related cirrhosis treated with direct-acting antivirals. *J Hepatol*. 2016;65(4):727-733.
74. Rutledge SM, Zheng H, Li DK, Chung RT. No evidence for higher rates of hepatocellular carcinoma after direct-acting antiviral treatment: a meta-analysis. *Hepatoma Res*. 2019;5.
75. Li DK, Ren Y, Fierer DS, et al. The short-term incidence of hepatocellular carcinoma is not increased after hepatitis C treatment with direct-acting antivirals: An ERCHIVES study. *Hepatology*. 2018;67(6):2244-2253.
76. Waziry R, Hajarizadeh B, Grebely J, et al. Hepatocellular carcinoma risk following direct-acting antiviral HCV therapy: A systematic review, meta-analyses, and meta-regression. *J Hepatol*. 2017;67(6):1204-1212.
77. Grandhe S, Frenette CT. Occurrence and Recurrence of Hepatocellular Carcinoma After Successful Direct-Acting Antiviral Therapy for Patients With Chronic Hepatitis C Virus Infection. *Gastroenterol Hepatol (N Y)*. 2017;13(7):421-425.
78. stanislas.pol@aphp.fr AcsgohcEa. Lack of evidence of an effect of direct-acting antivirals on the recurrence of hepatocellular carcinoma: Data from three ANRS cohorts. *J Hepatol*. 2016;65(4):734-740.
79. Verna EC. The dynamic landscape of liver transplant in the era of effective hepatitis C virus therapy. *Hepatology*. 2017;65(3):763-766.
80. Sharma P, Welch K, Hussain H, et al. Incidence and risk factors of hepatocellular carcinoma recurrence after liver transplantation in the MELD era. *Digestive diseases and sciences*. 2012;57(3):806-812.

81. Vagefi PA, Dodge JL, Yao FY, Roberts JP. Potential role of the donor in hepatocellular carcinoma recurrence after liver transplantation. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2015;21(2):187-194.
82. Liang W, Wu L, Ling X, et al. Living donor liver transplantation versus deceased donor liver transplantation for hepatocellular carcinoma: a meta-analysis. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2012;18(10):1226-1236.
83. Goldaracena N, Gorgen A, Doyle A, et al. Live donor liver transplantation for patients with hepatocellular carcinoma offers increased survival vs. deceased donation. *J Hepatol*. 2019;70(4):666-673.
84. Croome KP, Lee DD, Burns JM, et al. The Use of Donation After Cardiac Death Allografts Does Not Increase Recurrence of Hepatocellular Carcinoma. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2015;15(10):2704-2711.
85. Roayaie S, Schwartz JD, Sung MW, et al. Recurrence of hepatocellular carcinoma after liver transplant: patterns and prognosis. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2004;10(4):534-540.
86. Sapisochin G, Goldaracena N, Astete S, et al. Benefit of Treating Hepatocellular Carcinoma Recurrence after Liver Transplantation and Analysis of Prognostic Factors for Survival in a Large Euro-American Series. *Annals of surgical oncology*. 2015;22(7):2286-2294.
87. Fernandez-Sevilla E, Allard MA, Selten J, et al. Recurrence of hepatocellular carcinoma after liver transplantation: Is there a place for resection? *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2017;23(4):440-447.
88. Roh YN, David Kwon CH, Song S, et al. The prognosis and treatment outcomes of patients with recurrent hepatocellular carcinoma after liver transplantation. *Clinical transplantation*. 2014;28(1):141-148.
89. Azoulay D, Audureau E, Bhangui P, et al. Living or Brain-dead Donor Liver Transplantation for Hepatocellular Carcinoma: A Multicenter, Western, Intent-to-treat Cohort Study. *Annals of surgery*. 2017;266(6):1035-1044.

90. Sapisochin G, Bruix J. Liver transplantation for hepatocellular carcinoma: outcomes and novel surgical approaches. *Nature reviews Gastroenterology & hepatology*. 2017;14(4):203-217.
91. de Ataide EC, Garcia M, Mattosinho TJ, Almeida JR, Escanhoela CA, Boin IF. Predicting survival after liver transplantation using up-to-seven criteria in patients with hepatocellular carcinoma. *Transplantation proceedings*. 2012;44(8):2438-2440.
92. Decaens T, Roudot-Thoraval F, Badran H, et al. Impact of tumour differentiation to select patients before liver transplantation for hepatocellular carcinoma. *Liver international : official journal of the International Association for the Study of the Liver*. 2011;31(6):792-801.
93. Kneuert PJ, Cosgrove DP, Cameron AM, et al. Multidisciplinary management of recurrent hepatocellular carcinoma following liver transplantation. *Journal of gastrointestinal surgery : official journal of the Society for Surgery of the Alimentary Tract*. 2012;16(4):874-881.
94. Costentin CE, Amaddeo G, Decaens T, et al. Prediction of hepatocellular carcinoma recurrence after liver transplantation: Comparison of four explant-based prognostic models. *Liver international : official journal of the International Association for the Study of the Liver*. 2017;37(5):717-726.
95. Clavien PA, Lesurtel M, Bossuyt PM, Gores GJ, Langer B, Perrier A. Recommendations for liver transplantation for hepatocellular carcinoma: an international consensus conference report. *The Lancet Oncology*. 2012;13(1):e11-22.
96. Northen A, Asendorf T, Walson PD, Oellerich M. Diagnostic value of alpha-1-fetoprotein (AFP) as a biomarker for hepatocellular carcinoma recurrence after liver transplantation. *Clinical biochemistry*. 2018;52:20-25.
97. Notarpaolo A, Layese R, Magistri P, et al. Validation of the AFP model as a predictor of HCC recurrence in patients with viral hepatitis-related cirrhosis who had received a liver transplant for HCC. *Journal of hepatology*. 2017;66(3):552-559.
98. Mehta N, Heimbach J, Harnois DM, et al. Validation of a Risk Estimation of Tumor Recurrence After Transplant (RETREAT) Score for Hepatocellular Carcinoma Recurrence After Liver Transplant. *JAMA Oncol*. 2017;3(4):493-500.
99. Mehta N, Dodge JL, Roberts JP, Yao FY. Validation of the prognostic power of the RETREAT score for hepatocellular carcinoma recurrence using the UNOS database. *Am J Transplant*. 2018;18(5):1206-1213.
100. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144(5):646-674.

101. Pecchi A, Besutti G, De Santis M, et al. Post-transplantation hepatocellular carcinoma recurrence: Patterns and relation between vascularity and differentiation degree. *World journal of hepatology*. 2015;7(2):276-284.
102. Guba M, von Breitenbuch P, Steinbauer M, et al. Rapamycin inhibits primary and metastatic tumor growth by antiangiogenesis: involvement of vascular endothelial growth factor. *Nature medicine*. 2002;8(2):128-135.
103. Xu X, Ye L, Araki K, Ahmed R. mTOR, linking metabolism and immunity. *Seminars in immunology*. 2012;24(6):429-435.
104. Wang Y, Speeg KV, Washburn WK, Halff G. Sirolimus plus sorafenib in treating HCC recurrence after liver transplantation: a case report. *World journal of gastroenterology*. 2010;16(43):5518-5522.
105. Zhang J, Li H, Wang GS, Jiang N, Yang Y, Chen GH. [Effects of sirolimus on the growth of transplanted hepatocellular carcinoma]. *Zhonghua gan zang bing za zhi = Zhonghua ganzangbing zazhi = Chinese journal of hepatology*. 2009;17(6):413-416.
106. Menon KV, Hakeem AR, Heaton ND. Meta-analysis: recurrence and survival following the use of sirolimus in liver transplantation for hepatocellular carcinoma. *Alimentary pharmacology & therapeutics*. 2013;37(4):411-419.
107. Zhou J, Wang Z, Wu ZQ, et al. Sirolimus-based immunosuppression therapy in liver transplantation for patients with hepatocellular carcinoma exceeding the Milan criteria. *Transplantation proceedings*. 2008;40(10):3548-3553.
108. Kneteman NM, Oberholzer J, Al Saghier M, et al. Sirolimus-based immunosuppression for liver transplantation in the presence of extended criteria for hepatocellular carcinoma. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2004;10(10):1301-1311.
109. Zimmerman MA, Trotter JF, Wachs M, et al. Sirolimus-based immunosuppression following liver transplantation for hepatocellular carcinoma. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2008;14(5):633-638.
110. Gaumann A, Schlitt HJ, Geissler EK. Immunosuppression and tumor development in organ transplant recipients: the emerging dualistic role of rapamycin. *Transplant international : official journal of the European Society for Organ Transplantation*. 2008;21(3):207-217.

111. Liang W, Wang D, Ling X, et al. Sirolimus-based immunosuppression in liver transplantation for hepatocellular carcinoma: a meta-analysis. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2012;18(1):62-69.
112. Shiah HS, Chen CY, Dai CY, et al. Randomised clinical trial: comparison of two everolimus dosing schedules in patients with advanced hepatocellular carcinoma. *Alimentary pharmacology & therapeutics*. 2013;37(1):62-73.
113. Villanueva A, Chiang DY, Newell P, et al. Pivotal role of mTOR signaling in hepatocellular carcinoma. *Gastroenterology*. 2008;135(6):1972-1983, 1983.e1971-1911.
114. Zhu AX, Abrams TA, Miksad R, et al. Phase 1/2 study of everolimus in advanced hepatocellular carcinoma. *Cancer*. 2011;117(22):5094-5102.
115. Watt KD, Dierkhising R, Heimbach JK, Charlton MR. Impact of sirolimus and tacrolimus on mortality and graft loss in liver transplant recipients with or without hepatitis C virus: an analysis of the Scientific Registry of Transplant Recipients Database. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2012;18(9):1029-1036.
116. Geissler EK, Schnitzbauer AA, Zulke C, et al. Sirolimus Use in Liver Transplant Recipients With Hepatocellular Carcinoma: A Randomized, Multicenter, Open-Label Phase 3 Trial. *Transplantation*. 2016;100(1):116-125.
117. Cholongitas E, Mamou C, Rodriguez-Castro KI, Burra P. Mammalian target of rapamycin inhibitors are associated with lower rates of hepatocellular carcinoma recurrence after liver transplantation: a systematic review. *Transplant international : official journal of the European Society for Organ Transplantation*. 2014;27(10):1039-1049.
118. McAlister VC, Peltekian KM, Malatjalian DA, et al. Orthotopic liver transplantation using low-dose tacrolimus and sirolimus. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2001;7(8):701-708.
119. Ventura-Aguiar P, Campistol JM, Diekmann F. Safety of mTOR inhibitors in adult solid organ transplantation. *Expert opinion on drug safety*. 2016;15(3):303-319.
120. Kim R, Emi M, Tanabe K. Cancer immunoediting from immune surveillance to immune escape. *Immunology*. 2007;121(1):1-14.

121. Zhang Q, Chen H, Li Q, et al. Combination adjuvant chemotherapy with oxaliplatin, 5-fluorouracil and leucovorin after liver transplantation for hepatocellular carcinoma: a preliminary open-label study. *Investigational new drugs*. 2011;29(6):1360-1369.
122. Chen ZN, Mi L, Xu J, et al. Targeting radioimmunotherapy of hepatocellular carcinoma with iodine (131I) metuximab injection: clinical phase I/II trials. *International journal of radiation oncology, biology, physics*. 2006;65(2):435-444.
123. Xu J, Shen ZY, Chen XG, et al. A randomized controlled trial of Licartin for preventing hepatoma recurrence after liver transplantation. *Hepatology (Baltimore, Md)*. 2007;45(2):269-276.
124. Shetty K, Dash C, Laurin J. Use of adjuvant sorafenib in liver transplant recipients with high-risk hepatocellular carcinoma. *Journal of transplantation*. 2014;2014:913634.
125. Teng CL, Hwang WL, Chen YJ, Chang KH, Cheng SB. Sorafenib for hepatocellular carcinoma patients beyond Milan criteria after orthotopic liver transplantation: a case control study. *World journal of surgical oncology*. 2012;10:41.
126. Toso C, Mentha G, Majno P. Integrating sorafenib into an algorithm for the management of post-transplant hepatocellular carcinoma recurrence. *Journal of hepatology*. 2013;59(1):3-5.
127. Satapathy SK, Das K, Kocak M, et al. No apparent benefit of preemptive sorafenib therapy in liver transplant recipients with advanced hepatocellular carcinoma on explant. *Clinical transplantation*. 2018;32(5):e13246.
128. Bruix J, Takayama T, Mazzaferro V, et al. Adjuvant sorafenib for hepatocellular carcinoma after resection or ablation (STORM): a phase 3, randomised, double-blind, placebo-controlled trial. *Lancet Oncol*. 2015;16(13):1344-1354.
129. Squibb B-M. A study of nivolumab in patients with hepatocellular carcinoma who are at high risk of recurrence after curative hepatic resection or ablation (CHECKMATE 9DX). Available from: <https://clinicaltrials.gov/ct2/show/NCT03383458>. Clinicaltrials.gov identifier NCT03383458. Accessed March 29, 2019.
130. Kornberg A, Kupper B, Tannapfel A, et al. Long-term survival after recurrent hepatocellular carcinoma in liver transplant patients: clinical patterns and outcome variables. *European journal of surgical oncology : the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology*. 2010;36(3):275-280.
131. Regalia E, Fassati LR, Valente U, et al. Pattern and management of recurrent hepatocellular carcinoma after liver transplantation. *Journal of hepato-biliary-pancreatic surgery*. 1998;5(1):29-34.

132. Bodzin AS, Lunsford KE, Markovic D, Harlander-Locke MP, Busuttil RW, Agopian VG. Predicting Mortality in Patients Developing Recurrent Hepatocellular Carcinoma After Liver Transplantation: Impact of Treatment Modality and Recurrence Characteristics. *Ann Surg*. 2017;266(1):118-125.
133. Bates MJ, Farkas E, Taylor D, McFadden PM. Pulmonary resection of metastatic hepatocellular carcinoma after liver transplantation. *The Annals of thoracic surgery*. 2008;85(2):412-415.
134. Han KN, Kim YT, Yoon JH, et al. Role of surgical resection for pulmonary metastasis of hepatocellular carcinoma. *Lung cancer (Amsterdam, Netherlands)*. 2010;70(3):295-300.
135. Hwang S, Kim YH, Kim DK, et al. Resection of pulmonary metastases from hepatocellular carcinoma following liver transplantation. *World journal of surgery*. 2012;36(7):1592-1602.
136. Zhai H, Liang P, Yu XL, et al. Microwave ablation in treating intrahepatic recurrence of hepatocellular carcinoma after liver transplantation: An analysis of 11 cases. *International journal of hyperthermia : the official journal of European Society for Hyperthermic Oncology, North American Hyperthermia Group*. 2015;31(8):863-868.
137. Ko HK, Ko GY, Yoon HK, Sung KB. Tumor response to transcatheter arterial chemoembolization in recurrent hepatocellular carcinoma after living donor liver transplantation. *Korean journal of radiology*. 2007;8(4):320-327.
138. Rivera L, Giap H, Miller W, et al. Hepatic intra-arterial infusion of yttrium-90 microspheres in the treatment of recurrent hepatocellular carcinoma after liver transplantation: a case report. *World journal of gastroenterology*. 2006;12(35):5729-5732.
139. Bhoori S, Toffanin S, Sposito C, et al. Personalized molecular targeted therapy in advanced, recurrent hepatocellular carcinoma after liver transplantation: a proof of principle. *Journal of hepatology*. 2010;52(5):771-775.
140. Gomez-Martin C, Bustamante J, Castroagudin JF, et al. Efficacy and safety of sorafenib in combination with mammalian target of rapamycin inhibitors for recurrent hepatocellular carcinoma after liver transplantation. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2012;18(1):45-52.
141. Sposito C, Mariani L, Germini A, et al. Comparative efficacy of sorafenib versus best supportive care in recurrent hepatocellular carcinoma after liver transplantation: a case-control study. *J Hepatol*. 2013;59(1):59-66.

142. Weinmann A, Niederle IM, Koch S, et al. Sorafenib for recurrence of hepatocellular carcinoma after liver transplantation. *Digestive and liver disease : official journal of the Italian Society of Gastroenterology and the Italian Association for the Study of the Liver*. 2012;44(5):432-437.
143. Yoon DH, Ryoo BY, Ryu MH, et al. Sorafenib for recurrent hepatocellular carcinoma after liver transplantation. *Japanese journal of clinical oncology*. 2010;40(8):768-773.
144. Mancuso A, Mazzola A, Cabibbo G, et al. Survival of patients treated with sorafenib for hepatocellular carcinoma recurrence after liver transplantation: a systematic review and meta-analysis. *Digestive and liver disease : official journal of the Italian Society of Gastroenterology and the Italian Association for the Study of the Liver*. 2015;47(4):324-330.
145. Martin RC, 2nd, Bruenderman E, Cohn A, et al. Sorafenib use for recurrent hepatocellular cancer after resection or transplantation: Observations from a US regional analysis of the GIDEON registry. *American journal of surgery*. 2017;213(4):688-695.
146. Iavarone M, Invernizzi F, Czauderna C, et al. Preliminary experience on safety of regorafenib after sorafenib failure in recurrent hepatocellular carcinoma after liver transplantation. *Am J Transplant*. 2019.
147. Ravaioli M, Cucchetti A, Pinna AD, et al. The role of metronomic capecitabine for treatment of recurrent hepatocellular carcinoma after liver transplantation. *Scientific reports*. 2017;7(1):11305.
148. Kittaï AS, Oldham H, Cetnar J, Taylor M. Immune Checkpoint Inhibitors in Organ Transplant Patients. *Journal of immunotherapy (Hagerstown, Md : 1997)*. 2017;40(7):277-281.
149. Buchbinder EI, Desai A. CTLA-4 and PD-1 Pathways: Similarities, Differences, and Implications of Their Inhibition. *American journal of clinical oncology*. 2016;39(1):98-106.
150. Tanaka K, Albin MJ, Yuan X, et al. PDL1 is required for peripheral transplantation tolerance and protection from chronic allograft rejection. *Journal of immunology (Baltimore, Md : 1950)*. 2007;179(8):5204-5210.
151. Boils CL, Aljadir DN, Cantafio AW. Use of the PD-1 Pathway Inhibitor Nivolumab in a Renal Transplant Patient With Malignancy. *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons*. 2016;16(8):2496-2497.
152. Friend BD, Venick RS, McDiarmid SV, et al. Fatal orthotopic liver transplant organ rejection induced by a checkpoint inhibitor in two patients with refractory, metastatic hepatocellular carcinoma. *Pediatric blood & cancer*. 2017;64(12).

153. Lipson EJ, Bodell MA, Kraus ES, Sharfman WH. Successful administration of ipilimumab to two kidney transplantation patients with metastatic melanoma. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2014;32(19):e69-71.
154. Biondani P, De Martin E, Samuel D. Safety of an anti-PD-1 immune checkpoint inhibitor in a liver transplant recipient. *Annals of oncology : official journal of the European Society for Medical Oncology*. 2018;29(1):286-287.
155. DeLeon TT, Salomao MA, Aqel BA, et al. Pilot evaluation of PD-1 inhibition in metastatic cancer patients with a history of liver transplantation: the Mayo Clinic experience. *Journal of gastrointestinal oncology*. 2018;9(6):1054-1062.
156. Lipson EJ, Bagnasco SM, Moore J, Jr., et al. Tumor Regression and Allograft Rejection after Administration of Anti-PD-1. *The New England journal of medicine*. 2016;374(9):896-898.
157. Gassmann D, Weiler S, Mertens JC, et al. Liver Allograft Failure After Nivolumab Treatment-A Case Report With Systematic Literature Research. *Transplantation direct*. 2018;4(8):e376.
158. Pillai A, Couri T, Charlton M. Liver Allocation Policies in the USA: Past, Present, and the Future. *Digestive diseases and sciences*. 2019;64(4):985-992.
159. Toso C, Meeberg G, Hernandez-Alejandro R, et al. Total tumor volume and alpha-fetoprotein for selection of transplant candidates with hepatocellular carcinoma: A prospective validation. *Hepatology (Baltimore, Md)*. 2015;62(1):158-165.
160. Agopian VG, Harlander-Locke M, Zarrinpar A, et al. A novel prognostic nomogram accurately predicts hepatocellular carcinoma recurrence after liver transplantation: analysis of 865 consecutive liver transplant recipients. *Journal of the American College of Surgeons*. 2015;220(4):416-427.
161. Brattsand R, Linden M. Cytokine modulation by glucocorticoids: mechanisms and actions in cellular studies. *Alimentary pharmacology & therapeutics*. 1996;10 Suppl 2:81-90; discussion 91-82.
162. Wei Q, Gao F, Zhuang R, et al. A national report from China Liver Transplant Registry: steroid avoidance after liver transplantation for hepatocellular carcinoma. *Chinese journal of cancer research = Chung-kuo yen cheng yen chiu*. 2017;29(5):426-437.
163. Yazawa H, Kato T, Nakada T, Sendo F. Glucocorticoid hormone suppression of human neutrophil-mediated tumor cell cytostasis. *International journal of cancer*. 1999;81(1):74-80.
164. Yi NJ, Suh KS, Cho JY, et al. Recurrence of hepatitis B is associated with cumulative corticosteroid dose and chemotherapy against hepatocellular carcinoma recurrence after liver transplantation.

- Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society.* 2007;13(3):451-458.
165. Rodriguez-Peralvarez M, Germani G, Papastergiou V, et al. Early tacrolimus exposure after liver transplantation: relationship with moderate/severe acute rejection and long-term outcome. *Journal of hepatology.* 2013;58(2):262-270.
166. Schumacher G, Oidtmann M, Rueggeberg A, et al. Sirolimus inhibits growth of human hepatoma cells alone or combined with tacrolimus, while tacrolimus promotes cell growth. *World journal of gastroenterology.* 2005;11(10):1420-1425.
167. Vivarelli M, Cucchetti A, La Barba G, et al. Liver transplantation for hepatocellular carcinoma under calcineurin inhibitors: reassessment of risk factors for tumor recurrence. *Annals of surgery.* 2008;248(5):857-862.
168. Vivarelli M, Cucchetti A, Piscaglia F, et al. Analysis of risk factors for tumor recurrence after liver transplantation for hepatocellular carcinoma: key role of immunosuppression. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society.* 2005;11(5):497-503.
169. Benlloch S, Berenguer M, Prieto M, et al. De novo internal neoplasms after liver transplantation: increased risk and aggressive behavior in recent years? *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons.* 2004;4(4):596-604.
170. Rodriguez-Peralvarez M, Manousou P, Lerut J, De la Mata M, Burroughs AK. How much immunosuppression is needed after liver transplantation? *Clinical transplantation.* 2014;28(1):6-7.
171. Vivarelli M, Dazzi A, Zanello M, et al. Effect of different immunosuppressive schedules on recurrence-free survival after liver transplantation for hepatocellular carcinoma. *Transplantation.* 2010;89(2):227-231.
172. Khorsandi SE, Heaton N. Optimization of immunosuppressive medication upon liver transplantation against HCC recurrence. *Translational gastroenterology and hepatology.* 2016;1:25.