

ABSTRACT (200 of 200)

Objective: To examine associations of maternal GDM with offspring cardiovascular biomarkers from late childhood through adolescence.

Methods: We used mixed-effects linear regression models to examine associations of maternal GDM ($n=92$ cases of 597) with average offspring levels of serum lipids (total cholesterol, HDL, LDL, triglycerides) and SBP across two research visits spanning ~10.6 and 16.9 years of age. In sex-stratified analysis, we evaluated the impact of adjustment for sociodemographic characteristics, pubertal status, physical activity and total energy intake, maternal BMI, GDM treatment, and child's BMI.

Results: After adjusting for child's age, pubertal status, race/ethnicity, and maternal education and smoking, GDM exposure was associated with higher total (0.38 [95% CI: 0.16, 0.61] mmol/L) and LDL cholesterol (0.34 [0.14, 0.53] mmol/L) in girls. These estimates were robust to adjustment for lifestyle characteristics and maternal BMI, but were attenuated after accounting for GDM treatment with no appreciable change following further adjustment for current BMI. In boys, maternal GDM corresponded with 4.50 (1.90, 7.10) mmHg higher SBP. This association persisted after accounting for sociodemographic/lifestyle characteristics, maternal BMI, and GDM treatment, but was attenuated after adjusting for current BMI.

Conclusions: Maternal GDM is related to offspring lipid profile and SBP in a sex-specific manner.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/ijpo.12611](https://doi.org/10.1111/ijpo.12611)

KEY WORDS: gestational diabetes mellitus, adolescence, total cholesterol, low-density lipoprotein, blood pressure, longitudinal

INTRODUCTION

In his famous 1980 lecture, “Of Pregnancy and Progeny,” Norbert Freinkel called gestational diabetes mellitus (GDM) a “teratogen” that leads to higher birthweight and congenital defects (1). A number of epidemiological studies have since noted detrimental effects of maternal GDM on offspring fat distribution starting at birth (2-6), and on glycemic regulation (insulin resistance, impaired glucose tolerance, low insulin secretion, and reciprocally altered adipokines (6-9)) by age 6 years. Fewer studies have evaluated associations of *in utero* exposure to GDM with biomarkers of cardiovascular risk, which are important to consider given the interrelations among cardiovascular and metabolic biomarkers, and the persistent burden of cardiovascular disease worldwide (10) – a non-negligible proportion of which may be traced back to the gestational metabolic milieu (11). The majority of publications that have assessed the relation between maternal GDM and cardiovascular health in offspring focused on the adolescent life stage (though one study was conducted in children aged 3 years (12)) and identified associations with higher offspring blood pressure (6, 12-16) and an adverse lipid profile (16). Many of these studies were able to establish associations independent of offspring adiposity (6, 12, 14), suggesting a unique influence of maternal GDM on *in utero* “programming” of cardiovascular risk.

In this prospective study of >500 multi-ethnic youth, we test the hypothesis that *in utero* exposure to maternal GDM with biomarkers of cardiovascular risk in offspring from late childhood through adolescence in the Exploring Perinatal Outcomes in Children (EPOCH)

cohort. The biomarkers of interest included serum lipid profile (total cholesterol, triglycerides, high-density lipoprotein [HDL], and low-density lipoprotein [LDL]) and blood pressure, each of which track from childhood/adolescence into adulthood, and are independent predictors of future cardiovascular and metabolic disease risk (17-21). This work extends current knowledge in this area of research in two key ways. First, we investigate associations of maternal GDM with biomarkers of cardiovascular risk longitudinally across a period that spans the adolescent transition via a repeated measures approach. Given that multiple measurements of health are more informative than single measurements, our analytical strategy is an improvement upon that of current literature, which has focused on evaluating associations of early life exposures with health outcomes at a single point-in-time later in life (including an earlier study in EPOCH that focused on data at the first research visit only (6)). Second, in all models, we consider sex-specific associations given documented differences in fetal (as well as later-life (2)) response to *in utero* exposures (22-24), and sex differences in metabolism during the adolescent life stage (25).

METHODS

Study population

Study participants were from a historical prospective cohort of youth whose mothers were members of the Kaiser Permanente of Colorado (KPCO) Health plan. Details on eligibility and recruitment are published (3). Of the original 604 participants enrolled, we excluded from the present analysis children of 7 women with type 1 diabetes. The analytic sample comprised 597 youth at the first research visit and 410 at the second one with information on at least one cardiovascular risk factor of interest. The Colorado Multiple Institutional Review Board approved this study.

Exposure

All pregnant women at KPCO are routinely screened for GDM at 24-28 weeks using the standard two-step protocol (26). A total of 92 women in the sample had a GDM diagnosis.

Outcomes

Using blood collected after an 8-hour fast, total cholesterol, high-density lipoprotein (HDL), and triglycerides were assayed on the Olympus AU400 advanced chemistry analyzer system. We calculated LDL as (total cholesterol) – (high-density lipoprotein cholesterol [HDL]) – (triglycerides/5). Research assistants measured participants' blood pressure twice in the sitting position using an oscillometric monitor (Dinamap ProCare V100). We used the average of the two values in the analysis and focused on systolic (SBP) rather than diastolic blood pressure

because it is more accurately measured in children and is a stronger determinant of future health outcomes (27).

Covariates

Perinatal and delivery characteristics

We calculated maternal pre-pregnancy body mass index (BMI; kg/m²) using clinically-recorded pre-pregnancy weight from KPCO medical records and measured height at the first research visit. The medical record also provided information on pregnancy complications such as preeclampsia, and birthweight (g) as well as delivery characteristics like gestational age at delivery (weeks) for the index birth.

At T1, the women reported on smoking habits during pregnancy, education level at time of birth, and total gestational weight gain (lbs) via a self-administered questionnaire. In the analysis, we categorized maternal education as a three-level variable (“<high school,” “high school diploma or equivalent,” and “>high school) and smoking as a dichotomous variable (smoked while pregnant with index child, yes vs. no). At this visit, the mothers also filled out a questionnaire regarding any treatment they received for GDM. GDM treatment was categorized as diet and/or exercise only, diet and/or exercise with insulin, and insulin only. Due to the small sample size for insulin only treatment ($n=5$), we combined this category with diet and/or exercise when using this variable as a covariate in the analysis.

Participant characteristics

We calculated participants' age as the difference between date of each research visit and delivery date. At both visits, we measured the participants' weight on a digital scale to the nearest 0.1 kg, and measured height (cm) via a calibrated stadiometer to the nearest 0.1 m. We calculated BMI and standardized it according to the age- and sex-specific World Health Organization (WHO) growth reference (28). Participants self-reported on their race/ethnicity as non-Hispanic White, non-Hispanic Black, Hispanic, and non-Hispanic other. Participants reported their pubertal development based on diagrams of the Tanner stages. We based pubertal status on pubic hair development in boys, and breast development in girls. For each characteristic, we classified a child as pre-pubertal (Tanner stage=1) or pubertal (Tanner stage \geq 2). We obtained information on the participants' physical activity levels using the 3-Day Physical Activity Recall (3DPAR) Questionnaire (29). Using 3DPAR, we derived average energy expenditure (metabolic equivalents; METs) and mean number of moderate- to-vigorous 30-minute blocks of physical activity (METs) over the three-day period. At both visits, participants completed the Block Kid's Food Frequency Questionnaire (FFQ) (30). We used these data in conjunction with the USDA Food Composition Database to estimate total energy intake (kcal/day).

Data analysis

First, we assessed bivariate associations of *in utero* GDM exposure with maternal, perinatal, and offspring characteristics to identify potential confounders. This step, in

conjunction with our *a priori* knowledge of determinants of cardiovascular health in youth informed our multivariable analysis.

Next, we examined associations of GDM exposure with cardiovascular biomarkers from late childhood through adolescence using a mixed-effects linear regression model wherein the outcome of interest was repeated measurements of each biomarker. The explanatory variables included longitudinal assessments of age at each research visit, a random effect for individual ID to account for the repeated outcomes, and an unstructured covariance matrix. We have previously used this approach as a way to characterize growth trajectories from longitudinal assessments of anthropometric data (31-33). A key advantage of this approach (as opposed to conventional linear regression to examine associations of an exposure with the outcome of interest at separate time-points, or with change in the outcome between specific time-points) is that it does not require that all participants have the same number of measurements assessed at exactly the same time. In this study, the participants may have a maximum of two measurements for each biomarker, but if a participant has biomarker data at one visit only, their information still contributes to estimation of standard errors. This approach most efficiently leverages our longitudinal data, both from an analytical standpoint as well as from the perspective of interpreting the biological relevance of results given that multiple assessments of health are likely more meaningful than single evaluations. Using this approach, the estimate of interest (β for GDM exposure) is interpreted as the association of GDM with average levels of a biomarker over the 6 years of follow-up. A key assumption of using mixed-effects models to generate

population-level estimates is homogeneity of associations in the face of variation in the age distribution of the population. Thus, we tested for birth cohort effects by including an interaction term between age at first research visit and GDM exposure; the interaction terms provided no evidence of birth cohort effects.

In multivariable analysis, we evaluated the impact of covariate adjustment on the β and 95% confidence interval (CI) for maternal GDM. Covariates for Model 1 included longitudinal assessments of the child's age in years and pubertal status (i.e., up to two data points for each variable per child, measured at the same time as the cardiovascular outcomes), as well as race/ethnicity, maternal education at the time of birth, and smoking habits during pregnancy. Model 2 further accounted for longitudinal assessments of mean energy expenditure and total energy intake. Model 3 included Model 1 covariates plus maternal pre-pregnancy BMI, given that a woman's pre-gravid weight status likely influences offspring health through shared pathways with hyperglycemia. Model 4 included Model 1 covariates plus GDM treatment to account for variability in maternal hyperglycemia following a diagnosis of GDM. In Model 5, we adjusted for Model 4 covariates plus the child's current BMI to assess for an independent association of maternal GDM (while adjusting for treatment) with offspring cardiovascular biomarkers independent of offspring adiposity.

In all models, we tested for interactions with sex (which tests for a difference in association of GDM with cardiovascular outcomes for girls vs. boys), pubertal status at the first visit (which tests for a difference in association of GDM with cardiovascular outcomes by

pubertal status), and repeated measurements of age across the two research visit (which tests for differences in slopes of biomarker trajectories over time for GDM exposed vs. unexposed). We considered P -interaction <0.05 as the threshold for a need to conduct stratified analysis. As expected, we found evidence of effect modification of association between GDM and cardiovascular risk factors by sex, so we conducted all models separately for boys and girls. We did not observe evidence of effect modification by the other above-mentioned variables.

Sensitivity analyses

In addition to the main analysis, we carried out some sensitivity analyses. First, because height is a strong determinant of BP in children and adolescents (34), and because clinical assessments of pediatric blood pressure are based on age-, sex-, and height-specific values (35), we re-ran models for blood pressure with SBP z-score as the outcome. Results for SBP and SBP z-score were similar, so we focus on interpretation of SBP in mmHg for interpretability. Second, given the co-occurrence of GDM and preeclampsia (36), we re-ran all models after excluding 27 mother-offspring pairs affected by preeclampsia. Exclusion of these individuals did not appreciably change our findings; thus, we included these participants in our analysis. Finally, we assessed the impact of adjustment for birthweight and gestational age at delivery, two mediating perinatal characteristics that are determinants of long-term cardiovascular health (37-39). Adjusting for these variables did not change the direction, magnitude, or precision of the results, so we did not include them in the final models.

We carried out all analyses using Statistical Analyses System software (version 9.3; SAS Institute Inc., Cary, NC).

Author Manuscript

RESULTS

Mean \pm SD age of participants at the first research visit was 10.4 \pm 1.5 years (range: 6.0 to 13.9 years), and 16.7 \pm 1.2 years (range: 12.6 to 19.6 years) at the second research visit, with an approximate follow-up time of 6 years. Fifty percent of the participants were female, and 48.1% ($n=287$) were non-Hispanic White. Approximately 15% ($n=92$) of participants were exposed to GDM *in utero*. **Table 1** shows descriptive characteristics for mother-child dyads stratified by GDM exposure, and **Supplemental Table 1** show the same characteristics stratified by sex. Associations of GDM exposure status with background characteristics were similar across sex, although we noted that a higher proportion of exposed girls were post-pubertal at the first visit (53.5% post-pubertal) than exposed boys (24.5% post-pubertal). **Table 2** shows mean \pm SD for the cardiovascular risk biomarkers of interest at each visit for all children and separately by sex, respectively.

Table 3 displays sex-specific associations of GDM exposure with cardiovascular risk factors over the entire age-range of study participants from both research visits. We interpret these results as the association of *in utero* GDM exposure on average levels of each cardiovascular risk factor across the two research visits (rather than average trajectories of change per year) as we did not find a significant interaction between GDM exposure and time, which suggests a consistent influence of GDM exposure on these health outcomes. After accounting for key confounders in Model 1, girls who were exposed to GDM *in utero* had 0.38 (95% CI: 0.16, 0.61) mmol/L and 0.34 (95% CI: 0.14, 0.53) mmol/L higher total cholesterol and

LDL, respectively, than their unexposed counterparts. This association was robust to adjustment for physical activity and total energy intake (Model 2) and maternal pre-pregnancy BMI (Model 3). However, adjustment for GDM treatment in Model 4 attenuated both estimates to approximately 2/3 the magnitude, with the lower CI crossing the null (**Table 3**). Further adjustment for the offspring's current BMI z-score in Model 5 did not change the estimates.

In boys, GDM exposure corresponded with 4.50 (95% CI: 1.90, 7.10) mmHg higher SBP from late childhood through adolescence (**Table 3, Model 1**). Accounting for physical activity and total energy intake in Model 2 did not change this estimate, although adjusting for maternal pre-pregnancy BMI in Model 3 attenuated the estimate by ~30%. In Model 4, adjustment for GDM treatment augmented the association to 6.27 (95% CI: 0.18, 12.36) mmHg for exposed vs. unexposed males. However, additional adjustment for the offspring's current BMI z-score attenuated the estimate such that the magnitude was similar to those detected in Models 1 and 2, and the lower CI crossed the null (4.60 [95% CI: -1.82, 11.02] mmHg). We also noted that GDM exposure was associated with 0.11 (95% CI: 0.02, 0.20) mmol/L lower HDL in boys, although the 95% CI included the null for all models except Model 3.

Supplemental Table 2 presents estimates of mean \pm SE for the cardiovascular risk factors with respect to GDM exposure after confounder adjustment (Model 1). We provide graphical depictions of these results in **Figure 1** (lipid profile) and **Figure 2** (blood pressure).

Supplemental Table 3 shows the same set of models as in Table 3 for sex-, age, and height-specific blood pressure z-scores. The results from analyses of z-scores are similar to those

of blood pressure in mmHg, indicating no concern regarding residual confounding by age or height in our primary analysis.

DISCUSSION

In this prospective study of 597 multi-ethnic youth in Colorado, we found sex-specific associations of *in utero* exposure to GDM with lipid profile and blood pressure in offspring from late childhood through adolescence. In girls, GDM was associated with higher total cholesterol and low-density lipoprotein (LDL). In boys, GDM corresponded with consistently higher systolic blood pressure. These associations were consistent across the age-range studied, and did not change over time.

Girls

In utero exposure to GDM corresponded with an average of 0.38 mmol/L (6.8 mg/dL) higher total cholesterol, and 0.34 mmol/L (6.1 mg/dL) higher LDL cholesterol levels. These findings are comparable to those of an analysis of 96 male and female Polish youth aged 7 to 16 years wherein Wilk et al. (40) detected unadjusted differences of 19.0 mg/dL and 14.2 mg/dL higher total and LDL cholesterol, respectively, among GDM-exposed ($n=50$) vs. unexposed participants ($n=46$). In a recent study of >1000 boys and girls in the Danish National Birth Cohort, Grunnet et al. reported higher serum triglycerides (5% [95% CI: 1%, 10%]) and lower HDL (-0.07 [95% CI: -0.11, -0.02] mmol/L), as well as marginally higher LDL (0.06 [95% CI: -0.01, 0.14] mmol/L) at 12 years of age among GDM-exposed youth (16). However, these associations became null after adjusting for maternal and child BMI.

In the present analysis, the influence of maternal GDM on the above offspring cardiovascular risk factors remained apparent after accounting for offspring characteristics known to impact lipid profile, including pubertal status, physical activity and energy intake, as well as maternal pre-pregnancy BMI, a proxy for maternal adiposity that is hypothesized to account for a portion of variability in the relationship between maternal hyperglycemia and offspring health. However, accounting for GDM treatment attenuated the estimates for cholesterol and LDL toward the null, thereby providing support for a specific effect of maternal hyperglycemia on these cardiovascular traits. In other words, if removing variability in maternal hyperglycemia during gestation by adjusting for GDM treatment in the analysis resulted in smaller and non-significant effect estimates for maternal GDM, then maternal hyperglycemia is likely a determinant of the offspring health outcomes.

Our finding that exposure to maternal GDM was associated with an adverse lipid profile is likely related to the occurrence of perturbed maternal lipid metabolism in the context of gestational hyperglycemia (41), which in turn, may affect offspring lipid metabolism (42). Mechanistic pathways are difficult to disentangle in humans, but animal models indicate an impact of maternal hyperglycemia on hepatic lipid content and metabolism through oxidative stress and inflammatory pathways (43). Although we adjusted for pubertal status in multivariable analyses, the female-specific nature of the relationship between exposure to maternal GDM and offspring lipid profile may be related to unmeasured confounding by tempo of sexual maturation (which exists on a spectrum that is likely not wholly captured by discrete Tanner staging) given

recent findings that GDM-exposed girls undergo puberty earlier than their male counterparts. Earlier puberty may in turn translate to more rapid increases in total cholesterol and LDL (44).

Boys

Boys who were exposed to GDM *in utero* had ~5 mmHg higher SBP than their unexposed counterparts. This relationship remained apparent after accounting for sociodemographic and lifestyle characteristics during follow-up, but was modestly attenuated after accounting for maternal pre-pregnancy BMI and current BMI. Interestingly, adjustment for GDM treatment did not attenuate the estimate, but rather, slightly increased the magnitude of effect of maternal GDM from ~5 mmHg (Models 1 and 2) to ~6 mmHg higher SBP for GDM exposed vs. unexposed boys (Model 4). This indicates the existence of mechanisms other than hyperglycemia linking *in utero* GDM exposure to offspring blood pressure. Indeed, when we further adjusted for the offspring's current BMI z-score (in addition to GDM treatment), the estimate for maternal GDM in relation to SBP was approximately 5 mmHg and attenuated to non-significance, suggesting that the relationship between maternal hyperglycemia and offspring SBP is mediated by the offspring adiposity.

In a previous study conducted in this cohort that examined consequences of maternal GDM on adiposity and cardiometabolic biomarkers at the first visit, West et al. (6) detected a marginally significant 2 mmHg higher SBP between GDM-exposed vs. unexposed participants at 6-13 years. At this time, the investigators reported no differences in the relationship between GDM and health outcomes by sex, and adjustment for maternal and child BMI attenuated the

associations to the null. Similarly, in an investigation of 1,238 children in the Project Viva cohort, Wright et al. (12) found that GDM exposure corresponded with 3.2 mmHg higher SBP at 3 years of age after adjusting for key covariates including for maternal pre-pregnancy BMI. As with the West et al. study, further adjustment for the child's current BMI markedly attenuated the estimate.

In addition to investigations that focused exclusively on GDM, smaller studies have explored offspring cardiovascular consequences of exposure to maternal diabetes during pregnancy based on a combination of pre-existing diabetes (type 1 or type 2) and GDM with offspring blood pressure. Cho et al. (15) noted 8 mm Hg Higher SBP ($P < 0.001$) in a multi-ethnic population of 99 offspring aged 10-16 years, as compared to 80 unexposed counterparts after controlling for height, weight, sex, age, and pubertal status. Similarly, Rostand et al. (13) reported 8 mmHg ($P < 0.05$) higher SBP among Black and White children born to diabetic mothers as compared to those of non-diabetic mothers ($n=10$ diabetic vs. 252 non-diabetic) after accounting for sex, child's height, and maternal socioeconomic factors. Finally, in the Pima Indian Study, Bunt et al. (14) investigated associations of maternal diabetes with offspring cardiovascular traits after accounting for % fat mass and sex, and detected 11 mmHg higher SBP among 22 Pima Indian children aged 7-11 years exposed to diabetes *in utero*, as compared to 20 offspring of non-diabetic women. Yet again, these studies did not assess the impact of any treatments or interventions to address hyperglycemia during pregnancy.

Author Manuscript

Taken together, the current body of evidence suggests that maternal GDM is associated with higher blood pressure in offspring. While the underlying physiological mechanisms are not known, our data and those of others (12) suggest that this relationship may be mediated by excess adiposity given the established influence of maternal hyperglycemia on offspring fat mass at birth and beyond (2-6). Animal studies also suggest the possibility of a direct effect of hyperglycemia on kidney structure and function (45) which can affect blood pressure. While this hypothesis has not specifically been tested in humans, a study in the Pima Indians found that exposure to maternal diabetes corresponded with nearly four times higher urinary albumin excretion in the offspring, pointing toward the capacity of maternal diabetes to affect fetal nephron development (46). Another possible mechanism linking maternal GDM to an adverse lipid profile involves the co-occurrence of maternal dyslipidemia, which may influence offspring blood pressure through other mechanisms (15). The fact that we detected associations of GDM with SBP in boys but not girls may be related to the more rapid increase and greater variability in SBP from late childhood through adolescence in males than females (25).

Strengths & limitations

Our study had several strengths. First, we were able to examine long-term consequences of exposure to maternal GDM on offspring cardiovascular risk factors across the adolescent transition, a sensitive period for development of cardiometabolic disease (47). Second, rich covariate data enabled us to adjust for variables that may confound or mediate associations of interest. Third, our longitudinal modeling strategy appropriately accounted for correlations

among repeated assessments of cardiovascular risk factors and efficiently leveraged the outcome data across two study visits spanning approximately 6 years of follow-up.

However, this study is not without limitations. A weakness of this study is the fact that we only had access to medical record data which provided information on GDM diagnosis, thereby precluding our ability to explore associations of the *in utero* glycemie milieu at a finer resolution. A second limitation is the slightly decreasing sample sizes in multivariable models due to missing data for covariates, although the sub-samples did not differ with respect to sociodemographic characteristics. A third limitation is the potential lack of generalizability due to over-selection of participants exposed to maternal GDM. Finally, we cannot discount the potential for false positive findings given the large number of models. However, our research focus was to describe and assess the direction, magnitude, and precision of the estimates rather than focus on statistical significance, especially in light of the fact that many of the biomarkers are correlated markers of the same biological phenomena.

Conclusions

In this prospective study of ~600 multi-ethnic youth, we extend current literature regarding the relationship between *in utero* GDM exposure and cardiovascular health in offspring in two key aspects. First, we were able to explore the long-term consequences of maternal GDM on offspring cardiovascular risk factors throughout late childhood and adolescence via a repeated measures technique. This approach is not only more efficient from a statistical standpoint, but also lends itself to more robust findings given that the biomarkers of

interest – including total cholesterol and LDL, which we found to differ by GDM exposure in females – fluctuate across the age range of our study sample (44). Second, we considered sex-specific associations to reflect not only biological differences metabolic and cardiovascular traits as youth progress through the adolescent transition (25), but also, potential discrepancies in fetal response to *in utero* exposures like maternal glycemic control (2).

While the effect sizes that we detected are modest (~0.35 mmol/L higher total cholesterol and 0.31 mmol/L higher LDL in girls; ~5 mmHg higher SBP in boys for GDM exposed vs. unexposed), they may have important long-term ramifications given that cardiometabolic risk factors track from childhood/adolescence into adulthood, and are independent predictors of cardiovascular disease risk (48). For example, in the Bogalusa Heart Study, a 1 mg/dL increment in LDL (equivalent to an increment of 0.026 mmol/L LDL) at 11 years of age corresponded with nearly two times higher odds of metabolic syndrome in young adulthood (49). Similarly, a 1 mmHg increment in childhood SBP predicted elevations in a cluster of metabolic risk factors in young adulthood including SBP, cholesterol, triglycerides, and fasting glucose among Bogalusa Heart Study participants (21). Given the increasing prevalence of GDM in developed countries like the U.S., our findings underscore the importance of efforts to prevent maternal diabetes in order to improve the health of future generations.

CONFLICT OF INTEREST. The authors have no conflicts of interest to report.

ACKNOWLEDGEMENTS:

Author contributions: WP and DD conceived research questions. WP conducted the analysis, wrote the initial draft of the paper, and incorporated co-author comments. CWH and KS contributed to the analytical approach. All authors were involved in writing the paper and approved the final version.

Funding: This study was supported by the National Institutes of Health (NIH), National Institute of Diabetes, Digestive, and Kidney Diseases (R01 DK068001). The funders had no role in the design, conduct, or reporting of this work. We are grateful for Brandy M. Ringham for data preparation, and Deborah Glueck for analytical support. We thank the EPOCH participants, as well as past and present research assistants.

Legends for tables and figures

Table 1 Bivariate associations of *in utero* gestational diabetes mellitus (GDM) exposure with background characteristics of 597 EPOCH participants

Table 2 Descriptive statistics for cardiovascular risk factors of 597 EPOCH participants at two research visits (age 6-14 y and age 12-19 y).

Table 3 Associations of *in utero* exposure to gestational diabetes mellitus (GDM) with cardiovascular (CV) traits across in 597 EPOCH participants across T1 (age 6-14 y) and T2 (age 12-19 y)

Figure 1 Mean \pm SE for lipid profile components among GDM-exposed vs. unexposed EPOCH participants †

† Estimates are adjusted for child's age (years), pubertal status (pre- vs. post-pubertal), race/ethnicity (non-Hispanic White, Hispanic, non-Hispanic Black, non-Hispanic other); and maternal smoking habits during pregnancy (yes vs. no), and education level at time of birth (< high school, high school, >high school).

* Statistically significant difference at alpha = 0.05.

Figure 2 Mean \pm SE for systolic blood pressure (SBP) among GDM-exposed vs. unexposed EPOCH participants

† Estimates are adjusted for child's age (years), pubertal status (pre- vs. post-pubertal), race/ethnicity (non-Hispanic White, Hispanic, non-Hispanic Black, non-Hispanic other); and maternal smoking habits during pregnancy (yes vs. no) and education level at time of birth (< high school, high school, >high school).

* Statistically significant difference at alpha = 0.05.

Supplemental Table 1 Bivariate associations of *in utero* gestational diabetes mellitus (GDM) exposure with background characteristics of 300 girls and 297 boys in EPOCH

Supplemental Table 2 Mean \pm SE for cardiovascular traits among GDM-exposed vs. unexposed EPOCH participants

Supplemental Table 3 Associations of in utero exposure to gestational diabetes mellitus (GDM) with age-, sex-, and height-standardized systolic blood pressure (SBP) z-score across adolescence among 597 EPOCH participants

REFERENCES

1. Freinkel N. Banting Lecture 1980. Of pregnancy and progeny. *Diabetes*. 1980;29(12):1023-35.
2. Regnault N, Gillman MW, Rifas-Shiman SL, Eggleston E, Oken E. Sex-specific associations of gestational glucose tolerance with childhood body composition. *Diabetes care*. 2013;36(10):3045-53.
3. Crume TL, Ogden L, West NA, et al. Association of exposure to diabetes in utero with adiposity and fat distribution in a multiethnic population of youth: the Exploring Perinatal Outcomes among Children (EPOCH) Study. *Diabetologia*. 2011;54(1):87-92.
4. Chang Y, Chen X, Zhang ZK. Intrauterine Exposure to Maternal Diabetes is Associated with Adiposity in Children at 6 Years of Age in China. *Biomedical and environmental sciences : BES*. 2015;28(2):140-2.
5. Kc K, Shakya S, Zhang H. Gestational diabetes mellitus and macrosomia: a literature review. *Annals of nutrition & metabolism*. 2015;66 Suppl 2:14-20.
6. West NA, Crume TL, Maligie MA, Dabelea D. Cardiovascular risk factors in children exposed to maternal diabetes in utero. *Diabetologia*. 2011;54(3):504-7.
7. Sauder KA, Hockett CW, Ringham BM, Glueck DH, Dabelea D. Fetal overnutrition and offspring insulin resistance and beta-cell function: the Exploring Perinatal Outcomes among Children (EPOCH) study. *Diabetic medicine : a journal of the British Diabetic Association*. 2017;34(10):1392-9.

8. Krishnaveni GV, Veena SR, Hill JC, Kehoe S, Karat SC, Fall CH. Intrauterine exposure to maternal diabetes is associated with higher adiposity and insulin resistance and clustering of cardiovascular risk markers in Indian children. *Diabetes care*. 2010;33(2):402-4.
9. Gautier JF, Wilson C, Weyer C, et al. Low acute insulin secretory responses in adult offspring of people with early onset type 2 diabetes. *Diabetes*. 2001;50(8):1828-33.
10. Roth GA, Johnson C, Abajobir A, et al. Global, Regional, and National Burden of Cardiovascular Diseases for 10 Causes, 1990 to 2015. *Journal of the American College of Cardiology*. 2017;70(1):1-25.
11. Oken E, Gillman MW. Fetal origins of obesity. *Obesity research*. 2003;11(4):496-506.
12. Wright CS, Rifas-Shiman SL, Rich-Edwards JW, Taveras EM, Gillman MW, Oken E. Intrauterine exposure to gestational diabetes, child adiposity, and blood pressure. *American journal of hypertension*. 2009;22(2):215-20.
13. Rostand SG, Cliver SP, Goldenberg RL. Racial disparities in the association of foetal growth retardation to childhood blood pressure. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*. 2005;20(8):1592-7.
14. Bunt JC, Tataranni PA, Salbe AD. Intrauterine exposure to diabetes is a determinant of hemoglobin A(1)c and systolic blood pressure in pima Indian children. *The Journal of clinical endocrinology and metabolism*. 2005;90(6):3225-9.

15. Cho NH, Silverman BL, Rizzo TA, Metzger BE. Correlations between the intrauterine metabolic environment and blood pressure in adolescent offspring of diabetic mothers. *The Journal of pediatrics*. 2000;136(5):587-92.
16. Grunnet LG, Hansen S, Hjort L, et al. Adiposity, Dysmetabolic Traits, and Earlier Onset of Female Puberty in Adolescent Offspring of Women With Gestational Diabetes Mellitus: A Clinical Study Within the Danish National Birth Cohort. *Diabetes care*. 2017;40(12):1746-55.
17. Lee JH, Kim HC, Kang DR, Suh I. The 23-year tracking of blood lipids from adolescence to adulthood in Korea: the Kangwha study. *Lipids Health Dis*. 2017;16(1):221-.
18. Clarke WR, Schrott HG, Leaverton PE, Connor WE, Lauer RM. Tracking of blood lipids and blood pressures in school age children: the Muscatine study. *Circulation*. 1978;58(4):626-34.
19. Chen X, Wang Y. Tracking of blood pressure from childhood to adulthood: a systematic review and meta-regression analysis. *Circulation*. 2008;117(25):3171-80.
20. Webber LS, Srinivasan SR, Wattigney WA, Berenson GS. Tracking of serum lipids and lipoproteins from childhood to adulthood. The Bogalusa Heart Study. *American journal of epidemiology*. 1991;133(9):884-99.
21. Rademacher ER, Jacobs DR, Jr., Moran A, Steinberger J, Prineas RJ, Sinaiko A. Relation of blood pressure and body mass index during childhood to cardiovascular risk factor levels in young adults. *Journal of hypertension*. 2009;27(9):1766-74.
22. de Zegher F, Devlieger H, Eeckels R. Fetal growth: boys before girls. *Hormone research*. 1999;51(5):258-9.

23. Clifton VL. Review: Sex and the human placenta: mediating differential strategies of fetal growth and survival. *Placenta*. 2010;31 Suppl:S33-9.
24. Eriksson JG, Kajantie E, Osmond C, Thornburg K, Barker DJ. Boys live dangerously in the womb. *American journal of human biology : the official journal of the Human Biology Council*. 2010;22(3):330-5.
25. Perng W, Rifas-Shiman SL, Hivert MF, Chavarro JE, Sordillo J, Oken E. Metabolic trajectories across early adolescence: differences by sex, weight, pubertal status and race/ethnicity. *Annals of human biology*. 2019:1-10.
26. Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. National Diabetes Data Group. *Diabetes*. 1979;28(12):1039-57.
27. Barker DJ, Bagby SP. Developmental antecedents of cardiovascular disease: a historical perspective. *Journal of the American Society of Nephrology : JASN*. 2005;16(9):2537-44.
28. de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ*. 2007;85(9):660-7.
29. Weston AT, Petosa R, Pate RR. Validation of an instrument for measurement of physical activity in youth. *Medicine and science in sports and exercise*. 1997;29(1):138-43.
30. Cullen KW, Watson K, Zakeri I. Relative reliability and validity of the Block Kids Questionnaire among youth aged 10 to 17 years. *J Am Diet Assoc*. 2008;108(5):862-6.

31. Perng W, Villamor E, Mora-Plazas M, Marin C, Baylin A. Alpha-linolenic acid (ALA) is inversely related to development of adiposity in school-age children. *European journal of clinical nutrition*. 2015;69(2):167-72.
32. Perng W, Mora-Plazas M, Marin C, Villamor E. Iron status and linear growth: a prospective study in school-age children. *European journal of clinical nutrition*. 2013;67(6):646-51.
33. Perng W, Mora-Plazas M, Marin C, Rozek LS, Baylin A, Villamor E. A prospective study of LINE-1DNA methylation and development of adiposity in school-age children. *PLoS One*. 2013;8(4):e62587.
34. Regnault N, Kleinman KP, Rifas-Shiman SL, Langenberg C, Lipshultz SE, Gillman MW. Components of height and blood pressure in childhood. *International journal of epidemiology*. 2014;43(1):149-59.
35. U.S. Department of Health and Human Services. *The Fourth Report on the Diagnosis, Evaluation, and Treatment of High Blood Pressure in Children and Adolescents*. 2005.
36. Weissgerber TL, Mudd LM. Preeclampsia and Diabetes. *Current diabetes reports*. 2015;15(3):9.
37. Crispi F, Miranda J, Gratacós E. Long-term cardiovascular consequences of fetal growth restriction: biology, clinical implications, and opportunities for prevention of adult disease. *American Journal of Obstetrics & Gynecology*. 2018;218(2):S869-S79.

38. Mercurio G, Bassareo PP, Flore G, et al. Prematurity and low weight at birth as new conditions predisposing to an increased cardiovascular risk. *European journal of preventive cardiology*. 2013;20(2):357-67.
39. Kerkhof GF, Breukhoven PE, Leunissen RW, Willemsen RH, Hokken-Koelega AC. Does preterm birth influence cardiovascular risk in early adulthood? *The Journal of pediatrics*. 2012;161(3):390-6.e1.
40. Wilk M, Horodnicka-Jozwa A, Moleda P, Petriczko E, Safranow K, Walczak M. Assessment of selected lipid parameters in children exposed to gestational diabetes (GDM) in utero. *Pediatric endocrinology, diabetes, and metabolism*. 2016;22(4):140-7.
41. Ryckman K, Spracklen C, Smith C, Robinson J, Saftlas A. Maternal lipid levels during pregnancy and gestational diabetes: a systematic review and meta-analysis. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2015;122(5):643-51.
42. Talton OO, Bates K, Salazar SR, Ji T, Schulz LC. Lean maternal hyperglycemia alters offspring lipid metabolism and susceptibility to diet-induced obesity in micedagger. *Biology of reproduction*. 2019;100(5):1356-69.
43. Alfaradhi MZ, Fernandez-Twinn DS, Martin-Gronert MS, Musial B, Fowden A, Ozanne SE. Oxidative stress and altered lipid homeostasis in the programming of offspring fatty liver by maternal obesity. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2014;307(1):R26-R34.

44. Cook S, Auinger P, Huang TT. Growth curves for cardio-metabolic risk factors in children and adolescents. *J Pediatr*. 2009;155(3):S6.e15-26.
45. Amri K, Freund N, Vilar J, Merlet-Benichou C, Lelievre-Pegorier M. Adverse effects of hyperglycemia on kidney development in rats: in vivo and in vitro studies. *Diabetes*. 1999;48(11):2240-5.
46. Nelson RG, Morgenstern H, Bennett PH. Intrauterine diabetes exposure and the risk of renal disease in diabetic Pima Indians. *Diabetes*. 1998;47(9):1489-93.
47. Schooling CM. Life course epidemiology: recognising the importance of puberty. *Journal of epidemiology and community health*. 2015;69(8):820.
48. Raitakari OT, Juonala M, Kahonen M, et al. Cardiovascular risk factors in childhood and carotid artery intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study. *Jama*. 2003;290(17):2277-83.
49. Camhi SM, Katzmarzyk PT, Broyles S, et al. Predicting adult body mass index-specific metabolic risk from childhood. *Metab Syndr Relat Disord*. 2010;8(2):165-72.

Table 1 Bivariate associations of *in utero* gestational diabetes mellitus (GDM) exposure with background characteristics of 597 EPOCH participants

	GDM exposed <i>n</i> = 92	Unexposed <i>n</i> = 505	<i>P</i> -value ^a
Maternal characteristics			
Maternal pre-pregnancy BMI (kg/m ²)	27.8 ± 6.3	25.4 ± 5.8	0.001
Maternal education level			0.50
<High school	5.4%	3.4%	
High school or equivalent	15.2%	18.4%	
>High school	79.4%	78.2%	
Mother smoked during pregnancy	14.1%	7.3%	0.03
GDM treatment			
Diet and/or exercise	66.3%	--	
Diet and/or exercise with insulin, or insulin only	27.2%	--	
None	6.5%	--	
Preeclampsia			0.33
Yes	6.5%	4.2%	
No	93.5%	95.8%	
Child's characteristics			
Female	46.70%	50.90%	0.46
Race/ethnicity			0.02
Non-Hispanic White	63.0%	45.4%	
Hispanic	28.3%	40.0%	
Non-Hispanic Black	4.4%	8.5%	
Non-Hispanic Other	4.4%	6.1%	
<i>Anthropometry & lifestyle at T1</i>			
BMI z-score	0.37 ± 1.33	0.23 ± 1.21	0.32
Post-pubertal (Tanner stage ≥2)	57.10%	37.40%	0.003
Mean energy expenditure (METs)	1.85 ± 0.32	1.90 ± 0.31	0.13
Mean moderate-to-vigorous physical activity (METs)	4.15 ± 2.94	4.26 ± 2.80	0.74
Total energy intake (kcal/per day)	1744 ± 458	1798 ± 562	0.38
<i>Anthropometry & lifestyle at T2</i>			
BMI z-score	0.54 ± 1.05	0.38 ± 1.13	0.26
Post-pubertal (Tanner stage ≥2)	100.00%	100.00%	--
Mean energy expenditure (METs)	1.85 ± 0.27	1.91 ± 0.32	0.12

Mean moderate-to-vigorous physical activity (METs)	4.81 ± 4.48	4.70 ± 3.77	0.82
Total energy intake (kcal/per day)	1683 ± 767	1653 ± 725	0.76

a From an independent t-test for continuous variables; from a Pearson chi-squared test for categorical variables.

Table 2 Descriptive statistics for cardiovascular risk factors of 597 EPOCH participants at two research visits (age 6-14 y and age 12-19 y)

	Visit 1	Visit 2
	<i>n</i> = 300	<i>n</i> = 206
Girls		
Age (years)	10.4 ± 1.4	16.7 ± 1.2
Cholesterol (mmol/L)	4.1 ± 0.7	3.8 ± 0.8
HDL (mmol/L)	1.3 ± 0.3	1.2 ± 0.3
LDL (mmol/L)	2.3 ± 0.6	2.1 ± 0.6
Triglycerides (mmol/L)	1.1 ± 0.5	1.0 ± 0.6
SBP (mmHg)	102.1 ± 9.6	112.0 ± 9.7
Boys		
	<i>n</i> = 297	<i>n</i> = 204
Age (years)	10.5 ± 1.5	16.6 ± 1.1
Cholesterol (mmol/L)	4.1 ± 0.8	3.6 ± 0.6
HDL (mmol/L)	1.3 ± 0.3	1.1 ± 0.2
LDL (mmol/L)	2.4 ± 0.7	2.0 ± 0.5
Triglycerides (mmol/L)	1.0 ± 0.4	1.0 ± 0.4
SBP (mmHg)	103.8 ± 10.3	119.4 ± 10.6

Abbreviations: HDL - high-density lipoprotein; LDL - low-density lipoprotein; SBP - systolic blood pressure.

Table 3 Associations of *in utero* exposure to gestational diabetes mellitus (GDM) with cardiovascular (CV) traits across in 597 EPOCH participants across T1 (age 6-14 y) and T2 (age 12-19 y)^a

	Model 1 <i>n</i> = 300	Model 2 <i>n</i> = 298	Model 3 <i>n</i> = 207	Model 4 <i>n</i> = 300	Model 5 <i>n</i> = 300
Girls (<i>n</i> = 300)					
Cholesterol (mmol/L)	0.38 (0.16, 0.61)	0.39 (0.16, 0.62)	0.34 (0.07, 0.61)	0.24 (-0.08, 0.57)	0.25 (-0.08, 0.57)
HDL (mmol/L)	0.01 (-0.06, 0.08)	0.01 (-0.06, 0.08)	-0.05 (-0.13, 0.02)	-0.01 (-0.12, 0.10)	-0.02 (-0.13, 0.10)
LDL (mmol/L)	0.34 (0.14, 0.53)	0.36 (0.16, 0.55)	0.33 (0.10, 0.55)	0.21 (-0.10, 0.51)	0.21 (-0.10, 0.52)
Triglycerides (mmol/L)	0.11 (-0.07, 0.29)	0.08 (-0.10, 0.26)	0.16 (-0.04, 0.35)	0.10 (-0.12, 0.31)	0.11 (-0.10, 0.32)
SBP (mmHg)	0.93 (-1.16, 3.03)	0.81 (-1.30, 2.93)	-1.52 (-4.04, 0.99)	-1.75 (-6.32, 2.83)	-1.10 (-4.94, 2.74)
	<i>n</i> = 296	<i>n</i> = 293	<i>n</i> = 230	<i>n</i> = 297	<i>n</i> = 297
Boys (<i>n</i> = 297)					
Cholesterol (mmol/L)	-0.06 (-0.28, 0.16)	-0.06 (-0.29, 0.16)	0.03 (-0.22, 0.27)	0.15 (-0.40, 0.70)	0.09 (-0.47, 0.65)
HDL (mmol/L)	-0.06 (-0.14, 0.02)	-0.06 (-0.14, 0.02)	-0.11 (-0.20, -0.02)	0.13 (-0.01, 0.27)	0.14 (-0.01, 0.28)
LDL (mmol/L)	0.01 (-0.18, 0.19)	-0.02 (-0.21, 0.17)	0.10 (-0.11, 0.30)	0.08 (-0.34, 0.51)	0.05 (-0.39, 0.48)
Triglycerides (mmol/L)	0.04 (-0.07, 0.15)	0.06 (-0.06, 0.17)	0.09 (-0.03, 0.21)	-0.12 (-0.54, 0.30)	-0.18 (-0.59, 0.22)
SBP (mmHg)	4.50 (1.90, 7.10)	4.56 (2.02, 7.10)	3.11 (0.28, 5.94)	6.27 (0.18, 12.36)	4.60 (-1.82, 11.02)

^a Estimates are from a linear mixed effects model where the dependent variable is repeated measurements of the CVD traits across T1 and T2, and the predictors include GDM (yes vs. no), covariates, and a random effect for individual intercept.

Model 1: Adjusted for child's age (years), pubertal status (pre vs. post-pubertal), race/ethnicity (non-Hispanic White, Hispanic, non-Hispanic Black, non-Hispanic other); and maternal smoking habits during pregnancy (yes vs. no) and education level at time of birth (<high school, high school, >high school).


Model 2: Model 1 + physical activity levels (average energy expenditure; METs) and total energy intake (kcal)

Model 3: Model 1 + maternal pre-pregnancy BMI.

Model 4: Model 1 + GDM treatment (diet/exercise only vs. none, diet/exercise with insulin vs. none).

Model 5: Model 4 + child's BMI z-score.


Figure 1 Mean \pm SE for lipid profile components among GDM-exposed vs. unexposed EPOCH participants †



† Estimates are adjusted for child's age (years), pubertal status (pre- vs. post-pubertal), race/ethnicity (non-Hispanic White, Hispanic, non-Hispanic Black, non-Hispanic other); and maternal smoking habits during pregnancy (yes vs. no), and education level at time of birth (< high school, high school, >high school).

* Statistically significant difference at $\alpha = 0.05$.

Figure 2 Mean \pm SE for systolic blood pressure (SBP) among GDM-exposed vs. unexposed EPOCH participants †



† Estimates are adjusted for child's age (years), pubertal status (pre- vs. post-pubertal), race/ethnicity (non-Hispanic White, Hispanic, non-Hispanic Black, non-Hispanic other); and maternal smoking habits during pregnancy (yes vs. no) and education level at time of birth (< high school, high school, >high school).

* Statistically significant difference at alpha = 0.05.

Supplemental Table 1 Bivariate associations of *in utero* gestational diabetes mellitus (GDM) exposure with background characteristics of 300 girls and 297 boys in EPOCH

	Girls			Boys		
	GDM exposed <i>n</i> = 43	Unexposed <i>n</i> = 257	<i>P</i> -value ^a	GDM exposed <i>n</i> = 49	Unexposed <i>n</i> = 248	<i>P</i> -value ^a
Maternal characteristics						
Maternal pre-pregnancy BMI (kg/m ²)	26.4 ± 6.6	25.2 ± 5.3	0.24	28.7 ± 6.1	25.5 ± 6.2	0.002
Maternal education level			0.50			0.12
<High school	4.7%	2.3%		12.2%	6.1%	
High school or equivalent	14.0%	19.5%		16.3%	17.3%	
>High school	81.4%	78.2%		77.6%	83.6%	
Mother smoked during pregnancy	16.3%	8.6%	0.11	14.1%	7.3%	0.03
GDM treatment						
Diet and/or exercise	58.1%	--		73.5%	--	
Diet and/or exercise with insulin, or insulin only	30.2%	--		24.5%	--	
None	11.6%	--		2.0%	--	
Mother had preeclampsia	2.3%	3.9%	0.61	10.2%	4.5%	0.11
Yes	2.3%	3.9%		10.2%	4.5%	
Child's characteristics						
Race/ethnicity			0.59			0.01
Non-Hispanic White	60.5%	49.8%		65.3%	40.7%	
Hispanic	30.2%	36.2%		26.5%	44.0%	
Non-Hispanic Black	4.7%	8.6%		4.1%	8.5%	

Non-Hispanic Other	4.7%	5.5%		4.1%	6.9%	
<i>Anthropometry & lifestyle at T1</i>						
BMI z-score	0.37 ± 1.25	0.21 ± 1.17	0.42	0.38 ± 1.42	0.26 ± 1.24	0.55
Post-pubertal (Tanner stage ≥2)	53.5%	65.8%	0.12	24.5%	48.2%	0.002
Mean energy expenditure (METs)	1.77 ± 0.23	1.89 ± 0.29	0.01	1.85 ± 0.32	1.90 ± 0.31	0.13
Mean MVPA blocks (METs)	3.64 ± 2.48	4.08 ± 2.69	0.32	4.64 ± 3.27	4.44 ± 2.91	0.69
Total energy intake (kcal/per day)	1649 ± 470	1687 ± 489	0.62	1827 ± 435	1913 ± 609	0.38
<i>Anthropometry & lifestyle at T2</i>						
BMI z-score	0.51 ± 0.97	0.39 ± 1.05	0.55	0.57 ± 1.13	0.37 ± 1.20	0.34
Post-pubertal (Tanner stage ≥2)	100.0%	100.0%	--	100.0%	100.0%	--
Mean energy expenditure	1.85 ± 0.36	1.90 ± 0.34	0.42	2.00 ± .046	1.96 ± 0.45	0.70
Mean MVPA blocks (METs)	4.43 ± 3.62	4.67 ± 3.70	0.74	5.10 ± 5.05	4.72 ± 3.85	0.82
Total energy intake (kcal/per day)	1589 ± 818	1461 ± 58	0.30	1759 ± 724	1859 ± 802	0.48

Abbreviations: MVPA: moderate-to-vigorous physical activity; **METs** - metabolic equivalents

^a From an independent t-test for continuous variables; from a Pearson chi-squared test for categorical variables.

Supplemental Table 2 Mean \pm SE for cardiovascular traits among GDM-exposed vs. unexposed EPOCH participants

	Mean \pm SE ^a for each outcome according to <i>in utero</i> exposure to GDM		β (95% CI) ^a for GDM exposed vs. unexposed
	Exposed (<i>n</i> = 43)	Unexposed (<i>n</i> = 257)	
Girls (<i>n</i> = 300)			
Cholesterol (mmol/L)	4.29 \pm 0.12	3.91 \pm 0.06	0.38 (0.16, 0.61)
HDL (mmol/L)	1.26 \pm 0.03	1.25 \pm 0.02	0.01 (-0.06, 0.08)
LDL (mmol/L)	2.55 \pm 0.10	2.21 \pm 0.05	0.34 (0.14, 0.53)
Triglycerides (mmol/L)	1.04 \pm 0.09	0.95 \pm 0.03	0.11 (-0.07, 0.29)
SBP (mmHg)	108.2 \pm 1.1	107.5 \pm 0.7	0.93 (-1.16, 3.03)
Boys (<i>n</i> = 297)			
	Exposed (<i>n</i> = 49)	Unexposed (<i>n</i> = 248)	
Cholesterol (mmol/L)	3.77 \pm 0.11	3.83 \pm 0.06	-0.04 (-0.26, 0.19)
HDL (mmol/L)	1.22 \pm 0.04	1.28 \pm 0.03	-0.06 (-0.14, 0.02)
LDL (mmol/L)	2.13 \pm 0.10	2.15 \pm 0.05	0.01 (-0.18, 0.19)
Triglycerides (mmol/L)	0.95 \pm 0.06	0.91 \pm 0.03	0.04 (-0.07, 0.15)
SBP (mmHg)	115.0 \pm 1.3	110.4 \pm 0.7	4.50 (1.90, 7.10)

^a Estimates are from a linear mixed effects model where the dependent variable is repeated measurements of the CVD traits across T1 and T2, and the predictors include GDM (yes vs. no), child's age (years), pubertal status (pre- vs. post-pubertal), race/ethnicity (non-Hispanic White, Hispanic, non-Hispanic Black, non-Hispanic other); and maternal smoking habits during pregnancy (yes vs. no) and education level at time of birth (<high school, high school graduate, >high school); and a random effect for individual intercept.

Supplemental Table 3 Associations of in utero exposure to gestational diabetes mellitus (GDM) with age-, sex-, and height-standardized systolic blood pressure (SBP) z-score across adolescence among 597 EPOCH participants

	β (95% CI) for GDM exposed vs. unexposed^a				
	Model 1 <i>n</i> = 300	Model 2 <i>n</i> = 298	Model 3 <i>n</i> = 207	Model 4 <i>n</i> = 300	Model 5 <i>n</i> = 300
Girls (<i>n</i> = 303)					
SBP z-score	0.08 (-0.15, 0.31)	0.11 (-0.12, 0.33)	-0.13 (-0.40, 0.14)	0.03 (-0.45, 0.50)	0.12 (-0.29, 0.52)
	<i>n</i> = 297	<i>n</i> = 293	<i>n</i> = 230	<i>n</i> = 297	<i>n</i> = 297
Boys (<i>n</i> = 301)					
SBP z-score	0.37 (0.12, 0.61)	0.38 (0.13, 0.63)	0.19 (-0.07, 0.46)	0.50 (-0.30, 1.31)	0.25 (-0.58, 1.08)

^a Estimates are from a linear mixed effects model where the dependent variable is repeated measurements of the CVD traits across T1 and T2, and the predictors include GDM (yes vs. no), covariates, and a random effect for individual intercept.

Model 1: Adjusted for child's age (years), pubertal status (pre- vs. post-pubertal), race/ethnicity (non-Hispanic White, Hispanic, non-Hispanic Black, non-Hispanic other); and maternal smoking habits during pregnancy (yes vs. no) and education level at time of birth (<high school, high school graduate, >high school).

Model 2: Model 1 + physical activity levels (average energy expenditure over 3 days).

Model 3: Model 1 + maternal pre-pregnancy BMI.

Model 4: Model 1 + GDM treatment (diet/exercise only vs. none, diet/exercise with insulin vs. none).

Model 5: Model 4 + child's BMI z-score.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.


Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.


You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.



IJPO_12611_figure1.tif



IJPO_12611_figure2.tif

Title: *In utero* exposure to gestational diabetes mellitus (GDM) and cardiovascular risk factors in youth: A longitudinal analysis in the EPOCH cohort

Running title: Gestational diabetes mellitus and offspring cardiovascular health

Author list and affiliations: Wei Perng PhD^{1,2,3}, Christine W. Hockett PhD¹, Katherine A. Sauder PhD^{1,4}, and Dana Dabelea MD, PhD^{1,2,4}

¹Lifecourse Epidemiology of Adiposity and Diabetes (LEAD) Center, University of Colorado Anschutz Medical Campus, Aurora CO, USA

²Department of Epidemiology, Colorado School of Public Health, Aurora CO, USA

³Department of Nutritional Sciences, University of Michigan School of Public Health, Ann Arbor, MI, USA

⁴Department of Pediatrics, University of Colorado School of Medicine, Aurora CO, USA

Correspondence: Wei Perng, wei.perng@ucdenver.edu | 303-724-4366 (Office)

Conflict of interest: The authors have no conflict of interest to report.

Abstract word count: 245

Manuscript word count: 3498

Tables: 3

Figures: 2

References: 45

Supplemental material: 3 tables

Abbreviations: HDL - high-density lipoprotein; LDL - low-density lipoprotein; SBP - systolic blood pressure; MVPA – moderate to vigorous physical activity; METs – metabolic equivalents, BMI – body mass index