

# Deconstructing Tectonics: Ten Animated Explorations

Christopher R Scotese – PALEOMAP Project/Northwestern University

Ben A van der Pluijm – University of Michigan

Last update: July 22, 2020

## Abstract

The configuration of continents and oceans of our tectonically-active planet are ever-changing. Using new, high-resolution paleogeographic basemaps we created a set of animations that examine key elements of plate tectonics. These time- and space-based paleoglobe reconstructions illustrate: Continental Rifting, Continental Breakup, Ocean Ridges and Fracture Zones, Hotspot Tracks, Arc-Backarc Systems, Continental Collision, Terrane Accretion, Opening-Closing of Ocean Basins, Supercontinent Formation, Plate Velocities, and Future Earth. Each animation is supported by a narrative that offers a brief topical overview, some observations to guide a user's exploration, and key references that formulated the main ideas and concepts that became the foundations of modern plate tectonics.

## Plate Tectonics

Earth is a dynamic planet and the configuration of continents and oceans are ever-changing through geologic time. The planet outermost shell consists of discrete segments, called lithospheric plates, which continuously move relative to one another on the order of cm/year. Lithospheric plates consist of the crust and the mechanically strong part of the upper mantle, ranging in thickness from kilometers in oceanic realms to 200+km in continental regions of the plates. A plate can be viewed as a cap on the surface of a sphere, much like the cracked shell of a boiled egg (McKenzie and Parker, 1967; Morgan, 1968). The contact between two adjacent plates is called a plate boundary, and is where tectonic deformation is concentrated. As plates move, the plate interior (the region away from the plate

**This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1029/2019EA000989**

26 boundaries) stays relatively coherent, although regional structures can develop (such as rift zones,  
27 intracratonic basins, fault-and-fold zones, and plateaus).

28

29 Today's Earth has seven large plates (Pacific, North American, South American, Eurasian, African, Indo-  
30 Australian and Antarctic plates), seven smaller plates (Juan de Fuca, Caribbean, Cocos, Nazca, Scotia,  
31 Arabian, Philippine plates) and several microplates (Figure 1 ; Bird, 2003). A plate can consist entirely of  
32 oceanic lithosphere (such as the Nazca Plate), but most plates consist of both oceanic and continental  
33 lithosphere. For example, the North American Plate consists of the continent of North America and the  
34 western half of the Atlantic Ocean floor. This also means that continental margins, where continental  
35 lithosphere transform into oceanic lithosphere, are not necessarily tectonic plate boundaries.

36




Figure 1. The seven major and several minor plates of Earth. Ocean ridges, transfers zones and trenches (lines with teeth on the overriding plate) mark the surface expression of today's plate boundaries.

37

38 We identify three types of plate boundaries.

39

40

41

42

43

44

1. At divergent plate boundaries, typically ocean ridges, two plates move apart due to seafloor spreading. This process produces new oceanic lithosphere. For example, the boundary between the North American and Eurasian plates, South American and African plates, and Antarctic and Pacific, Nazca, South American, African and Indo-Australian Pacific plates.
2. At convergent plate boundaries, the oceanic part of a plate sinks beneath an overriding plate into the deeper mantle. The overriding plate can be either a continental or oceanic plate. This

45 sinking process, called subduction, gradually consumes the oceanic plate. Volatiles (such as  
46 water and carbon dioxide) released from the subducted plate often trigger melting in the  
47 overlying material. The resulting magma erupts in a chain of volcanoes, called a volcanic arc,  
48 that formed near the edge of the overriding plate. Arcs can be floored by continental or oceanic  
49 crust, with characteristic properties. Examples are, the boundary between the Nazca and South  
50 American plates (with a continental arc), and the Pacific and Eurasian plates (with intraoceanic  
51 arcs).

52 3. The boundary where one plate moves past another is called a lateral-slip fault (or transform  
53 fault). No new plate is created nor consumed along a transform boundary fault. They can occur  
54 in continental or oceanic lithosphere and have comparatively little associated volcanism. For  
55 example, the boundary between the Pacific and North American plates in California, and the  
56 Indo-Australian and Pacific plates in New Zealand are large continental transforms, and many  
57 smaller transforms are found along ocean ridges.

58  
59 The movement of plates generates major geologic structures. Plates that consist of continental  
60 lithosphere, volcanic arcs, or oceanic plateaus are too buoyant to be subducted. Instead, they create  
61 broad belts of deformation, igneous activity and metamorphism, called collision zones. The buoyant  
62 blocks that collide form a larger contiguous continental block, with a zone, called a suture, that marks  
63 where they are joined. Larger continents occasionally form when smaller continents are sutured  
64 together creating a supercontinent. Rifting stretches a continent and splits it apart. As rifting proceeds,  
65 a new ocean ridge is formed that results in the production of oceanic lithosphere. At an unsuccessful (or  
66 failed) rift, rifting stops before the split is complete, and the rift remains as a permanent feature  
67 characterized by volcanic rock and partially filled with continental sediments.

68

69 We recognize two different reference frames for tectonic plate motion (Jurdy, 1990; Demets et al.,  
 70 2010). The absolute reference frame describes plate motions with respect to a fixed point in Earth’s  
 71 interior. In contrast, the relative reference frame describes the motion of one plate with respect to  
 72 another. To illustrate this distinction, consider the motion of two cars driving along a road. If we say  
 73 that Car A travels at 60 km/h and Car B travels at 40 km/h, we are specifying the absolute velocity of the  
 74 cars relative to a fixed point on the road. If we say that Car A drives 20 km/h faster than Car B, we are  
 75 specifying the relative velocity of Car A with respect to Car B.

76  
 77 In this contribution, we explore the fundamentals of plate tectonics in a series of 10 plate tectonic  
 78 vignettes and paleogeographic animations. Each of these animations is accompanied by a brief  
 79 description on tectonic principles, information about regional or local tectonic settings, and references  
 80 pioneering research. The target audiences for this educational contribution are introductory Earth  
 81 science courses, but it also offers foundational material for upper-level college courses in structure and  
 82 tectonics. The literature of plate tectonics since its formulation in the early 1960s is vast, representing  
 83 100,000s of studies. Here, we limit references to works that formulated the main ideas and concepts  
 84 that became the foundations of modern plate tectonics and that continue to stand as seminal works.  
 85 Tremendous progress has been made in the 50 years since these original contributions, which are  
 86 summarized in thousands of review papers and books on plate tectonics theory.




Figure 2. Global Tectonic Tree of the Phanerozoic (since ~450 Ma). Roots that join represent



continental collisions. Upward branching events represent the breakup of continents and the formation of new ocean basins. A branch that terminates represents an ocean basin that stopped opening. Abbreviations are: AFR = Africa; AMR = Amuria; ANT = Antarctica; ARB = Arabia; ARG = Argoland; AUS = Australia; BAJA = Baja California; BLT = Baltica; BSN = Basin and Range; CHL = Chile; CIM = Cimmeria; COR = Corsica and Sardinia; DNZ = Donetz Basin; EUR = Europe; GOND = Gondwana; GRN = Greenland; HON = Honduras; INC = Indochina; IND = India; ITL = Italy; JAP = Japan; KAZ = Kazakhstan; LAR = Laurentia; MEX = Mexico; MB = Marie Byrdland; NAM= North America; NCH = North China; NSL = North Slope of Alaska; NZ = New Zealand; OML = Omolon; PAT = Patagonia; SAM = South America; SBM = Sibumasu; SCH = South China; SCT = Scotia Arc; SEY = Seychelles; SIB = Siberia; SNM = Sonomia; SOM - Somalia; SPN = Spain; WRN = Wrangellia. (After Scotese, 2004)

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

## Making the Maps

102

103

104

Hundreds of new 0.25 to 1 million year interval paleoelevation basemaps were created to produce a set of animations that builds on several decades of plate tectonics and paleogeographic research (the PALEOMAP Project; Scotese & Baker, 1975; Scotese et al., 1979; Scotese & Sager, 1988; Scotese &

105 McKerrow, 1990; Golonka, Ross & Scotese, 1993; Scotese, 2001; Scotese, 2004; Scotese, 2009; Scotese  
106 & Elling, 2017; Scotese & Wright, 2018). These latest animations were created using the GPlates  
107 modeling software (Earthbyte; Scotese, 2016) and the resulting movies smoothed with Adobe® After  
108 Effects. The animations are presented without markup to allow a range of applications and use, except  
109 for the addition of geologic time in millions of years (Ma). The high-resolution of the animations permits  
110 screengrabs for annotation. Supporting figures in the vignettes offer geographic context.

111

112 1. Continental Rifting

113 Animation 1: Continental Rifting [[Vignette01.mp4](#); hotlink at Wiley production stage]

114 Continental rifting is the process by which continental lithosphere undergoes horizontal extension,  
115 creating a rift zone (Burke and Dewey, 1973). During rifting, the lithosphere stretches roughly  
116 perpendicular to the trend of the rift. We distinguish between active and inactive rifts, based on the  
117 timing of extension. Active rifts are places where extension currently takes place. We find an array of  
118 active normal faults that cuts the plate, which is accompanied by earthquakes and volcanic eruptions.  
119 The faulting in active rifts yields a distinctive topography that is characterized by the occurrence of linear  
120 ridges that border depressions.

121  
122 Inactive rifts are places where lithospheric extension ceased some time ago. Instead of earthquakes and  
123 eruptions, we find inactive normal faults and thick deposits of sandstones, conglomerates and volcanics  
124 in depressions. The preservation of an inactive rift means that rifting stopped before it succeeded in  
125 splitting a continent. Such inactive rifts are also known as aulacogens (from the Greek for “furrow”).

126



Figure 3. Map of Africa, showing the East African Rift, the Red Sea and the Gulf of Aden. The East African Rift consists of a belt of normal faults, bounding deep troughs, some filled with water (lakes). The Afar Triangle lies at the triple junction between the Red Sea, the Gulf of Aden, and the East African Rift. In the Red Sea and Gulf of Aden the rift has evolved to form a narrow ocean basin.

127

128 Eastern Africa and the Arabian Peninsula preserve a recent record of continental rifting and incipient  
129 ocean basin formation (Figure 3). We see the evolution of a triple junction (today's Afar Triangle) since  
130 ~35Ma, where one arm, the East African Rift, accommodates limited extension, possibly resulting in a  
131 failed continental rift in the future. Such failed rifts are found around the world, including the Paleozoic  
132 Donets Basin in Ukraine and the Proterozoic Midcontinent Rift of the USA.

133  
134 The other two arms of the triple junction in Africa show the full rift-to-drift cycle, resulting in the  
135 formation of oceanic lithosphere in the Red Sea and Gulf of Aden. Opening of the northern arms of the  
136 triple junction continues today at a rate of 2-3 cm/y. Further ocean spreading in the region may be  
137 limited, however, as the African and Arabian plates move northward and collide with the Eurasian  
138 continent.

139

140 2. Continental Breakup, Ocean Ridges and Fracture Zones

141 Animation 2: Continental Breakup [[Vignette02.mp4](#) ; hotlink at Wiley production stage]

142 Following initial rifting, the successful breakup of continents results in the formation of an oceanic ridge,  
143 which generates new oceanic lithosphere. (Vine, 1966; MacDonald, 1982). Ocean ridges are long, linear  
144 mountain ranges that divide the abyssal plain of the ocean floor. They occur in all modern oceans,  
145 though not necessarily in the middle, and can tower several kilometers above the ocean floor. In some  
146 places, the ridge even rises above sealevel, such as Iceland. Stretched end-to-end, today's ocean ridges  
147 make a submarine mountain chain that encircles the planet (ca. 40,000 km long).

148  
149 At slow spreading ridges, such as the Mid-Atlantic Ridge, plates move apart at rates of <4 cm/y, whereas  
150 at fast spreading ridges, such as the East Pacific Rise, plates move apart at rates of > 8 cm/y. Slow  
151 spreading ridges are steep, relatively narrow (100s km) and have deep axial troughs bordered by step-  
152 like escarpments. Fast spreading ridges have gentle slopes, are up to 1500 km wide, and do not have  
153 axial troughs.

154  
155 The separation of the North and South American plates from the Eurasian and African plates since the  
156 Triassic Period resulted in the formation of the Mid-Atlantic Ridge. The trajectory of an ocean ridge is  
157 accommodated by a special type of strike-slip faults, called ridge-to-ridge transform faults (Wilson,  
158 1965). Unlike typical continental strike-slip faults (also called transcurrent faults), the sense of motion of  
159 a transform fault is opposite to the ridge offset (see Figure 4), as it reflects the relative plate motion on  
160 either side.




Figure 4. (a) Regional setting of the Mid-Atlantic Ridge in the South Atlantic Ocean. SAP = South American Plate, NAP = North American Plate, AFP = African Plate, ANP = Antarctic Plate, NZP = Nazca Plate, CS = Caribbean Sea, SS = Scotia Sea, MAR = Mid-Atlantic Ridge. (b) Evolution of an oceanic transform fault (red). At time 1, the transform is 10 km long, which does not change over time as spreading at the ridges continues. (c) At time 2, the amount of displacement on the fault exceeds the length of the fault. The inactive continuation of a transform fault is marked by a topographic lineament on the ocean floor, called a fracture zone.

161

162

163

164

165

166

167

168

169

170

171

172

173

Once a plate grows beyond the location of the offset ridge segments, active faulting and earthquake activity stop. The traces of the once active transform faults form long, linear oceanic fracture zones, which, like railroad tracks, track the motions of the plates. Given a difference in age of the ocean floor on either side of fracture zone, these regions produce steep ocean floor scarps with the older, denser side sitting lower.

The S-shaped margins of Africa and South America require major curvature of the Mid-Atlantic Ridge, which is accommodated by steps with transfer faults, some as long as hundreds of km. These strike-slip faults link straight ridge segments, with spreading of about 3cm/y. The animation also shows the evolution of the Caribbean plate that migrates eastward, closing of the Atlantic-Pacific seaway and connecting North and South America via Central America.

### 174 3. Hotspots and Hotspot Tracks

175 Animation 3: Hotspots and Hotspot Tracks [[Vignette03.mp4](#) ; hotlink at Wiley production stage]

176 Anomalously hot mantle can create continental and oceanic regions at the surface that are  
177 characterized by extensive volcanic activity. These volcanic regions are thought to be due to hot,  
178 buoyant, rising plumes of mantle, called mantle plumes, and the associated volcanic activity is called  
179 hotspot volcanism (Morgan, 1971). Hotspots are characterized by the deposition of a large volume of  
180 volcanic rocks with geochemical signatures that are distinct from subduction-related and ocean ridge  
181 or rift-related volcanism (Sleep, 1992). Especially voluminous hotspots produce oceanic island chains  
182 and plateaus (Figure 5).

183

184 One modern example of hotspot volcanism is the Hawaiian volcanic island chain in the Pacific Ocean.  
185 The mantle is more stationary than the overlying lithospheric plate (moving faster by at least one order  
186 of magnitude), so the location of active volcanism changes as the plate moves over the mantle plume  
187 with time. This creates a trail of volcanic islands that become older as they move away from the active  
188 hotspot (Wilson, 1963). Today, Hawaii's Big Island is the active part of one such hot spot track, while a  
189 series of extinct volcanic islands and submarine volcanoes extend off to the northwest and north,  
190 forming the Hawaiian-Emperor island chain.

191




Figure 5. Hotspots of the modern world (data from Whitaker et al. 2015).

192

193 Some mantle plumes are long-lived, lasting 100 million years or more, while others are short-lived,  
194 lasting less than 10 million years. The orientation of a hotspot track provides the direction of plate  
195 motion relative to a mantle reference frame, so absolute motion, and the age of volcanic rocks along the  
196 track records the velocity of overlying plates.

197  
198 The opening of the Atlantic Ocean along the Mid-Atlantic Ridge was accompanied by several mantle  
199 plumes under the late Paleozoic supercontinent Pangea, including Iceland in the North Atlantic and  
200 Tristan da Cunha in the South Atlantic. The South Atlantic preserves a trail of progressively older  
201 hotspot volcanism from the southern Mid-Atlantic Ridge, marked today by Tristan da Cunha, to northern  
202 Namibia in Africa and southern Brazil in South America.

203  
204 Closing the Atlantic Ocean shows these two continental areas coming together at ~140 Ma and marks  
205 the location of a mantle plume under Pangea at the start of ocean spreading. The Tristan da Cunha  
206 hotspot track is especially pronounced in the eastern South Atlantic and is mirrored by a track to the  
207 west of the mid-Atlantic Ridge. Notice also that the mirror-image tracks are not parallel to the ridge's  
208 spreading direction, showing that the ridge and plates were together moving in a northerly direction  
209 relative to the hotspot (absolute) framework.

210

211



#### 212 4. Volcanics Arcs and Backarc Basins

213 Animation 4: Volcanic Arcs and Backarc Basins [[Vignette04.mp4](#); hotlink at Wiley production stage]

214 Trenches are linear or curvilinear troughs that mark the boundary, at Earth's surface, between the  
215 downgoing (or subducting) plate and the overriding plate (Mitchell and Reading, 1971). The floor of the  
216 Mariana Trench in the western Pacific reaches a depth of over 11 km (Figure 6), which is deeper than  
217 the highest mountains (Mt. Everest is ~ 9 km high). Trenches form as the downgoing oceanic  
218 lithosphere pulls the ocean floor into the mantle.

219  
220 A volcanic arc is a chain of volcanoes that forms along the edge of the overriding plate about 200-300  
221 km from the trench and about 100–150 km above the surface of the subducted oceanic lithosphere.  
222 Most of the magma that rises to feed the volcanic arc forms by partial melting of mantle above the  
223 downgoing oceanic slab. Partial melting takes place primarily because volatiles (H<sub>2</sub>O) released from the  
224 downgoing plate reduce the melting point of the overlying mantle rock.

225  
226 The region on the other side of the volcanic arc, away from the subducting oceanic lithosphere, is the  
227 back-arc region (Karig, 1970; Uyeda and Kanamori, 1979). Its character varies with tectonic setting, and  
228 can be contractional, extensional, or tectonically stable. Extensional back-arc settings contain rift zones  
229 that may evolve into basins floored by oceanic lithosphere. One example of a large back-arc basin with  
230 active seafloor spreading occurs behind the Mariana Volcanic Arc in the western Pacific Ocean. Another  
231 example of a back-arc basin in the northwestern Pacific is the Japan Sea, which formed about 30 million  
232 years ago by rifting of the Japan Volcanic Arc from eastern Asia, due to subduction along the Japan  
233 Trench.

234

235 Back-arc regions may also be contractional, producing fold-thrust belts and basement uplifts, such as  
236 those found along the Andean margin of South America. Lastly, back-arc region where no deformation  
237 occurs are called stable back-arcs, such as the Bering Sea that is located north of the Aleutian Islands,  
238 which connects Alaska and Kamchatka.

239




Figure 6. Trenches (heavy lines), arc volcanoes (black dots) and backarc basins related to subduction in the western Pacific Ocean Basin. The depth to the subducted plate interface is shown by contour lines, given in multiples of 50 km (e.g., 2 means 100 km).

240

241 5. Continental Collision

242 Animation 5: Continental Collision [[Vignette05.mp4](#); hotlink at Wiley production stage]

243 As subduction progressively consumes an oceanic plate, a piece of buoyant crust on the downgoing  
244 plate eventually reaches the trench. Examples of buoyant crust include large continents, but also  
245 smaller continental fragments, island arcs, oceanic plateaus (regions of anomalously thick oceanic crust  
246 formed by hotspot volcanism) and spreading ridges. Buoyant lithosphere cannot be subducted, so when  
247 it impinges on the overriding plate, the area becomes a tectonic collision zone. The boundary between  
248 the once separate plates, called a suture zone, often preserves slivers of oceanic crust and mantle (a  
249 rock suite called ophiolites) that have become trapped and exposed in the collision zone (Dewey and  
250 Bird, 1970; Coleman, 1971).

251



Figure 7. Tectonic elements that accreted in South-central Asia during the Late Paleozoic, Mesozoic, culminating with India's collision in Early Cenozoic time. The sutures between fragments are shown by barbed lines with the barbs indicating the upper plate of the former subduction zone. The suture between the Qangtang and Songpan Ganze (SG) fragments is of early Mesozoic age representing the closure of the Paleotethys Ocean. All sutures to the north are Paleozoic. The southern boundary of the Lhasa block is the Indus-Tsangpo suture and represents the closure of Neotethys. Dark regions are variably deformed, Cenozoic sedimentary rocks.

252

253 The collision of India with Southern Asia is the modern culmination of a long history of convergence and  
254 successive collisions in the region since the Paleozoic (Figure 7; Molnar and Tapponnier, 1975).

255 Northward motion of India continues today, resulting in the tectonically active Himalayan Mountain

256 range and Tibetan Plateau (which includes the Qangtang and Lhasa blocks). The region's latest collision

257 started around 50 Ma and formed the Indus-Tsangpo suture zone. The closure of smaller ocean basins  
258 preceded the terminal India-Asia collision, recorded by accretion of volcanic arcs (Şengör, 1979).

259

## 260 6. Terrane Accretion

261 Animation 6: Terrane Accretion [[Vignette06.mp4](#) ; hotlink at Wiley production stage]

262 The parts of plates that are relatively buoyant (meaning less dense than surrounding material) cannot be  
263 subducted. Therefore, such pieces eventually collide with, detach and suture to other buoyant plate  
264 segments (see Continental Collision). After each collision, a new convergent margin forms on the  
265 oceanic side of the collision zone, and the continent grows. This process is called accretion and the  
266 pieces that are added to larger continental blocks by this process are called accreted (or exotic) terranes  
267 (Coney et al., 1980).

268  
269 The recognition that part of a continent is an accreted terrane comes from analysis of the geologic  
270 histories of terranes and adjacent regions. If the rocks and structures of these fault-bounded blocks do  
271 not readily correlate with adjacent regions, then that block was likely accreted. In addition,  
272 paleomagnetism is used to test whether an accreted block and the continent to which it is now attached  
273 occupied different paleogeographic locations. Fossils are also used to determine if a terrane originated  
274 at a different location than its neighboring continent.

275  
276 Based on geologic mapping, geochronology, paleomagnetism and paleontology, we learn that, during  
277 the Mesozoic and Cenozoic, the North American Cordillera grew westward by as much as 1500 km due  
278 to accretion of exotic terranes. Much of the land that now comprises California, Oregon, Washington  
279 and Alaska in the United States and British Columbia and the Northwest Territories in Canada originated  
280 as accreted terranes (Figure 8).

281  
282 During the Jurassic and Cretaceous (150 Ma – 100 Ma) two major terranes, Stikine and Wrangellia,  
283 collided with the western margin of North America (Monger et al., 1982). These terranes originated at a

284 more southerly latitude than their current location. Convergence and collision were oblique, so right-  
 285 lateral strike-slip faults cut and moved slivers of these accreting terranes northward along the North  
 286 American plate margin.

287




Figure 8. Map of western North America that shows the regions consisting of accreted terranes. The oblique convergence of terranes resulted in strike-slip faulting along the western North American continental margin that sliced the original tectonic terranes into separate blocks. For example, the accreted Wrangellia terrane (dark brown) is found today as dismembered blocks along the North American margin, from Idaho to Alaska.

288

289 7. Closing and Opening of Ocean Basins

290 Animation 7: Closing and Opening of Ocean Basins [[Vignette07.mp4](#); hotlink at Wiley production stage]

291 The formation of the Appalachians and Caledonides mountains in eastern North America and western  
292 Europe, respectively, resulted from closure of a vanished ocean basin, called the Iapetus Ocean. Today, a  
293 modern ocean separates these continental blocks, the Atlantic Ocean. In late Proterozoic to early  
294 Paleozoic times, the ancient Iapetus Ocean opened and grew by rifting and seafloor spreading, and  
295 closed in the late Paleozoic when the bordering continents collided, forming the Appalachians-  
296 Caledonides collisional mountain belt. In Mesozoic times, the Atlantic Ocean opened in about the same  
297 relative position between the continental blocks of North and South America on one side, and Europe  
298 and Africa on the other side. Today, small subduction zones exist in the Caribbean region and the South  
299 Georgia-South Sandwich Islands of the Atlantic Ocean, which may lead to the future demise of the  
300 Atlantic Ocean and new America-Europe/Africa collision (see #10, Future Earth).

301  
302 The repetition of successive stages of rifting, seafloor spreading, convergence and collision is called the  
303 “Wilson Cycle”, in recognition of the person who first highlighted this pattern (Figure 9; Wilson, 1966).  
304 As a result of this process, we do not find oceanic lithosphere in today’s ocean basins that is older than  
305 about 200 Ma. All older oceanic lithosphere has been subducted, except for a few slivers preserved as  
306 ophiolites in ancient mountain ranges, which mark the tectonic sutures between colliding continents  
307 (Dewey and Bird, 1970). In contrast, older continental lithosphere is widely preserved on today’s  
308 surface, because it was too buoyant to be subducted. This explains why Proterozoic and Archean rocks,  
309 some as old as 4 billion years, are found on most of today’s continents.




Figure 9. (a-b) A continent rifts, such that the crust stretches, faults and subsides. (c) Seafloor spreading begins, forming a new ocean basin. (d) The ocean widens and is flanked by passive margins. (e) Subduction of oceanic lithosphere begins on one of the passive margins, closing the ocean basin by subduction of the oceanic part of plate and the ridge. (f-g) The closure of the ocean basin culminates with continental collision. At a later time, continental rifting begins again and the process repeats, in what is called the Wilson Cycle. (Modified from S. Stein, used with permission).

310

311

312

313

314

315

316

317

318

319

320

321

322

The Paleozoic evolution of the Appalachians of North America and the Caledonides of Western Europe records the progressive closure of the Iapetus Ocean (Dewey, 1969). This past ocean basin contained several terranes that were accreted during convergence, including volcanic arcs and ocean islands (MacNiocaill et al., 1997). The Iapetus Ocean formed in the same relative position as today's Atlantic Ocean, but was rimmed by the E-W oriented continental margins of North America (Laurentia) and Europe-Africa (Gondwana). The Iapetus Ocean first opened around the late Precambrian and fully closed by the Late Carboniferous, forming the Appalachian-Caledonide collisional mountain belt of the supercontinent Pangea. The area's later history is marked by counter-clockwise rotation into today's N-S configuration of the successor ocean basin, the Atlantic Ocean. The opening of the Atlantic Ocean started in the north and continued to the south, as part of the Tethys Ocean to the east closed.



## 323 8. Formation of Supercontinents

324 Animation 8: Formation of the supercontinent Pangea [[Vignette08.mp4](#); hotlink at Wiley production  
325 stage]

326 The closure of the Iapetus Ocean led to the formation of a large continental landmass, or  
327 supercontinent, called Pangea (meaning “all-land”; Wegener, 1929/1966). At various times in the  
328 geologic past, plate movements and continental collisions produced similar supercontinents that lasted  
329 for several tens of millions of years before they broke apart.

330  
331 The supercontinent Pangea was formed in the late Paleozoic (260 Ma) and rifted apart during the early  
332 Mesozoic (Figure 10). An older supercontinent Rodinia was formed at the end of the Mesoproterozoic  
333 (~1.1 Ga) and dispersed by ~750 Ma (Hoffman, 1991). There is also evidence that supercontinents  
334 formed even earlier in Earth history, such as the supercontinent Nuna that formed at the end of the  
335 Paleoproterozoic (~1.8 Ga), indicating that supercontinents repeatedly formed and subsequently broke  
336 up. This mega-tectonic process is called the Supercontinent Cycle or the “Wegener Cycle” after the  
337 German scientist who championed the idea that the landmasses were once joined together in a large  
338 continent that he called “Pangea”.

339



Figure 10. The Pangea Supercontinent formed after the closure of ancient ocean basins in approximately the same relative location as today’s Atlantic Ocean. The subducted ocean basins included the ancient Rheic, Theic and Iapetus Oceans. Final closure of these ocean basins resulted in the Appalachian-Caledonian mountain chain of North America and western Europe. Pangea

contained all major Paleozoic landmasses. The large internal ocean, Tethys, is later subducted as the Indian plate moved northward and collided with the China block of Asia.

340

341 The supercontinent cycle may be related to long-term (>100 million years) convection patterns in the  
342 deeper mantle. The relative motion of a continent is controlled by plate forces, particularly the  
343 downward pull from negative buoyancy of cold and dense plate material (Forsyth and Uyeda, 1975), but  
344 over longer periods of time the continental components accumulate over a mantle downwelling to form  
345 a supercontinent. Once formed, the thermal structure of the mantle beneath it changes, because the  
346 supercontinent acts as a giant insulator that blocks escaping heat from the mantle. In response, the  
347 mantle below the supercontinent heats up, and is no longer a region of downwelling. Instead, hot  
348 mantle begins to upwell beneath the supercontinent, causing it to heat up and weaken the overlying  
349 plate (Anderson, 1982). Eventually, a rift system forms that evolves into a new ocean basin, dispersing  
350 the supercontinent.

351

352 With the last supercontinent ~250 my old, we are still in the dispersal stage of the supercontinent cycle,  
353 but in the next 200 million years or so, the continents may once again coalesce to form a new  
354 supercontinent that we call "Pangea Proxima" (see #10, Future Earth).

355

## 356 9. Plate Velocities

357 Animation 9: Plate Velocities [[Vignette09.mp4](#); hotlink at Wiley production stage]

358 Linear magnetic anomalies on the ocean floor, which preserve the reversing polarity of the Earth's  
359 magnetic field over time, and geochronologic data offer a robust geologic estimate of plate velocities  
360 during the last 200 million years. Latitudinal velocities measured by paleomagnetic measurements offer  
361 estimates of plate velocities for times before that, but with greater uncertainty. For these reasons, we  
362 use the history of plate motions from 200 Ma to today to explore tectonic plate velocities. Plate  
363 velocities are given in cm/year, which is equivalent to 10's of km per million years (Figure 11).

364




Figure 11. Global (black) and continent-only (brown) plate velocities. The continental curve represents the motion of continents alone, without the oceanic portion of a plate. Plates and continents with large areas and/or high velocities dominate fast velocity intervals. Horizontal scale is age in million years; vertical scale is speed in cm/y. From Zahirovic et al. (2015).

365

366 The paleogeographic evolution of the oceans and continents since 200Ma shows that some plates move  
367 fast and others slower, in the range of 1-20 cm/year, and that accelerations and decelerations are  
368 common. Opening of the Atlantic was relatively slow, whereas opening of the Pacific was fast. The  
369 subduction of the Tethys Ocean was very fast at times, until collision between India and Asia which  
370 greatly slowed India's northward movement.

371

372 Using the PALEOMAP model of plate motions, Zahirovic et al. (2015) tracked the velocities of about a  
373 dozen plates, which traveled at an average global plate velocity of ~4 cm/year. Notably, plates with  
374 continental components tend to be relatively slow. Given that continental lithosphere is thicker and

375 more buoyant than oceanic lithosphere, the continental component of a plate introduces drag on plate  
376 motion. Plates do not move at constant velocity over time and may move as fast as 18 cm/year, as  
377 exemplified by the Cenozoic northward motion of India. The plate velocities estimated from the global  
378 plate model match modern plate velocities that are based on geodetic measurements.

379

## 380 10. Future Earth

381 Animation 10: Future Earth [[Vignette10.mp4](#); hotlink at Wiley production stage]

382 There is no certainty about the future plate geometry of the Earth, but using current plate motions and  
383 tectonic principles we can make educated guesses about future plate configurations. We surmise that  
384 the Atlantic and Indian Oceans will continue to widen until a new subduction zone brings these  
385 continents back together, eventually forming a new supercontinent, “Pangea Proxima”. The northward  
386 trajectory of Africa suggests that the Red Sea and Gulf of Aden will close, and that the East African Rift  
387 will not grow into an ocean basin. Africa will collide with Europe and Arabia closing the Mediterranean  
388 Sea and the Red Sea. A Himalayan-scale mountain range will extend from Spain, across Southern  
389 Europe, through the Middle East and into Asia. Similarly, Australia will collide with Southeast Asia and a  
390 new subduction zone will encircle Australia, extending westwards across the Central Indian Ocean.

391




Figure 12. The tectonic setting of a future supercontinent, Pangea Proxima. After 100 my into the future, the Atlantic Ocean becomes subducted, while the Pacific Ocean continues to grow and a small remnant of the Indian Ocean eventually becomes land-locked after the northward convergence and accretion of Antarctica.

392

393 One of the key events in this geography of the future is the beginning of subduction along the eastern  
394 coasts of North America and South America (Figure 12). As the Atlantic Ocean widens, the Puerto Rican  
395 Trough and Scotia Arc propagate northward and southward along the east coast of North and South  
396 America. In time, this new westward dipping subduction zone will consume the Atlantic Ocean.

397  
398 The Atlantic Ocean, 100 million years in the future, will begin to narrow by subduction beneath the  
399 Americas. The Indian Ocean will also contract due to northward subduction into the Central Indian  
400 trench. Antarctica will collide along the southern margin of Australia, and the Mid-Atlantic Ridge, the  
401 last vestige of seafloor spreading in the Atlantic Ocean basin, will be subducted beneath eastern North  
402 America. Once the last bit of the Atlantic Ocean's spreading ridge is subducted beneath the Americas,  
403 the Atlantic Ocean will rapidly close and a new supercontinent will begin to form. The rocks that contain  
404 the remains of ancient New York City, Boston and Washington DC will be sitting atop high mountain  
405 ranges.

406  
407 About 250 million years in the future, the Atlantic and Indian oceans will have closed. North America  
408 will have collided with Africa, but in a more southerly position than from where it rifted. In this new  
409 supercontinent, South America is wrapped around the southern tip of Africa, with Patagonia in contact  
410 with Antarctica and Indonesia, enclosing a remnant of the Indian Ocean. The Pacific has grown much  
411 wider, encircling more than half the Earth. We call this future supercontinent, "Pangea Proxima",  
412 because it would be the next Pangea, but not necessarily the last!

413

## 414 Acknowledgements

415 The project was created with the PALEOMAP dataset (<https://www.earthbyte.org/paleomap-paleoatlas->  
416 [for-gplates/](https://www.earthbyte.org/paleomap-paleoatlas-for-gplates/)) from new, high-resolution maps with greater time and space resolution than earlier  
417 models, and developed with the open-source GPlates software (<https://www.gplates.org/>) and Adobe®  
418 software. We thank journal reviewers John Encarnacion and David Mogk and the editor for thoughtful  
419 comments that helped to improve the presentation. The project was partially supported by funds from  
420 Northwestern University (CRS) and the University of Michigan-Ann Arbor (BvdP).

421

## 422 References

- 423 Anderson, D., 1982. Hotspots, polar wander, Mesozoic convection and the geoid. *Nature*, 297, 391–393.  
424 <https://doi.org/10.1038/297391a0>
- 425 Bird, P., 2003. An updated digital model of plate boundaries. *Geochemistry Geophysics Geosystems*, 4,  
426 1027, doi:10.1029/2001GC000252.
- 427 Burke K. & Dewey J.F. 1973. Plume generated triple junctions: key indicators in applying plate tectonics  
428 to old rocks. *Journal of Geology*, 81, 406–433.
- 429 Coleman, R. G., 1971. Plate tectonic emplacement of upper mantle peridotites along continental edges,  
430 *J. Geophys. Res.*, 76, 1212-1222.
- 431 Coney, P.J., Jones, D.L., Monger, J.W.H., 1980. Cordilleran suspect terranes. *Nature*, 288, 329–333.
- 432 DeMets, D.C., Gordon, R.G., Argus, D.F., 2010. Geologically current plate motions. *Geophys. J. Int.*, 181,  
433 1–80, doi:10.1111/j.1365-246X.2010.04491.x.
- 434 Dewey, J. F., J. M. Bird, 1970. Mountain belts and the new global tectonics, *J. Geophys. Res.*, 75, 2625-  
435 2647.
- 436 Dewey, J.F., 1969. Evolution of the Appalachian/Caledonian Orogen. *Nature*, 222, 124-129.

437 Forsyth, D, Uyeda, S., 1975. On the Relative Importance of the Driving Forces of Plate Motion.  
438 Geophysical Journal International, 43, 163–200, <https://doi.org/10.1111/j.1365->  
439 246X.1975.tb00631

440 Golonka, J., Ross, M.I., and Scotese, C.R., 1994. Phanerozoic Paleogeographic and Paleoclimatic  
441 Modeling Maps, in A. F. Embry, B. Beauchamp, and D.J. Glass (editors), Pangea, Global  
442 Environments and Resources, Canadian Society of Petroleum Geologists Memoir, 17, 1-47.

443 Hoffman, P.E., 1991. Did the breakout of Laurentia turn Gondwana inside out? *Science*, 252, 1409-1412.

444 Jurdy, D.M., 1990. Reference frames for plate tectonics and uncertainties. *Tectonophysics*, 182, 373-382,  
445 [https://doi.org/10.1016/0040-1951\(90\)90173-6](https://doi.org/10.1016/0040-1951(90)90173-6).

446 Karig, D., 1970. Ridges and Trenches of the Tonga-Kermadec Island Arc System. *Journal of Geophysical*  
447 *Research*, 75, 239-254.

448 Macdonald, K. C., 1982. "Mid-Ocean Ridges: Fine Scale Tectonic, Volcanic and Hydrothermal Processes  
449 within the Plate Boundary Zone". *Annual Review of Earth and Planetary Sciences*, 10, 155–190.

450 McKenzie, D., Parker, R.L., 1967. The North Pacific: An example of tectonics on a sphere. *Nature*, 216,  
451 1276-1280.

452 Mac Niocaill, C., van der Pluijm, B.A., Van der Voo, R., 1997. Ordovician paleogeography and the  
453 evolution of the Iapetus Ocean. *Geology*, 25, 159-162.

454 Mitchell, A.H, Reading, H.G., 1971. Evolution of Island Arcs. *Journal of Geology*, 79, 253-284.

455 Molnar, P., and P. Tapponnier, 1975. Cenozoic tectonics of Asia: Effects of a continental collision,  
456 *Science*, 189, 419–426, doi:10.1126/science.189.4201.419.

457 Monger, J.W.H., Price, R.A., Tempelman-Kluit, D.J., 1982. Tectonic accretion and the origin of the two  
458 major metamorphic and plutonic belts in the Canadian Cordillera. *Geology*, 10, 70-75.

459 Morgan, J., 1971. Convection plumes in the lower mantle: *Nature*, 230, 42-43.

460 Morgan, W.J., 1968. Rises, trenches, great faults, and crustal blocks. *Journal Geophys. Res.*, 73, 1959–  
461 1982.

462 Scotese, C.R., 2001. *Atlas of Earth History, Volume 1, Paleogeography*, PALEOMAP Project, Arlington,  
463 Texas, 52 pp.

464 Scotese, C.R., 2004. A continental drift flipbook, *Journal of Geology*, 112, 729-741.

465 Scotese, C.R., 2009. Late Proterozoic plate tectonics and paleogeography: A tale of two supercontinents,  
466 Rodinia and Pannotia, in *Global Neoproterozoic petroleum systems: The emerging potential in*  
467 *North Africa*, J. Craig, J. Thurow, A. Whitman, and Y. Abutarruma (editors), Geological Society of  
468 London Special Publication, 326, 67-83.

469 Scotese, C.R., 2016. Tutorial: PALEOMAP Paleoatlas for GPlates and the PaleoData Plotter Program,  
470 <http://www.earthbyte.org/paleomap-paleoatlas-for-gplates/>

471 Scotese, C.R., and Elling, R., 2017. Plate Tectonic Evolution during the Last 1.5 Billion Years: The Movie,  
472 William Smith Meeting, “Plate Tectonics at 50”, Geological Society of London (abstract).  
473 <https://www.youtube.com/watch?v=CnVGFv-1Wqc&feature=youtu.be>.

474 Scotese, C.R., and Wright, N., 2018. PALEOMAP Paleodigital Elevation Models (PaleoDEMS) for the  
475 Phanerozoic, PALEOMAP Project, [https://www.earthbyte.org/paleodem-resource-scotese-and-](https://www.earthbyte.org/paleodem-resource-scotese-and-wright-2018/)  
476 [wright-2018/](https://www.earthbyte.org/paleodem-resource-scotese-and-wright-2018/)

477 Scotese, C.R., Baker, D.W., 1975. Continental drift reconstructions and animation, *Journal of Geological*  
478 *Education*, 23, 167-171.

479 Scotese, C.R., Bambach, R.K., Barton, C., Van der Voo, R., Ziegler, A.M., 1979. Paleozoic base maps,  
480 *Journal of Geology*, 87, 217-277.

481 Scotese, C.R., McKerrow, W.S., 1990. Revised world maps and introduction, in *Paleozoic Paleogeography*  
482 *and Biogeography*, W.S. McKerrow and C.R. Scotese (editors), Geological Society of London  
483 *Memoir*, 12, 1-21.



484 Scotese, C.R., Sager, W.W., 1988. 8th Geodynamics Symposium, Mesozoic and Cenozoic Plate  
485 Reconstructions. *Tectonophysics*, 155, 1-399.

486 Şengör, AMC, 1979. Mid-Mesozoic closure of Permo–Triassic Tethys and its implications. *Nature*, 279,  
487 590–593.

488 Sleep, N.H., 1992. Hotspot Volcanism and Mantle Plumes. *Annual Review of Earth and Planetary  
489 Sciences*, 20, 19-43.

490 Uyeda, S., Kanamori, H., 1979. Back-arc opening and the mode of subduction. *Journal of Geophysical  
491 Research: Solid Earth*, 84 1049–1061, doi:10.1029/JB084iB03p01049.

492 Vine, F., 1966. Spreading of the Ocean Floor: New Evidence. *Science*, 154, 1405-1415.

493 Wegener, A., 1966. *The Origin of Continents and Oceans* (translated from Wegener, A., 1929, *Die  
494 Entstehung der Kontinente und Ozeane*). New York: Dover. ISBN 0-486-61708-4.

495 Wilson, J.T. , 1963. A possible origin of the Hawaiian Islands. *Canadian Journal of Physics*, 41, 863–870.


496 Wilson, J.T., 1965. A New Class of Faults and their Bearing on Continental Drift. *Nature*, 207, 343–347.

497 Wilson, J.T., 1966. Did The Atlantic Close And Then Re-Open? *Nature*, 211, 676-681.

498 Zahirovic, S., Dietmar Müller, R., Seton, M., Flament, N., 2015. Tectonic speed limits from plate  
499 kinematic reconstructions. *Earth and Planetary Science Letters*, 418, 40-52.  
500 doi.org/10.1016/j.epsl.2015.02.037

Figure 1.

Author Manuscript



This article is protected by copyright. All rights reserved.

Figure 2.

Author Manuscript








Figure 3.


Author Manuscript




Figure 4.

Author Manuscript






(a)



(b)



(c)

This article is protected by copyright. All rights reserved.

Figure 5.

Author Manuscript




Figure 6.

Author Manuscript




Figure 7.

Author Manuscript




This article is protected by

Figure 8.

Author Manuscript





- Alaska
- NS North Slope
  - Kv Kagvik
  - En Endicott
  - R Ruby
  - Sp Seaward Peninsula
  - I Innoko
  - NF Nixon Fork
  - PM Pingston and McKinley
  - YT Yukon-Tanana
  - Cl Chuiitna
  - P Peninsular
  - W Wrangellia
  - Cg Chugach and Prince William
  - TA Tracy Arm
  - T Taku
  - Ax Alexander
  - G Goodnews

- Canada
- Ch Cache Creek
  - St Stikine
  - BR Bridge River
  - E Eastern assemblages

- Washington, Oregon, and California
- Ca Northern Cascades
  - SJ San Juan
  - O Olympic
  - S Siletzia
  - BL Blue Mountains
  - Trp Western Triassic and Paleozoic of Klamath Mountains
  - KL Klamath Mountains
  - Fh Foothills belt
  - F Franciscan and Great Valley
  - C Calaveras
  - Si Northern Sierra
  - SG San Gabriel
  - Mo Mohave
  - Sa Salinia
  - Or Orocopia

- Nevada
- S Sonomia
  - RM Roberts Mountains
  - GL Golconda

- Mexico
- B Baja
  - V Vizcaino











Figure 9.

Author Manuscript





Early Triassic 237 Ma




- Ancient Landmass
- Modern Landmass
- Subduction Zone (triangles point in the direction of subduction)
- Sea Floor Spreading Ridge

This article is protected by copyright. All rights reserved.

© 2000 C. R. Scotese, PALEOMAP Project


Figure 11.

Author Manuscript











Author Manuscript





ESS2\_650\_2019EA000989-f01-z-.png




ESS2\_650\_2019EA000989-f02-z-.png


Author Manuscript




ESS2\_650\_2019EA000989-f03-z-.png



(a)




(b)




(c)


ESS2\_650\_2019EA000989-f04-z-.png





ESS2\_650\_2019EA000989-f05-z-.png






ESS2\_650\_2019EA000989-f07-z-.png



Author Manuscript



Alaska

- NS North Slope
- Kv Kagvik
- En Endicott
- R Ruby
- Sp Seaward Peninsula
- I Innoko
- NF Nixon Fork
- PM Pingston and McKinley
- YT Yukon-Tanana
- Cl Chuiitna
- P Peninsular
- W Wrangellia
- Cg Chugach and Prince William
- TA Tracy Arm
- T Taku
- Ax Alexander
- G Goodnews

Canada

- Ch Cache Creek
- St Stikine
- BR Bridge River
- E Eastern assemblages

Washington, Oregon, and California


- Ca Northern Cascades
- SJ San Juan
- O Olympic
- S Siletzia
- BL Blue Mountains
- Trp Western Triassic and Paleozoic of Klamath Mountains
- KL Klamath Mountains
- Fh Foothills belt
- F Franciscan and Great Valley
- C Calaveras
- Si Northern Sierra
- SG San Gabriel
- Mo Mohave
- Sa Salinia
- Or Orocopia

Nevada

- S Sonomia
- RM Roberts Mountains
- GL Golconda


Mexico

- B Baja
- V Vizcaino




ESS2\_650\_2019EA000989-f09-z-.png


Early Triassic 237 Ma



ESS2\_650\_2019EA000989-f10-z-.png



ESS2\_650\_2019EA000989-f11-z-.png



ESS2\_650\_2019EA000989-f12-z-.png