

# Author Manuscript

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/cyt.12885](https://doi.org/10.1111/cyt.12885)

This article is protected by copyright. All rights reserved

[DR GUIDO FADDA \(Orcid ID : 0000-0001-9736-0544\)](#)  
[DR ESTHER DIANA ROSSI \(Orcid ID : 0000-0003-3819-4229\)](#)

Article type : Original article

## Relevance of Rosette Patterns in Variants of Papillary Thyroid Carcinoma

Marco Dell'Aquila<sup>1</sup>, Teresa Musarra<sup>1</sup>, Vincenzo Fiorentino<sup>1</sup>, Chiara Brunelli<sup>1</sup> Costanza De Marco<sup>1</sup>, Marco Raffaelli<sup>2</sup>, Emanuela Traini<sup>2</sup>, Celestino Pio Lombardi<sup>2</sup>, Guido Fadda<sup>1</sup>, Luigi Maria Larocca<sup>1</sup>, Liron Pantanowitz<sup>3</sup>, Esther Diana Rossi<sup>1</sup>°.

<sup>1</sup>Division of Anatomic Pathology and Histology; <sup>2</sup>Division of Endocrine-Surgery-Fondazione Policlinico "Agostino Gemelli"-IRCCS, Rome Italy, <sup>2</sup> Division of Endocrine surgery-Fondazione Policlinico "Agostino Gemelli"-IRCCS-Rome Italy, <sup>3</sup>Department of Pathology, University of Pittsburgh Medical Center, Pittsburgh, PA, USA

° These authors share the senior authorship

**Running title:** Rosettes in thyroid carcinoma

**KEY WORDS:** Thyroid, rosette pattern, microfollicular pattern, papillary pattern, thyroid malignancy, PTC variant

**Corresponding author:** Dr. Esther Rossi, MD, PhD, MIAC- Division of Anatomic Pathology and Histology – Fondazione Policlinico Universitario "Agostino Gemelli"-IRCCS, Università Cattolica

del Sacro Cuore, Largo Francesco Vito, 1 – 00168 Rome (Italy) Phone: +3906-3015-4433 Fax: +3906-3015-7008, e-mail: esther.rossi@policlinicogemelli.it

Conflicts of interests: the authors have no conflicts of interest to declare

## ABSTRACT

**BACKGROUND:** The detection of rosette-like clusters (RLC) of follicular cells in thyroid carcinoma has been reported mostly in the columnar cell variant of papillary thyroid carcinoma (CCV-PTC). Despite the fact that diagnosing variants of PTC is no longer encouraged by The Bethesda System for Reporting Thyroid Cytopathology (TBSRTC), the identification of cytomorphological features such as RLC linked with these tumors might help reduce possible misinterpretation in thyroid fine needle aspiration cytology (FNAC). We accordingly investigated the potential correlation of architectural patterns including RLC with PTC variants.

**MATERIALS AND METHODS:** We analyzed 225 thyroid FNAC cases diagnosed as suspicious for malignancy (SFM) and positive for malignancy (M) over a 1 year time where all samples had corresponding histology. We also included 150 benign lesions (BL) from the same time period. The presence of RLC versus similar appearing solid clusters, papillary structures, and microfollicles were evaluated. We also performed immunocytochemistry (ICC) and molecular testing for BRAFV600E.

**RESULTS.** We included 100 (44.4%) SFM favoring PTC and 125 (55.6%) M cases with cyto-histological correlation. On histology, all SFM and M cases showed malignancy including 140 (62.2%) classic PTC and 85 (37.8%) PTC variants. The cytomorphological patterns in all FNA samples included solid (74%), papillary (89%), microfollicular (70%), and pseudo-RLC morphology (25.7%). We identified only pseudo-RLC in 33 FNA specimens from PTC variant cases that included tall cell variant (42.4%), hobnail variant (21.2%) and miscellaneous variants (36.3%) of PTC. No

definitive RLC were detected in our series. ICC and BRAFV600E were not specifically linked with a RLC pattern.

**CONCLUSIONS:** These findings demonstrate that in our dataset the architectural pattern of RLC was not recognized within PTC variants. However, we did identify a pseudo-RLC pattern that was observed in association with TCV and hobnail variant cases of PTC.

## INTRODUCTION

Rosette and rosette-like patterns can be found in different tumors, particularly those from the nervous system<sup>1</sup>. Histologically, a typical rosette is characterized by a small round group or cluster of cells organized in a spoke-wheel or halo arrangement around a central lumen. The name is derived from their morphological resemblance to rose windows found in gothic churches, such as those found in the Cathedral of Notre Dame in Paris<sup>2-4</sup>. Whilst different authors have described rosette and rosette-like patterns as a manifestation of certain tumors, this architectural pattern alone is not pathognomonic of any specific tumor type<sup>1-2</sup>. Their presence, in fact, may be associated with various forms of tumor differentiation<sup>1-10</sup>. True rosettes are a characteristic finding in neurological tumors<sup>1</sup>. However, rosettes have also been identified in other neoplasms such as medullary thyroid carcinoma (MTC), osteosarcoma, fibromyxoid sarcoma, rhabdomyosarcoma, endometrial stromal sarcoma and some non-Hodgkin lymphomas<sup>1-11</sup>. Pseudo-rosettes tend to be associated with neuroblastomas, medulloblastomas, malignant melanoma, Merkel cell carcinoma, other neuroendocrine tumors of the skin, primitive neuroendocrine tumor (PNET), retinoblastoma, ependymoma, neurocytoma, glioblastoma, Wilm tumor and pheochromocytoma<sup>1-11</sup>.

To the best of our knowledge, the histological identification of a rosette pattern in thyroid histological samples has not been well published. However, rosettes have been previously documented in cytologic samples from benign and malignant lesions such as adenomatous goiter,

insular carcinoma and MTC)<sup>11-17</sup>. Sen et al also reported the presence of a rosette-like pattern in a fine needle aspiration cytology (FNAC) specimen of a well-differentiated thyroid follicular carcinoma diagnosed as columnar cell variant of papillary thyroid carcinoma (CCV-PTC)<sup>12</sup>. Only a few other papers have reported this finding associated with thyroid carcinoma in which a rosette-like pattern was linked with aggressive variants of PTC including the tall cell variant (TCV) of PTC.<sup>13-17</sup> Therefore, whether a rosette-like pattern is indeed associated with CCV-PTC and its potential diagnostic value in thyroid cytology remains unknown. In order to assess any possible correlation of rosette-like morphology with specific PTC variants, we accordingly, examined a retrospective series of thyroid FNAC cases for such specific features.

## **MATERIALS AND METHODS**

We received institutional (Catholic University of the Sacred Heart) ethical approval for this study. A retrospective search was performed for all thyroid FNACs diagnosed as SFM and M over a one year period (January 2018 to December 2018) at the Catholic University “Agostino Gemelli” Hospital in Rome, Italy. Furthermore, in order to determine the presence of a rosette-like pattern in benign versus malignant thyroid entities, we included another 150 benign sequential lesions diagnosed during the same study period. The institutional electronic medical record system Armonia-Metafora, Italy (CU) was also searched for thyroidectomy specimens during the same study period. The patient’s age, gender, FNAC diagnoses and follow-up information were recorded. All available pathology slides were reviewed. The majority of the thyroid nodules were evaluated and biopsied under ultrasound guidance by clinicians and radiologists.

### **Thyroid FNAC Specimens:**

All aspirations (with usually two passes performed for each lesion) were performed with 25 to 27G needles. No rapid on-site assessment for adequacy of material was done. All patients consented to

their procedure. All FNAC specimens were processed using a ThinPrep 5000™ processor (Hologic Co., Marlborough, MA, USA). Prepared slides were fixed in 95% methanol and stained with a Papanicolaou stain. Any remaining material was stored in Preservcyt solution for potential ancillary studies.

Specimen adequacy was determined according to the Bethesda and British RCPATH classification schemes.<sup>18-19</sup> The cytology cases were classified and diagnosed according to the new Italian Working Group SIAPEC-IAP classification.<sup>20-21</sup> All of the cases were re-evaluated and then re-classified according to The Bethesda System for Reporting Thyroid Cytology II (TBSRTC, 2017).<sup>19, 36</sup> For this retrospective study, analyses were conducted using TBSRTC terminology. This case series included the following distribution of diagnoses: 5.9% non-diagnostic (ND) including cystic cases; 77.8% benign lesions (BL); 3% atypia of undetermined significance/follicular lesion of undetermined significance (AUS/FLUS); 6.1% follicular neoplasms (FN); 2.2% SFM and 5% malignant (M) cases. All cytology and histology cases were reviewed by two cytopathologists whilst the re-classification according to TBSRTC was done by one cytopathologist (EDR). Cases with an equivocal interpretation were subject to consensus review. The concordance between SIAPEC-IAP and TBSRTC classification systems was 95.9%.

According to the literature, rosettes are defined as a group of 15 or fewer follicular cells with a circular arrangement and central lumen that may/may not be filled with colloid.<sup>18-19</sup> Most of the cytoplasm of the cells in a rosette tends to be located towards the center (i.e. nuclei are peripheral), implying that these cells are tall/columnar in shape. At our institution we utilize LBC preparation versus the conventional slide (CS) method. Figures 1A-1C and Table 1 compares the rosette-like clusters (RLC) as reported by Sen to microfollicular and papillary structures. By comparison, a microfollicle is also comprised of a group of 15 or fewer follicular cells with a circular arrangement and central area usually filled with colloid. However, these follicular cells are shorter (cuboidal) and

thus exhibit less cytoplasm in the microfollicle center. A micropapillary cluster is comprised of 15 or more follicular cells arranged around a central fibrovascular core. The individual follicular cells in a papillary cluster are also shorter (more cuboidal) and have minimal cytoplasm that typically does not merge toward the center of the cellular cluster. Nonetheless, we only found a pseudorosette pattern in some cases. A pseudorosette pattern resembles rosettes, but does not exhibit the features depicted in Figure 1a and as described in the paper by Sen et al.<sup>12</sup> The presence/absence of these different follicular structures was recorded in our cohort of cases.

#### **Immunocytochemistry (ICC) Analysis:**

In our routine practice we perform ancillary techniques including ICC and BRAF<sup>V600E</sup> analysis in all suspicious and malignant thyroid cytological cases. Specifically, we perform ICC on all liquid-based cytology (LBC) samples. HBME-1 and Galectin-3 ICC is performed using a protocol previously described by our group.<sup>22-24</sup> The minimal percentage of adequate lesional cells for the performance of ICC evaluation was defined at 30% in LBC samples. Lesional cells were interpreted to be positive when at least 50% of the cells demonstrated strong cytoplasmic staining. To avoid false negative and/or false positive yields, this 50% ICC cutoff value was also used for histological tissue sections. A case was considered to be overall positive for malignancy when there was concomitant expression of both immunomarkers. Adequate Galectin-3 immunoreactivity was represented by cytoplasmic staining, and suitable HBME-1 staining was chiefly within the cytoplasm with accentuation on the cytoplasmic membrane and within the lumen. Positive controls included mesothelioma for HBME-1 (membranous positivity) and histiocytes for Galectin-3 (cytoplasmic staining). Lymphocytes identified in the majority of the thyroid slides were used as a negative control.

**Molecular analysis for BRAF mutation:** DNA was extracted from both LBC stored material and paraffin embedded tissues, with 100% concordance. BRAF<sup>V600E</sup> mutational analysis was performed on DNA extracted from cytological and surgical specimens containing at least 70% tumor. Details of the protocol employed have been previously published by our group.<sup>25-28</sup>

#### **Histology Specimens:**

All surgical specimens were fixed in 10% buffered formaldehyde, embedded in paraffin and 5 micron-thick sections then stained with hematoxylin-eosin. The diagnosis of classical variant of PTC was based on the presence of true papillary structures and distinctive nuclear features. The diagnosis of FVPTC relied upon the detection of follicular architecture coupled with the nuclear features of PTC in multiple foci. Encapsulated tumors with either lympho-vascular invasion (within the capsule or beyond) or capsular penetration were diagnosed as invasive FVPTC. All cases were classified according to the seventh edition of the tumor-node-metastasis (TNM)-based staging system recommended by the American Joint Commission on Cancer (AJCC)<sup>29</sup>. The different PTC variants were classified according to the WHO 2017<sup>30-31</sup>. For the definition of TCV of PTC, we adopted the standard definition of at least 30% of tall cell component. However, in our analysis we also included cases of PTC with a less than 30% TCV component. The histological diagnosis of non-invasive follicular thyroid neoplasm with papillary like nuclear features (NIFTP) was rendered according to the criteria described by Nikiforov et al.<sup>24</sup> However, in our institution NIFTP terminology was used for follicular neoplasms without any overt papillary structures.

#### **Statistical Analysis:**

Statistical analysis was performed using GraphPad-Prism 5 software (Graph Pad Software, San Diego, CA) and MedCalc version 10.2.0.0 (MedCalc Software, Mariakerke, Belgium). Statistical comparison of continuous variables was performed using the Mann-Whitney U-test or Paired t-test,



as appropriate. Comparison of categorical variables was performed using the chi-square statistic, and the Fisher's exact test. P-values less than 0.05 were considered as statistically significant.

## RESULTS

Table 2 summarizes the clinic-pathological features for our retrospective series. In order to validate the search for a rosette-like pattern in malignant lesions as reported by Sen et al<sup>12</sup>, we selected SFM and M cases with a with cyto-histological correlation resulting in a histological diagnosis of PTC and its variants. Furthermore, in order to determine if there was a difference of a rosette-like pattern for benign and malignant thyroid entities, we also included 150 sequential benign lesions diagnosed during the same study period. The entire series was composed of 239 female and 136 male patients with an age range of 9 to 82 years (mean: 36.5 years). The size of thyroid lesions subject to FNAC ranged from 5 mm to 65 mm in greatest dimension (median size = 14 mm). All sub-centimeter lesions were discovered during radiologic screening for causes unrelated to thyroid disease. There was no significant difference in the size of these thyroid nodules among the diagnostic entities. The cytological series included 150 BL (40%), 100 SFM (26.6%) and 125 positive for malignancy (M) (33.3%). Only 20 (13.3%) out of 150 BL had histological follow-up, showing 15 goiter and 5 follicular adenoma (FA) cases. The histological diagnoses for SFM and positive for malignant cases included: 140 classical PTC (cPTC-62.2%) and 85 PTC variants (37.8%).

The distribution of the PTC variants, analyzed in Table 3, was characterized by 140 classic PTC, 19 pure tall cell variant (TCV), 27 PTC with a less than 30% TCV component, 13 hobnail PTC variant, 8 follicular variant of PTC (FVPTC), 6 Warthin-like PTC, 5 columnar cell variant of PTC (CCV-PTC), 4 cases with both TCV plus CCV-PTC, and 3 solid variant of PTC (Table 3). For the TCV cases, we included both pure TCV (19 cases) and also PTC cases with a TCV component (27 cases, 58.7%) that contained less than 30% tall cells. In Table 3 we further evaluated the 85 PTC variants

distributed according to their cytology diagnosis of SFM (33 cases) and positive for malignancy (52 cases). On histology, the 33 SFM were diagnosed as 6 pure TCV and 14 PTC with a less than 30% TCV component, 6 hobnail PTC variant, 1 Warthin-like PTC, 4 FVPTC and 2 CCV-PTC. The 52 malignant cases were diagnosed as 13 TCV and 13 PTC with less than 30% TCV, 7 hobnail PTC variant, 5 Warthin-like PTC, 4 FVPTC, 3 CCV-PTC, 4 combined TCV and CCV and 3 as solid variant of PTC (Table 3).

Tables 4 and 5 compare the distribution of the different follicular architectural patterns encountered in the different samples according to FNAC and final histopathology diagnoses, respectively. In the 150 benign cases we confirmed the presence of large sheets of follicular cells, few follicular clusters, and lack of any papillary, rosette or pseudorosette patterns. Regardless of the FNAC diagnostic category, Table 4 shows that solid (without either a rosette or a pseudo-rosette pattern) clusters were present in all PTC cases. The different cytologic architectural patterns (in order of decreasing frequency) showed that there were 225 cases with solid clusters of neoplastic cells (100%), isolated follicular cells in 201 cases (89.3%), papillary structures in 200 cases (88.8%), microfollicular clusters in 154 cases (68.4%). None of our cases had an unequivocal RLC pattern. Nonetheless, we did identify a pseudorosette pattern in 33 cases (14.6%, Figures 2A-C and Figure 3A-B). Comparing these different cytomorphological patterns stratified according to the FNAC diagnosis of SFM and positive for malignancy, we found that the number of cases with pseudorosettes was documented in 21 cases with M and 12 in SFM without any significant statistical correlation ( $P>0.001$ ). The other cytomorphological patterns had a similar distribution in both SFM and malignant cases. In Table 5 we analyzed these patterns specifically according to the different histological variants of PTC. A pseudorosette pattern was found in cases with a histological diagnosis of pure TCV (14 cases-42.4%) and hobnail variant (7 cases-21.2%). Further, seven out of nine cases with a CCV component (including three pure CCV-PTC and 4 mixed TCV and CCV-PTC) had a focal pseudorosette pattern.

None of our cPTC cases showed a RLC component or a pseudorosette pattern. Of note, we did not find any pseudorosette cluster within the histological samples of our series.

None of our BL cases had been diagnosed with the support of ancillary techniques. ICC stain results are reported in Table 6. ICC results showed that the majority of our SFM and M cases (209 cases, 92.8%) had a concordant positive immunopanel for HBME-1 and Galectin-3 with only 16 cases (7.1%) showing positivity for a single immunomarker (i.e. HBME-1). For cases grouped in the SFM category, ICC showed 9 cases with negative staining for galectin-3 and revealed 100% positivity for HBME-1. The cases interpreted as being positive for malignancy had positive ICC staining in all cases except for seven that only showed immunopositivity for HBME-1. None of our malignant lesions had a negative concordant panel.

The different cytomorphological patterns did not correlate with any of the analyzed immunomarkers. BRAF<sup>V600E</sup> was expressed in 47 cases out of 225 malignancies (21%). We found a 100% concordance between cytology and histology for BRAF<sup>V600E</sup> expression. The results of molecular testing for BRAF<sup>V600E</sup> demonstrated a lack of any statistical correlation between BRAF<sup>V600E</sup> and architectural patterns. In fact, we found that 88 cases (70.4%) were of BRAF<sup>V600E</sup> wild type phenotype whilst 37 M cases (29.6%) had a BRAF mutation (Table 6). Among the BRAF<sup>V600E</sup> mutated cases, we found 14 cPTC, 7 hobnail PTC and 16 TCV. Of note, we did not find any specific significant statistical correlation between BRAF and the pseudorosette pattern ( $p>0.001$ ).

## DISCUSSION

It is well established that specific morphological patterns may be characteristic of particular tumors.<sup>1-</sup>  
<sup>2</sup> Such patterns often play an important role in the pathological identification of these specific entities. However, they are usually not pathognomonic of any one tumor. One such pattern observed in various tumors is the formation of rosettes or rosette-like structures. The term “rosette” has been attributed to a pattern resembling a flower, characterized by cells radially organized around a center. Four main kinds of rosette have been described with different morphological features including: 1) Homer Wright rosette; 2) the Flexner-Wintersteiner rosette; 3) the ependymal rosette and 4) perivascular pseudo-rosettes.<sup>1-10</sup> Whilst these rosettes may not always be seen, their recognition is often helpful in the diagnosis of neural entities such as medulloblastoma/PNET, retinoblastoma, ependymoma, central neurocytoma and pineocytoma.

To the best of our knowledge, the definition of a rosette pattern has not been reported in thyroid lesions based on histological samples. Nevertheless, whilst a few books and atlases have mentioned the possibility of focal rosette-like features, this has not been clearly linked with either a specific entity or PTC variant. Of note, a few papers have described the presence of a rosette-like component representing a minor or focal feature in cytological samples from both benign and malignant thyroid lesions, even though they lacked pictures demonstrating this pattern.<sup>11-17</sup> In our series of 150 BL we did not find any true rosette or pseudo-rosette follicular structures, confirming that this pattern is likely linked more with malignant thyroid nodules. The case report by Sen et al is, to the best of our knowledge, the only paper reporting the presence of rosettes in a thyroid FNAC with CCV-PTC, which triggered the reason to undertake our study.<sup>12</sup> To the best of our knowledge, no prior large

series has corroborated a correlation of a specific PTC variant with a RLC pattern. A few authors previously described microfollicles as “rosettes” in cases of insular thyroid carcinoma and/or benign adenomatous lesions (14-18). However, their “rosette-like pattern” did not meet its strict definition.<sup>17</sup>

Since the first edition of TBSRTC, several new variants of thyroid carcinoma have been described.<sup>32-</sup>  
<sup>33</sup> The revised second edition of TBSRTC even included a detailed definition with particular criteria for some of these PTC variants.<sup>33</sup> However, there was no mention of a RLC pattern in any of these PTC variants. Analyzing published data from the literature, we found that very few authors mentioned the presence of rosettes and/or a pseudo-rosette architectural pattern in association with thyroid carcinoma<sup>31</sup>. Importantly, these patterns were not linked with specific diagnostic or prognostic features. Honambirajan et al, however, did emphasize that rosettes may superficially resemble the microfollicles of FVPTC and that their presence should thus raise suspicion for an aggressive PTC variant.<sup>13</sup>

According to TBSRTC, the recognition of PTC variants is generally unnecessary on FNAC given that precise subtyping is rarely possible or reliable due to difficulties in sampling the predominant pattern, due to the rarity of some PTC variants, and because of the morphological overlap with conventional PTC.<sup>32-33</sup> Nevertheless, the recognition of certain PTC variants (e.g. TCV, CCV-PTC, and hobnail variant) that manifest with an aggressive behavior may warrant recognizing them on FNAC because they often require more aggressive therapy. In a recent review article, Rossi et al discussed the morphological features that may be associated with PTC variants.<sup>31</sup> However, there were no details discussed regarding the relevance of rosettes in PTC variants.<sup>31</sup> By definition, variants of PTC must have some of the essential nuclear features of typical PTC including nuclear grooves, clearing and pseudoinclusions. In contrast to the relatively easy identification of TCV findings on FNAC, the recognition of CCV-PTC is more elusive since the typical features of PTC are not constantly found in this variant. TBSRTC defines CCV-PTC as a variant characterized by columnar

cells with hyperchromatic, oval and pseudostratified nuclei with supranuclear or subnuclear cytoplasmic vacuoles resembling colonic epithelium or secretory-type endometrium. With regard to the architectural patterns seen in CCV-PTC, TBSRTC mentions that neoplastic follicular cells may be arranged as papillae, clusters, flat sheets, and sometimes, small tubular structures but not RLC as observed by Sen and colleagues<sup>1</sup>. As reported by several authors, our LBC series confirms some morphological differences with CS. In particular, we confirmed a lack of unequivocal features of rosette clusters. However, we did appreciate that a few of the cases had a pseudorosette pattern.

Our series, involving a review of one year's worth of thyroid FNAC cases diagnosed as SFM or malignant, found that a few pseudorosette clusters were present in 14 TCV and 7 hobnail variant cases of PTC. Apart from the case report by Sen et al, we did not uncover articles mentioning an association with rosettes and either the TCV or hobnail variant. Such pseudo-rosette pattern may be attributed to peculiar morphological features in these variants (tall cells and columnar features of neoplastic cells), and as such when detected could suggest a possible aggressive variant. Concerning CCV-PTC, we did not find a obvious RLC pattern but did document a few pseudorosette clusters in 7 out of 9 cases, including four cases where there was a combination of TCV and CCV-PTC. Furthermore, a few pseudorosette clusters were also recognized in a minority of the other PTC variants including one solid variant of PTC, 2 FVPTC and 2 Warthin-like variant cases. None of the conventional cPTC cases had either a RLC or pseudorosette pattern identified, but did frequently contain papillary and microfollicular cell clusters. While ICC, especially HBME-1, demonstrated strong and diffuse positivity in all cases of PTC supporting this diagnosis, such staining did not show any correlation with RLC. For BRAF<sup>V600E</sup>, we found an association with the aggressive variants of PTC but no correlation with cytomorphological pattern. Compared to the literature, we found a low number of BRAF<sup>V600E</sup> mutated cases 47 (21% for SFM plus M) which might be attributed to the fact that cPTC and several of the PTC variants are frequently wild type BRAF<sup>V600E</sup>.

In conclusion, our analysis confirmed the presence of a RLC pattern in several PTC variants, and that such “rosettes” were not associated with cPTC cases. Our data shows that a pseudo-rosette pattern is possibly identify in the TCV, hobnail variant of PTC and CCV-PTC. Furthermore, none of our BL cases had a rosette or pseudorosette pattern, suggesting that their presence is most likely associated with malignant thyroid lesions. Another interesting point is the absence of pseudo-rosette clusters in histological samples, confirming that this architectural pattern is mostly seen in cytological samples. The detection of a pseudo-rosette architectural pattern combined with the typical cytological findings of classical nuclear and cellular features of conventional PTC might offer an additional feature in those cases of thyroid FNAC with an equivocal interpretation or scant neoplastic component.

**Acknowledgement** : EDR, LP, LMR GF contributed to the idea and preparation of the manuscript for the first draft and revisions; MDA, TM, VF, CB, CDM contributed to the data and revisions; MR, ET, CPL, GF contributed to the revisions of the manuscript.

**Data availability statement:** The Data used for this paper can be shared

## REFERENCES

- 1) Wippol FJ, Perry A. Neuropathology for neuroradiologist: rosette and pseudorosette. *Am J Neurorad* 2006; 27:488-92.
- 2) Rosette. Pathology discussion forum. Available at: <http://patho.medtut.net/2016/01/07/rosette>. Last accessed March 2016
- 3) Krafts K. The four main type rosettes in pathology. In pathology student. Available at: <http://www.pathologystudent.com/?p=5400>. Last accessed February 2016.
- 4) Das D, Bhattacharjee K, Barthkur SS, Tahiliuani PS; DeKa P, Bhattacharjee H et al. A new rosette in retinoblastoma. *Ind J ophtalm* 2014; 62: 636-41.
- 5) Ishida M, Iwai M, Yoshida K, Kazotani A, Okabe H. A distinct histopathological variant of a malignant melanoma with perivascular pseudorosettes: a case report. *Oncology Letters* 2013; 6: 673-75.
- 6) Aral A, Sasavama T, Tamaki M, Sakagami Y, Enoki E, Ohbavashi C et al. Rosette-forming glioneural tumor of fourth ventricle-case report. *Neurol Med Chir* 2010; 50: 224-28.
- 7) Ferri N, Martinez-Lage F, Almangro MJ et al. Embryonal tumor with abundant neuropil and true rosette (ETANTR): a new distinctive variety of pediatric PNET; a case based update. *Chil Nervous Syst* 2010; 26: 1003-8.
- 8) Harding MJ, Mcgrew HF, Nechiporuk A. The roles and regulation of multicentric rosette structures during morphogenesis. *Development* 2014; 141: 2549-58.
- 9) Miller K, Hall RC, Brenn T. Spitz nevus with homer wright rosette like structures. *Am J Dermpath* 2012; 34: 457-59.
- 10) Fukudome N, Kimura F, Arita S et al. Significance of cytological findings of neuroblastoma: Rosette arrangement and neuropil structure. *J Cytol Pathol* 2015 Aug 26. Available at : <http://www.omicssonline.org/open-access/significance-of-cytology>.



- 11) Desai SS, Sankar S Borges AM. A study of histopathological features of medullary carcinoma of the thyroid: cases from a single institute in India. *Indian J Cancer* 2005; 42: 25-29
- 12) Sen A, Nalwa A, Mathur SR, Jain D, Iyer VK. Cytomorphology of columnar cell variant of papillary carcinoma thyroid: A case report and review of the literature. *CytoJournal* 2014;11:27
- 13) Nambirajan A, Batool S, Mandal P et al. Rosettes in papillary carcinoma of the thyroid: a rare cytomorphological indicator of aggressive variants. *Cytopathol* 2017; 28: 560-62
- 14) Oertel YC, Miyahara-Felice L. Cytologic features of insular carcinoma of the thyroid: A case report. *Diagn Cytopathol* 2006; 34: 572-5
- 15) El Hg IA, Kollur SM. Benign follicular thyroid lesions versus follicular variant of papillary carcinoma: differentiation by architectural pattern. *Cytopathol* 2004; 15: 200-5
- 16) Mesonero CE, Jugle JE, Wilbur DC, Nayar R. Fine needle aspiration of the macrofollicular subtypes of the follicular variant of papillary carcinoma of the thyroid. *Cancer* 1998; 84: 235-44
- 17) Oertel YC. Fine needle aspiration in the evaluation of thyroid neoplasms. *Endocr Pathol* 1997; 8: 215-24
- 18) Gharib H, Goellner RJ, Johnson DA. Fine needle aspiration of the thyroid: a 12 years experience with 11,000 biopsies. *Clin Lab Med* 1993; 13: 699-709
- 19) British Thyroid Association, Royal College of Physicians. Guidelines for the management of thyroid cancer (Perros P, ed) 2<sup>nd</sup> edition. Report of the Thyroid Cancer Guidelines Update Group. London: Royal College of Physicians, 2007, Perros P 2<sup>nd</sup> edition
- 20) Fadda G, Basolo F, Bondi A, Bussolati G, Crescenzi A, Nappi O, et al. Cytological classification of thyroid nodules. Proposal of the SIAPEC-IAP Italian consensus working group. *Pathologica*. 2010;102:405-408

- 21) Nardi F, Basolo F, Crescenzi A, Fadda G, Frasoldati A, Orlandi F, et al. . Italian consensus for the classification and reporting of thyroid cytology. *J Endocrinol Invest*. 2014; 37:593–9
- 22) Rossi ED, Martini M, Capodimonti S et al. Analysis of immunocytochemical and molecular BRAF expression in thyroid carcinomas: a cyto-histological institutional experience. *Cancer Cytopathol* 2014; 122: 527-35
- 23) Rossi ED, Mule' A, Russo R et al. Application of liquid based preparation to non-gynecological exfoliative cytology. *Pathologica* 2008; 100: 461-5
- 24) Rossi ED, Raffaelli M, Minimo C et al. Immunocytochemical evaluation of thyroid neoplasms on thin-layer smears from fine-needle aspiration biopsies. *Cancer* 2005; 105: 87-95
- 25) Rossi ED, Martini M, Capodimonti S et al. Diagnostic and prognostic value of immunocytochemistry and BRAF mutation analysis on liquid based biopsies of thyroid neoplasms suspicious for carcinoma. *Eur J Endocrin* 2014; 168: 853-9
- 26) Rossi ED, Martini M, Capodimonti S et al. BRAF (v600e) mutation analysis on LBC-processed aspiration biopsies predicts bilaterality and nodal involvement in papillary thyroid microcarcinoma. *Cancer Cytopathol* 2013; 121: 291-7
- 27) Rossi ED, Martini M, Capodimonti S et al. Morphology combined with ancillary techniques: An algorithm approach for thyroid nodules. *Cytopathol* 2018; 29: 418-27
- 28) Rossi ED, Larocca LM, Pantanowitz L. Ancillary molecular testing of indeterminate thyroid nodules. *Cancer Cytopathol* 2018; 126: 654-71
- 29) American Joint Commission on Cancer (AJCC) Cancer Staging Atlas, 2017 8th Edition.
- 30) Lloyd R. V., Osamura R. Y., Kloppel G., Rosai J. (Eds): WHO Classification of Tumours of Endocrine Organs (4th edition). IARC: Lyon 2017

- 31) Rossi ED, Faquin WC, Pantanowitz L. Cytologic Features of Aggressive Variants of Follicular-Derived Thyroid Carcinoma. *Cancer Cytopathol* 2019; 127(7):432-446.
- 32) Ali S, Cibas ES. *The Bethesda system for reporting thyroid cytopathology*. 1st Edition, Springer, New York. 2010.
- 33) Ali S, Cibas ES. *The Bethesda system for reporting thyroid cytopathology*. 2<sup>nd</sup> Edition, Springer, Berlin. 2018.

**Table 1.** Comparison of cytomorphological patterns

Follicular structure	<b>Rosette pattern</b>	<b>Microfollicles</b>	<b>Papillary clusters</b>
<b>Definition</b>	<ul style="list-style-type: none"> <li>• Cells with circular arrangement</li> <li>• 15 or less cells</li> <li>• Cytoplasm merges toward center</li> <li>• Most of the cytoplasm is located within center (i.e. nuclei peripheral), implying the cells are tall/columnar</li> <li>• Central area can have a lumen (+/- filled with</li> </ul>	<ul style="list-style-type: none"> <li>• Cells with circular arrangement</li> <li>• 15 or less cells</li> <li>• Cytoplasm merges toward center</li> <li>• Minimal cytoplasm/cell causes less to be located within center</li> <li>• Cells shorter (cuboidal)</li> <li>• Central area usually has colloid</li> </ul>	<ul style="list-style-type: none"> <li>• Cells with more elongated arrangement</li> <li>• 15 or more cells</li> <li>• Cytoplasm need not merge toward center</li> <li>• Minimal cytoplasm/cell</li> <li>• Cells shorter (cuboidal)</li> <li>• Central</li> </ul>

	colloid)		fibrovascular core
--	----------	--	-----------------------

**Table 2.** Clinicopathological features of study cases

Parameter	Quantity (N=375)
Female	239 (63.7%)
Male	136 (36.2%)
Thyroid nodule size	6-65 mm
Cytological Diagnosis	BL=150 (40%) SFM=100 (26.6%) PM=125 (33.3%)
Histological Diagnosis (245 cases)	15 Goiter (6.1%) 5 FA (2%) cPTC=140 (48.9%)

	PTC variants=85(34.6%)
ICC*	
H+/G+	209 (92.8%)
H+/G-	16 (7.1%)
H-/G+	0
H-/G-	0
BRAF <sup>V600E</sup> *	47(21%)

BL= Benign Lesions; FA Follicular Adenoma; cPTC = classical papillary thyroid carcinoma; PM = positive for malignancy; PTC = papillary thyroid carcinoma SFM = suspicious for malignancy; ICC: Immunocytochemistry; H: HBME-1; G: Galectin-3.

- \*ICC and BRAF<sup>V600E</sup> only in SFM and M categories

**Table 3. Distribution of variants of PTC in the suspicious for malignancy and positive for malignancy cytological categories**

Histology	CCV-PTC	FVPTC	Hobnail-PTC	Classic PTC	Solid-PTC	TCV*	TCV+CCV	Warthin-like PTC
SFM	2	4	6	44	0	6 (14)	0	1
PM	3	4	7	96	3	13 (13)	4	5

CCV-PTC = columnar cell variant of papillary thyroid carcinoma; FVPTC = follicular variant of PTC; PM = positive for malignant; PTC = papillary thyroid carcinoma; SFM = suspicious for malignancy; TCV = tall cell variant of PTC

- Including pure TCV and in brackets , cases of PTC with TCV component less than 30%

**Table 4. Distribution of cases according to follicular architectural pattern based upon FNAC diagnoses**

Growth pattern	BL (N=150)	SFM (N=100)	PM (N=125)	SFM+PM (N=225)
Solid clusters	150 (100%)	100 (100%)	125 (100%)	225 (100%)
Papillary	0	88 (88%)	112 (89.6%)	200 (88.8%)
Microfollicular	0	56 (56%)	98 (78.4%)	154 (68.4%)
Isolated follicular cells	0	94 (94%)	107 (85.6%)	201 (89.3%)
<b>Pseudorosette- like clusters</b>	0	12 (12%)	21 (16.8%)	33 (14.6%)

PM: positive for malignancy; SFM: suspicious for malignancy; BL benign lesions

**Table 5. Distribution of cases according to follicular architectural pattern stratified by histological diagnoses**

Growth pattern	CCV-PTC (N=5)	FVPTC (N=8)	H-PTC (N=13)	S-PTC (N=3)	TCV (N=46)	TCV+CCV- PTC (N=4)	W-PTC (N=6)	cPTC (N=140)
Solid clusters	2 (40%)	8 (100%)	13 (100%)	3 (100%)	27 (58.6%)	4 (100%)	6 (100%)	140 (100%)
Papillary	2 (40%)	2 (25%)	8 (61.5%)	2 (66.6%)	29 (63%)	4 (100%)	6 (100%)	130 (92.8%)
Microfollicles	2 (40%)	3 (37.5%)	10 (76.9%)	2 (66.6%)	20 (43.4%)	4 (100%)	0 (%)	126 (90%)
Isolated cells	2 (40%)	8 (100%)	12 (92.3%)	3 (100%)	27 (58.6%)	4 (100%)	6 (100%)	140 (100%)
<b>Pseudorosette- like clusters</b>	3 (60%)	3 (25%)	7 (53.8%)	2 (33.3%)	14 (30.4%)	2 (50%)	2 (33.%)	0

CCV-PTC = columnar cell variant of papillary thyroid carcinoma; FVPTC: follicular variant of PTC;  
 PTC = papillary thyroid carcinoma; TCV = tall cell variant of PTC; W-PTC: Warthin-like PTC;  
 cPTC: Classic PTC; S-PTC: Solid variant PTC; H-PTC: Hobnail variant PTC;

**Table 6. Correlation between diagnostic categories and ancillary techniques**

Cytology diagnosis	H+	G+	BRAF <sup>V600E</sup>
SFM (100 cases)	100 (100%)	91 (91%)	10 (10%)
PM (125 cases)	125 (100%)	118 (94.4%)	37 (29.6%)

SFM = suspicious for malignancy; PM: Positive for malignancy; H: HBME-1; G: Galectin-3.





**Figure Legends**

**Figure 1 A, B and C:** Comparison papillary, rosette and microfollicular structures. Drawing of the different patterns in Figure 1 A) **Classic rosette structure**; Figure 1 B) **Cluster of follicular neoplastic cells**; Figure 1 C) **Papillary structure pap stain, 200x, liquid based cytology-LBC)**

**Figure 2 A and B and C:** Figure 2 A and B Follicular cells in a pseudorosette-like pattern from a case of tall cell variant of papillary thyroid carcinoma (pap stain, 200x, liquid based cytology-LBC); Figure 2 C details of another pseudorosette-like patten from the same case (pap stain, 400x, liquid based cytology-LBC)

**Figure 3 A and B:** Follicular cells in a pseudorosette-like pattern from a case of hobnail cell variant of papillary thyroid carcinoma (Pap stain, 400x liquid based cytology-LBC); Figure 3B details from the same case with a pseudorosette-like cluster in the left bottom and a neoplastic cluster in the upper right (Pap stain, 200x liquid based cytology-LBC)

Author Manuscript





Author Manuscript

This article is protected by


Author Manuscript



This article is protected by


Author Manuscript

This article is protected by




Author Manuscript

This article is protected by copyright. All rights reserved.



Author Manuscript



This article is protected by