

DR ALICIA B LICHVAR (Orcid ID : 0000-0003-3804-6517)

DR JEONG M. PARK (Orcid ID : 0000-0002-7961-494X)

Article type : TI on Transplantation

Full Title: Use of Direct-acting Oral Anticoagulants in Solid Organ Transplantation: A Systematic Review

Running Title: DOAC in SOT

Authors:

¹Alexandra L. Bixby; ²Alicia B. Lichvar, ³David Salerno, ⁴Jeong M. Park

Affiliations:

¹Department of Pharmacy Services, University of Hospitals Cleveland Medical Center, Cleveland, Ohio

²Department of Pharmacy Practice, University of Illinois at Chicago College of Pharmacy, Chicago, Illinois; Department of Surgery, University of Illinois at Chicago College of Medicine, Chicago, Illinois

³Department of Pharmacy, NewYork-Presbyterian Hospital, New York, New York

⁴Department of Clinical Pharmacy, College of Pharmacy, University of Michigan, Ann Arbor, Michigan; Department of Pharmacy Services, Michigan Medicine, Ann Arbor, Michigan

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/PHAR.2485](https://doi.org/10.1002/PHAR.2485)

This article is protected by copyright. All rights reserved

Corresponding Author:

Jeong M. Park
College of Pharmacy
University of Michigan
1111 Catherine St., VVH 302
Ann Arbor, MI 48109-2435
Email: jeongp@med.umich.edu
Phone: 734-647-4711
Fax: 734-615-2314

Conflict of Interest: The authors declare no conflicts of interest.

ABBREVIATIONS:

4F-PCC	4-factor prothrombin complex concentrate
AHA/ACC/HRS	American Heart Association/American College of Cardiology/Heart Rhythm Society
ANNEXA-4	The Andexanet Alfa, a Novel Antidote to the Anticoagulation Effects of Factor Xa Inhibitors
aPCC	activated prothrombin complex concentrate
aPTT	activated partial thromboplastin time
ATC	American Transplant Congress
AUC	area under the curve
BID	bis in die
BMI	body mass index
CI	confidence interval
C _{max}	maximum concentration
CNI	calcineurin inhibitor
CrCl	creatinine clearance
CYP	cytochrome P450
DDI	drug-drug interaction
DOAC	direct-acting oral anticoagulant
dTT	dilute thrombin time

ECA	ecarin chromogenic assay
ECT	ecarin clotting time
eGFR	estimated glomerular filtration rate
FDA	Food and Drug Administration
GMR	geometric mean ratio
HR	hazard ratio
INR	international normalized ratio
IQR	interquartile range
ISHLT	International Society for Heart & Lung Transplantation
ISTH	International Society on Thrombosis and Haemostasis
IV	intravenous
NVAF	non-valvular atrial fibrillation
OR	odds ratio
PAUSE	Perioperative Anticoagulation Use for Surgery Evaluation
P-gp	P-glycoprotein
PT	prothrombin time
RE-VERSE AD	RE-VERSal Effects of Idarucizumab on Active Dabigatran
ROTEM	rotational thromboelastometry
RR	relative risk
SOT	solid organ transplantation
TEG	thromboelastography
TT	thrombin time
VKA	vitamin K antagonist
VTE	venous thromboembolism

ABSTRACT

The use of direct-acting oral anticoagulants (DOACs) has increased secondary to the mounting evidence for comparable efficacy and potentially superior safety to vitamin K antagonists (VKAs) in the general population. However, insufficient data regarding DOAC use in solid organ transplant (SOT) recipients and numerous pharmacokinetic and pharmacodynamic considerations limit their use in this highly selected patient population. A systematic review of recent clinical evidence on the safety and efficacy of DOACs compared to VKAs in SOT recipients was conducted. Additional considerations including transplant-specific strategies for

DOAC reversal and common pharmacokinetic/pharmacodynamic concerns were also reviewed. Although current evidence is limited to single-center retrospective analyses, DOACs, especially apixaban, appear to be a safe and effective alternative to VKAs for SOT recipients with stable graft function and without drug-drug interactions. Reliable data on DOAC reversal at the time of transplant surgery are lacking and clinicians should consider idarucizumab, andexanet alfa, and other non-specific reversal agents on an individual patient basis. There is no evidence supporting deviations from the Food and Drug Administration labeling recommendations for DOAC dosing in the setting of drug-drug interactions, obesity and renal function especially in patients on hemodialysis.

KEY WORDS: direct acting anticoagulant, organ transplantation, apixaban, dabigatran, edoxaban, rivaroxaban

1 | INTRODUCTION

Since the Food and Drug Administration (FDA) approval of dabigatran in 2010, direct-acting oral anticoagulants (DOACs) have been rapidly adopted for treatment of venous thromboembolism (VTE) and stroke prevention in non-valvular atrial fibrillation (NVAF). (1, 2) Numerous studies have demonstrated comparable or superior efficacy of four DOACs in the aforementioned indications with a significantly lower risk of bleeding complications compared with warfarin. (3-10) While DOACs are the preferred anticoagulants over vitamin K antagonists (VKA) in the general population, the data examining safety and efficacy of DOACs in special patient populations are limited. In particular, solid organ transplant (SOT) recipients demonstrate unique pharmacokinetic considerations regarding renal and hepatic function as well as drug-drug interactions (DDIs).

Managing anticoagulation in patients receiving DOACs who are undergoing an unplanned transplant surgery or urgent allograft biopsy can also be challenging. (11) In more recent years, DOAC-specific reversal agents such as idarucizumab and andexanet alfa became available to manage life-threatening or uncontrolled bleeding from anticoagulation with DOACs. (12, 13) Idarucizumab can also be considered to reverse the anticoagulant effect of dabigatran prior to emergency surgery and urgent procedures; however, safety and efficacy of other reversal strategies have not been well established for this indication. (14)

To provide insight into the practice trends of DOAC use in pre-, peri- and post-transplant settings, a national survey was conducted among transplant pharmacists in the United States in 2019. (15) Out of the 115 transplant programs responding to the survey, only 43 programs (37.4%) allowed DOACs in patients on the transplant waitlist. There were heterogeneous approaches to perioperative management of DOACs. While DOAC use was permissive in the post-transplant setting by 94.3% of the responding programs, patient's renal function and concomitant DDIs were among major factors that influenced prescribing decisions. Pharmacist perceptions on bleeding risk of DOACs in the SOT population were split between similar (38.3%) vs. increased (33.0%) compared to the non-transplant population. The majority of survey respondents (64.3%) recognized that transplant-specific data on DOAC therapy are lacking.

To address the primary clinical questions identified from these survey results, this review will focus on: 1) whether DOACs are safer and/or more effective than VKAs in SOT recipients; 2) how DOACs should be reversed in the setting of SOT; and 3) how DOAC doses should be modified for special populations of SOT recipients.

2 | LITERATURE SEARCH AND REVIEW

In March 2020, a systematic search was conducted to identify studies evaluating the safety and efficacy of DOACs in adult (age >18 years old) SOT recipients including kidney, liver, pancreas, heart and lung. PubMed was searched for English-language, full-text articles using the following combination of search terms: (DOAC OR "direct acting anticoagulant" OR "apixaban" OR "dabigatran"[MeSH] OR "edoxaban" OR "rivaroxaban"[MeSH]) AND ("Organ Transplantation"[MeSH] OR "Transplantation"[MeSH] OR "kidney transplant" OR "liver transplant" OR "lung transplant" OR "heart transplant"). Articles were excluded if the studies were not conducted in SOT recipients or did not evaluate safety or efficacy outcomes of DOAC. Additional studies were identified by reviewing references of relevant articles and searching abstracts presented at the American Transplant Congress (ATC) and the International Society for Heart & Lung Transplantation (ISHLT) between 2010 and 2020. Search terms for abstracts included: DOAC, direct acting anticoagulant, apixaban, dabigatran, edoxaban, and rivaroxaban. When evaluating the safety and efficacy of DOAC, preference was given to studies employing VKA as control. Eligibility assessment was performed independently by two reviewers. Disagreements were resolved by consensus. The following data were extracted from the included studies: study design, organ type, DOAC agents, bleeding complications, and thromboembolic events.

This study design followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement. (16) All four investigators (AB, AL, DS, and JP) independently assessed and rated study quality following the Newcastle-Ottawa Scale (NOS; http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp) for cohort studies.

For the DOAC reversal strategies and pharmacokinetic/pharmacodynamics considerations, references were qualitatively evaluated for relevance to peri-transplant settings. When transplant-specific evidence was lacking, data were supplemented from the general population. Based on the available literature, expert recommendations were formulated on criteria for appropriate use of DOAC reversal agents prior to transplant surgery and DOAC dosing in special patient populations.

3 | SAFETY AND EFFICACY OF DOAC IN SOLID ORGAN TRANSPLANT

A total of 120 publications were identified through the systematic search between January 1, 2010 and May 31, 2020. After applying the inclusion and exclusion criteria, nine English-language full-text articles were included in the content review (Figure 1). Seventeen abstracts reporting DOAC use in SOT recipients were identified from the meeting archives of ATC and ISHLT and were also summarized for discussion.

3.1 | Single-arm Studies of DOACs

Prior to 2020, reports on the DOAC use among SOT recipients were limited to single-center, retrospective case series without control arms (Table 1). (17-42) Among the abstracts from ATC and ISHLT, the incidences of major bleeding (0% to 23%) and composite bleeding (0% to 37%) ranged widely, and the incidence of thrombotic events was also variable (0% to 21%).

Five full-text articles described the safety and/or efficacy outcomes of DOACs in various SOT recipients in greater detail. The first study by Ambrosi and colleagues was a case series of 11 heart transplant recipients taking rivaroxaban, of which one patient experienced severe bleeding. (21) This patient had creatinine clearance (CrCl) of 25-30 mL/min, leading to the authors to conclude that rivaroxaban use should be discouraged in those with moderate renal impairment. Of the 42 kidney transplant recipients reviewed by Bukari and colleagues, 7% experienced a major bleed. (22) Apixaban was the most frequently prescribed DOAC (69%). Interestingly, 53% of patients received a DOAC dose reduction despite only 26.2% of patients being older than 75 years and an average estimated glomerular filtration rate (eGFR) of

62.9±18.9 mL/min/1.73 m². Vanhove and colleagues examined the interaction between calcineurin inhibitors (CNIs) and DOACs in 39 transplant recipients (kidney, lung, and heart). (24) The authors reported 2 bleeding events during the average follow up of 34 months. Wannhoff and colleagues investigated the differences in rivaroxaban blood levels among 9 liver transplant recipients on CNIs. (25) Three of 5 patients on cyclosporine experienced a bleeding complication during a median follow up of 11 months. The most robust single-arm study was a retrospective single-center cohort study of 37 thoracic transplant recipients by Lichvar and colleagues. Bleeding occurred in 19% of patients (major bleeding in 3%) during a median follow up of 3 months. (23) Based on a center specific protocol, 60% of patients required a 50% DOAC dose reduction due to concomitant azole antifungal use.

Given the small sample sizes, heterogeneity in study designs, and lack of controls, conclusions could not be made from these single-arm cohort studies regarding the safety and efficacy of DOACs in SOT recipients.

3.2 | Comparative Studies of DOACs to Vitamin K Antagonists

In early 2020, four single-center retrospective cohort studies were published comparing DOACs to VKAs in SOT recipients. (17-20) Table 2 summarizes the study population, anticoagulants, and efficacy and safety outcomes of the studies. (17-20) The NOS quality assessments for each study were listed in the Supplemental Table. (17-20)

Liver Transplantation

Santeusanio and colleagues compared 27 liver transplant recipients on DOACs to 20 matched controls on warfarin from 2014 to 2018. (20) Patients were excluded with active malignancy, those that stopped anticoagulation prior to 2 months for reasons other than bleeding or thrombosis, and those without appropriate laboratory follow up. DOAC patients were matched with warfarin controls based on type of transplant, age, history of hepatocellular carcinoma, indication for anticoagulation, HAS-BLED score, timing of anticoagulation with regards to transplant, and duration of anticoagulation. The primary endpoint was the incidence of clinically relevant major or non-major bleeding, defined by the International Society on Thrombosis and Haemostasis (ISTH) criteria for bleeding.

Apixaban was the most frequently prescribed DOAC (55.6%), followed by dabigatran (25.9%) and rivaroxaban (18.5%). There were no significant differences in baseline characteristics. Less patients in the DOAC group experienced a clinically relevant bleeding event (15% vs 45%, p=0.01) over a similar follow up period (356 vs 434 days, p=0.23). No

major bleeding events occurred in patients on DOACs and 4 major bleeding events occurred in the warfarin group (2 requiring transfusion, 1 retroperitoneal bleed, and 1 acute subdural hematoma). There was no difference in the incidence of thrombotic events (20% vs 15%, $p=0.67$). Five patients received inappropriate DOAC dose reductions and 1 patient was prescribed a higher dose than recommended. However, all of the patients that experienced a bleeding event were on appropriate dosing. One patient that was on lower than recommended dose experienced a thrombotic event. When the authors performed a univariate analysis of this matched cohort, warfarin use and a baseline eGFR < 30 mL/min or total bilirubin \geq 3 mg/dL was associated with clinically relevant bleeding. The final multivariable logistic regression showed that warfarin use (odds ratio [OR] 6.9, 95% confidence interval [CI] 1.1-44.6) and baseline eGFR < 30 mL/min (OR 10.4, 95% CI 1.3-81.3) were associated with higher odds of clinically relevant bleeding. Based on these results, the authors concluded that DOAC use appeared relatively safe compared with warfarin in liver transplant recipients. (20)

Kidney Transplantation

Leon and colleagues compared kidney transplant recipients taking DOACs ($n=52$) to a control group taking warfarin or fluindione ($n=50$) between 2013 and 2018. (19) Patients were excluded if eGFR < 30 mL/min/1.73 m² or had a mechanical valve. The co-primary efficacy outcomes were arterial thromboembolic events and venous thromboembolic events. The primary safety outcome was any bleeding event. Major bleeding was defined using the ISTH definition. The only significant differences between baseline characteristics included a higher HAS-BLED score, a shorter time post-transplant, and a lower baseline hemoglobin in the warfarin group. Twenty-five patients (48%) were on reduced doses of DOAC.

Over a mean follow-up of 14 ± 13 months in the DOAC group vs 22 ± 20 months in the VKA group ($p=0.08$), there was no difference in VTE (0% vs 8%, $p=0.054$). However, fewer patients in the DOAC group experienced a bleeding event (13% vs 42%, $p=0.003$; hazard ratio [HR] 0.39, 95% CI 0.19-0.85). None of the bleeding events in the DOAC group were considered major. Of the 4 patients that were on a strong cytochrome P450 (CYP) 3A4 and/or P-glycoprotein (P-gp) inhibitor or inducer, 3 patients experienced a bleeding event. Three of the 36 (8.3%) patients on apixaban and 4 of the 15 (26.7%) patients on rivaroxaban experienced a bleeding event. Characteristics of patients that experienced bleeding and severity of bleeding events were only reported for patients in the DOAC group (7 non-major bleeding events: 1

gastrointestinal, 2 urogenital, 3 subcutaneous hematoma, and 1 hemoptysis). The authors concluded that DOACs appeared to be effective and safe anticoagulants in kidney transplant recipients with stable graft function. (19)

Bixby and colleagues compared kidney transplant recipients taking DOACs (n=99) to those taking warfarin (n=98) between 2011 and 2018. (17) Patients who required anticoagulation for mechanical valve, had inadequate follow up, multiorgan transplant, gastrointestinal bleeding in the last 6 months, anticoagulation for less than 30 days (unless stopped early due to bleed), or indication for anticoagulation other than VTE and NVAf were excluded. The primary outcome was incidence of major bleeding defined using the ISTH definition. Baseline characteristics were similar with the exception of year anticoagulation started, antiplatelet use, and baseline hemoglobin. Reduced doses of DOACs were utilized in 19% of patients.

Over a median follow-up of 11.0 (interquartile range [IQR] 5.3-18.5) months in the DOAC group vs 13.1 (IQR 5.0-36.8) months in the warfarin group (p=0.15), there was no difference in new onset stroke (1.0% vs 5.1%, p=0.13) or VTE (2.0% vs 3.1%, p=0.50). Over the follow up period, 9.1% of DOAC vs 21.4% of warfarin patients experienced a major bleed and 24.2% vs 34.7% experienced a composite bleeding event. When accounting for time, there was no statistical difference in major bleeding (Mantel-Cox p=0.15) or composite bleeding. After multivariable Cox regression, DOAC use was not associated with an increased risk of bleeding (HR 0.73, 95% CI 0.27-1.95). When stratified by agent, there was a lower incidence of major bleeding with apixaban compared to all other anticoagulants (6.7% vs 19.0%, p=0.027). Similarly, there was a lower incidence of major bleeding when apixaban was compared with warfarin (6.7% vs 21.4%, p=0.014). The authors concluded that further research is warranted to definitively determine whether DOACs are safe and effective alternatives to warfarin for kidney transplant recipients. (17)

Heart Transplantation

Henricksen and colleagues evaluated heart transplant recipients taking a DOAC (n=51) or warfarin (n=22) between January 2012 and July 2019. (18) Except for more patients in the warfarin group on dialysis (22.7% vs 3.9%, p=0.01), there were no significant differences in baseline characteristics. There was no difference in duration of follow-up between the groups (DOAC 130 days [IQR 84.5-203] vs warfarin 93.5 days [IQR 59-183], p=0.15). Two recurrent VTE events occurred in the DOAC group (1 progressive upper extremity VTE and 1 fatal pulmonary embolism) and both patients were taking reduced dose apixaban due to concomitant

itraconazole use. Bleeding occurred in 23% of warfarin patients and 10% of DOAC patients ($p=0.08$). From both groups, there were 3 gastrointestinal bleeds, 3 intracranial bleeds, 2 abdominal hematomas, and 2 patients with epistaxis. In their Cox regression model, female sex was the only variable predictive of bleeding (HR 6.7, 95% CI 1.8-25.2). DOAC use was not associated with bleeding (HR 0.34, 95% CI 0.10-1.20). However, DOAC use was associated with a lower rate of bleeding requiring transfusion ($p=0.04$). The authors of this study concluded that anticoagulation with a DOAC demonstrated a trend toward a lower rate of bleeding when compared to warfarin, but a larger cohort is needed to validate these findings. (18)

3.3 | Clinical Implications

In this review, four studies were discussed comparing the safety and efficacy of DOACs to VKAs among SOT recipients. (17-20) There was no significant difference in efficacy with regards to VTE or stroke in any of the studies. Three of the four studies used the ISTH definitions to report bleeding outcomes. Consistently, all four studies found a numerically lower incidence of bleeding with DOACs than that with warfarin. Higher rates of bleeding were observed with both DOACs and warfarin in SOT recipients compared to the general population. (3-10)

Apixaban was the most commonly prescribed DOAC which is consistent with the national survey of transplant center practices. (15) Henricksen and colleagues did not find a significant difference in bleeding events between warfarin and apixaban in heart transplant recipients ($p=0.09$). (18) However, Bixby and colleagues found a lower incidence of major bleeding with apixaban compared to any other oral anticoagulants in kidney transplant recipients ($p=0.027$). (17) Similarly, a recent abstract comparing various DOACs in SOT recipients reported that apixaban was associated with less bleeding than dabigatran and rivaroxaban (12.5% vs 33.3%, $p=0.017$). (41)

Differences in dosing regimens may lead to differences in the bleeding incidence seen among different transplant centers. Santeusano and colleagues reported that 18.5% of DOAC patients were on doses not compliant with the prescribing guidelines. (20) Leon and colleagues reported that 48% of DOAC users were on reduced dose but did not report whether these dose adjustments were appropriate. (19) Bixby and colleagues reported that 19% of DOAC patients were on a reduced dose and 13% of patients were inappropriately on a reduced dose of a DOAC. (17) Although Henricksen and colleagues did not report the number of patients requiring dose adjustments, they did describe that dose adjustments were made according to renal function, weight, age, and DDIs. (18) Inconsistent practices in DOAC dose reduction is a major

limitation of the current data. In the national survey of transplant centers, 51.5% of respondents answered that they follow the recommended dosing in the prescribing information. (15) Of those who deviated, 56.3% reduced the DOAC dose even with only one concomitant DDI. Until more consistent dose reduction criteria are utilized in transplant recipients, comparing bleeding rates between institutions will remain a challenge.

Additionally, combining patients taking anticoagulation for various indications is a limitation in the present studies. However, Bixby and colleagues included DOAC use for atrial fibrillation into their multivariable cox regression and found that it did not increase risk for major bleeding (HR 2.11, 95% CI 0.94-4.11, $p=0.071$). (17) Based on the current data, DOACs, particularly apixaban, appear to be a safe and effective alternative to warfarin for transplant recipients with stable graft function and without DDIs.

4 | PERIPROCEDURAL MANAGEMENT AND REVERSAL OF DOAC THERAPY

The periprocedural management of DOACs and reversal strategies remain controversial and heterogeneous across the transplant landscape. In the recent survey of transplant programs, 37.4% of respondents allow patients to be maintained on DOAC therapy while they are on the transplant waitlist. (15) The varied nature of DOAC utilization during the pre-transplant period drives the need to understand the clinical management, monitoring, and reversal of DOACs in the setting of surgical procedures.

4.1 | Clinical Monitoring of DOACs

DOACs have variable effects on coagulation assays, and there are different clinical implications of these effects (Table 3). (43, 44) Despite the availability of laboratory assessments, only 29.4% of transplant centers report using routine laboratory monitoring to assess safety prior to transplant surgery in the setting of a DOAC therapy. (15) Limitations behind their limited use is delayed turnaround time, lack of individual agent calibration, and lack of quantitative measures that clinically relate to anticoagulation expectations and bleeding in the setting of surgery.

Dabigatran, a direct thrombin inhibitor, can alter activated partial thromboplastin time (aPTT), thrombin time (TT), dilute thrombin time (dTT), ecarin clotting time (ECT), prothrombin time (PT) and international normalized ratio (INR). (43) Out of these markers, dTT and ECT, and the plasma drug concentration itself are all thought to be sensitive markers that have some quantitative utility. The dTT is a coagulation test that measures coagulation by diluting plasma to better measure the activity of thrombin and has a linear dose relationship with dabigatran. (44)

The dTT can indicate anticoagulation intensity, and in the setting of a normal value, is indicative of no clinically relevant anticoagulation effect from dabigatran. The ECT and plasma drug concentration, while sensitive, are infrequently available in clinical practice.

Factor Xa inhibitors can alter PT, aPTT (with the exception of apixaban) and anti-factor Xa levels. (43, 44) The PT is not sensitive enough to exclude clinically relevant anticoagulation effect; furthermore, it is less sensitive to apixaban compared to edoxaban and rivaroxaban. (43) Secondly, aPTT is less sensitive than PT and does not have a role in factor Xa inhibitor clinical monitoring, as a normal aPTT does not exclude on-therapy levels. After specific agent calibration, anti-factor Xa levels can be used for quantitative assessment and are the best monitoring parameter for this class of DOACs. Anti-factor Xa levels can be utilized to assess on-therapy or supratherapeutic levels if this calibration is performed (Table 4). (43, 45)

Thromboelastography (TEG) and rotational thromboelastometry (ROTEM) are visual assessments of clot formation, kinetics, and strength and have been assessed in the setting DOAC therapy. (46) With apixaban and dabigatran, the kaolin test reaction time (R time) and the time to maximum rate of thrombus generation were prolonged versus control samples. The utility of this laboratory monitoring parameter rests in its ability to guide intraoperative transfusion in the transplant surgery setting. (47)

4.2 | Perioperative Management of DOACs

Transplant surgery and allograft biopsies are generally categorized as high bleed risk procedures that require specific management in the setting of DOAC therapy. (14) In the setting of living donor transplantation or scheduled allograft biopsy, holding DOAC therapy is the most realistic and feasible clinical choice. The American College of Cardiology recommendations stratify duration of DOAC interruption based on the bleeding risk of the procedure and also CrCl as many DOACs have some degree of renal elimination (Table 5). (14, 45)

Furthermore, the Perioperative Anticoagulation Use for Surgery Evaluation (PAUSE) cohort study analyzed 3007 patients with atrial fibrillation on apixaban, dabigatran, and rivaroxaban. (48) In this particular assessment, perioperative DOAC interruption strategy was based on procedure-related bleeding risk and also CrCl. The 30-day postoperative rate of major bleeding was 1.35% (95% CI, 0%-2.00%) in the apixaban cohort, 0.90% (95% CI, 0%-1.73%) in the dabigatran cohort, and 1.85% (95% CI, 0%-2.65%) in the rivaroxaban cohort. (48) This study demonstrates that procedural pauses in DOAC therapy can result in low incidences of bleeding in the setting of elective procedures.

4.3 | Reversal of DOAC Therapy

A major limitation to widespread DOAC utilization was a lack of specific reversal therapy, although specific agents are now available in the United States. In the recent survey, only 7.8% of transplant programs reported routine DOAC reversal at the time of transplant surgery, and of those programs using DOAC reversal, idarucizumab was most common for dabigatran reversal and 4-factor prothrombin complex concentrate (4F-PCC) for oral factor Xa inhibitors. (15)

Non-specific DOAC reversal agents can be used but are sometimes impractical in the setting of unexpected procedures and are used more so in the setting of acute ingestion or overdose. These modalities include activated charcoal, high-flow hemodialysis modalities, and plasmapheresis. (49) Both 4F-PCC and activated prothrombin complex concentrate (aPCC) can also be used off-label for DOAC reversal, despite not being specific reversal therapy agents.

Reversal of Direct Thrombin Inhibitor (Dabigatran)

Dabigatran reversal is routinely performed with idarucizumab in clinical practice. Idarucizumab is a human monoclonal antibody fragment that binds to dabigatran and its metabolites, neutralizing their anticoagulation effect (Table 6). (50-53) In the RE-VERSE AD trial, 213 patients were analyzed and 68% had cessation of bleeding within 24 hours with a median time to hemostasis of 2.5 hours. (12) A sub-analysis of surgical patients demonstrated that complete reversal of dabigatran effect occurred rapidly in 91% of the cohort, with a median time of drug administration to surgery to be 2 hours in almost all groups. (54)

The use of idarucizumab in the setting of transplant recipients has limited data including case reports and small cohort experiences. There have been several case reports using this reversal modality in the setting of transplant surgery with success. (55-61) In a series of 10 patients receiving idarucizumab in the setting of heart transplantation, 2 patients (20%) required re-intervention because of bleeding which was similar to a historical control group. (60) In a multicenter observational study using idarucizumab reversal for 53 heart transplant recipients, 7.5% required re-operation in the immediate postoperative period to control bleeding, 66% required blood product transfusion, and 92.4% demonstrated 30-day survival. (59) One patient had surgical problems with right ventricular failure resulting in a death, which was the only mortality associated with bleeding. In the setting of inadequate reversal and suboptimal hemostasis, there have been reports of repeat idarucizumab dosing. (62) Per the prescribing information, in the setting of clinically relevant bleeding together with elevated coagulation parameters, then an additional dose of idarucizumab 5 g may be considered. (52)

In a prospective cohort study assessing the effect of aPCC for dabigatran reversal in the setting of major bleeding, hemostasis was determined to be “good” in 9 (64%), “moderate” in 5 (36%) and “poor” in none with no thrombotic events. (63) However, there is no data for this reversal strategy in the setting of urgent surgery in the absence of bleeding, and much of this use is extrapolated from data of *in vitro*, animal, healthy volunteer, and bleeding patients. (64) Likewise, there is no peer-reviewed published data for this agent in a transplant population.

Currently, guidance from the Anticoagulation Forum and 2020 American College of Cardiology (ACC) Expert Consensus Decision Pathway on Management of Bleeding in Patients on Oral Anticoagulants recommends using idarucizumab 5 g intravenous (IV) for first-line dabigatran reversal with aPCC 50 units/kg IV as an alternative therapy. (45, 64) Based on currently available literature, in the setting of the need for DOAC reversal in transplant recipients, idarucizumab should be first-line with aPCC as a secondary substitute for reversal. At this time, there is no specific data to guide whether repeat dosing is more effective than dosing aPCC, but given the achievement of hemostasis with primary idarucizumab dosing, this should be used before aPCC if there is enough supply to accommodate additional doses.

Reversal of Oral Factor Xa Inhibitors (Apixaban, Edoxaban, and Rivaroxaban)

The approach to oral factor Xa reversal is more heterogeneous compared to dabigatran. Specific reversal can be utilized with andexanet alfa, a recombinant variant of human factor Xa which binds to oral factor Xa inhibitors and inhibits the activity of tissue factor pathway inhibitor (Table 6). The Andexanet Alfa, a Novel Antidote to the Anticoagulation Effects of Factor Xa Inhibitors (ANNEXA-4) Study analyzed 254 patients (100 on rivaroxaban, 134 on apixaban, 16 on edoxaban, and 4 on enoxaparin) and 352 for safety outcomes (128 on rivaroxaban, 194 on apixaban, 20 on edoxaban, and 3 on enoxaparin). (13) There was a 92% reduction in anti-factor Xa activity after the andexanet alfa bolus with excellent or good hemostasis occurring in 82% of patients on apixaban and 80% of rivaroxaban treated patients. A total of 14% died and 10% had thrombotic events within 30 days. Currently, no data exists for the use of andexanet alfa in the setting of urgent surgical procedural reversal or in the setting of transplant recipients.

Another option for DOAC reversal is 4F-PCC. In a study by Majeed and colleagues, 84 patients received 4F-PCCs to reverse major bleeds from rivaroxaban and apixaban. (65) Patients weighing < 65 kg received 1500 units, and those weighing > 65 kg received 2000 units. A total of 69% of patients achieved hemostasis. Furthermore, a prospective cohort study in Canada analyzed 66 patients who presented with major bleeds on rivaroxaban and apixaban and received 4F-PCC reversal. (66) In total hemostasis was categorized as “good” in 65%,

“moderate” in 20%, and “poor” in 15%. In *post-hoc* analysis, reversal was effective in 68% of patients according to the ISTH criteria. A total of 8% of patients had thrombotic events within 30 days of their bleed reversal. The utilization of PCCs at the time of transplant surgery is limited to case reports. (67)

Guidance from the Anticoagulation Forum and 2020 ACC Expert Consensus Decision Pathway on Management of Bleeding in Patients on Oral Anticoagulants recommends using andexanet alfa for first-line oral factor Xa reversal with 4F-PCC utilization (2000 units fixed dose) as an alternative therapy. (45, 64) Utilization of andexanet alfa in the setting of transplant reversal should be carefully considered and perhaps avoided based on the current paucity of literature and potential thrombosis risk. Figure 2 illustrates an overview of DOAC reversal in the setting of bleeding and urgent procedures.

5 | TRANSPLANT-SPECIFIC CONSIDERATIONS FOR DOAC USE

Pharmacokinetic/pharmacodynamic properties of individual DOAC agents that are pertinent to the SOT population have been examined in a previous comprehensive review. (11) Since the publication, there have been significant additions to the literature with regard to DDIs with common transplant medications, concomitant use with antiplatelet agents, and pharmacokinetics of DOACs in other specialized populations including obesity and hemodialysis. Approaches to DOAC dosing in these specific conditions will be discussed here.

5.1 | Drug Interactions between DOACs and Calcineurin Inhibitors

Cyclosporine and tacrolimus are known substrates of CYP3A4 and P-gp. Both are inhibitors of P-gp, and cyclosporine is a moderate inhibitor of CYP3A4. (68) Both rivaroxaban and apixaban undergo metabolism via CYP3A4 and all of the DOACs are substrates of P-gp. (11) As such, CNIs and DOACs have the potential of either being the perpetrator or target drug in a DDI. Although the FDA labeling of DOACs specify contraindications and dosing recommendations in concomitant use with a strong CYP3A4 and/or P-gp inhibitor, pharmacokinetic interactions with moderate CYP3A4 and/or P-gp inhibitors and the associated risk of bleeding are not well defined. (69, 70)

The potential DDI between rivaroxaban and CNIs has been reported by three different groups in heterogenous patient populations. Each case series noted that rivaroxaban may accumulate in the setting of CNI use but other confounders such as decreases in renal function may also play a role. (19, 21, 25) Bashir and colleagues investigated the DDIs between CNIs and apixaban in healthy volunteers. (71) Co-administration of 100 mg of cyclosporine resulted in

an increase in the apixaban C_{max} by the geometric mean ratio (GMR) of 1.43 [90% CI 1.12-1.83] but not in the apixaban area of the curve (AUC) (GMR 1.20, 90% CI 0.97-1.48). Co-administration with tacrolimus 5 mg resulted in reduced apixaban AUC (GMR 0.78, 90%CI 0.63-0.97) but no change in C_{max} (GMR 0.87, 90% CI 0.69-1.12). Although this study was unable to comment on the mechanism for the above variability, the magnitude of differences in AUC and C_{max} are not clinically significant. Parasrampur and colleagues, studied healthy subjects receiving a single oral dose of 60 mg edoxaban with or without cyclosporine 500 mg as a single dose. (72) Co-administration with cyclosporine increased the edoxaban AUC by 73% and the C_{max} by 74%. In each of these cases, the relative changes in drug exposure were thought not to be clinically significant to warrant a dose adjustment. Even though cyclosporine has been implicated as having the greater potential for a DDI with the DOACs, registry data from the Taiwan National Health Insurance database did not find an association with increased risk of bleeding in patients receiving cyclosporine (n=567) versus those without. (69)

Vanhove and colleagues investigated whether DOACs can be considered the perpetrator drug and CNIs the victim drug in a single-center retrospective analysis including 39 organ recipients (18 kidney, 13 lung, and 5 heart) treated with the combination of a CNI and rivaroxaban (n=29) or apixaban (n=10). (24) Rivaroxaban may cause a clinically insignificant (<20%) increase in CNI trough concentration, making it practical to repeat the CNI trough concentration measurement 5-7 days after DOAC initiation in select patients. Santeusano and colleagues reported no significant difference in the mean CNI trough/dose ratio in liver transplant recipients at 30 days following DOAC initiation regardless of tacrolimus (p=0.53) or cyclosporine use (p=0.50). (20) Further, Bukhari and colleagues reported no changes in serum tacrolimus levels three days after the initiation of DOACs among kidney transplant recipients treated with tacrolimus (pre- and post-DOAC serum tacrolimus concentration was 7.35 and 7.80 ng/mL, p=0.55). (22)

5.2 | Concomitant Use of Azole Antifungals

Azole antifungals are routinely used in SOT recipients for fungal prophylaxis and are considered either moderate inhibitors (fluconazole) or strong inhibitors (itraconazole, posaconazole, and voriconazole) of CYP3A4. (68) Currently, no controlled studies with azole antifungals exist to assess the risk of bleeding in patients concomitantly receiving DOACs and there was no uniform clinical approach to managing these interactions in the studies reviewed herein (17-25). Chang and colleagues, in a large retrospective cohort study found that fluconazole increased the risk of major bleeding in patients concomitantly receiving a DOAC

(dabigatran, rivaroxaban, and apixaban). (69) Given the paucity of data, no empiric adjustment to DOAC dose can be recommended for concomitant use of an azole antifungal alone in patients with preserved renal function. However, the cumulative impact of this potential DDI in a patient with impaired renal function warrants individualized dose considerations. The apixaban prescribing information recommends a 50% dose decrease when coadministered with drugs that are strong dual-inhibitors of CYP3A4 and P-gp including ketoconazole and itraconazole, but it is unclear if this recommendation extends to posaconazole, voriconazole and isavuconazole. (73) Drug-specific monitoring may be beneficial in patients with DDI and/or impaired clearance.

5.3 | Concomitant Use of Antiplatelet Agents

Mounting data from large registries have revealed an increased risk of bleeding in patients concomitantly receiving DOACs and an add-on antiplatelet agent versus those without. (70, 74) Although, a meta-analysis of the DOAC NVAF trials and registry data in patients with anticoagulation and concomitant antiplatelet agents suggests the use of DOACs with aspirin has less bleeding than VKA plus the addition of aspirin. (75, 76) Further, randomized data from patients with NVAF who underwent percutaneous coronary intervention demonstrates that the use of DOACs and P2Y12 inhibitors together has a similar efficacy and potentially reduced risk of bleeding comparatively to warfarin plus P2Y12 inhibitors. (76-79)

It should be noted that in all of the peer-reviewed literature of DOAC use post-transplant there is a significant amount of empiric dose-adjustments being made (6-100%). (17) This appears to be based on a variety of factors including potential pharmacokinetic and pharmacodynamic interactions, renal and hepatic dysfunction, and other clinical indicators of bleeding risk. In the absence of controlled clinical trials, empiric dose-adjustments for presumed pharmacokinetic or pharmacodynamic interactions is not recommended. Rather, DOAC prescribing in SOT recipients should be congruent with the FDA labeling recommendations.

5.4 | DOAC Dosing in Obese Patients

Obesity in SOT recipients is common, as many patients with end-stage organ disease have a component of the metabolic syndrome. No large randomized controlled trial exists investigating the efficacy and safety of the DOACs in obese or morbidly obese patients. Therefore, clinicians are left to make treatment decisions from discordant pharmacokinetic/pharmacodynamic studies, retrospective studies, *post-hoc* analyses of phase III trials, and meta-analyses. (80-92)

Given that routine monitoring of DOAC concentrations is difficult to interpret and may be subject to variability, DOACs are not recommended in obese patients defined as a body mass index (BMI) ≥ 40 kg/m² and weight ≥ 120 kg. (93) In-depth considerations in morbidly obese patients and the data since the Scientific and Standardization Committee (SSC)/ISTH recommendation have been reviewed previously in this Journal. (94) Since this publication, there have been two additional meta-analyses assessing the rate of stroke or systemic embolism (SSE) and major bleeding in NVAf patients with obesity and morbid obesity. (92)

Zhou and colleagues, stratified NVAf patients based on BMI category and whether patients received a DOAC or warfarin and found no difference in SSE (relative risk [RR] 0.87, 95% CI 0.73-1.04) or major bleeding (RR 0.90, 95% CI 0.81-1.01) in patients receiving a DOAC or warfarin with BMI category ≥ 30 kg/m². (92) Kido and colleagues, specifically focused the meta-analyses on patients with morbid obesity (BMI > 40 kg/m² or weight > 120 kg) and found no difference between DOACs and warfarin for SSE (OR 0.85, 95% CI 0.60-1.19), but less major bleeding (OR 0.63, 95% CI 0.43-0.94) in patients receiving DOACs. Of note, there were no differences in major bleeding between those who received apixaban versus rivaroxaban. (95) Given the lack of pharmacokinetic/pharmacodynamic data in patients who are both transplant recipients and obese, the use of DOACs in obese SOT patients should be used with caution.

5.5 | DOAC Dosing in Hemodialysis Patients

The 2019 update to the American Heart Association (AHA)/ACC/Heart Rhythm Society (HRS) 2014 guideline suggests the use of warfarin or apixaban for patients with atrial fibrillation on dialysis may be reasonable if a CHA₂DS₂-VASc score is ≥ 2 in men or ≥ 3 in women. (96) Despite patients on hemodialysis being excluded from registration trials, the use of DOACs in patients on hemodialysis has been attempted. With the exception of apixaban, the FDA labeling for each DOAC recommends against use in patients on hemodialysis. (97)

Both a 10 mg and 15 mg dose of rivaroxaban has been evaluated in single-dose studies in hemodialysis patients, but meta-analyses suggest that there are higher rates of bleeding when rivaroxaban is used for NVAf patients on hemodialysis. (98-102) Apixaban on the other hand, undergoes significantly less renal clearance and has pharmacokinetic data to support its use in hemodialysis. (103) Additionally since the publication of the guideline two additional meta-analyses support the use of apixaban in hemodialysis. (101, 102) Unfortunately, the much anticipated RENAL-AF trial that randomized hemodialysis patients to either apixaban 5 mg twice a day (BID) or warfarin was stopped early due to slow enrollment. (104) While the available data

point at a relative benefit in safety of apixaban versus warfarin in patients on hemodialysis, a definitive clinical trial is lacking to determine the best anticoagulant in this patient population awaiting transplant. Moreover, heterogeneity exists between the data supporting apixaban 2.5 mg BID versus 5 mg BID for patients on hemodialysis without other reasons for a dose-adjustment. Pharmacokinetic data from patients taking apixaban 2.5 mg BID on hemodialysis demonstrated comparable drug exposure to that of the 5 mg BID dose in patients with preserved renal function. Further, after a washout period, the same patients received apixaban 5 mg BID and a further increase was noted above the 90th percentile compared with the reference value generated in patients with preserved renal function. (105) This is in contrast to clinical data from retrospective cohort study exploring the United States Renal Data System that analyzed hemodialysis patients taking either apixaban or warfarin for NVAf. In a sensitivity analysis, the 5 mg BID dose was associated with significantly lower risks of stroke or systemic embolism and death as compared to either the reduced dose of 2.5 mg BID or warfarin. (103) Recently, a meta-analysis of observational studies compared DOACs versus warfarin versus no anticoagulation and found no benefit of either DOACs or warfarin versus no anticoagulation for NVAf patients on hemodialysis. However, in the sensitivity analysis, the 5 mg BID dose had no difference in major bleeding and a reduction in mortality when compared to the 2.5 mg BID dose. (101)

As most patients on hemodialysis with atrial fibrillation will have an elevated CHA₂DS₂-VASc score, it is reasonable to follow the FDA labeling recommendation of apixaban 5 mg BID, as this has shown benefit in available registry data and a meta-analysis studies. (97, 101, 103) However, each patient's risk of bleeding and clotting should be scrutinized prior to choosing either 2.5 mg or 5 mg BID. Drug-specific monitoring may be beneficial to avoid complications of supra-therapeutic anticoagulation.

6 | CONCLUSION

Given the absence of prospective, randomized, controlled trials investigating safety and efficacy of DOACs in SOT recipients, this review highlights four important retrospective observational studies comparing DOACs to VKAs in SOT recipients. In these studies, the incidence of bleeding with DOACs was similar, if not lower, to that of VKAs. Based on these findings, DOACs, particularly apixaban, are safe and effective alternatives to VKAs. In the studies discussed, DOAC doses were often reduced based on clinical factors. This review discusses pharmacokinetic considerations for DOACs relating to dose adjustments, DDIs, weight, and renal function. Given that none of these factors alone warrant DOAC dose

adjustments, FDA labeling for DOAC dose adjustments should be followed. In cases where multiple factors are present that may increase DOAC concentrations, clinical discretion should be utilized. Although novel agents for DOAC reversal in the setting of life-threatening bleeding have emerged, DOAC use in the pre-transplant setting remains controversial given the absence of a standardized approach for periprocedural DOAC reversal. Utilization of andexanet alfa in the setting of pre-transplant reversal should be carefully considered on an individual patient basis and a universal use of reversal agents in this setting cannot be recommended based on the current paucity of literature and potential thrombosis risk. Clinicians must consider the phase of transplant, DDIs, weight, and renal function in order to determine the optimal anticoagulant for a transplant candidate or recipient.

REFERENCES

1. Lutsey PL, Walker RF, MacLehose RF, Alonso A, Adam TJ, Zakai NA. Direct oral anticoagulants and warfarin for venous thromboembolism treatment: Trends from 2012 to 2017. *Res Pract Thromb Haemost.* 2019;3(4):668-73. Epub 2019/10/19. doi: 10.1002/rth2.12222. PubMed PMID: 31624786; PubMed Central PMCID: PMC6782014.
2. Zhu J, Alexander GC, Nazarian S, Segal JB, Wu AW. Trends and Variation in Oral Anticoagulant Choice in Patients with Atrial Fibrillation, 2010-2017. *Pharmacotherapy.* 2018;38(9):907-20. Epub 2018/06/20. doi: 10.1002/phar.2158. PubMed PMID: 29920705; PubMed Central PMCID: PMC6448138.
3. Connolly SJ, Ezekowitz MD, Yusuf S, Eikelboom J, Oldgren J, Parekh A, et al. Dabigatran versus warfarin in patients with atrial fibrillation. *N Engl J Med.* 2009;361(12):1139-51. Epub 2009/09/01. doi: 10.1056/NEJMoa0905561. PubMed PMID: 19717844.
4. Schulman S, Kearon C, Kakkar AK, Mismetti P, Schellong S, Eriksson H, et al. Dabigatran versus warfarin in the treatment of acute venous thromboembolism. *N Engl J Med.* 2009;361(24):2342-52. Epub 2009/12/08. doi: 10.1056/NEJMoa0906598. PubMed PMID: 19966341.
5. Bauersachs R, Berkowitz SD, Brenner B, Buller HR, Decousus H, Gallus AS, et al. Oral rivaroxaban for symptomatic venous thromboembolism. *N Engl J Med.* 2010;363(26):2499-510. Epub 2010/12/07. doi: 10.1056/NEJMoa1007903. PubMed PMID: 21128814.

6. Granger CB, Alexander JH, McMurray JJ, Lopes RD, Hylek EM, Hanna M, et al. Apixaban versus warfarin in patients with atrial fibrillation. *N Engl J Med*. 2011;365(11):981-92. Epub 2011/08/30. doi: 10.1056/NEJMoa1107039. PubMed PMID: 21870978.
7. Patel MR, Mahaffey KW, Garg J, Pan G, Singer DE, Hacke W, et al. Rivaroxaban versus warfarin in nonvalvular atrial fibrillation. *N Engl J Med*. 2011;365(10):883-91. Epub 2011/08/13. doi: 10.1056/NEJMoa1009638. PubMed PMID: 21830957.
8. Agnelli G, Buller HR, Cohen A, Curto M, Gallus AS, Johnson M, et al. Oral apixaban for the treatment of acute venous thromboembolism. *N Engl J Med*. 2013;369(9):799-808. Epub 2013/07/03. doi: 10.1056/NEJMoa1302507. PubMed PMID: 23808982.
9. Buller HR, Decousus H, Grosso MA, Mercuri M, Middeldorp S, Prins MH, et al. Edoxaban versus warfarin for the treatment of symptomatic venous thromboembolism. *N Engl J Med*. 2013;369(15):1406-15. Epub 2013/09/03. doi: 10.1056/NEJMoa1306638. PubMed PMID: 23991658.
10. Giugliano RP, Ruff CT, Braunwald E, Murphy SA, Wiviott SD, Halperin JL, et al. Edoxaban versus warfarin in patients with atrial fibrillation. *N Engl J Med*. 2013;369(22):2093-104. Epub 2013/11/21. doi: 10.1056/NEJMoa1310907. PubMed PMID: 24251359.
11. Salerno DM, Tsapepas D, Papachristos A, Chang JH, Martin S, Hardy MA, et al. Direct oral anticoagulant considerations in solid organ transplantation: A review. *Clin Transplant*. 2017;31(1). Epub 2016/11/20. doi: 10.1111/ctr.12873. PubMed PMID: 27859621; PubMed Central PMCID: PMC5490493.
12. Pollack CV, Jr., Reilly PA, van Ryn J, Eikelboom JW, Glund S, Bernstein RA, et al. Idarucizumab for Dabigatran Reversal - Full Cohort Analysis. *N Engl J Med*. 2017;377(5):431-41. Epub 2017/07/12. doi: 10.1056/NEJMoa1707278. PubMed PMID: 28693366.
13. Connolly SJ, Crowther M, Eikelboom JW, Gibson CM, Curnutte JT, Lawrence JH, et al. Full Study Report of Andexanet Alfa for Bleeding Associated with Factor Xa Inhibitors. *N Engl J Med*. 2019;380(14):1326-35. Epub 2019/02/08. doi: 10.1056/NEJMoa1814051. PubMed PMID: 30730782; PubMed Central PMCID: PMC6699827.
14. Doherty JU, Gluckman TJ, Hucker WJ, Januzzi JL, Jr., Ortel TL, Saxonhouse SJ, et al. 2017 ACC Expert Consensus Decision Pathway for Periprocedural Management of Anticoagulation in Patients With Nonvalvular Atrial Fibrillation: A Report of the American College of Cardiology Clinical Expert Consensus Document Task Force. *J Am Coll Cardiol*. 2017;69(7):871-98. Epub 2017/01/14. doi: 10.1016/j.jacc.2016.11.024. PubMed PMID: 28081965.

15. Lichvar AB, Pierce DR, Salerno D, Klem P, Waldman G, Park JM. Utilization of direct-acting oral anticoagulation in solid organ transplant patients: A national survey of institutional practices. *Clin Transplant*. 2020:e13853. Epub 2020/03/13. doi: 10.1111/ctr.13853. PubMed PMID: 32163212.
16. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Health Care Interventions: Explanation and Elaboration. *PLOS Medicine*. 2009;6(7):e1000100. doi: 10.1371/journal.pmed.1000100.
17. Bixby AL, Shaikh SA, Naik AS, Cotiguala L, McMurry K, Samaniego-Picota MD, et al. Safety and efficacy of direct-acting oral anticoagulants versus warfarin in kidney transplant recipients: a retrospective single-center cohort study. *Transplant international : official journal of the European Society for Organ Transplantation*. 2020. Epub 2020/02/29. doi: 10.1111/tri.13599. PubMed PMID: 32107804.
18. Henricksen EJ, Tremblay-Gravel M, Moayedi Y, Yang W, Lee R, Ross HJ, et al. Use of direct oral anticoagulants after heart transplantation. *The Journal of heart and lung transplantation : the official publication of the International Society for Heart Transplantation*. 2020;39(4):399-401. Epub 2020/02/03. doi: 10.1016/j.healun.2019.12.007. PubMed PMID: 32007373.
19. Leon J, Sabbah L, Aubert O, Anglicheau D, Delavenne X, Zuber J, et al. Efficacy and safety of direct oral anticoagulants in kidney transplantation: a single-center pilot experience. *Transplantation*. 2020. Epub 2020/02/12. doi: 10.1097/tp.0000000000003168. PubMed PMID: 32044892.
20. Santeusanio AD, Weinberg AD, Florman SS, Schiano TD. Safety of direct-acting oral anticoagulants relative to warfarin in a matched cohort of liver transplant recipients. *Clin Transplant*. 2020;34(1):e13756. Epub 2019/11/19. doi: 10.1111/ctr.13756. PubMed PMID: 31738454.
21. Ambrosi P, Kreitmann B, Cohen W, Habib G, Morange P. Anticoagulation with a new oral anticoagulant in heart transplant recipients. *International journal of cardiology*. 2013;168(4):4452-3. Epub 2013/07/31. doi: 10.1016/j.ijcard.2013.06.148. PubMed PMID: 23890900.
22. Bukhari MA, Al-Theaby A, Tawhari M, Al-Shaggag A, Pyrke R, Gangji A, et al. Efficacy and safety of non-vitamin K antagonist oral anticoagulants post-kidney transplantation. *World journal of transplantation*. 2019;9(6):134-44. Epub 2019/11/22. doi: 10.5500/wjt.v9.i6.134. PubMed PMID: 31750090; PubMed Central PMCID: PMC6851500.
23. Lichvar AB, Moore CA, Ensor CR, McDyer JF, Teuteberg JJ, Shullo MA. Evaluation of Direct Oral Anticoagulation Therapy in Heart and Lung Transplant Recipients. *Progress in transplantation (Aliso*

Viejo, Calif). 2016;26(3):263-9. Epub 2016/09/07. doi: 10.1177/1526924816661951. PubMed PMID: 27597772.

24. Vanhove T, Spriet I, Annaert P, Maertens J, Van Cleemput J, Vos R, et al. Effect of the Direct Oral Anticoagulants Rivaroxaban and Apixaban on the Disposition of Calcineurin Inhibitors in Transplant Recipients. *Therapeutic drug monitoring*. 2017;39(1):77-82. Epub 2016/11/20. doi: 10.1097/ftd.0000000000000356. PubMed PMID: 27861314.

25. Wannhoff A, Weiss KH, Schemmer P, Stremmel W, Gotthardt DN. Increased levels of rivaroxaban in patients after liver transplantation treated with cyclosporine A. *Transplantation*. 2014;98(2):e12-3. Epub 2014/07/16. doi: 10.1097/tp.0000000000000223. PubMed PMID: 25022236.

26. Bellam N, Kamath M, DePasquale EC, Smith JG, Fuentes J, Ardehali R, et al. Direct Acting Oral Anticoagulant Utilization, Safety and Efficacy in Heart Transplant (HT). *The Journal of Heart and Lung Transplantation*. 2018;37(4):S112-S3. doi: 10.1016/j.healun.2018.01.269.

27. Bixby AL, Shaikh SA, Naik AS, Cotiguala L, McMurry K, Samaniego-Picota MD, et al. Safety and Efficacy of Direct-Acting Oral Anticoagulants versus Warfarin in Kidney Transplant Recipients . *Am J Transplant*. 2020; 20 (suppl 3). <https://atcmeetingabstracts.com/abstract/safety-and-efficacy-of-direct-acting-oral-anticoagulants-versus-warfarin-in-kidney-transplant-recipients/>. Accessed June 1, 2020.

28. Brown A, Palkimas S, Ally W. Apixaban Therapy Is Safe And Effective In Most Organ Transplant Populations . *Am J Transplant*. 2019; 19 (suppl 3). <https://atcmeetingabstracts.com/abstract/apixaban-therapy-is-safe-and-effective-in-most-organ-transplant-populations/>. Accessed May 28, 2020.

29. Do V, Haakinson D, Belfield K. When to hold: apixaban in renal transplant recipients. *Am J Transplant*. 2019;19(suppl 3). <https://atcmeetingabstracts.com/abstract/when-to-hold-apixaban-in-renal-transplant-recipients/>.

30. Hazelcorn J, Yau R, Sabagha N, Moadda M, Manson M. Risk of major bleeding in solid organ transplant recipients taking calcineurin inhibitors concomitantly with direct oral anticoagulants compared to warfarin . *Am J Transplant*. 2017;17:640. doi:10.1111/ajt.14306.

31. Kim M, Gabardi S, Townsend KR, Page DS, Woodcome EL, Givertz MM. Post Transplant Thrombosis and Atrial Arrhythmia May Be Safely Managed by Direct Oral Anticoagulants in Cardiac Transplant Patients. *The Journal of Heart and Lung Transplantation*. 2016;35(4):S123. doi: 10.1016/j.healun.2016.01.338.

32. Leon J, Zuber J, Amrouche L, Anglicheau D, Divard G, Legendre C, et al. Novel Oral Anticoagulants in Kidney Transplantation: Results from a Pilote Study . *Am J Transplant*. 2019; 19 (suppl

- 3). <https://atcmeetingabstracts.com/abstract/novel-oral-anticoagulants-in-kidney-transplantation-results-from-a-pilote-study/>. Accessed May 27, 2020.
33. Loosier T, Gutierrez K, James A, Dodson A. A Comparison of the Safety and Efficacy of Apixaban Use in Renal Transplant vs. Non-Renal Transplant Recipients . Am J Transplant. 2019; 19 (suppl 3). <https://atcmeetingabstracts.com/abstract/a-comparison-of-the-safety-and-efficacy-of-apixaban-use-in-renal-transplant-vs-non-renal-transplant-recipients/>. Accessed May 27, 2020.
34. McMurry KA, Shuster J, Bain K, Horwedel T, Hartupee J. Safety and Efficacy of Anti Xa Inhibitors Post-Transplant with Concomitant Calcineurin Inhibitors . Am J Transplant. 2019; 19 (suppl 3). <https://atcmeetingabstracts.com/abstract/safety-and-efficacy-of-anti-xa-inhibitors-post-transplant-with-concomitant-calcineurin-inhibitors/>. Accessed May 28, 2020.
35. Pasley T, Logan A, Brueckner A, Bowman L, Silverman A, Rumore A. Direct Oral Anticoagulants: An Evaluation of the Safety and Efficacy in Solid Organ Transplant Recipients . Am J Transplant. 2019; 19 (suppl 3). <https://atcmeetingabstracts.com/abstract/direct-oral-anticoagulants-an-evaluation-of-the-safety-and-efficacy-in-solid-organ-transplant-recipients/>. Accessed May 27, 2020.
36. Pasley T, Logan A, Brueckner A, Bowman-Anger L, Silverman A, Rumore A. Direct Oral Anticoagulants: An Evaluation of the Safety and Efficacy in Cardiothoracic Transplant Recipients. The Journal of Heart and Lung Transplantation. 2019;38(4):S304. doi: 10.1016/j.healun.2019.01.765.
37. Rakouki L, Yau R, Tunwar C. Evaluation of Apixaban Usage in Solid Organ Transplant Patients with Renal Impairment . Am J Transplant. 2020; 20 (suppl 3). <https://atcmeetingabstracts.com/abstract/evaluation-of-apixaban-usage-in-solid-organ-transplant-patients-with-renal-impairment/>. Accessed June 1, 2020.
38. Santeusano A, Weinberg A, Schiano T. The Comparative Safety and Efficacy of Direct Acting Oral Anticoagulants Relative to Warfarin Following Liver Transplantation . Am J Transplant. 2019; 19 (suppl 3). <https://atcmeetingabstracts.com/abstract/the-comparative-safety-and-efficacy-of-direct-acting-oral-anticoagulants-relative-to-warfarin-following-liver-transplantation/>. Accessed May 28, 2020.
39. Shaikh S, Samaniego M, Park J. Direct Oral Anticoagulation Use in Kidney Transplant Recipients . <https://atcmeetingabstracts.com/abstract/direct-oral-anticoagulation-use-in-kidney-transplant-recipients/>. Accessed May 28, 2020.
40. Shuster JE, LaRue SJ, Vader JM. Dabigatran May Have More Significant Drug Interactions with Calcineurin Inhibitors Than Oral Anti-Xa Inhibitors. The Journal of Heart and Lung Transplantation. 2016;35(4):S417. doi: 10.1016/j.healun.2016.01.1204.

41. Thornberg M, Scheffert J, Jennings D, Hedvat J, Lange N, Robbins H, et al. Less Bleeding Associated with Apixaban versus Other Direct Acting Oral Anticoagulation in Solid Organ Transplant Recipients . *Am J Transplant*. 2020; 20 (suppl 3). <https://atcmeetingabstracts.com/abstract/less-bleeding-associated-with-apixaban-versus-other-direct-acting-oral-anticoagulation-in-solid-organ-transplant-recipients/>. Accessed June 6, 2020.
42. Tremblay-Gravel M, Alexander KM, Czobor P, Lee R, Foroutan F, Yang WA, et al. Directly to a DOAC? Safety of Alternatives to Warfarin for Anticoagulation in Heart Transplantation. *The Journal of Heart and Lung Transplantation*. 2019;38(4):S389. doi: 10.1016/j.healun.2019.01.991.
43. Conway SE, Hwang AY, Ponte CD, Gums JG. Laboratory and Clinical Monitoring of Direct Acting Oral Anticoagulants: What Clinicians Need to Know. *Pharmacotherapy*. 2017;37(2):236-48. Epub 2016/12/17. doi: 10.1002/phar.1884. PubMed PMID: 27983747.
44. Samuelson BT, Cuker A. Measurement and reversal of the direct oral anticoagulants. *Blood reviews*. 2017;31(1):77-84. Epub 2016/09/15. doi: 10.1016/j.blre.2016.08.006. PubMed PMID: 27625113; PubMed Central PMCID: PMC5296289.
45. Tomaselli GF, Mahaffey KW, Cuker A, Dobesh PP, Doherty JU, Eikelboom JW, et al. 2020 ACC Expert Consensus Decision Pathway on Management of Bleeding in Patients on Oral Anticoagulants. *Journal of the American College of Cardiology*. 2020;76(5):594. doi: 10.1016/j.jacc.2020.04.053.
46. Dias JD, Norem K, Doorneweerd DD, Thurer RL, Popovsky MA, Omert LA. Use of Thromboelastography (TEG) for Detection of New Oral Anticoagulants. *Archives of pathology & laboratory medicine*. 2015;139(5):665-73. Epub 2015/05/01. doi: 10.5858/arpa.2014-0170-OA. PubMed PMID: 25927150.
47. Hawkins RB, Raymond SL, Hartjes T, Efron PA, Larson SD, Andreoni KA, et al. Review: The Perioperative Use of Thromboelastography for Liver Transplant Patients. *Transplantation proceedings*. 2018;50(10):3552-8. Epub 2018/12/24. doi: 10.1016/j.transproceed.2018.07.032. PubMed PMID: 30577236.
48. Douketis JD, Spyropoulos AC, Duncan J, Carrier M, Le Gal G, Tafur AJ, et al. Perioperative Management of Patients With Atrial Fibrillation Receiving a Direct Oral Anticoagulant. *JAMA Intern Med*. 2019. Epub 2019/08/06. doi: 10.1001/jamainternmed.2019.2431. PubMed PMID: 31380891; PubMed Central PMCID: PMC6686768.
49. Ruff CT, Giugliano RP, Antman EM. Management of Bleeding With Non-Vitamin K Antagonist Oral Anticoagulants in the Era of Specific Reversal Agents. *Circulation*. 2016;134(3):248-61. Epub 2016/07/21. doi: 10.1161/circulationaha.116.021831. PubMed PMID: 27436881.

50. KCentra® [prescribing information] Kankakee ICBL.
51. FIEBA [prescribing information] Lexington MBUI.
52. Praxbind® [package insert] Ridgefield CBIP, Inc.; 2017.
53. Andexxa® [package Insert]. South San Francisco CPP. 2018.
54. Levy JH, van Ryn J, Sellke FW, Reilly PA, Elsaesser A, Glund S, et al. Dabigatran Reversal With Idarucizumab in Patients Requiring Urgent Surgery: A Subanalysis of the RE-VERSE AD Study. *Annals of surgery*. 2019. Epub 2019/10/11. doi: 10.1097/sla.0000000000003638. PubMed PMID: 31599808.
55. Rimsans J, Rhoten M, Sylvester K, Singh SK, Connors JM. Idarucizumab for urgent reversal of dabigatran for heart transplant: A case report. *American journal of hematology*. 2017;92(3):E34-e5. Epub 2016/12/30. doi: 10.1002/ajh.24638. PubMed PMID: 28033655.
56. Tralhão A, Aguiar C, Ferreira J, Rebocho MJ, Santos E, Martins D, et al. Dabigatran reversal with idarucizumab in a patient undergoing heart transplantation: first European report. *Thrombosis journal*. 2017;15:23. Epub 2017/09/08. doi: 10.1186/s12959-017-0147-z. PubMed PMID: 28878572; PubMed Central PMCID: PMC5583973.
57. Józwiak A, Lisik W, Czerwiński J, Kosieradzki M. Simultaneous Pancreas-Kidney Transplantation in a Patient with Heparin-Induced Thrombocytopenia on Dabigatran Therapy. *Annals of transplantation*. 2018;23:232-5. Epub 2018/04/07. doi: 10.12659/aot.905868. PubMed PMID: 29622761; PubMed Central PMCID: PMC6248287.
58. Lopez-Vilella R, Sanz-Sanchez J, Sanchez-Lazaro I, Marques-Sule E, Rueda-Soriano J, Almenar-Bonet L. Idarucizumab in High-risk Thoracic Surgery. *International journal of organ transplantation medicine*. 2018;9(2):97-100. Epub 2018/01/01. PubMed PMID: 30834094; PubMed Central PMCID: PMC6390979.
59. Crespo-Leiro MG, López-Vilella R, López Granados A, Mirabet-Pérez S, Díez-López C, Barge-Caballero E, et al. Use of Idarucizumab to reverse the anticoagulant effect of dabigatran in cardiac transplant surgery. A multicentric experience in Spain. *Clin Transplant*. 2019;33(12):e13748. Epub 2019/11/02. doi: 10.1111/ctr.13748. PubMed PMID: 31670852.
60. Van Keer JM, Vanassche T, Droogne W, Rex S, Rega F, Van Cleemput J, et al. Idarucizumab for the reversal of dabigatran in patients undergoing heart transplantation. *European journal of heart failure*. 2019;21(1):129-31. Epub 2018/12/07. doi: 10.1002/ejhf.1356. PubMed PMID: 30520546.
61. Herrera-Escandon A, Castano-Cifuentes O, Plata-Mosquera CA. Use of Idarucizumab to Revert the Anticoagulant Effect of Dabigatran in Heart Transplant Surgery: An Institutional Experience. *Case*

reports in cardiology. 2020;2020:6927423. Epub 2020/03/25. doi: 10.1155/2020/6927423. PubMed PMID: 32206355; PubMed Central PMCID: PMC7081017.

62. Marino KK, Santiago RA, Dew RB, 3rd, Berliner N, Connors JM, Connell NT, et al. Management of Dabigatran-Associated Bleeding with Two Doses of Idarucizumab Plus Hemodialysis. *Pharmacotherapy*. 2016;36(10):e160-e5. Epub 2016/09/02. doi: 10.1002/phar.1830. PubMed PMID: 27581709.

63. Schulman S, Ritchie B, Nahirniak S, Gross PL, Carrier M, Majeed A, et al. Reversal of dabigatran-associated major bleeding with activated prothrombin concentrate: A prospective cohort study. *Thrombosis research*. 2017;152:44-8. Epub 2017/02/22. doi: 10.1016/j.thromres.2017.02.010. PubMed PMID: 28222322.

64. Cuker A, Burnett A, Triller D, Crowther M, Ansell J, Van Cott EM, et al. Reversal of direct oral anticoagulants: Guidance from the Anticoagulation Forum. *American journal of hematology*. 2019;94(6):697-709. Epub 2019/03/28. doi: 10.1002/ajh.25475. PubMed PMID: 30916798.

65. Majeed A, Ågren A, Holmström M, Bruzelius M, Chaireti R, Odeberg J, et al. Management of rivaroxaban- or apixaban-associated major bleeding with prothrombin complex concentrates: a cohort study. *Blood*. 2017;130(15):1706-12. Epub 2017/08/25. doi: 10.1182/blood-2017-05-782060. PubMed PMID: 28835439.

66. Schulman S, Gross PL, Ritchie B, Nahirniak S, Lin Y, Lieberman L, et al. Prothrombin Complex Concentrate for Major Bleeding on Factor Xa Inhibitors: A Prospective Cohort Study. *Thrombosis and haemostasis*. 2018;118(5):842-51. Epub 2018/03/23. doi: 10.1055/s-0038-1636541. PubMed PMID: 29564837.

67. Schultz NH, Lundblad R, Holme PA. Activated prothrombin complex concentrate to reverse the factor Xa inhibitor (apixaban) effect before emergency surgery: a case series. *Journal of medical case reports*. 2018;12(1):138. Epub 2018/05/17. doi: 10.1186/s13256-018-1660-9. PubMed PMID: 29764497; PubMed Central PMCID: PMC5954448.

68. US Food and Drug Administration. [cited 2020 May 15 2020]. Available from: <https://www.fda.gov/drugs/drug-interactions-labeling/drug-development-and-drug-interactions-table-substrates-inhibitors-and-inducers#table3-2>.

69. Chang SH, Chou IJ, Yeh YH, Chiou MJ, Wen MS, Kuo CT, et al. Association Between Use of Non-Vitamin K Oral Anticoagulants With and Without Concurrent Medications and Risk of Major Bleeding in Nonvalvular Atrial Fibrillation. *Jama*. 2017;318(13):1250-9. Epub 2017/10/04. doi: 10.1001/jama.2017.13883. PubMed PMID: 28973247; PubMed Central PMCID: PMC5818856 Form for Disclosure of Potential Conflicts of Interest and none were reported.

70. Zhang Y, Souverein PC, Gardarsdottir H, van den Ham HA, Maitland-van der Zee AH, de Boer A. Risk of major bleeding among users of direct oral anticoagulants combined with interacting drugs: A population-based nested case-control study. *British journal of clinical pharmacology*. 2020. Epub 2020/02/06. doi: 10.1111/bcp.14227. PubMed PMID: 32022295.
71. Bashir B, Stickle DF, Chervoneva I, Kraft WK. Drug-Drug Interaction Study of Apixaban with Cyclosporine and Tacrolimus in Healthy Volunteers. *Clinical and translational science*. 2018;11(6):590-6. Epub 2018/07/05. doi: 10.1111/cts.12580. PubMed PMID: 29972633; PubMed Central PMCID: PMC6226116.
72. Parasrampur DA, Mendell J, Shi M, Matsushima N, Zahir H, Truitt K. Edoxaban drug-drug interactions with ketoconazole, erythromycin, and cyclosporine. *British journal of clinical pharmacology*. 2016;82(6):1591-600. Epub 2016/08/18. doi: 10.1111/bcp.13092. PubMed PMID: 27530188; PubMed Central PMCID: PMC5099547.
73. Lempers VJ, van den Heuvel JJ, Russel FG, Aarnoutse RE, Burger DM, Brüggemann RJ, et al. Inhibitory Potential of Antifungal Drugs on ATP-Binding Cassette Transporters P-Glycoprotein, MRP1 to MRP5, BCRP, and BSEP. *Antimicrobial agents and chemotherapy*. 2016;60(6):3372-9. Epub 2016/03/24. doi: 10.1128/aac.02931-15. PubMed PMID: 27001813; PubMed Central PMCID: PMC4879353.
74. Sotomi Y, Hirata A, Amiya R, Kobayashi T, Hirayama A, Sakata Y, et al. Bleeding Risk of Add-On Anti-Platelet Agents to Direct Oral Anticoagulants in Patients With Nonvalvular Atrial Fibrillation (From 2216 Patients in the DIRECT Registry). *The American journal of cardiology*. 2019;123(8):1293-300. Epub 2019/02/06. doi: 10.1016/j.amjcard.2019.01.027. PubMed PMID: 30717887.
75. Bennaghmouch N, de Veer A, Bode K, Mahmoodi BK, Dewilde WJM, Lip GYH, et al. Efficacy and Safety of the Use of Non-Vitamin K Antagonist Oral Anticoagulants in Patients With Nonvalvular Atrial Fibrillation and Concomitant Aspirin Therapy: A Meta-Analysis of Randomized Trials. *Circulation*. 2018;137(11):1117-29. Epub 2017/11/05. doi: 10.1161/circulationaha.117.028513. PubMed PMID: 29101289.
76. Douros A, Renoux C, Yin H, Filion KB, Suissa S, Azoulay L. Concomitant Use of Direct Oral Anticoagulants with Antiplatelet Agents and the Risk of Major Bleeding in Patients with Nonvalvular Atrial Fibrillation. *The American journal of medicine*. 2019;132(2):191-9.e12. Epub 2019/01/30. doi: 10.1016/j.amjmed.2018.10.008. PubMed PMID: 30691551.
77. Cannon CP, Bhatt DL, Oldgren J, Lip GYH, Ellis SG, Kimura T, et al. Dual Antithrombotic Therapy with Dabigatran after PCI in Atrial Fibrillation. *N Engl J Med*. 2017;377(16):1513-24. Epub 2017/08/29. doi: 10.1056/NEJMoa1708454. PubMed PMID: 28844193.

78. Lopes RD, Heizer G, Aronson R, Vora AN, Massaro T, Mehran R, et al. Antithrombotic Therapy after Acute Coronary Syndrome or PCI in Atrial Fibrillation. *N Engl J Med*. 2019;380(16):1509-24. Epub 2019/03/19. doi: 10.1056/NEJMoa1817083. PubMed PMID: 30883055.
79. Vranckx P, Valgimigli M, Eckardt L, Tijssen J, Lewalter T, Gargiulo G, et al. Edoxaban-based versus vitamin K antagonist-based antithrombotic regimen after successful coronary stenting in patients with atrial fibrillation (ENTRUST-AF PCI): a randomised, open-label, phase 3b trial. *Lancet (London, England)*. 2019;394(10206):1335-43. Epub 2019/09/08. doi: 10.1016/s0140-6736(19)31872-0. PubMed PMID: 31492505.
80. Barsam SJ, Patel JP, Roberts LN, Kavarthapu V, Patel RK, Green B, et al. The impact of body weight on rivaroxaban pharmacokinetics. *Res Pract Thromb Haemost*. 2017;1(2):180-7. Epub 2018/07/27. doi: 10.1002/rth2.12039. PubMed PMID: 30046688; PubMed Central PMCID: PMC6058267.
81. Bertaglia-Calderara D, Kröll D, Gerschheimer C, Nicolas N, Nett P, Stirnimann G, et al. Effect of Rivaroxaban on thrombin generation in vivo. A study in obese patients. *International journal of laboratory hematology*. 2018;40(1):e11-e4. Epub 2017/12/15. doi: 10.1111/ijlh.12767. PubMed PMID: 29239121.
82. Hohnloser SH, Fudim M, Alexander JH, Wojdyla DM, Ezekowitz JA, Hanna M, et al. Efficacy and Safety of Apixaban Versus Warfarin in Patients With Atrial Fibrillation and Extremes in Body Weight. *Circulation*. 2019;139(20):2292-300. Epub 2019/02/19. doi: 10.1161/circulationaha.118.037955. PubMed PMID: 30773022.
83. Kido K, Ngorsuraches S. Comparing the Efficacy and Safety of Direct Oral Anticoagulants With Warfarin in the Morbidly Obese Population With Atrial Fibrillation. *The Annals of pharmacotherapy*. 2019;53(2):165-70. Epub 2018/08/23. doi: 10.1177/1060028018796604. PubMed PMID: 30130979.
84. Kubitzka D, Becka M, Zuehlsdorf M, Mueck W. Body weight has limited influence on the safety, tolerability, pharmacokinetics, or pharmacodynamics of rivaroxaban (BAY 59-7939) in healthy subjects. *Journal of clinical pharmacology*. 2007;47(2):218-26. Epub 2007/01/25. doi: 10.1177/0091270006296058. PubMed PMID: 17244773.
85. Kushnir M, Choi Y, Eisenberg R, Rao D, Tolu S, Gao J, et al. Efficacy and safety of direct oral factor Xa inhibitors compared with warfarin in patients with morbid obesity: a single-centre, retrospective analysis of chart data. *The Lancet Haematology*. 2019;6(7):e359-e65. Epub 2019/05/28. doi: 10.1016/s2352-3026(19)30086-9. PubMed PMID: 31133411.

86. Peterson ED, Ashton V, Chen YW, Wu B, Spyropoulos AC. Comparative effectiveness, safety, and costs of rivaroxaban and warfarin among morbidly obese patients with atrial fibrillation. *American heart journal*. 2019;212:113-9. Epub 2019/04/14. doi: 10.1016/j.ahj.2019.02.001. PubMed PMID: 30981035.
87. Piran S, Traquair H, Chan N, Bhagirath V, Schulman S. Peak plasma concentration of direct oral anticoagulants in obese patients weighing over 120 kilograms: A retrospective study. *Res Pract Thromb Haemost*. 2018;2(4):684-8. Epub 2018/10/24. doi: 10.1002/rth2.12146. PubMed PMID: 30349887; PubMed Central PMCID: PMC6178753.
88. Reilly PA, Lehr T, Haertter S, Connolly SJ, Yusuf S, Eikelboom JW, et al. The effect of dabigatran plasma concentrations and patient characteristics on the frequency of ischemic stroke and major bleeding in atrial fibrillation patients: the RE-LY Trial (Randomized Evaluation of Long-Term Anticoagulation Therapy). *J Am Coll Cardiol*. 2014;63(4):321-8. Epub 2013/10/01. doi: 10.1016/j.jacc.2013.07.104. PubMed PMID: 24076487.
89. Sandhu RK, Ezekowitz J, Andersson U, Alexander JH, Granger CB, Halvorsen S, et al. The 'obesity paradox' in atrial fibrillation: observations from the ARISTOTLE (Apixaban for Reduction in Stroke and Other Thromboembolic Events in Atrial Fibrillation) trial. *European heart journal*. 2016;37(38):2869-78. Epub 2016/04/14. doi: 10.1093/eurheartj/ehw124. PubMed PMID: 27071819.
90. Upreti VV, Wang J, Barrett YC, Byon W, Boyd RA, Pursley J, et al. Effect of extremes of body weight on the pharmacokinetics, pharmacodynamics, safety and tolerability of apixaban in healthy subjects. *British journal of clinical pharmacology*. 2013;76(6):908-16. Epub 2013/03/16. doi: 10.1111/bcp.12114. PubMed PMID: 23488672; PubMed Central PMCID: PMC3845314.
91. Yin OQ, Tetsuya K, Miller R. Edoxaban population pharmacokinetics and exposure-response analysis in patients with non-valvular atrial fibrillation. *European journal of clinical pharmacology*. 2014;70(11):1339-51. Epub 2014/08/30. doi: 10.1007/s00228-014-1736-4. PubMed PMID: 25168620.
92. Zhou Y, Ma J, Zhu W. Efficacy and Safety of Direct Oral Anticoagulants Versus Warfarin in Patients with Atrial Fibrillation Across BMI Categories: A Systematic Review and Meta-Analysis. *American journal of cardiovascular drugs : drugs, devices, and other interventions*. 2020;20(1):51-60. Epub 2019/07/26. doi: 10.1007/s40256-019-00362-4. PubMed PMID: 31342343.
93. Martin K, Beyer-Westendorf J, Davidson BL, Huisman MV, Sandset PM, Moll S. Use of the direct oral anticoagulants in obese patients: guidance from the SSC of the ISTH. *Journal of thrombosis and haemostasis : JTH*. 2016;14(6):1308-13. Epub 2016/06/15. doi: 10.1111/jth.13323. PubMed PMID: 27299806; PubMed Central PMCID: PMC4936273.

94. Kido K, Lee JC, Hellwig T, Gulseth MP. Use of Direct Oral Anticoagulants in Morbidly Obese Patients. *Pharmacotherapy*. 2020;40(1):72-83. Epub 2019/12/14. doi: 10.1002/phar.2353. PubMed PMID: 31834939.
95. Kido K, Shimizu M, Shiga T, Hashiguchi M. Meta-Analysis Comparing Direct Oral Anticoagulants Versus Warfarin in Morbidly Obese Patients With Atrial Fibrillation. *The American journal of cardiology*. 2020. Epub 2020/04/30. doi: 10.1016/j.amjcard.2020.03.048. PubMed PMID: 32345473.
96. January CT, Wann LS, Calkins H, Chen LY, Cigarroa JE, Cleveland JC, Jr., et al. 2019 AHA/ACC/HRS Focused Update of the 2014 AHA/ACC/HRS Guideline for the Management of Patients With Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol*. 2019;74(1):104-32. Epub 2019/02/01. doi: 10.1016/j.jacc.2019.01.011. PubMed PMID: 30703431.
97. Eliquis[®] (apixaban) [prescribing information]. Princeton, NJ: Bristol-Myers Squibb Company; June 2019.
98. Chan KE, Edelman ER, Wenger JB, Thadhani RI, Maddux FW. Dabigatran and rivaroxaban use in atrial fibrillation patients on hemodialysis. *Circulation*. 2015;131(11):972-9. Epub 2015/01/18. doi: 10.1161/circulationaha.114.014113. PubMed PMID: 25595139; PubMed Central PMCID: PMC4363265.
99. De Vriese AS, Caluwé R, Bailleul E, De Bacquer D, Borrey D, Van Vlem B, et al. Dose-finding study of rivaroxaban in hemodialysis patients. *American journal of kidney diseases : the official journal of the National Kidney Foundation*. 2015;66(1):91-8. Epub 2015/03/26. doi: 10.1053/j.ajkd.2015.01.022. PubMed PMID: 25804678.
100. Dias C, Moore KT, Murphy J, Ariyawansa J, Smith W, Mills RM, et al. Pharmacokinetics, Pharmacodynamics, and Safety of Single-Dose Rivaroxaban in Chronic Hemodialysis. *American journal of nephrology*. 2016;43(4):229-36. Epub 2016/04/22. doi: 10.1159/000445328. PubMed PMID: 27100875.
101. Kuno T, Takagi H, Ando T, Sugiyama T, Miyashita S, Valentin N, et al. Oral Anticoagulation for Patients With Atrial Fibrillation on Long-Term Hemodialysis. *J Am Coll Cardiol*. 2020;75(3):273-85. Epub 2020/01/25. doi: 10.1016/j.jacc.2019.10.059. PubMed PMID: 31976865.
102. Wang X, Tirucherai G, Marbury TC, Wang J, Chang M, Zhang D, et al. Pharmacokinetics, pharmacodynamics, and safety of apixaban in subjects with end-stage renal disease on hemodialysis. *Journal of clinical pharmacology*. 2016;56(5):628-36. Epub 2015/09/04. doi: 10.1002/jcph.628. PubMed PMID: 26331581.

103. Siontis KC, Zhang X, Eckard A, Bhave N, Schaubel DE, He K, et al. Outcomes Associated With Apixaban Use in Patients With End-Stage Kidney Disease and Atrial Fibrillation in the United States. *Circulation*. 2018;138(15):1519-29. Epub 2018/06/30. doi: 10.1161/circulationaha.118.035418. PubMed PMID: 29954737; PubMed Central PMCID: PMC6202193.
104. Pokorney SD. AHA 2019 Presentation Slides for the RENAL-AF Trial. [cited April 27 2020]. Available from: <https://www.acc.org/education-and-meetings/image-and-slide-gallery/media-detail?id=469e5bd88fff4d2bb8ec183e71521637>.
105. Mavrakanas TA, Samer CF, Nessim SJ, Frisch G, Lipman ML. Apixaban Pharmacokinetics at Steady State in Hemodialysis Patients. *Journal of the American Society of Nephrology : JASN*. 2017;28(7):2241-8. Epub 2017/03/18. doi: 10.1681/asn.2016090980. PubMed PMID: 28302754; PubMed Central PMCID: PMC5491286.
106. Schulman S, Angerås U, Bergqvist D, Eriksson B, Lassen MR, Fisher W. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in surgical patients. *Journal of thrombosis and haemostasis : JTH*. 2010;8(1):202-4. Epub 2009/11/03. doi: 10.1111/j.1538-7836.2009.03678.x. PubMed PMID: 19878532.

Table 1. Studies Analyzing DOAC Use in Solid Organ Transplant Recipients

First Author (Year)	N (for DOAC)	Organ	Major bleed (%)	Composite bleed (%)	Thrombotic event (%)
Full-text Comparative Studies					
Bixby (2020)† (17)	99	Kidney	9%	24%	3%
Henricksen (2020) (18)	51	Heart	-	10%	4%
Leon (2020) (19)	52	Kidney	0%	13%	0%
Santeusanio (2020)† (20)	27	Liver	0%	11%	15%
Full-text Single-arm Studies					
Ambrosi (2013) (21)	11	Heart	10%	-	0%
Bukhari (2019) (22)	42	Kidney	2%	7%	0%
Lichvar (2016) (23)	37	Heart or Lung	3%	19%	5%
Vanhove (2017) (24)	39	Any	0%	5%	-
Wannhoff (2014) (25)	9	Liver	-	33%	0%
Abstracts					
Bellam (2018) (26)	23	Heart	0%	0%	0%
Bixby (2020)† (27)	99	Kidney	9%	24%	3%
Brown (2019) (28)	162	Any	3%	17%	4%
Do (2019) (29)	52	Kidney	4%	12%	4%
Hazelcorn (2017) (30)	37	Any	11%	-	-
Kim (2016) (31)	7	Heart	0%	14%	0%
Leon (2019) (32)	33	Kidney	0%	12%	0%
Loosier (2019) (33)	22	Kidney	0%	0%	5%

McMurry (2019) (34)	172	Any	-	6%	1%
Pasley (2019) (36)	38	Heart or Lung	0%	16%	21%
Pasley (2019) (35)	109	Heart or Lung	1%	14%	9%
Rakouki (2020) (37)	77	Any	-	13%	6%
Santeusanio (2019)† (38)	27	Liver	-	11%	15%
Shaikh (2017) (39)	62	Kidney	23%	37%	0%
Shuster (2016) (40)	27	Heart	9%	-	-
Thornberg (2020) (41)	94	Any	-	19%	7%
Tremblay-Gravel (2019) (42)	62	Heart	-	-	-

† Published both in a full-text article and an abstract

Abbreviation: DOAC, direct-acting oral anticoagulant

Author Manuscript

Table 2. Summary of Studies Comparing DOAC to Vitamin K Antagonists in Solid Organ Transplant Recipients (Single-center Retrospective Cohort Studies)

First Author, Year (NOS Study Quality Rating)	Study Treatments	Outcomes	Baseline Characteristics	Efficacy Outcome Results	Safety Outcome Results
<i>Liver Transplant</i>					
Santeusanio, 2020 (NOS: S 3 stars, C 2 stars, and O 3 stars) (20)	DOAC (n=20): apixaban, dabigatran, rivaroxaban Warfarin (n=20)	Primary efficacy endpoint: thrombotic event Primary safety endpoint: incidence of clinically relevant major or non-major bleeding	DOAC vs warfarin: Age 60.3 ± 11.3 vs 59.4 ± 11.1 years (p=0.79) NVAf 25% vs 25% VTE 55% vs 35% (p=0.24) Time post-transplant 676 (IQR 39-4023) vs 1916 (IQR 24-4786) days (p=0.54) HAS-BLED score 2 (1.5- 3) vs 2 (1.5-2.5) (p=1.00) CHADS ₂ 1 (1-3) vs 3 (2-3) (p=0.40)	DOAC vs warfarin: Thrombotic events 20% vs 15% (p=0.67)	DOAC vs warfarin: clinically relevant bleeding 15% vs 45% (p=0.01)
<i>Kidney Transplant</i>					

<p>Bixby, 2020 (NOS: S 4 stars, C 1 stars, O 2 stars) (19)</p>	<p>DOAC (n=99): apixaban (n=60), dabigatran (n=6), rivaroxaban (n=33) Warfarin (n=98)</p>	<p>Primary efficacy endpoint: new onset stroke, new onset stroke or recurrent VTE Primary safety endpoint: incidence of major bleeding</p>	<p>DOAC vs Warfarin: Age 62.0 (IQR 56.5-70.3) vs 62.3 (IQR 54.4-68.4) (p=0.52) Time post-transplant 6.5 (IQR 2.8-10.8) vs 6.4 (IQR 1.6-12.5) years (p=0.98) NVAf 54% vs 42% VTE 47% vs 58% (p=0.10) ATRIA 3.2 ± 1.9 vs 3.6 ± 2.1 (p=0.21) CHA₂DS₂-VASc 3.5 ± 1.7 vs 3.5 ± 1.4 (p=0.27)</p>	<p>DOAC vs Warfarin: Stroke 1.0% vs 5.1% (p=0.13) VTE 2.0% vs 3.1% (p=0.50)</p>	<p>DOAC vs Warfarin Major bleeding 9.1% vs 21.4% (p=0.15) Composite bleeding 24.2% vs 34.7% (NS in time to event analysis)</p>
--	---	--	--	---	--

<p>Leon, 2020 (NOS: S 3 stars, C 0 stars, O 3 stars) (17)</p>	<p>DOAC (n=52): apixaban (n=36), dabigatran (n=1), rivaroxaban (n=15) vs VKA (n=50): warfarin (n=20), fluindione (n=30)</p>	<p>Primary efficacy endpoint: arterial thromboembolism event or VTE event Primary safety endpoint: any bleeding</p>	<p>DOAC vs VKA: Age 62 ± 13 vs 60 ± 11 years (p=0.45) Time post-transplant 87 ± 95 vs 65 ± 90 months (p=0.049) NVAf 60% vs 42% (p=0.11) VTE 29% vs 44% (p=0.15) HAS-BLED score 2.9 ± 1.3 vs 2.4 ± 1.1 (p=0.044) CHA₂DS₂-VASc 3.4 ± 1.7 vs 3.2 ± 1.5 (p=0.81)</p>	<p>DOAC vs VKA: Arterial thromboembolism 0% vs 0% (p=1.0) VTE 0% vs 8% (p=0.054)</p>	<p>DOAC vs VKA: Any bleeding event 13% vs 42% (p=0.002)</p>
<p><i>Heart Transplant</i></p>					

Henricksen, 2020 (NOS: S 3 stars, C 1 stars, O 2 stars) (18)	DOAC (n=51): apixaban (n=35), dabigatran (n=14), rivaroxaban (n=2) Warfarin (n=22)	Primary safety endpoint: any bleeding event requiring intervention	DOAC vs Warfarin: Age: 58 (IQR 51-65) vs 54 (IQR 46-62) (p=0.28) Time post-transplant 42 vs 75 days (p=0.29) NVAf 13.7% vs 13.6% VTE 84.3% vs 77.3% (p=0.60)	VTE 4% vs 0%	DOAC vs Warfarin Bleeding event 10% vs 23% (p=0.08)
---	--	--	---	--------------	--

Abbreviations: DOAC, direct-acting oral anticoagulant; IQR, interquartile range; NVAf, non-valvular atrial fibrillation; NOS, Newcastle-Ottawa Scale; VKA, vitamin K antagonist; VTE, venous thromboembolism

Table 3. Impact of DOACs on Anticoagulation Monitoring Parameters (43, 44)

Assay	Function	Sensitivity in Direct Thrombin Inhibitors	Effect in Direct Thrombin Inhibitors	Utility in Direct Thrombin Inhibitors	Sensitivity in Factor Xa Inhibitors	Effect in Factor Xa Inhibitors	Utility in Factor Xa Inhibitors
Activated partial thromboplastin time (aPTT)	Activity and presence of FII, FV, and FIII/FXII	Not sensitive	Increased	Qualitative assessment, normal does not rule out effect	Low	Increased for all but apixaban	No utility in routine clinical practice
Anti-Xa	Concentration -dependent inhibition of FXa	-	-	-	High	Increased (Trough dependent on FXa inhibitor)	Quantitative, if calibrated to specific anticoagulant
Dilute thrombin time (dTT)	Diluted plasma to make a more sensitive test than TT	Sensitive	Increased Trough > 200 ng/mL dTT ≥ 65 seconds associated with bleeds	Quantitative assessment	-	-	-
Ecarin clotting time (ECT)	Measures thrombin generation	Sensitive	Increased Trough 3x the upper limit of	Quantitative assessment	-	-	-

			normal associated with bleeds				
--	--	--	-------------------------------	--	--	--	--

Table 4. Limited Utility of Anticoagulation Monitoring Parameters for DOAC Monitoring (43, 45)

DOAC	Exclude Clinically Relevant* Drug Levels		Measure On-therapy or Above On-Therapy Levels	
	Suggested test	Interpretation	Suggested test	Interpretation
Dabigatran	TT aPTT	<p>Normal TT: excludes clinically relevant* levels.</p> <p>Prolonged TT: does not discriminate between clinically significant and insignificant levels.</p> <p>Normal aPTT: usually excludes clinically relevant* levels if a sensitive reagent is used.</p>	aPTT	<p>Prolonged aPTT: suggests that on-therapy or above on-therapy levels are present.</p> <p>Normal aPTT: may not exclude on-therapy levels, particularly if a relatively insensitive aPTT reagent is used.</p>

Apixaban	UFH or LMWH anti-FXa	<p>Normal PT and aPTT: do not exclude clinically relevant* levels.</p> <p>UFH or LMWH anti-FXa: below the lower limit of quantitation probably excludes clinically relevant* levels</p>	PT	<p>Prolonged PT: suggests that on-therapy or above on-therapy levels are present.</p> <p>Normal PT: may not exclude on-therapy or above on-therapy levels, particularly if a relatively insensitive PT reagent is used.</p>
Betrixaban, edoxaban, rivaroxaban	UFH or LMWH anti-FXa	<p><u>Normal PT and aPTT:</u> does not exclude clinically relevant* levels.</p> <p><u>UFH or LMWH anti-FXa</u> below the lower limit of quantitation probably excludes clinically relevant* levels.</p>	PT	<p>Prolonged PT: suggests that on-therapy or above on-therapy levels are present.</p> <p>Normal PT: may not exclude on-therapy levels, particularly if a relatively insensitive PT reagent is used.</p>

*The term “clinically relevant” refers to DOAC levels that may contribute to bleeding or surgical bleeding risk. The minimum DOAC level that may contribute to bleeding or surgical bleeding risk is unknown. The International Society on Thrombosis and Hemostasis recommends consideration of anticoagulant reversal for patients with serious bleeding and a

DOAC level >50 ng/mL, and for patients requiring an invasive procedure with high bleeding risk and a DOAC level >30 ng/mL. (106)

Abbreviations: aPTT, activated partial thromboplastin time; DOAC, direct-acting oral anticoagulant; dTT, dilute thrombin time; ECA, ecarin chromogenic assay; ECT, ecarin clotting time; INR, international normalized ratio; PT, prothrombin time; TT, thrombin time

Table 5. Recommendations for Pre-procedural Interruption of DOAC therapy (adapted from the 2017 ACC Expert Consensus Discussion Pathway for Periprocedural Management of Anticoagulation in Patients with Nonvalvular Atrial Fibrillation and the 2020 ACC Expert Consensus Decision Pathway on Management of Bleeding in Patients on Oral Anticoagulants) (14, 45)

	Rivaroxaban (R), Edoxaban (E), Apixaban (A), Betrixaban (B)			Dabigatran		
	t _{1/2} (hrs)	Low Risk	Uncertain, Intermediate or High Risk	t _{1/2} (hrs)	Low Risk	Uncertain, Intermediate or High Risk
CrCl ≥80 mL/min	6-15 (R, E, A) 19 – 27 (B)	≥ 24 hrs	≥ 48 hrs	13	≥ 24 hrs	≥ 48 hrs
CrCl 50-79 mL/min	6-15 (R, E, A) 19 – 27 (B)	≥ 24 hrs	≥ 48 hrs	15	≥ 36 hrs	≥ 72 hrs
CrCl 30-49 mL/min	6-15 (R, E, A) 19 – 27 (B)	≥ 24 hrs	≥ 48 hrs	18	≥ 48 hrs	≥ 96 hrs
CrCl 15-29 mL/min	R: 9	≥ 36 hrs	Not indicated.	27	≥ 72 hrs	≥ 120 hrs

	E: 17 A: 17		No data. Consider measuring agent-specific anti-Xa level and/or withholding \geq 72 h.			
CrCl <15 mL/min	R: 13 (ff dialysis) E: 10-17 (off dialysis) A: 17	Not indicated. No data. Consider measuring agent-specific anti-Xa level and/or withholding \geq 72 h.	30 (off dialysis)	Not indicated. No data. Consider measuring dTT and/or withholding \geq 96 h (low risk procedures only)		

Abbreviations: DOAC, direct-acting oral anticoagulant; CrCl, creatinine clearance

Table 6. Specific and Non-specific DOAC Reversal Agents

	KCentra® (4 factor Prothrombin Complex Concentrate, 4F-PCC) (50)	FEIBA® (Anti-Inhibitor Coagulant Complex, aPCC) (51)	Praxbind® (Idarucizumab) (52)	Andexxa® (Andexanet Alfa) (53)
Classification	Non-specific prohemostatic agent		Specific antidote (humanized monoclonal antibody fragment)	Specific antidote (recombinant variant of human factor Xa)
Mechanism of action	II, VII, IX, X , Proteins C and S, heparin	II, VIIa, IX, X, VIII inhibitor bypassing activity	Binding to dabigatran and its metabolites neutralizing the anticoagulation effect	Binds oral factor Xa inhibitors and binds/ inhibits tissue factor pathway inhibitor
Half-life ($t_{1/2}$)	Dependent on half-lives of individual clotting factors Elevated levels of clotting factors persistent for ~24h		Pharmacodynamic: 45 min Terminal: 4 to 8 h	Pharmacodynamic: 30 to 60 min (anti-Xa rebound) Terminal: 5 to 7 h
Elimination	Hepatic		Renal	Unknown
Dose	Potency based on FIX content 2000 units x 1 dose (fixed dose) Alternative: 50 units/kg x 1 dose	Potency based on FVIII inhibitor bypassing activity in units 50 units/kg x 1 dose	2.5 g IV over 5 min x 2 doses	High dose bolus + infusion: 800 mg at 30mg/min then 8mg/min for up to 120 min Low dose bolus + infusion:

				400 mg at 30mg/min then 4mg/min for up to 120 min
Onset	15 minutes (warfarin data)	15 minutes (warfarin data)	< 5 mon	2 – 5 min

FIGURE LEGENDS

Figure 1. PRISMA Flow Diagram of Studies Included

Figure 2. Summary of the Reversal of Direct Oral Anticoagulants (adapted Guidance from the Anticoagulation Forum and 2020 ACC Expert Consensus Decision Pathway on Management of Bleeding in Patients on Oral Anticoagulants) (45)

Figure 1: PRISMA Flow Diagram of Studies Included





Figure 2



Author Manuscript