

## Early Enteral Nutrition in Mechanically Ventilated Patients with SARS-CoV-2 Infection

Nicholas Farina, PharmD, BCCCP<sup>1</sup>; Sarah Nordbeck, PharmD, BCNSP<sup>1</sup>; Michelle Montgomery, RDN<sup>2</sup>; Laura Cordwin, MMN, RDN; Faith Blair, RDN<sup>2</sup>; Jill Cherry-Bukowiec, MD, MS, FACS, FCCM<sup>3</sup>; Michael D. Kraft, PharmD, BCNSP<sup>1,4</sup>; Melissa R. Pleva, PharmD, BCPS, BCNSP, BCCCP<sup>1</sup>; Erica Raymond, MS, RDN<sup>2</sup>

1. Michigan Medicine, Department of Pharmacy, Ann Arbor MI
2. Michigan Medicine, Department of Nutrition Services, Ann Arbor MI
3. University of Michigan, Department of Surgery, Ann Arbor MI
4. University of Michigan College of Pharmacy, Ann Arbor MI

Corresponding Author:

Nicholas Farina, PharmD, BCCCP

Clinical Pharmacy Specialist – Surgery

Michigan Medicine

1111 E Catherine Street

Victor Vaughan Building Room 301

Ann Arbor, MI 48109

[nfarina@med.umich.edu](mailto:nfarina@med.umich.edu)

Conflict of Interest: Study authors do not have any conflicts of interest to disclose

Financial Disclosure: None declared

Abstract

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as [doi: 10.1002/ncp.10629](https://doi.org/10.1002/ncp.10629).

This article is protected by copyright. All rights reserved.

Background: Nutrition therapy is essential in critically ill adults. Little is known about appropriate nutrition therapy in patients with SARS-CoV-2 (COVID-19) infection.

Methods: This was a retrospective, observational study in adult patients with confirmed COVID-19 infection receiving mechanical ventilation. Data regarding patient demographics and nutrition therapy were collected. Patients that received enteral nutrition within 24 hours of starting mechanical ventilation were compared to patients started on enteral nutrition later. The primary outcome was inpatient length of stay. Propensity score matching was conducted to control for baseline differences in patient groups.

Results: 155 patients were included in final analysis. Patients who received enteral nutrition within 24 hours received a significantly greater daily amount of calories (17.5 vs. 15.2 kcal/kg,  $p=0.015$ ) and protein (1.04 vs. 0.85 gm/kg,  $p=0.003$ ). There was no difference in length of stay (18.5 vs. 23.5 days,  $p=0.37$ ). The propensity score analysis included 100 patients. Following propensity scoring, significant differences in daily calorie (17.7 (4.6) vs 15.1 (5.1) kcal/kg/day,  $p=0.009$ ) and protein (1.03 (0.35) vs. 0.86 (0.38) g/kg/day,  $p=0.014$ ) provision remained. No differences in length of stay or other outcomes were noted in the propensity score analysis.

Conclusion: Initiation of enteral nutrition within 24 hours was not associated with improved outcomes in mechanically ventilated adults with COVID-19. No harm was detected either. Future research should seek to clarify optimal timing of enteral nutrition initiation in patients with COVID-19 requiring mechanical ventilation.

Keywords: COVID-19, SARS-CoV-2, critical care, enteral nutrition, length of stay, mechanical ventilation

## Introduction

Nutrition therapy plays an important role in the management of critically ill patients. Significant delay in initiation of enteral nutrition with accumulation of caloric deficit has been associated with more nosocomial infections, longer length of stay, and higher mortality.<sup>1-4</sup> Enteral nutrition is the delivery route of choice, as it promotes gut integrity and reduces systemic inflammation when compared to parenteral nutrition.<sup>5,6</sup> Guidelines recommend initiation of enteral nutrition within 24-48 hours of onset of critical illness in patients who cannot take in nutrition orally on their own.<sup>7,8</sup>

Although early nutrition intervention is widely accepted as standard of care, much is unknown about optimal nutrition therapy practices. Although some studies have found that early, aggressive calorie provision via enteral nutrition is beneficial, other studies have demonstrated that trophic or hypocaloric feeding may yield similar outcomes.<sup>9,10</sup> Early use of PN to meet 100% of caloric goals has not been shown to significantly improve outcomes.<sup>11,12</sup> The optimal amount of nutrition provision is especially controversial in patients with Acute Respiratory Distress Syndrome (ARDS) and Acute Lung Injury (ALI). One randomized controlled trial in patients with ALI found no benefit to full enteral nutrition as opposed to trophic feeding for the first 6 days of mechanical ventilation.<sup>13</sup> Another randomized controlled trial in patients with ALI was terminated early because full nutrition was associated with higher mortality.<sup>14</sup>

Like many other therapeutic modalities, the role of nutrition intervention in critically ill patients with SARS-CoV-2 (COVID-19) is poorly understood. Previous literature in ALI suggests that early, goal enteral nutrition does not improve outcomes. However, others have suggested that due to the severity of inflammation and illness observed in patients with COVID-19, risk of acquiring malnutrition may be higher.<sup>15</sup> Additionally, patients with COVID-19 may have had poor oral intake for days prior to admission. Statements from nutrition societies have recommended early consideration of parenteral nutrition in patients with COVID-19 that cannot tolerate enteral nutrition.<sup>16,17</sup> The objective of this study was primarily to associate provision of early enteral nutrition with outcomes in mechanically ventilated patients with COVID-19 and to secondarily describe practical nutrition practices in mechanically ventilated patients with COVID-19.

## Methods

This retrospective observational study was conducted at a single tertiary academic medical center. Patients were included if primarily admitted for COVID-19 pneumonia between March 1, 2020 - May 1, 2020 and required mechanical ventilation. Active COVID-19 infection was confirmed by positive polymerase chain reaction test on admission. Patients who were mechanically ventilated for greater than 48 hours at an outside institution prior to transfer were excluded. Patients that were mechanically ventilated for less than 48 hours and patients in which nutrition provision was unable to be accurately collected retrospectively from the medical record were also excluded. **This study was approved by the University of Michigan Institutional Review Board with waiver of informed consent (HUM 00181276).**

Institutional recommendations for nutrition support in critically ill patients with COVID-19 were established and disseminated early during the response to the pandemic. However, critically ill patients with COVID-19 were not managed via a nutrition support team or universal protocol. Decisions to initiate or hold nutrition therapy and how much enteral nutrition to provide were at the discretion of the multidisciplinary medical team caring for the patient. Each multidisciplinary team included a registered dietitian who evaluated the patients and made nutrition recommendations. Open label remdesivir was not readily available at the study institution during the study period. Additionally, early corticosteroids were not routinely administered for respiratory indications during the study period.

Patient demographic information, including age, sex, and relevant baseline comorbidities were collected. Dietitians assessed the nutrition status of all patients at baseline, when possible. A diagnosis of non-severe or severe malnutrition was made based on a local assessment tool adapted from American Society for Parenteral and Enteral Nutrition (ASPEN)/Academy of Nutrition and Dietetics criteria.<sup>18</sup> Presentation with gastrointestinal symptoms of COVID-19, including nausea, diarrhea, and lack of appetite was recorded. Days of fever in the first 7 days of mechanical ventilation were likewise recorded. To assess severity of illness, Acute Physiology and Chronic Health Evaluation (APACHE II) scores, Sequential Organ Failure Assessment (SOFA) scores, and PaO<sub>2</sub>: FiO<sub>2</sub> (P:F) ratios were

calculated for each patient using the most aberrant data points in the 24 hours following start of mechanical ventilation at the study institution.<sup>19,20</sup> Other therapies used in the treatment of COVID-19 were recorded including prone positioning, neuromuscular blockade, tocilizumab, inhaled nitric oxide, and vevo-venous extracorporeal membrane oxygenation (ECMO). Enrollment in a clinical trial for an experimental therapy for COVID-19 was also recorded.

Nutrition provision was recorded for the first 7 days following start of mechanical ventilation. Each day was defined as an individual calendar day from midnight to midnight. Total calorie intake from enteral nutrition, protein supplements, intravenous (IV) lipids via propofol, IV continuous dextrose infusions, and parenteral nutrition was totaled each day. Total protein intake was also calculated each day. Concomitant use of enteral nutrition and vasopressor administration was recorded, in addition to instances of bowel ischemia within 24 hours of these concomitant therapies. Due to staffing limitations and limits of staff exposure to patients, calorie intake was not recorded in patients that were able to tolerate a diet within the first 7 days following mechanical ventilation. Use of a pro-kinetic medication, defined as metoclopramide or erythromycin, was recorded. Days of severe hypophosphatemia (defined as a serum phosphorous level below 2 mg /dL) were collected. Days with hyperglycemia (defined as a blood glucose reading greater than 180 mg/d) and days with hypoglycemia (defined as a blood glucose reading less than 70 mg/dL) were recorded. Blood glucose levels were obtained from serum values, point of care readings, or blood gases. For patients receiving IV vitamin C, point of care readings were disregarded due to inaccuracies in readings. Ideal body weight was calculated using the Hamwi method. For this analysis, weight-based daily provision of calories and protein was calculated using ideal body weight for all patients. In clinical practice, registered dietitians recommended weight-based protein and calorie goals using ideal, actual, or adjusted weight based on clinical judgement. Modified NUTRIC scores were also retrospectively calculated for all patients using laboratory values drawn within 24 hours after initiation of mechanical ventilation; all modified NUTRIC scores were conducted without incorporating interleukin-6.<sup>21</sup>

Outcomes were compared in patients who received enteral nutrition within the first 24 hours of starting mechanical ventilation at the study institution to patients initiated on enteral nutrition at a later time. The primary outcome of the study was to compare

inpatient length of stay in patients that were initiated on enteral nutrition within the first 24 hours of mechanical ventilation compared later initiation. Secondary outcomes were measured at 60 days after admission and included inpatient mortality, days alive and free of mechanical ventilation, tracheostomy, and discharge on oxygen in patients that were initiated on enteral nutrition within the first 24 hours of mechanical ventilation compared later initiation.<sup>16</sup> Additionally, patients were compared based on receipt of at least 15 kcal/kg/day (using ideal body weight) of nutrition versus less than 15 kcal/kg/day. The value of 15 kcal/kg/day was chosen because has been suggested as a possible minimum goal calorie target for patients in the acute phase of critical illness.<sup>22</sup> Inpatient length of stay, inpatient mortality, days alive and free of mechanical ventilation, tracheostomy, and discharge on oxygen were also compared in patients who received at least 15 kcal/kg/day compared to less than 15 kcal/kg/day.

All statistical analyses were conducted using SPSS 26 (Armonk, NY). Baseline characteristics were analyzed using descriptive statistics. Continuous variables were evaluated using Student's T test and Mann-Whitney U test as appropriate based on normalcy of distribution. Nominal variables were compared using a Chi-Square test or Fisher's exact test. To control for baseline differences between groups, Propensity Score Matching was performed. Propensity score matching controlled for any of the following variables if the p-value was less than 0.1 when groups were compared at baseline: APACHE II, SOFA, age, BMI, neuromuscular blockade, tocilizumab, prone positioning, and ECMO. These variables were selected because differences either likely pre-disposed patients to a poor outcome or indicated a higher severity of illness. Patients were included in the final propensity score analysis based on a match tolerance of 0.1. After propensity score matching was completed, 100 patients were included for analysis. Following propensity score matching, no differences were found for standardized mean differences and each covariate assessed.

## Results

A total of 162 patients were included in the study. Six patients were excluded because they were mechanically ventilated for less than 48 hours and one patient was excluded due to missing data, leaving 155 patients for final analysis. Patient demographics

are represented in Table 1. A clear diagnosis of malnutrition was rarely made at the time of initiating mechanical ventilation due to lack information regarding prior energy intake or nutrition focused physical findings. The majority (72.9%, n=113) of patients had a modified NUTRIC score of  $\geq 5$ , indicating that they were at a high nutrition risk during admission. Data regarding treatment location and admission type can be found in the supplementary appendix.

Practices describing nutrition provision are displayed in Table 2. Enteral nutrition was started within 24 hours in 38.7% (n=60) of patients and within 48 hours in 69.7% (n=108) of patients. While mechanically ventilated, patients received a mean of 16.1 (SD 5.5) kcal/kg/day from all sources and 0.93 (SD 0.38) gm/kg/day of protein. Enteral nutrition was administered to 75.8% (n=116) of patients while a vasopressor infusion was concurrently administered, with no documented instances of bowel ischemia or emergent operations due to suspicion of bowel ischemia. No patients were initiated on parenteral nutrition within the first 7 days of mechanical ventilation. Hyperglycemia was common, occurring in 78.7% (n=122) of patients.

Patient outcomes are shown in Table 3. During the admission, 27.1% (n=42) of patients died. Median inpatient length of stay was 22.0 (IQR 23.8) days. A tracheostomy was performed in 16.8% (n=26) of patients and 26.5% (n=41) of patients were discharged on a new oxygen requirement. Only 40% (n=62) of patients were discharged home.

To assess the role of early nutrition, patients were divided by receipt of enteral nutrition within 24 hours of starting mechanical ventilation or after. The results of this comparison are shown in Table 4. Patients who received enteral nutrition within 24 hours of starting mechanical ventilation received significantly higher amounts of weight-based calories (17.5 (4.7) vs. 15.2 (5/8) kcal/kg/day, p=0.015) and protein (1.04 (0.35) vs. 0.85 (0.39) g/kg/day, p=0.003) daily. In patients who were initiated on enteral nutrition within 24 hours of mechanical ventilation, APACHE II scores (23.4 (6.4) vs. 26.8 (7.5), p=0.005) and SOFA scores (9.0 (2.8) vs. 10.3 (3.1), p=0.006) were significantly lower. The results of a propensity-matched analysis are shown in Table 5. Significant differences in daily calorie (17.7 (4.6) vs 15.1 (5.1) kcal/kg/day, p=0.009) and protein (1.03 (0.35) vs. 0.86 (0.38) g/kg/day, p=0.014) provision were still present after propensity score matching. Initiating

enteral nutrition within 24 hours of mechanical ventilation was not associated with a difference in length of stay (18.5 (25.0) vs. 24.6 (20.6) days,  $p=0.136$ ). Inpatient mortality (28% vs. 22%,  $p=0.644$ ), days alive and free of mechanical ventilation (31.4 (22.4) vs. 35.3 (20.0) days,  $p=0.36$ ), and the proportion of patients discharged home (40% vs. 50%,  $p=0.422$ ) also did not differ between groups. Moreover, administration of enteral nutrition within 24 hours was not associated with more days of hyperglycemia (4.3 (2.5) vs. 3.5 (2.7) days,  $p=0.129$ ) or number of patients that developed severe hypophosphatemia (24% vs. 12%,  $p=0.192$ ). A similar analysis was conducted comparing patients that received at least 15 kcal/kg/day of total calories or less, with no difference in outcomes found. Results of this analysis can be found in supplementary appendix.

## Discussion

In this retrospective study of mechanically ventilated patients with COVID-19 pneumonia, patients presented with a high severity of illness as evidenced by high baseline APACHE II and SOFA scores in addition to low P:F ratios. Over 70% of patients had a modified NUTRIC score  $\geq 5$ , which indicates that they may particularly benefit from aggressive nutrition therapy. Despite this, initiation of nutrition support within 24 hours of mechanical ventilation was not associated with improved clinical outcomes.

The results of this study conflict with previously published data. A meta-analysis conducted by Marik and colleagues demonstrated that early enteral nutrition was associated with fewer infectious complications and a reduction in inpatient length of stay.<sup>2</sup> Similarly, Doig, et al. also demonstrated a reduction in pneumonia and mortality when enteral nutrition was started within 24 hours in a separate meta-analysis.<sup>1</sup> Both of these meta-analyses primarily included studies conducted in surgical and trauma patients. Medical patients, such as those with COVID-19, may not derive a similar benefit from very early nutrition. Although less than half of the overall cohort was fed within 24 hours of mechanical ventilation, nearly 70% of patients were started on enteral nutrition within 48 hours, which aligns with current guideline recommendations. Thus, the relative delay in initiation of nutrition observed in this study may not have been long enough to contribute to patient outcomes. Future studies could investigate if delaying enteral nutrition for 48 hours in COVID patients affects patients outcomes. In this study, patients who were initiated



on enteral nutrition within 24 hours of mechanical ventilation compared to a later initiation of enteral nutrition. This selection of the 24 hour cutoff may be controversial. Current recommendations in COVID patients are to initiate enteral nutrition within 12 hours of starting mechanical ventilation. In the experience of the authors, the 12 hour threshold can be challenging to meet in the setting of surge staffing and the significance of a 24 hour threshold was thus studied instead.

Patients who received enteral nutrition within 24 hours did receive a greater number of weight-based calories and protein per day. However, the provision of both was still well below goal requirements and absolute differences in daily calorie and protein provision were small. These small differences may in part explain why early nutrition did not impact outcomes. However, similar outcomes were observed when patients were compared by those who received a daily calorie provision of at least 15 kcal/kg. The PERMIT trial compared permissive underfeeding to standard feeding in critically ill adults. Patients in the permissive underfeeding arm received 46% of goal calories compared to 71% in the standard care arm, with no differences in outcomes observed.<sup>23</sup> The EDEN trial compared initial trophic compared to full enteral feeding in patients with ALI and found that a greater calorie provision did not increase the number of ventilator-free days or reduce mortality.<sup>13</sup> Early parenteral nutrition in patients with contraindications to enteral nutrition has also been examined in a randomized control trial. Patients receiving early parenteral nutrition received more calories and protein per day, yet no differences mortality, length of stay, or complications were observed.<sup>11</sup>

Aggressive nutrition has also been shown to potentially harm outcomes in patients with ALI. The INTACT trial compared early intensive nutrition therapy to standard nutrition therapy. Patients in the intensive nutrition therapy arm received significantly more calories and protein. Mortality was 40% in the intensive nutrition group compared to 15.8% in the standard nutrition therapy group, a significant difference.<sup>14</sup> In the current study, there was no signal that early initiation of enteral nutrition or greater calorie provision was harmful in COVID-19 patients. Early nutrition could result in more hyperglycemia, which has been demonstrated to increase incidence of nosocomial complications.<sup>24,25</sup> However, patients who received nutrition within 24 hours of mechanical ventilation did not experience more days of hyperglycemia after propensity score analysis was incorporated.

To the knowledge of the authors, this is one of the first reports describing nutrition support for COVID-19 patients in the United States. The results of this study have implications on the care of critically ill COVID-19 patients. Despite a high severity of illness, it was still possible to deliver enteral nutrition to these patients often within 48 hours of being intubated. The presence of gastrointestinal symptoms on admission did not appear to limit the ability to provide enteral nutrition once mechanically ventilated in this cohort. However, it is unclear to determine the severity of these gastrointestinal symptoms given the retrospective nature of this study. Over 70% of patients received enteral nutrition despite also receiving vasopressor medications. This practice was associated with an excellent safety profile as no patients developed bowel ischemia requiring intervention. Although greater calorie provision was not associated with improved outcomes in this study, very few patients received 100% of goal calories. Previous trials have demonstrated that it is very challenging to provide 100% of goal calories with conventional enteral nutrition strategies alone.<sup>12,13</sup> Volume-based enteral nutrition strategies or supplemental parenteral nutrition may be necessary to provide 100% of caloric goals in this patient population. More research is necessary to clarify the role of parenteral nutrition in patients with COVID-19. An ongoing prospective, observational trial examining caloric needs in mechanically ventilated COVID-19 patients using indirect calorimetry should provide more information.<sup>26</sup>

This study does have several limitations. Nutrition assessment was limited by an inability to examine patients and perform nutrition-focused physical assessments. Oral intake prior to admission was likewise often unclear. It is possible that many patients presented after several days of suffering from COVID-19 with poor oral intake and they were already at a significant calorie deficit prior to the initiation of mechanical ventilation. A supplemental multivitamin was not routinely administered to patients during the first seven days of mechanical ventilation. Due to the retrospective nature of this project and documentation inconsistencies, gastric residual volumes, vomiting, and the decision to stop enteral nutrition based on gastric residual volumes could not be assessed. Glucose control was challenging in these patients. Insulin infusions were generally avoided due to the need to frequently enter patient rooms to check glucose levels and adjust infusion rates. Overall incidence of hyperglycemia in these patients was nevertheless higher than ideal. Due to risk of infection spread and surge staffing, daily calorie counts were not recorded on patients

that were extubated and eating. This study is subject to the limitations inherent to retrospective evaluations (bias, confounding variables, etc.). However, we attempted to minimize the impact of these limitations through the use of propensity score matching. Following propensity score matching, differences in calorie and protein provision still existed despite no apparent differences in severity of illness. These differences in enteral nutrition provision could be attributed to heterogeneity in practice among different providers. However, other undetermined variables that were not collected may have contributed to nutrition provision as well.

The natural history of COVID-19 infections and the complexity of presentation is not yet well understood. Nutritional intake may only play a small role in survival for the most critically ill patients with COVID-19 pneumonia. The results of this study suggest that initiation of enteral nutrition within 24 hours of starting mechanical ventilation may not improve outcomes in COVID-19 patients. Additional studies are necessary to further clarify the ideal time to initiate enteral nutrition in critically ill patients with COVID-19.

## References

1. Doig GS, Heighes PT, Simpson F, Sweetman EA, Davies AR. Early enteral nutrition, provided within 24 h of injury or intensive care unit admission, significantly reduces mortality in critically ill patients: A meta-analysis of randomised controlled trials. *Intensive Care Med.* 2009;35(12):2018-2027. doi:10.1007/s00134-009-1664-4
2. Marik PE, Zaloga GP. Early enteral nutrition in acutely ill patients: A systematic review. *Crit Care Med.* 2001;29(12):2264-2270. <https://doi.org/10.1097/00003246-200112000-00005>.
3. Artinian V, Krayem H, DiGiovine B. Effects of Early Enteral Feeding on the Outcome of Critically Ill Mechanically Ventilated Medical Patients. *Chest.* 2006;129:960-967. <https://doi.org/10.1378/chest.129.4.960>.
4. Elke G, van Zanten ARH, Lemieux M, et al. Enteral versus parenteral nutrition in critically ill patients: An updated systematic review and meta-analysis of randomized controlled trials. *Crit Care.* 2016;20(1). doi:10.1186/s13054-016-1298-1
5. Jabbar A, Chang WK, Dryden GW, McClave SA. Gut Immunology and the Differential Response to Feeding and Starvation. *Nutr Clin Pract.* 2003;18(6):461-482. doi:10.1177/0115426503018006461
6. Windsor ACJ, Kanwar S, Li AGK, et al. Compared with parenteral nutrition, enteral feeding attenuates the acute phase response and improves disease severity in acute pancreatitis. *Gut.* 1998;42(3):431-435. doi:10.1136/gut.42.3.431
7. McClave SA, Taylor BE, Martindale RG, et al. Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *J Parenter Enter Nutr.* 2016;40(2):159-211. doi:10.1177/0148607115621863
8. Singer P, Blaser AR, Berger MM, et al. ESPEN guideline on clinical nutrition in the intensive care unit. *Clin Nutr.* 2019;38(1):48-79. doi:10.1016/j.clnu.2018.08.037
9. Barr J, Hecht M, Flavin KE, Khorana A, Gould MK. *Outcomes in Critically Ill Patients Before and After the Implementation of an Evidence-Based Nutritional Management Protocol\**. Vol 125.; 2004. <https://doi.org/10.1378/chest.125.4.1446>.
10. Dickerson RN, Boschert KJ, Kudsk KA, Brown RO. Hypocaloric Enteral Tube Feeding in Critically Ill Obese Patients. *Nutrition.* 2002;18:241-246. [https://doi.org/10.1016/S0899-9007\(01\)00793-6](https://doi.org/10.1016/S0899-9007(01)00793-6).
11. Doig GS, Simpson F, Sweetman EA, et al. Early parenteral nutrition in critically ill patients with short-term relative contraindications to early enteral nutrition: A randomized controlled trial. *JAMA - J Am Med Assoc.* 2013;309(20):2130-2138. doi:10.1001/jama.2013.5124
12. Casaer MP, Mesotten D, Hermans G, et al. Early versus late parenteral nutrition in critically ill adults. *N Engl J Med.* 2011;365(6):506-517. doi:10.1056/NEJMoa1102662

13. Rice TW, Wheeler AP, Thompson BT, et al. Initial trophic vs full enteral feeding in patients with acute lung injury: The EDEN randomized trial. *JAMA - J Am Med Assoc.* 2012;307(8):795-803. doi:10.1001/jama.2012.137
14. Braunschweig CA, Sheean PM, Peterson SJ, et al. Intensive nutrition in acute lung injury: A clinical trial (INTACT). *J Parenter Enter Nutr.* 2015;39(1):13-20. doi:10.1177/0148607114528541
15. Yu PJ, Cassiere H, DeRosa S, Bocchieri K, Yar S, Hartman A. Hypermetabolism and COVID-19. *JPEN J Parenter Enteral Nutr.* Published online June 19, 2020. doi:10.1002/jpen.1948
16. Martindale R, McClave Md -Professor SA. *Nutrition Therapy in the Patient with COVID-19 Disease Requiring ICU Care.* Available from: [https://www.nutritioncare.org/Guidelines\\_and\\_Clinical\\_Resources/Resources\\_for\\_Clinicians\\_Caring\\_for\\_Patients\\_with\\_Coronavirus/](https://www.nutritioncare.org/Guidelines_and_Clinical_Resources/Resources_for_Clinicians_Caring_for_Patients_with_Coronavirus/)
17. Barazzoni R, Bischoff SC, Breda J, et al. ESPEN expert statements and practical guidance for nutritional management of individuals with SARS-CoV-2 infection. *Clin Nutr.* Published online 2020. doi:10.1016/j.clnu.2020.03.022
18. White J V., Guenter P, Jensen G, Malone A, Schofield M. Consensus statement: Academy of nutrition and dietetics and American society for parenteral and enteral nutrition: Characteristics recommended for the identification and documentation of adult malnutrition (undernutrition). *J Parenter Enter Nutr.* 2012;36(3):275-283. doi:10.1177/0148607112440285
19. Knaus W, Draper E, Wagner D, Zimmerman J. APACHE II: a severity of disease classification system. *Crit Care Med.* 1985;13(10):818-829.
20. Vincent J, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. *Intensive Care Med.* 1996;22:707-710. <https://doi.org/10.1007/bf01709751>.
21. Heyland DK, Dhaliwal R, Jiang X, Day AG. Identifying critically ill patients who benefit the most from nutrition therapy: the development and initial validation of a novel risk assessment tool. *Crit Care.* 2011;15:R268. doi.org/10.1016/j.clnu.2015.01.015.
22. Wischmeyer PE. Tailoring nutrition therapy to illness and recovery. *Crit Care.* 2017;21. doi:10.1186/s13054-017-1906-8
23. Arabi YM, Aldawood AS, Haddad SH, et al. Permissive underfeeding or standard enteral feeding in critically ill adults. *N Engl J Med.* 2015;372(25):2398-2408. doi:10.1056/NEJMoa1502826
24. Van den Berghe G, Wilmer A, Meersseman W, et al. Intensive Insulin Therapy in the Medical ICU. *NEJM.* 2006;354(2):449-461. doi.org/10.1056/NEJMoa052521.
25. Van den Berge G, Wouters P, Weekers F, et al. Intensive Insulin Therapy in Critically Ill Patients. *N Engl J Med.* 2001;345(19):1359-1367. doi.org/10.1056/NEJMoa011300.

26. Longitudinal Energy Expenditure and Metabolic Effects in Patients With COVID-19 (LEEP-COVID). Accessed July 19, 2020.  
<https://clinicaltrials.gov/ct2/show/NCT04350073>

Author Manuscript

**Table 1.** Baseline Characteristics

Characteristic	n=155
Age, mean (SD)	60.3 (13.8)
Male, n (%)	103 (66.5)
Weight kg, median (IQR)	98.5 (34.3)
BMI, median (IQR)	33.2 (12.6)
Comorbidities	
Chronic Obstructive Pulmonary Disease, n (%)	17 (11.0)
Asthma, n (%)	23 (14.8)
Diabetes, n (%)	85 (54.9)
Pre-Diabetes, n (%)	8 (5.2)
Hypertension, n (%)	113 (72.9)
Chronic Kidney Disease, n (%)	36 (23.2)
End Stage Renal Disease, n (%)	4 (2.6)
Immunocompromised, n (%)	21 (13.5)
Presentation with GI symptoms, n (%)	44 (28.3)
Temperature $\geq 38.3^{\circ}\text{C}$ for at least one day, n (%)	105 (67.7)
PaO <sub>2</sub> /FiO <sub>2</sub> Ratio, median (IQR)	128 (80)
PaO <sub>2</sub> /FiO <sub>2</sub> Ratio <100, n (%)	44 (28.4)
APACHE II, mean (SD)	25.5 (7.3)
SOFA, mean (SD)	9.9 (3.1)
Albumin (gm/dL), median (IQR)	3.2 (0.6)
Modified NUTRIC $\geq 5$ , n (%)	113 (72.9)
Unclear nutrition status	152 (98.1)

Diagnosis of non-severe malnutrition	3 (1.9)
Diagnosis of severe malnutrition	0 (0)

APACHE, Acute Physiology and Chronic Health Evaluation; BMI, Body Mass Index; FiO<sub>2</sub>, Fraction of Inspired Oxygen; GI, Gastrointestinal; IQR, Interquartile Range; NUTRIC, Nutrition Risk in Critically Ill; PaO<sub>2</sub>, Partial Pressure of Oxygen; SOFA, Sequential Organ Failure Assessment; SD, Standard Deviation

Author Manuscript



**Table 2.** Nutrition Provision

Characteristic	n=155
Daily Calories (kcal/kg IBW), mean (SD)	16.1 (5.5)
Daily Protein (grams/kg IBW), mean (SD)	0.93 (0.38)
Time to Starting Enteral Nutrition (days), median (IQR)	1.16 (0.52)
Started Enteral Nutrition Within 24 Hours, n (%)	60 (38.7)
Started Enteral Nutrition Within 48 Hours, n (%)	108 (69.7)
Received Enteral Nutrition While on Vasopressor, n (%)	116 (74.8)
Developed Hyperglycemia, n (%)	122 (78.7)
Days with Hyperglycemia, mean (SD)	3.86 (2.69)
Developed Hypoglycemia, n (%)	18 (11.6)
Developed Severe Hypophosphatemia, n (%)	24 (15.5)
Enteral Tube Location	
Gastric Only, n (%)	140 (93.5)
Post-Pyloric Only, n (%)	2 (1.4)
Both Gastric and Post-Pyloric, n (%)	2 (1.4)
Unclear, n (%)	6 (3.9)
Received Prokinetic Medication, n (%)	10 (6.5)
Received Multivitamin, n (%)	19 (12.3)
Enteral Nutrition Formula	
Nutren 1.5 <sup>®</sup> , n (%)	118 (76.1)
Nutren 2.0 <sup>®</sup> , n (%)	8 (5.2)
Novasource Renal <sup>®</sup> , n (%)	62 (40)
Replete <sup>®</sup> , n (%)	2 (1.3)
Nepro <sup>®</sup> , n (%)	2 (1.3)

IBW, Ideal Body Weight; IQR, Interquartile Range; SD, Standard Deviation

# Author Manuscript

**Table 3.** Patient Outcomes

Outcome	n=155
Inpatient Mortality, n (%)	42 (27.1)
Inpatient Length of Stay, median (IQR)	22.0 (23.8)
ICU Length of Stay, median (IQR)	14.3 (14.4)
Days Alive and Ventilator Free at 60 days, mean (SD)	32.5 (21.3)
Required tracheostomy, n (%)	26 (16.8)
Re-admission to ICU following ICU discharge, n (%)	7 (4.5)
Discharge on new oxygen requirement	41 (26.5)
Discharge Location	
Home, n (%)	62 (40)
Rehab Facility, n (%)	26 (16.7)
Skilled Nursing Facility, n (%)	21 (13.6)
Long-term acute care hospital, n (%)	1 (0.6)
Still admitted to hospital, n (%)	4 (2.6)
Deceased, n (%)	42 (27.1)

ICU, Intensive Care Unit; IQR, Interquartile Range; SD, Standard Deviation

Table 4. Comparison By Initiation Time of Enteral Nutrition (n=155)

	Started Within 24 Hours of Starting Mechanical Ventilation (n=60)	Started After 24 Hours of Starting Mechanical Ventilation (n=95)	P value
Age (years), mean (SD)	58.3 (14.1)	61.5 (13.6)	0.166
Weight (kg), median (IQR)	100 (33.3)	96 (36)	0.481
BMI (median)	33.2 (12.8)	33.2 (11.2)	0.530
APACHE II, mean (SD)	23.4 (6.4)	26.8 (7.5)	0.005
SOFA, mean (SD)	9.0 (2.8)	10.4 (3.1)	0.006
NUTRIC $\geq$ 5, n (%)	38 (63.3)	75 (78.9)	0.042
PaO <sub>2</sub> /FiO <sub>2</sub> Ratio, median (IQR)	142 (81.5)	118 (83)	0.594
Neuromuscular Blockade, n (%)	19 (31.7)	29 (30.5)	1.00
Inhaled Nitric Oxide, n (%)	10 (16.7)	10 (10.5)	0.327
Prone Positioning, n (%)	15 (25)	26 (27.4)	0.852
ECMO, n (%)	3 (5)	3 (3.2)	0.677
Tocilizumab, n (%)	25 (41.7)	45 (47.4)	0.512
Daily Calories (kcal/kg/day), mean (SD)	17.5 (4.7)	15.2 (5.8)	0.015
Daily Protein (gm/kg/day), mean (SD)	1.04 (0.34)	0.85 (0.39)	0.003
Time to Starting Enteral Nutrition (days), median (IQR)	0.65 (0.47)	1.99 (2.43)	0.006
Inpatient Length of Stay (days), median (IQR)	18.5 (24.4)	23.5 (21.5)	0.37
Mortality, n (%)	17 (28.3)	25 (26.3)	0.842

ICU Length of Stay (days), median (IQR)	12.9 (10.1)	14.7 (17.9)	0.07
Days Alive and Ventilator Free at 60, mean (SD)	31.3 (22.3)	33.3 (20.7)	0.743
Tracheostomy, n (%)	7 (11.7)	19 (20)	0.194
Discharge to Home, n (%)	24 (40)	38 (40)	1.00
Days of Hyperglycemia, mean (SD)	4.4 (2.6)	3.5 (2.7)	0.048
Hypoglycemia, n (%)	8 (13.3)	10 (10.5)	0.615
Severe Hypophosphatemia, n (%)	14 (23.3)	10 (10.5)	0.04

APACHE, Acute Physiology and Chronic Health Evaluation; BMI, Body Mass Index; ECMO, Extracorporeal Membrane Oxygenation; FiO<sub>2</sub>, Fraction of Inspired Oxygen; ICU, Intensive Care Unit; IQR, Interquartile Range; NUTRIC, Nutrition Risk in Critically Ill; PaO<sub>2</sub>, Partial Pressure of Oxygen; SOFA, Sequential Organ Failure Assessment; SD, Standard Deviation

Author Manuscript

Table 5. Propensity-Matched Comparison By Start of Enteral Nutrition Within 24 Hours (n=100)

	Started within 24 Hours (n=50)	Started After 24 Hours (n=50)	P value
Age (years), mean (SD)	58.9 (14.1)	59.6 (13.8)	0.544
Weight (kg), median (IQR)	100 (36.2)	95.8 (37.5)	0.341
BMI (median)	34.5 (13.8)	33.3 (8.7)	0.424
APACHE II, mean (SD)	23.5 (6.4)	25.3 (6.0)	0.160
SOFA, mean (SD)	9.4 (2.9)	10.0 (2.6)	0.232
NUTRIC $\geq$ 5, n (%)	31 (62)	36 (72)	0.395
PaO <sub>2</sub> /FiO <sub>2</sub> Ratio, median (IQR)	141 (75.9)	132 (88.8)	0.758
Neuromuscular Blockade, n (%)	18 (36)	17 (34)	1.00
Inhaled Nitric Oxide, n (%)	8 (16)	4 (8)	0.357
Prone Positioning, n (%)	11 (22)	13 (26)	0.815
ECMO, n (%)	2 (4)	2 (4)	1.00
Tocilizumab, n (%)	22 (44)	24 (48)	0.841
Daily Calories (kcal/kg/day), mean (SD)	17.7 (4.6)	15.1 (5.1)	0.009
Daily Protein (gm/kg/day), mean (SD)	1.03 (0.35)	0.86 (0.38)	0.014
Time to Starting Enteral Nutrition (days), median (IQR)	0.65 (0.47)	1.97 (2.68)	<0.001
Inpatient Length of Stay (days), median (IQR)	18.5 (25.0)	24.6 (20.6)	0.136
Mortality, n (%)	4 (28)	11 (22)	0.644
ICU Length of Stay (days),	13.0 (11.4)	15 (18.6)	0.389

median (IQR)			
Days Alive and Ventilator Free at 60, mean (SD)	31.4 (22.4)	35.3 (20.0)	0.36
Tracheostomy, n (%)	6 (12)	11 (22)	0.287
Discharge to Home, n (%)	20 (40)	25 (50)	0.422
Days of Hyperglycemia, mean (SD)	4.3 (2.5)	3.5 (2.7)	0.129
Hypoglycemia, n (%)	5 (10)	4 (8)	1.00
Severe Hypophosphatemia, n (%)	12 (24)	6 (12)	0.192

APACHE, Acute Physiology and Chronic Health Evaluation; BMI, Body Mass Index; ECMO, Extracorporeal Membrane Oxygenation; FiO<sub>2</sub>, Fraction of Inspired Oxygen; ICU, Intensive Care Unit; IQR, Interquartile Range; NUTRIC, Nutrition Risk in Critically Ill; PaO<sub>2</sub>, Partial Pressure of Oxygen; SOFA, Sequential Organ Failure Assessment; SD, Standard Deviation