

REVIEW ARTICLE

Axon–glia interactions in the ascending auditory system

David C. Kohrman | Beatriz C. Borges | Luis R. Cassinotti |
Lingchao Ji | Gabriel Corfas 

Department of Otolaryngology - Head and Neck Surgery, Kresge Hearing Research Institute, University of Michigan, Ann Arbor, MI, USA

Correspondence

Gabriel Corfas, Department of Otolaryngology - Head and Neck Surgery, Kresge Hearing Research Institute, The University of Michigan, Medical Sciences I Building, Rm. 5428, 1150 West Medical Center Drive, Ann Arbor, MI 48109-5616, USA.

Email: corfas@med.umich.edu

Funding information

National Institute on Deafness and Other Communication Disorders, Grant/Award Number: R01 DC018500 and R21 DC017916; Lynn and Ruth Townsend Professorship of Communication Disorders

Abstract

The auditory system detects and encodes sound information with high precision to provide a high-fidelity representation of the environment and communication. In mammals, detection occurs in the peripheral sensory organ (the cochlea) containing specialized mechanosensory cells (hair cells) that initiate the conversion of sound-generated vibrations into action potentials in the auditory nerve. Neural activity in the auditory nerve encodes information regarding the intensity and frequency of sound stimuli, which is transmitted to the auditory cortex through the ascending neural pathways. Glial cells are critical for precise control of neural conduction and synaptic transmission throughout the pathway, allowing for the precise detection of the timing, frequency, and intensity of sound signals, including the sub-millisecond temporal fidelity is necessary for tasks such as sound localization, and in humans, for processing complex sounds including speech and music. In this review, we focus on glia and glia-like cells that interact with hair cells and neurons in the ascending auditory pathway and contribute to the development, maintenance, and modulation of neural circuits and transmission in the auditory system. We also discuss the molecular mechanisms of these interactions, their impact on hearing and on auditory dysfunction associated with pathologies of each cell type.

KEYWORDS

astrocytes, auditory system, glia, hearing loss, neuron–glia interactions, oligodendrocytes, satellite glial cells, Schwann cells, supporting cells

Abbreviations: ABR, auditory brainstem response; ALDH1L1, aldehyde dehydrogenase 1 family member L1; AMPA, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid; AN, auditory nerve; AP, action potential; APD, auditory processing disorder; ASD, autism spectrum disorder; AVCN, anteroventral cochlear nucleus; BDNF, brain-derived neurotrophic factor; CMT, Charcot–Marie–Tooth; CN, cochlear nucleus; CNS, central nervous system; Cx, connexin; dB SPL, decibels sound pressure level; DTI, diffusor tensor imaging; ErbBR, ErbB receptor; FGF, fibroblast growth factor; GABA, gamma aminobutyric acid; GBC, globular bushy cell; GBS, Guillain–Barre syndrome; GFAP, glial fibrillary acidic protein; GLAST, glutamate/aspartate transporter (also known as EAAT1); GLT, glutamate transporter; HC, hair cell; HHL, hidden hearing loss; IC, inferior colliculus; IGF1, insulin-like growth factor-1; IHC, inner hair cell; IID, interaural intensity difference; IP3, inositol trisphosphate; ITD, interaural time difference; KCC4, K/Cl cotransporter 4; KCNQ4, potassium voltage-gated channel subfamily Q member 4; LSO, lateral superior olive; MBP, myelin basic protein; MGN, medial geniculate nucleus; MIF, macrophage migration inhibitory factor; MNTB, medial nucleus of the trapezoid body; MPZ, myelin protein zero; MRI, magnetic resonance imaging; MS, multiple sclerosis; MSO, medial superior olive; NL, nucleus laminaris; NMDA, n-methyl-d-aspartate; NoR, nodes of Ranvier; NRG1, neuregulin 1; NT3, neurotrophin 3; OHC, outer hair cell; OL, oligodendrocyte; OPC, oligodendrocyte precursor cell; PLP1, proteolipid protein 1; PMD, Pelizaeus–Merzbacher disease; PMP22, peripheral myelin protein 22; PNS, peripheral nervous system; SBC, spherical bushy cell; SGC, satellite glial cell; SGN, spiral ganglion neuron; SOC, superior olivary complex; Sox10, SRY-Box Transcription Factor 10; SuppC, supporting cell; TNF- α , tumor necrosis factor alpha; TRK, tropomyosin receptor kinase; VCN, ventral cochlear nucleus.

David Kohrman, Beatriz C. Borges, Luis Cassinotti, and Lingchao Ji denotes equal contribution.

1 | INTRODUCTION

The vertebrate auditory system is the part of the nervous system dedicated to capturing environmental sounds and processing them to extract information about the surroundings and allow for communication between individuals. Using unique mechanosensory cells and specialized synaptic connections, the auditory system is exceptionally proficient at encoding and transmitting a high-fidelity representation of sound frequency and intensity. In mammals, the auditory pathway starts with the organ of Corti located within the cochlea in the inner ear (Figure 1a). There, inner hair cells (IHCs) transduce sound waves into neural activity through excitatory synapses onto the primary auditory neurons, the type I spiral ganglion neurons (SGNs). These neurons then propagate auditory information to the auditory cortex through an ascending pathway consisting

of several nuclei. At every single stage of the pathway, neural cells are in close contact with glia, including the glia-like supporting cells (SuppCs) of the organ of Corti that surround hair cells (HCs) and their synapses with SGNs, Schwann cells that myelinate the peripheral portion of the auditory nerve (AN), satellite glial cells (SGCs) that surround SGN cell bodies, oligodendrocytes (OLs) that myelinate the central portion of the AN as well as axons in central auditory pathways, and astrocytes that wrap around synapses in the central portion of the ascending auditory pathway (Figure 1b). This diverse set of glial cells is critical for precise control of axonal conduction and synaptic transmission at specialized synapses throughout the pathway, facilitating the detection of sub-millisecond changes in the timing of sound signals. This extraordinary temporal fidelity is necessary for tasks such as sound localization, and in humans, for processing complex sounds including speech and music.




FIGURE 1 The auditory pathway. (a) Schematic representation of the auditory periphery, including glial cells associated with SGNs (supporting cells, Schwann cells, satellite glial cells, and oligodendrocytes). Oligodendrocyte myelination of SGN axons begins at the CNS–PNS transition zone (arrows). (b) Schematic representation of the central auditory pathway highlighting ITD and IID circuitry at the level of the pons. VCN SBCs (blue) extend excitatory projections to the LSO and MSO in the SOC. VCN GBCs (green) extend excitatory projections to the MNTB, which then send inhibitory projections (red) to the LSO and MSO. Auditory information then passes through several subsequent nuclei including the IC before reaching the MGN and projecting to the auditory cortex. (c) Auditory system function can be measured using auditory brainstem responses (ABR). This method uses electrodes to detect evoked electrical potentials along the early auditory pathway, from the cochlea through the midbrain. The summating potential (SP) reflects activation of IHCs, peak I corresponds to evoked potentials in SGNs while later peaks generally correspond to evoked potentials in CN (peak II), SOC (peak III), and IC (peaks IV and V) in the mouse (Henry, 1979; Land et al., 2016). The magnitude of peak I amplitudes correlates with the number and synchronous firing rate of the SGN fibers. (d) Calyx of held synapse in MNTB is formed by a highly fenestrated calyx ending of the presynaptic GBC axon and the postsynaptic soma of the MNTB principal cell (principal cell projections not shown). Astrocytes are in close proximity to the principal cell soma and extend processes into the spaces between calyceal elements (Ford et al., 2009; Reyes-Haro et al., 2010). Figure 1b is adapted from (Cope et al., 2015). Figure 1c is adapted from (Kohrman et al., 2019). AVCN, anteroventral cochlear nucleus; GBC, globular bushy cell; IHC, inner hair cell; LSO, lateral superior olive; MGN, medial geniculate nucleus; MNTB, medial nucleus of the trapezoid body; MSO, medial superior olive; SBC, spherical bushy cell; SGN, spiral ganglion neuron; SOC, superior olivary complex

This review aims at providing glial biologists information about the auditory system, and auditory scientists information about the roles of glia in the auditory system. We focus on the interactions that glia have with sensory HCs and ascending auditory neurons; the roles these interactions play in the development, maintenance, and modulation of auditory neural circuits; and their impact on auditory disorders. We also discuss our current understanding of the reciprocal neural–glial interactions that influence development and maintenance of hearing and review the auditory dysfunction associated with pathologies of each cell type.

2 | AUDITORY SYSTEM GLIA

2.1 | Supporting cells, the glia of the organ of Corti

The sensory epithelium of the cochlea, the organ of Corti, contains two types of mechanoreceptive HCs—IHC and outer (OHC) hair cells. IHCs are the principal mechanoreceptor cells for sound detection, whereas OHCs are electromotile cells that amplify the vibrational motion of the organ of Corti and control the gain of auditory responses (Fettiplace, 2017). IHCs form ribbon-type glutamatergic synapses with the peripheral axons of type I SGNs, bipolar neurons which give rise to approximately ninety percent of myelinated axons in both the central and peripheral parts of the AN. Specialized exocytosis and endocytosis machinery present at the presynaptic ribbon support sound-evoked synaptic release that is robust and possesses high temporal fidelity (Moser et al., 2019). The remaining ten percent of AN-myelinated axons belong to brainstem efferent neurons (Brown, 1987; Campbell & Henson, 1988; Ehret, 1979). Each type I SGN innervates a single IHC, while each IHC makes synaptic contact with approximately 10 to 30 type I SGN fibers (Coate et al., 2019). The AN also contains unmyelinated axons that belong to type II SGNs, smaller caliber afferent neurons that innervate OHC. While the roles of type II afferents remain poorly understood, it has been proposed that they signal cochlear damage (Liu, Glowatski, et al., 2015).

Hearing sensitivity is often assessed by measuring auditory brainstem responses (ABRs) to transient sounds. ABRs are auditory evoked far-field responses detected by head-mounted electrodes that represent the synchronous electrical activity of the auditory system at multiple locations along the pathway. A typical ABR waveform from the mouse (Figure 1c) consists of five major peaks, with peak I corresponding to activity generated by the SGNs within 2 msec following the sound stimulus, and the later peaks II through V corresponding to activity generated by neurons in successive nuclei of the auditory hindbrain. Thresholds are

determined by the lowest intensity of sound that produces a recognizable waveform. The peak I amplitude correlates with the number and synchronous firing rate of the type I SGN fibers and can be influenced by changes in IHC-AN synapse numbers and by myelin alterations (see below for details and references). Reductions in ABR peak I amplitudes are considered to reflect a reduction in the efficiency of transmission of information from the cochlea to the CNS.

SuppCs in the organ of Corti were traditionally considered to be connective tissue providing structural and homeostatic support. However, recent work has shown that these cells exhibit many characteristics of glia and play glia-like roles (Wan et al., 2013). The IHCs and their synapses are surrounded by SuppCs known as inner border and inner phalangeal cells, which express molecules that regulate synaptic development and may function in a manner analogous to that of perisynaptic Schwann cells (Wan et al., 2013). Below, we discuss the diverse roles that SuppCs play in development and function of the cochlea.

2.1.1 | Spontaneous cochlear activity prior to the onset of hearing

While sound-evoked responses are not present in early postnatal cochlea of rodents, type I SGNs display spontaneous electrical activity (Jones et al., 2007; Walsh & Romand, 1992). Although IHC intrinsic firing properties likely contribute to early spontaneous activity of SGNs (Johnson et al., 2011, 2012), several studies indicate a role for SuppCs in modulating this spontaneous activity via ATP-dependent signaling. Beginning at birth (P0), ATP released from SuppCs within the developing cochlea activates purinergic receptors on IHCs, leading to IHC depolarization and glutamate release at ribbon synapses, which in turn triggers coordinated bursts of action potentials (APs) in the AN (Tritsch & Bergles, 2010; Tritsch et al., 2007, 2010). The released ATP also activates purinergic autoreceptors on SuppCs, inducing K⁺ efflux that potentiates IHC depolarization (Babola et al., 2020; Wang et al., 2015). Later in postnatal development, ATP release declines along with spontaneous activity levels in SGNs (Tritsch & Bergles, 2010). Spontaneous activity is important for the maturation of auditory neurons (Seal et al., 2008; Shrestha et al., 2018; Sun et al., 2018) as well as refinement of synaptic connections in central auditory pathways (Clause et al., 2014; Erazo-Fischer et al., 2007; McKay & Oleskevich, 2007).

2.1.2 | Neuronal survival and outgrowth

HCs have traditionally been considered critical for SGN survival, at least in part due to their expression of

neurotrophins such as brain-derived neurotrophic factor (BDNF) and neurotrophin-3 (NT3) (Green et al., 2012). Indeed, loss of either BDNF or NT3, or their respective cognate receptors *trkB* and *trkC*, result in embryonic loss of SGNs, with NT3 KOs exhibiting virtually complete loss of type I SGNs while loss of BDNF results in loss of type II SGNs (Bianchi et al., 1996; Ernfors et al., 1995; Fariñas et al., 2001; Schimmang et al., 1995). Nonetheless, a large body of evidence suggests that cochlear SuppCs play an equally important role in SGN biology, including acting as an additional source of BDNF and NT3 in the cochlea (Sugawara et al., 2007; Wiechers et al., 1999). Furthermore, data from aged or injured animal indicate a strong correlation between SGN survival and the number of remaining SuppCs, rather than HCs (Leake & Hradek, 1988; McFadden et al., 2004; Sugawara et al., 2005; Xu et al., 1993). Notably, experimental ablation of IHCs in two independent mouse model systems, both of which left SuppCs intact, demonstrated survival of SGNs for at least 3 to 4 months post-ablation, suggesting that the intact SuppCs are capable of providing neurotrophic support in the absence of HCs (Tong et al., 2015; Zilberstein et al., 2012). Several studies of human temporal bones have also reported a positive correlation between SuppC and SGN survival in the cochlea (Johnsson, 1974; Suzuka & Schuknecht, 1988). Finally, developmental expression gradients of BDNF and NT3 observed along the length of the cochlea in rodents, including SuppCs, is likely to influence the extension and refinement of fibers toward target HCs (Huang et al., 2007; Sugawara et al., 2007; Wiechers et al., 1999). These neurotrophin gradients may also modulate synaptic and ion channel protein expression in SGNs and thereby contribute to differences in firing properties in basal versus more apical SGNs (Adamson et al., 2002; Flores-Otero et al., 2007; Zhou et al., 2005).

The ability of SuppCs to impact SGN survival also appears to depend on Neuregulin 1-ErbB receptor (NRG1-ErbBR) signaling, as blockade of SuppC ErbBR signaling results in SGN loss in spite of the presence of intact HCs and SuppCs (Stankovic et al., 2004). Although SGN loss in this model was originally attributed to an associated decrease in cochlear NT3 expression, more recent work using conditional knockout mice demonstrated normal SGN survival following postnatal loss of NT3 or BDNF in either HCs or SuppCs (Wan et al., 2014). These results suggest that residual neurotrophin expression from either cell type is sufficient for maintenance of SGNs. Other trophic factors that are expressed in postnatal SuppCs, such as insulin-like growth factor-1 (IGF1) or macrophage migration inhibitory factor (MIF), may also contribute to neuronal preservation, as knockout mice for either IGF1 or MIF exhibit SGN loss or altered innervation (Bank et al., 2012; Camarero et al., 2001; Elkon et al., 2015; Okano et al., 2011).

2.1.3 | Regulation of synapse formation and regeneration

While postnatal NT3 expression by SuppCs is not required for SGN survival, it is nevertheless critical for normal IHC-AN synaptogenesis. Neonatal knockout of NT3 in cochlear SuppCs results in a significant decrease in the number of IHC ribbon synapses and impaired hearing sensitivity. Conversely, overexpression of NT3 in cochlear SuppCs during the neonatal period results in increases in the number of synapses and better hearing sensitivity (Wan et al., 2014).

Interestingly, studies in numerous animal models have demonstrated that moderate noise exposures can result in the loss of a subset of IHC-AN synapses and an associated phenotype known as “hidden hearing loss” (HHL), named for the presence of normal hearing thresholds but decreases in ABR peak I amplitudes at higher stimulus intensities (Kohrman et al., 2019; Liberman & Kujawa, 2017). In adult mice, SuppC overexpression of NT3 promotes regeneration of ribbon synapses and functional recovery after moderate noise overstimulation (Wan et al., 2014). These results have suggested the therapeutic potential of neurotrophins for repair of IHC-AN synapses following noise damage. Indeed, local delivery of NT3 into the cochlea was able to reduce synapse loss following moderate noise exposure (Suzuki et al., 2016). The efficacy of NT3 for synapse protection/repair, however, appears to be sensitive to dosage and/or timing of delivery. Although overexpression of NT3 in IHCs via infection with adeno-associated virus prior to noise exposure was able to protect synapses, viral delivery following noise was unable to reproduce the protective effect of direct NT3 injection (Hashimoto et al., 2019).

2.1.4 | Regulation of neurotransmission

Sound-evoked release of glutamate by IHC ribbon synapses activates AMPA receptors on SGN postsynaptic terminals (Fuchs et al., 2003). Glutamate accumulation and overactivation of glutamate receptors can degrade the fidelity of glutamatergic signaling (Barbour & Häusser, 1997) and potentially lead to excitotoxicity-induced withdrawal of the axon terminal and cell death of SGNs (Duan et al., 2006; Jäger et al., 2000; Kostandy, 2012; Oestreicher et al., 2002; Pujol & Puel, 1999). Similar to traditional glial cell-neuronal interactions that act to replenish glutamate neurotransmitter levels, the glutamate-glutamine cycle also occurs in the organ of Corti. SuppCs mediate glutamate removal at the IHC afferent synapse through the activity of glutamate/aspartate transporter (GLAST, also known as EAAT1). Glutamate uptake is then followed by conversion of glutamate to glutamine by glutamine synthetase (Usami et al., 1992). GLAST expression in SuppCs is closely matched to the amount of

glutamate released (Furness & Lawton, 2003) and the time course of GLAST expression in SuppCs correlates with the electrophysiological onset and maturation of mouse auditory function (Jin et al., 2003). GLAST-deficient mice experience more pronounced swelling of afferent terminals and exacerbated hearing loss after noise exposure than wild types, consistent with enhanced glutamate excitotoxicity (Glowatzki et al., 2006; Hakuba et al., 2000).

2.1.5 | Regulation of organ of Corti ionic environment

Ion recycling mechanisms are required in the cochlea to maintain homeostasis, and SuppCs are key players in these pathways. For example, potassium ions (K^+) that flow through mechanotransduction channels in HCs during sound stimulation are released through KCNQ4 channels in the HC basolateral membranes and K^+ transporter proteins are required for further recycling (Zdebik et al., 2009). Expression of the K -Cl cotransporter protein KCC4 is restricted to SuppCs in the mature cochlea (Boettger et al., 2002). Mice with deletion of the *Slc12a7* gene that encodes KCC4 exhibit early progressive degeneration of HCs, SuppCs and SGNs, leading to profound hearing loss, consistent with a critical role for KCC4 in uptake of K^+ into SuppCs (Boettger et al., 2002).

All cochlear SuppCs are connected by gap junctions that permit small molecule exchange, and this is believed to impact K^+ homeostasis in the inner ear (Zdebik et al., 2009). Gap junctions are formed by connexin (Cx) protein hemichannel complexes on adjacent cells. Cx26 and Cx30 are the major connexin proteins expressed by cochlear SuppCs and mutations in the genes encoding these isoforms have been associated with hearing loss in mice and in humans (Mammano, 2018). Most prominently, mutations in the human Cx26 gene (*GJB2*) account for up to 50% of autosomal recessive non-syndromic deafness in many populations (Chan & Chang, 2014). Potassium ion homeostasis, however, is unlikely to be the sole critical function for connexins in the cochlea, as several known *GJB2* mutations that are associated with hearing loss appear to form channels capable of small ion passage (Wingard & Zhao, 2015). Connexins also mediate passage of larger molecules such as ATP and inositol trisphosphate (IP₃), both of which have been linked to intercellular Ca^{2+} signaling in the organ of Corti (Mammano & Bortolozzi, 2018). SuppCs release ATP through apical connexin hemichannels, which results in purinergic receptor activation on SuppCs and leads to both the influx of external Ca^{2+} and release of internal Ca^{2+} stores. ATP release from SuppCs has been implicated in the regulation of spontaneous activity in the early postnatal cochlea (see above). Intercellular passage of the second messenger IP₃ also occurs through gap junctions, which similarly activates Ca^{2+}

release from internal stores (Mammano & Bortolozzi, 2018). The importance of this pathway is supported by the identification of a human deafness-associated variant in Cx30 that retains permeability to small ions but is deficient in intercellular IP₃ transfer and the associated Ca^{2+} response (Schütz et al., 2010). In addition, reduction of connexin-dependent Ca^{2+} signaling has been associated with an immature IHC phenotype, consistent with a role for SuppCs in regulating early sensory cell development (Johnson et al., 2017).

2.2 | Schwann cells

Schwann cells are derived from precursor cells in the neural crest that migrate to the periphery during embryogenesis, proliferate, associate with axons and adopt myelinating or non-myelinating fates (Salzer, 2015). Myelinating Schwann cells elaborate specialized membranes that wrap single axons and form the myelin sheath. Non-myelinating Schwann cells typically bundle multiple smaller caliber unmyelinated axons. A number of extracellular signaling cues and intrinsic transcriptional pathways have been demonstrated to guide these steps from neural crest progenitor to mature Schwann cell in the peripheral nervous system (PNS) (Stolt & Michael, 2016). The myelin sheath supports saltatory conduction of neural signals by regulating the assembly and maintenance of neuronal domains that are required for precise localization of voltage-gated Na^+ and K^+ channels at nodes of Ranvier (NoR) and at terminal heminodes (Rasband, 2016).

2.2.1 | Schwann cell development in the cochlea

Schwann cell development occurs across an extended time period in the vertebrate cochlea. In humans, immature Schwann cells are associated with AN axons by week 9 of gestation (Locher et al., 2014). Myelin formation, as evidenced by myelin basic protein (MBP) staining, is present within the cochlear portion of the AN by week 22 (Locher et al., 2014), consistent with earlier electron microscopy studies that indicate myelin formation is first evident in the peripheral AN segment between weeks 20 and 22 (Lavigne-Rebillard & Pujol, 1988; Moore & Linthicum, 2001; Ray et al., 2005). In the mouse, Schwann cell progenitors arrive in the developing inner ear by embryonic day 10.5, when neural progenitor cells begin to differentiate into SGNs (Breuskin et al., 2010). Myelination by Schwann cells initiates near birth at the cell body of SGNs and progresses centrally toward the PNS-CNS transition zone and peripherally toward the axon terminals between P8 and P10, just days prior to the onset of hearing (Anniko, 1983; Wang et al., 2013). Immunolocalization studies in rat indicate that, like in other peripheral nerves,

maturation of AN myelin continues throughout the first post-natal month, with progressively tighter clustering of Na⁺ and K⁺ channels, along with associated proteins, at NoRs and terminal heminodes (Kim & Rutherford, 2016).

2.2.2 | Regulation of SGN migration, maturation, and survival

Several studies suggest that pre-myelinating Schwann cells produce a “stop signal” required for appropriate SGN migration and fiber extension in the cochlea, and that they might contribute to SGN survival during development. Morris and coworkers found that in mice with loss of ErbB2, a receptor tyrosine kinase critical for Schwann cell development, SGN cell bodies migrate to unusual positions in the cochlea and their processes overshoot the sensory epithelia during early development (Morris et al., 2006). Similarly, in mice that lack Schwann cells due to Sox10 deletion in neural crest cells, SGN fibers extend peripherally to the lateral part of the cochlea past their normal IHC targets (Mao et al., 2014). Interestingly, ErbB2 KO but not Sox10 KO mice have significant SGN loss, suggesting that Schwann cells are not essential for SGN survival during this developmental stage, and that the neuronal loss in the ErbB2 KO reflects SuppC dysfunction (Stankovic et al., 2004).

There is also evidence that growth factor pathways mediate critical axon-Schwann cell interactions in the AN, including NRG1-ErbBR signaling, which controls myelin thickness in other peripheral nerves (Chen et al., 2006; Garratt et al., 2000; Michailov et al., 2004). All SGNs express NRG1 (Hansen et al., 2001; Morley, 1998; Stankovic et al., 2004), and AN Schwann cells express ErbBRs (Hansen et al., 2001). We have found that loss of ErbBR signaling in myelinating Schwann cells due to expression of a dominant negative ErbBR results in HHL (Cassinotti, Kohrman, et al., 2020; Kohrman et al., 2019). This dysfunction is associated with deficits in the structure of the terminal heminodes of the AN, consistent with a critical role for the heminodes in ensuring fast, synchronous firing of APs in response to neurotransmitter release from HCs.

Another example is the fibroblast growth factor (FGF) signaling pathway, which has been shown to regulate multiple functions of Schwann cells, including proliferation, differentiation, regeneration of peripheral nerves following injury, and recovery following demyelination (Davis & Stroobant, 1990; Grothe & Nikkhah, 2001; Hansen et al., 2001). SGNs express FGF-1 and FGF-2 (Lin et al., 1993; Pirvola et al., 1995; Silva et al., 2005), while Schwann cells and OLs express a number of FGF receptors, including FGFR1, FGFR2, and FGFR3 (Bansal et al., 2002; Grothe & Nikkhah, 2001). Wang and coworkers showed that inactivation of FGF signaling in myelinating cells resulted in significant loss of SGNs, accompanied

by age-related hearing impairment in adulthood, without apparent loss of glia (Wang et al., 2009). Although myelination was not quantitatively assessed in this study, apparent decreases in amplitudes of the sound-evoked responses and increases in their latencies are suggestive of subtle myelination defects. However, since the loss of FGF function occurred in both OLs and Schwann cells, the relative contribution of each myelinating cell type to the observed phenotypes remains to be determined.

2.2.3 | Myelin-related pathologies in the peripheral auditory system

There is increasing evidence that peripheral myelin disorders can lead to auditory neuropathy, a type of hearing loss in which sound detection in the cochlea is normal but transmission of auditory activity to the brain is impaired. This is the case in Guillain–Barre syndrome (GBS), an acute and transient peripheral neuropathy caused by Schwann cell damage due to autoimmune processes (Stathopoulos et al., 2015). It has been reported that a portion of GBS patients exhibit abnormal ABRs (Nelson et al., 1988; Ropper & Chiappa, 1986; Schiff et al., 1985; Takazawa et al., 2012; Ueda & Kuroiwa, 2008). Although hearing thresholds gradually recover in some patients, their ABR wave forms show persistent increases in interpeak latencies (Takazawa et al., 2012), consistent with permanent auditory neuropathy. Using a genetic approach to ablate myelinating Schwann cells in mice, we recently demonstrated that transient AN demyelination leads to permanent HHL that is associated with a disruption of the terminal AN heminodes (Wan & Corfas, 2017). These defects are consistent with previous modeling and immunostaining studies that support the heminodes as the initial spike generators in type I SGNs (Kim & Rutherford, 2016; Rutherford et al., 2012) and indicate that loss of the normal compacted heminode structure significantly impairs synchronized initiation and conduction of APs (Budak et al., 2019; Wan & Corfas, 2017). Thus, heminodal disruption may be the underlying cause of the auditory neuropathy observed in some GBS patients.

Another Schwann cell-related disorder is demyelinating Charcot–Marie–Tooth (CMT), also known as CMT1, in which some patients have been reported to present classic auditory neuropathy characterized by normal OHC function but substantial decreases in hearing sensitivity, often with ABR latency alterations (Rance, 2005). CMT is a genetically heterogeneous disorder, and the auditory phenotype in CMT patients is quite variable, likely due to the nature of the affected gene, the mutation type, age, and other genetic modifiers (Morelli et al., 2017). For example, evaluation of a cohort of children genetically diagnosed with CMT1, mostly with dominant peripheral myelin protein 22 (PMP22) gene duplication (CMT1A), exhibited relatively normal hearing thresholds but

decreased speech understanding in background noise (Rance et al., 2012). ABR studies from these patients show altered peak latencies consistent with demyelination and prolonged conduction times in the peripheral AN (Rance et al., 2012). Similarly, a large cohort of adult CMT1A patients with normal hearing thresholds was recently found to exhibit HHL, with deficits in speech understanding in background noise that were also associated with other deficits in temporal processing of sound (Choi et al., 2018). On the contrary, older patients carrying a dominant missense mutation in myelin protein zero (MPZ) gene (CMT1B) were found to exhibit significant threshold increases associated with gross defects in ABR peaks (Starr, 2003). Postmortem analysis of the cochlea of one of these patients demonstrated a significant loss of SGNs and demyelination of remaining peripheral fibers, compared with age matched controls (Starr, 2003).

Our recent studies on mouse models of CMT1 support the notion that peripheral myelin disorders lead to auditory neuropathy with distinct phenotypes for each mutation (Cassinotti, Ji, et al., 2020). CMT1A mice, which carry three copies of the human PMP22 gene (Verhamme et al., 2011), exhibit normal hearing thresholds at 4 months of age but decreases in ABR peak I amplitudes, indicating an HHL phenotype. This phenotype is associated with defects in the terminal heminode of auditory nerve fibers. In contrast, Trembler-J mice, which carry a dominant missense mutation in PMP22 and serve as a model of CMT1E, exhibit large threshold elevations at 4 months that are associated with hypomyelination and severe loss of peripheral AN fibers and SGN. These features are consistent with the auditory neuropathy previously reported for Trembler-J mice and with the peripheral demyelination caused by direct ablation of Schwann cells with diphtheria toxin at birth (Zhou, Abbas, et al., 1995; Zhou, Assouline, et al., 1995). Severe to profound threshold elevations are also observed in CMT1E patients (Boerkoel et al., 2002; Kovach et al., 2002). These CMT phenotypes highlight the critical role of Schwann cell myelination in neurotransmission in the cochlea and in long-term AN survival.

Loss of other Schwann cell proteins, including connexins, also result in peripheral neuropathy and auditory dysfunction. Hearing impairment is observed in a subset of individuals with an X-linked form of demyelinating CMT disease (CMT1X), which is caused by any one of a large number of mutations in the gap junction protein Cx32 (*GJB1*) (Kleopa & Scherer, 2006). Electrophysiological and pathological studies of peripheral nerves in these patients provide evidence of demyelinating neuropathy with prominent axonal degeneration. Cx32 has been implicated in the gap junction-mediated intracellular transfer of small molecules across myelin layers, as well as in hemichannel passage of ATP from activated neurons, and is expressed in both Schwann cells and OLs (Bortolozzi, 2018). Alterations in ABR peak I as well as in central peaks have been observed in different CMT1X

patients, consistent with a role for Cx32 in both the peripheral and central auditory systems (Bähr et al., 1999; Giuliani et al., 2019; Nicholson & Corbett, 1996). Although not associated with general neuropathy, loss of the gap junction protein Cx29 in mice causes AN myelin defects along with a delay in the maturation of auditory thresholds and the prolongation of auditory latencies (Tang et al., 2006).

Knowledge of the mechanisms of axon–glial communication provides potential targets for treating neuropathy and myelin disorders. For example, based on the importance of NRG1/ErbB signaling in modulating myelin development, experimental manipulation of this pathway has been tested for efficacy in treating demyelinating neuropathies. Delivery of soluble type I NRG1 via either injected recombinant protein or through expression of a neuronal transgene was found to normalize-myelinated axon counts in the sciatic nerve and to improve muscle function in a rat model of CMT1 (Fledrich et al., 2014). Similarly, transgenic expression of type III NRG1 in peripheral neurons resulted in partial correction of conduction defects and hypomyelination in the sciatic nerves of two independent mouse models of CMT1 (Belin et al., 2018; Scapin et al., 2018). Manipulation of this pathway must be carefully controlled, however, as overactivation of NRG1/ErbB signaling can lead to hypermyelination and associated neuropathy (Belin et al., 2018; Michailov et al., 2004).

2.2.4 | Acquired and age-related hearing loss

While noise overexposure and some ototoxic drugs have direct damaging effects on HCs, these traumas have also been associated with alterations in Schwann cells and myelin structure. Studies in young rats demonstrated reduced AN myelin thickness and changes in nodal and paranodal dimensions within 4 days of exposure to intense noise sufficient to cause HC damage and permanent threshold shifts (Tagoe et al., 2014). Similarly, high intensity noise exposure to young adult mice (8–16 kHz at 106 or 112 dB SPL for 2 hr)-induced disruption of AN myelin lamellae and heminode structures within 1 day of exposure, also suggesting direct effects of noise on myelin structure (Panganiban et al., 2018). Myelin pathologies in AN have been observed in other animal models following either prolonged low level or acute intense sounds, yet, in most cases the effects on myelin were evaluated at later time periods following substantial loss of SGN, and thus, hampering interpretation (Coyat et al., 2019; Rossi et al., 1976).

Cochlear myelin defects have also been noted in animal models within days following exposure to chemotherapeutic compounds such as cisplatin and carboplatin (Ding et al., 2002; van Ruijve et al., 2004), consistent with the effects of chemotherapeutic drugs on Schwann cells observed in other peripheral nerves (Carozzi et al., 2015). Chemotherapeutic compounds

also cause CNS damage, including within the central auditory system (Meyers, 2008). For example, studies indicate that 5-fluoro-uracil is toxic to OLs in vitro, damages central myelin and axons in the corpus callosum, and results in increased ABR interpeak latencies, consistent with myelin dysfunction in auditory pathways in mice (Han et al., 2008).

In addition to HCs, aminoglycoside antibiotic toxicity may directly affect SGNs and peripheral fibers in the cochlea soon after local or systemic delivery, with residual AN fibers often lacking myelin sheaths (Dodson & Mohuiddin, 2000; Jiang et al., 2017; Wise et al., 2017). Finally, myelin deficits have also been associated with aging in animal models, including thinning and degeneration of myelin on cochlear AN fibers, which correlated with decreases in MBP levels and in ABR peak I amplitudes (Xing et al., 2012). Similar age-associated decreases in MBP levels were also observed in AN from human temporal bone samples (Xing et al., 2012), suggesting that myelin loss could contribute to the temporal processing abnormalities described in aging humans (Harris & Dubno, 2017; Plack et al., 2014).

2.2.5 | Satellite glial cells

Like most peripheral neurons, SGN cell bodies in the cochlea are enclosed by SGCs. Remarkably, and distinct from all other peripheral ganglia, SGCs create a thin myelin sheath around type I SGN in most vertebrates (Felix, 2002). Notably, this is not the case in humans (Liu, Edin, et al., 2015; Tylstedt et al., 1997). The mechanisms that mediate this cell body myelination and its precise physiological impact remain unknown, although biophysical modeling studies have suggested that human type I SGNs have lower spike conduction velocity due to lack of myelin in the soma (Rattay et al., 2001, 2013). Interestingly, high resolution 3D transmission electron microscopy of human cochleae has demonstrated the presence of membrane specializations between groups of SGN soma clustered within common SGCs (Tylstedt & Rask-Andersen, 2001). These membrane areas were speculated to contribute to electrotonic cross-transmission that could be related to coding of speech signals.

Several studies have provided evidence that SGCs are important for proper auditory function. SGCs express glutamine synthetase, which is important for metabolizing excess glutamate and preventing excitotoxicity (Eybalin et al., 1996). In addition, deletion of the Saposin B (Sap B) gene, which plays an essential role in the regulation of myelin lipid biosynthesis, results in selective loss of cochlear SGCs in mice and is associated with progressive degeneration of SGNs and hearing loss, implicating SGCs in AN maintenance (Akil et al., 2015). In line with the regenerative ability of SGCs in other sensory ganglia following nerve injury (Donegan et al., 2013; Elson et al., 2004; Nascimento et al., 2014), SGCs in the cochlea also regenerate following genetic ablation (Wan & Corfas, 2017).

2.3 | Oligodendrocytes

Oligodendrocytes (OLs) derive from neuroepithelial precursor cells (OPCs) that originate in the ventricular zone of the developing CNS. Intrinsic transcriptional pathways together with extracellular signaling cues guide the migration, proliferation, and differentiation of OPCs into mature, myelin-producing OLs (Elbaz & Popko, 2019). In contrast to the single nerve fibers wrapped by mature Schwann cells in the periphery, myelinating OLs send out cellular extensions that are capable of wrapping and myelinating multiple CNS axons.

2.3.1 | Oligodendrocyte development in the auditory CNS

Myelination of AN central projections and auditory brainstem axons by OLs starts around the time of hearing onset in rodents and humans. In mice, in which hearing onset occurs around P12-P14, OPCs are present in both the AN (Bojrab et al., 2017) and the auditory hindbrain (Dinh et al., 2014) at birth, while myelination of the central projections of AN are first detectable by P8 near the PNS-CNS transitional zone (Wang et al., 2013) (Figure 1a,b). Myelin formation in the auditory hindbrain and cortex initiates approximately 1 week later (Hackett et al., 2015; Kolson et al., 2015). Hearing onset in humans occurs between 24 and 25 weeks of gestation (Birnholtz & Benacerraf, 1983), with myelination of the central projection of AN as well as the auditory brainstem and midbrain initiating soon after, at 26 to 29 weeks (Moore & Linthicum, 2001; Moore et al., 1995). In both humans and mice, myelin maturation in the auditory CNS progresses after hearing onset, coinciding with functional maturation. For example, in the murine auditory brainstem, myelin thickness and axon diameter double in size during the 2 weeks following hearing onset. Importantly, during this time, conduction velocity in the auditory brainstem also doubles (Sinclair et al., 2017). Similarly, in the auditory brainstem and midbrain neurons of humans, increases in myelination are observed up to 1 year of age (Moore et al., 1995). In the auditory cortex, myelin signals detected by magnetic resonance imaging (MRI) rapidly increase until 1.5 years of age, then, progress at a slower rate until adulthood (Su et al., 2008).

2.3.2 | Oligodendrocyte function in the auditory CNS

The precise coding and maintenance of timing information of sound stimuli are critical for normal auditory tasks, including spatial hearing and sound localization. Mammals and birds localize sounds in the horizontal plane by comparing

the arrival times and intensities through auditory pathways originating at each ear (Grothe & Pecka, 2014). Humans are able to discriminate interaural time differences (ITDs) as small as 10 to 20 microseconds (Klumpp & Eady, 1956) and interaural intensity differences (IIDs) down to 1 dB SPL (Mills, 1960). The circuits in the mammalian central auditory pathway that subserve these sensory processes are depicted in Figure 1b. Neurons in the superior olivary complex (SOC) integrate sound input from each ear. Medial superior olive (MSO) neurons in the SOC play a key role in the ITD detection pathway by comparing the timing of excitatory input from spherical bushy cells (SBCs) in the ipsilateral and contralateral cochlear nuclei (Cant & Casseday, 1986; Lindsey, 1975; Stotler, 1953; Warr, 1966) and firing when these binaural inputs arrive within a short time window (Goldberg & Brown, 1969; Yin & Chan, 1990). In addition to this excitatory input, MSO neurons also receive inhibitory input, most importantly from principal neurons in the medial nucleus of the trapezoid body (MNTB), which fire in response to excitation from single axons of globular bushy cells (GBC) in the contralateral ventral cochlear nucleus (VCN). This contralateral information is transmitted through a large, specialized synapse known as the calyx of Held directly onto the cell bodies of the MNTB principal neurons (Figure 1d). The fast and precise nature of this synapse is critical for sound processing in the auditory hindbrain (Baydyuk et al., 2016). Similar integration of excitatory and inhibitory binaural inputs also occurs in lateral superior olive (LSO) neurons in the SOC to detect IIDs (Grothe & Pecka, 2014).

Evidence from several model systems suggests that specialized properties of OL-based myelination and their associated axons play a key role in fine tuning the timing of neural transmission in the auditory CNS. Comparisons of ipsilateral and contralateral pathways involved in sound localization suggest that these specializations are likely to influence conduction speeds to compensate for inherent differences in fiber projection lengths and synaptic connection numbers between pathways. For instance, in the ITD pathway of gerbils, excitatory projections from SBCs to the contralateral MSO exhibit longer internodes and increased axon diameters relative to those in the shorter ipsilateral projections and are expected to elevate conduction velocity and compensate for the longer contralateral projection distance (Seidl & Rubel, 2016). Systematic differences in internode lengths and axon diameters that are associated with conduction velocity variation have also been demonstrated in rodent GBCs, which are part of both the IID and the ITD pathways, and thus, seem to compensate for the additional calyx of Held synapse in the contralateral inhibitory circuit in SOC (Ford et al., 2015). Notably, a recent comparative study between gerbils, which depend upon ITD detection for localization of low frequency sounds, and mice, which rely instead upon IID, demonstrated unique structural specializations of axons and myelin in the

localization circuit of each species that correlated with ITD requirements (Stange-Marten et al., 2017).

The importance of myelination for temporal processing of sound information in the CNS is highlighted by electrophysiological defects found in Long-Evans shaker rats, which carry an insertional mutation of the MBP gene and exhibit little or no condensed myelin in CNS neurons (O'Connor et al., 1999). Increased peak I to peak V latencies and decreased peak amplitudes of ABRs to click sound stimuli in the MBP mutants indicated an increased central conduction time and lack of neural synchrony in the auditory hindbrain (Kim, Renden, et al., 2013; Kim, Turkington, et al., 2013). Additionally, brain slice recordings in the mutants indicated delayed and less reliable AP generation in MNTB principal neurons following high frequency synaptic activation at the calyx of Held. The delays and loss of AP fidelity were associated with altered Na⁺ and K⁺ channel distribution at NoR and the last heminodes of presynaptic GBC axons (Berret et al., 2016). These results support previous studies indicating the requirement of normal myelin structure for tight clustering of ion channels at nodes and heminodes in the CNS (Rasband et al., 1999; Susuki et al., 2013) and demonstrate its critical role for high-fidelity transmission along axons and at nerve terminals in the auditory hindbrain.

2.3.3 | Myelin plasticity in the auditory CNS

A large body of evidence supports a role for neuronal activity and experience in modulating myelin structure in the CNS during development and in the mature organism through multiple mechanisms (Almeida & Lyons, 2017; Monje, 2018). Many studies have demonstrated that such activity-dependent myelin plasticity impacts behaviors and cognitive function. The coincident timing of myelination and development of auditory processing capabilities, including language acquisition in humans, has suggested that auditory experience impacts myelination to refine sensation during auditory system maturation.

Imaging studies in humans support a relationship between sensory-evoked activity and myelin levels in the auditory CNS. MRI studies showed that white matter to gray matter ratios are lower in the Heschl's gyrus and superior temporal gyrus of congenitally deaf individuals relative to normal hearing controls (Emmorey et al., 2003; Hribar et al., 2014). Changes in white matter microstructure in the temporal lobes of deaf subjects relative to hearing controls have also been detected by diffusor tensor imaging (DTI), which measures water diffusion properties to estimate brain structural features (Hribar et al., 2014; Kim et al., 2009). Increasing sensory input, (auditory enrichment), has been associated with enhanced levels of CNS myelination. In rats, age-related decreases in myelin gene expression in the auditory cortex

were partially reversed by auditory enrichment in an operant conditioning paradigm (de Villers-Sidani et al., 2010). Similarly, DTI evaluations in humans have demonstrated a positive correlation between more extensive white matter connections in the corpus callosum and both the extent of musical training and the age of training initiation (Bengtsson et al., 2005; Steele et al., 2013). Using mice, Sinclair and colleagues found that eliciting a hearing loss (50 dB SPL threshold elevation) with ear plugs at the time of normal hearing onset results in thinner myelin wrapping axons projecting to the MNTB of the auditory brainstem (Sinclair et al., 2017). Similar deficits were observed following ear plugging of mature mice, suggesting that activity is also important in the maintenance of myelin.

Several mechanisms have been proposed to underlie the impact of neuronal activity and experience on CNS myelination, including impacts on OPC proliferation and differentiation (Foster et al., 2019). Trophic factors expressed by active neurons such as BDNF and NRG1 have been shown to stimulate myelination by OLs (Roy et al., 2007; Taveggia et al., 2007; Venkatesh et al., 2015; Wong et al., 2013; Xiao et al., 2010) and experience-dependent changes in the levels of NRG1 expression have been shown to underlie the impact of social isolation on prefrontal cortex myelin maturation (Makinodan et al., 2012). Likewise, sustained genetic activation in OLs of ERK1/2 kinases, which are part of the intracellular signaling pathway initiated by NRG1-ErbBR engagement (Mei & Nave, 2014), results in increased myelin thickness in the CNS and decreased latencies in ABR brainstem peaks (Jeffries et al., 2016).

Neurotransmitters such as ATP and glutamate released by active neurons may also impact myelination. In vitro studies indicated an indirect effect of neuronal ATP release, which induces the release of leukemia inhibitory factor (a cytokine with pro-myelinating activity), from astrocytes, which in turn stimulates myelin formation by OLs (Ishibashi et al., 2006). OPCs express a variety of voltage-gated ion channels and neurotransmitter receptors, including AMPA, NMDA, and GABA receptors (de Biase et al., 2010). In multiple regions of the CNS, subsets of these progenitors also receive synapse-like input from either glutamatergic excitatory and GABAergic inhibitory neurons (Bergles et al., 2000; Lin & Bergles, 2003; Ziskin et al., 2007). A recent study provided evidence that OPC excitability contributes to their own maturation and myelination (Berret et al., 2017). This article showed that a subset of pre-OLs in the MNTB of P7-P14 rats fire APs in response to glutamatergic transmission from nearby neurons, and these APs depend upon the voltage-gated Na^+ channel Nav1.2. Moreover, shRNA-mediated knockdown of Nav1.2 blocks this Na^+ current and results in altered OL cellular extensions and a decrease in MBP levels, consistent with an impact of activity on pre-OL development and myelination.

In addition to modulating neural transmission through effects on myelination, OL-nerve interactions also appear to impact synaptic plasticity at the calyx of Held synapse via BDNF-trkB signaling (Jang et al., 2019). Building on prior studies that demonstrated release of BDNF from OLs in response to glutamate in the cortical CNS (Bagayogo & Dreyfus, 2009), Jang and coworkers verified BDNF expression by OLs in the MNTB and demonstrated decreased glutamate release from presynaptic neurons in slice preparations from mice that lack BDNF expression in pre-myelinating OLs (Jang et al., 2019). These conditional *Bdnf* knockouts also exhibited defects in neural synchrony in the auditory brainstem. The rescue of glutamatergic transmission in the knockouts by local delivery of either BDNF or trkB agonists suggests a mechanism by which BDNF secretion from OLs near the calyx of Held modulates neurotransmitter release from the GBC terminal and supports a modulating role for activity-dependent regulation of presynaptic properties by OLs in the central auditory system.

2.3.4 | Oligodendrocyte pathology and hearing impairments

CNS myelin disorders can affect auditory function, as might be expected given the impact of myelin on conduction velocity and timing in neuronal circuits. Studies of these disorders have provided insights into how myelin impacts processing of auditory signals and the nature of hearing defects that arise due to abnormal CNS myelination. While a number of case reports indicate increases in hearing thresholds in multiple sclerosis (MS) patients that present with lesions in CNS auditory regions (Bergamaschi et al., 1997; Drulovic et al., 1993), larger controlled studies failed to validate those observations (Doty et al., 2012). Sudden sensorineural hearing loss has also been described in MS patients, but typically resolves over time (Hellmann et al., 2011). Although MS does not appear to result in consistent decreases in auditory sensitivity, subtler defects in hearing that involve losses in temporal auditory processing have been associated with the disorder. Common abnormalities in MS patients include increases in ABR I to V peak latencies that are consistent with myelin deficits in the auditory brainstem. MS patients often exhibit decreased abilities in binaural hearing tasks such as detection of ITDs and IIDs, suggesting impairment of sound localization pathways (Furst & Levine, 2015).

Defects in CNS myelin development may also underlie auditory dysfunction. Auditory processing disorder (APD) is a heterogeneous developmental disorder characterized by perceptual problems including deficits in speech comprehension, spatial hearing, and attention. The normal cochlear thresholds often present in individuals diagnosed with APD has suggested that the source of the disorder lies within

defects in the central auditory system. Associations between APD and abnormalities in the temporal characteristics of speech-evoked ABRs (Rocha-Muniz et al., 2012, 2014) have led to the hypothesis that decreases in the precision of AP firing and conduction in auditory pathways contribute to perceptual dysfunction (Kopp-Scheinflug & Tempel, 2015). Consistent with this notion, myelin abnormalities have been observed in APD patients in brain regions involved in auditory processing. For example, an increased mean diffusion in auditory cortex was observed by DTI in children diagnosed with APD, consistent with a potential disruption of thalamocortical communication (Farah et al., 2014). The same DTI study also found white matter structural abnormalities in the prefrontal cortex and anterior cingulate, which are areas of the brain closely tied to attention and top-down information processing. This suggests that altered myelination in these structures may underlie difficulties in attending to auditory stimuli in APD.

The transmembrane proteolipid protein 1 (PLP1) is a major constituent of myelin in OL. Loss of function mutations in the human X-linked *PLP1* gene generally result in spastic paraplegia, a progressive neuronal degenerative disorder affecting mainly spinocerebellar and lateral corticospinal tracts, while missense mutations with apparent toxic gain of function effects underlie Pelizaeus–Merzbacher Disease (PMD), a more severe leukodystrophy disorder characterized by hypomyelination (Garbern, 2007). PMD patients typically have speech difficulties that are associated with normal cochlear function (ABR thresholds and peak I amplitudes) but dyssynchrony of later ABR peaks and delayed and/or reduced myelin levels in the auditory brainstem and higher auditory pathways (Coticchia et al., 2011; Kuan et al., 2009; Morlet et al., 2018). Defects in CNS myelination may also influence the auditory dysfunction observed in other neurological disorders. For example, many individuals diagnosed with autism spectrum disorder (ASD) exhibit alterations in several aspects of auditory perception including loudness detection and complex speech comprehension (O'Connor, 2012). ASD patients also often exhibit reduced detection of temporal features of auditory stimuli such as gaps in sounds (Foss-Feig et al., 2017) as well as decreased white matter properties in several brain regions, including the corpus callosum and temporal lobes (Travers et al., 2012). Defects in CNS myelin and auditory processing have also been associated with psychiatric disorders such as schizophrenia, bipolar disorder, and chronic depression (Fields, 2008; Kahkonen et al., 2007; McLachlan et al., 2013; Zenisek et al., 2015).

2.4 | Astrocytes

Astrocytes account for approximately 20%–40% of the glial population across different regions in the CNS and play a

range of supportive and regulatory roles that contribute to neuronal development, synaptic plasticity, and responses to neuronal damage (Bartheld et al., 2016). These cells elaborate highly branched processes that interact with neural structures, including synapses, and express ion channels, extracellular matrix proteins, growth factors and cytokines that appear to mediate their influence on neurons (Schiweck et al., 2018). Of note, astrocytes are chemically excitable via expression of a variety of neurotransmitter receptors (Nimmerjahn & Bergles, 2015).

2.4.1 | Astrocytes and development in the central auditory pathway

Astrocytes have been linked to the postnatal development of the central auditory system (Cramer & Rubel, 2016). Immunostaining of astrocyte-associated marker proteins including the intermediate filament glial fibrillary acidic protein (GFAP), the calcium-binding protein S100 β , and aldehyde dehydrogenase 1 family member L1 (ALDH1L1), revealed the emergence of a diverse pattern of astrocyte cell types in VCN and MNTB of rodents across the 3 week postnatal developmental period (Dinh et al., 2014; Saliu et al., 2014). At P14, just after the time of hearing onset and when auditory brainstem pathways are approaching maturity, all astrocyte markers were expressed in the VCN and MNTB, with GFAP-positive processes found in both locations while GFAP-positive cell bodies remained outside of these nuclei (Dinh et al., 2014). In the MNTB, cell bodies of ALDH1L1-positive and S100 β -positive astrocytes were present near principal neurons and calyces of Held, the excitatory terminal of GBC axons on principal neurons, while GFAP-positive processes were often present near the calyces (Dinh et al., 2014) (Figure 1d). This heterogeneous population of astrocytes present during auditory hindbrain maturation has suggested that distinct classes of astrocytes carry out different developmental functions, with early emerging astrocytes participating in neuronal refinement and synaptogenesis and later emerging astrocytes influencing neurotransmission (Cramer & Rubel, 2016; Dinh et al., 2014).

It has been suggested that interactions between astrocytic processes and the pre- and postsynaptic membranes of the calyx of Held influence the development and function of this critical auditory synapse. Spontaneous Ca²⁺ transient currents in astrocytes appear to elicit slow inward currents in postsynaptic principal neurons in early postnatal brain slice recordings in MNTB (Reyes-Haro et al., 2010). These astrocyte-mediated currents, known as gliotransmission, correlate with astrocytic Ca²⁺ activity and seem to be NMDA receptor mediated. Glutamate release from the presynaptic GBCs mediates fast excitatory synaptic transmission at the calyces of Held. Group II metabotropic glutamate receptors

(mGluRs) are expressed in astrocytic processes at the maturing calyces of Held, suggesting that synaptic transmission may also influence astrocyte function during this time (Elezgarai et al., 2001).

Astrocytes play key homeostatic roles in the CNS, including biosynthetic recycling of the neurotransmitter glutamate at synapses (Verkhatsky et al., 2015). Studies in the developing postnatal MNTB of rats and gerbils demonstrated that the glutamate transporters GLAST and GLT are selectively localized in astrocyte processes ensheathing the calyx of Held (Ford et al., 2009; Renden et al., 2005). This is likely to aid in limiting glutamate diffusion between adjacent terminals during this period of structural and functional remodeling of the calyx. Similar to SuppCs in the cochlea and other glia, glutamate taken up by astrocytes in the MNTB is converted to glutamine; release of glutamine by astrocytes provides a precursor for biosynthetic replenishment of glutamate in adjacent neurons (Uwechue et al., 2012).

Astrocytes appear to influence neuronal development in the avian auditory hindbrain through release of modulatory factors (Cramer & Rubel, 2016). Neurons in nucleus laminaris (NL), which are involved in ITD pathways of sound localization in chicks, display a gradient of dendritic arborization that systematically changes during development of the auditory circuits (Parks & Rubel, 1975). Morphological alterations in dendrites coincide with the appearance of GFAP-positive astrocytes, which led researchers to assess the contribution of glial cells to branch order and dendritic morphometric features in NL neurons (Korn et al., 2011). Studies in embryonic brainstem slices indicated that conditioned media from cultured embryonic astrocytes promoted the formation of a normal gradient of dendritic morphology in NL neurons, suggesting that astrocytes are an important source of molecules capable of modulating dendritic branching and reorganization during development (Korn et al., 2011). Similar brainstem slice experiments also support a role for astrocyte-derived factors in regulating the final numbers of inhibitory synapses on NL neurons, which in turn facilitate more precise binaural coincidence detection (Korn et al., 2012).

2.4.2 | Astrocyte involvement in auditory system damage

Glial cells, including astrocytes and microglia, participate in complex adaptive responses to neurological injury and disease, including local inflammation as well as degenerative and neural protective activities (Schiweck et al., 2018). CNS injuries often lead to an increase in reactive astrocytes, which serve to dampen inflammatory responses and increase neuronal viability (Faulkner et al., 2004). Glial responses to injury often produce a “scar” composed of astrocytes, microglia,

and associated extracellular matrix material. Such glial scars, while containing the extent of injury, also appear to limit the ability of neural regeneration (Bradbury & Burnside, 2019). A number of studies support the involvement of astrocytes and other glia following damage or peripheral sensory loss in the auditory system.

In a model of acute injury by compression of the central portion of the AN in rats, Sekiya and colleagues observed GFAP-positive reactive astrocytes along with markers of glial scar formation in the vicinity of the injured site, which also extended beyond the PNS-CNS transitional zone boundary into the cochlea (Sekiya et al., 2015). The injury resulted in partial degeneration of SGNs that was associated with ABR threshold elevations. Interestingly, the authors reported that transplantation of auditory neuroblast cells onto the surface of the injured nerves resulted in neurons capable of forming glutamatergic synapses with HCs in the organ of Corti and extending processes into the auditory hindbrain. Rats receiving the neuroblasts exhibited modest recovery of central ABR peak thresholds relative to sham-treated controls. The transplanted cells were closely associated with GFAP-positive processes, suggesting that the reactive astrocytes may have provided neural guidance cues, similar to the glia-guided migration of neurons in the developing cortex (Marín & Rubenstein, 2003).

Disruption of cochlear function decreases afferent AN signaling in the cochlear nucleus (CN) and leads to activity-dependent alterations in central auditory pathways, including neural degeneration and rewiring of circuits (Gold & Bajo, 2014). Activated glial responses involving astrocytes and microglia have often been observed following loss of AN nerve input in mammals and birds during development and in the adult, consistent with a damage response (de Waele et al., 1996; Janz & Illing, 2014; Lurie & Rubel, 1994). For example, the neural degeneration in ipsilateral CN that follows unilateral cochlear ablation leads to loss of normal VCN connections to the contralateral MNTB (Hashisaki & Rubel, 1989; Mostafapour et al., 2000; Trune, 1982). Interestingly, following cochlear ablation during early postnatal development, neurons of the intact VCN have been shown to send branches that form new calyceal contacts with the denervated MNTB (Hsieh et al., 2007; Kitzes et al., 1995; Moore & Kowalchuk, 1988). Dinh and coworkers observed that these new synapses exhibited closely apposed astrocytes, suggesting that glial cells are involved in reorganizing the new calyces (Dinh et al., 2014). In contrast to manipulations during development, loss of afferent synapses in VCN following cochlear ablation in the mature animal leads to reinnervation of VCN cells by axon collateral sprouting from SOC-derived neurons (Kraus & Illing, 2004). Molecular studies of astrocyte markers during this reorganization suggested two phases of glial cell responses, one associated with AN and synaptic degeneration and the second associated

with reinnervation and new synapse formation (Fredrich et al., 2013).

A long-lasting activated glial response (increased number of astrocytes and microglia processes) has been observed in the CN of mature rats following deafening by loud noise exposure, prior to any evidence of overt AN degeneration (Fuentes-Santamaria et al., 2017). Glial processes were found in close apposition to VCN neurons and synapses, along with increased expression of pro-inflammatory cytokines, including tumor necrosis factor alpha (TNF- α). This glial response may be related to the increased spontaneous firing rate in CN neurons that is frequently observed following noise trauma (Kaltenbach & Afman, 2000; Vogler et al., 2011). Following deafening of neonatal rats, glial activation has also been correlated with increased neuronal activity in the inferior colliculus (IC), a central auditory nucleus that receives input from both the periphery and higher level pathways (Rosskothenkühl et al., 2018). These findings suggest that glial remodeling contributes to auditory system plasticity following sensory loss.

Deafness-associated plasticity, including increased spontaneous activity and synchrony in central auditory system nuclei, has also been implicated in tinnitus, an often debilitating condition characterized by the sensation of phantom sounds (Shore et al., 2016). A recent study has suggested an association of microglial activation and increased TNF- α expression with excitatory-inhibitory synaptic imbalance in the auditory cortex following noise-induced deafness in mice (Wang et al., 2019). Pharmacological and genetic inhibition of TNF- α resulted in a decrease in microglial activation and normalized synaptic firing in the cortex, together with reduced behavioral evidence of tinnitus in deafened animals. Although the use of behavioral studies to evaluate tinnitus in animal models remains controversial (Jones & May, 2017), this study suggests that neuroinflammatory responses to noise could influence auditory system plasticity, and thus, may also impact development of tinnitus.

3 | CONCLUSIONS

The auditory system provides a tractable experimental model to investigate the roles of glia and neuron–glia interactions in health and disease. The accessibility of the auditory system for in vivo recordings at the physiological and behavioral levels, in conjunction with the growing toolbox for genetic manipulation of sensory HCs, neurons and glia, provide opportunities to interrogate the molecular and cellular mechanisms of neuron–glia interactions. As a note of caution, however, a number of studies have underscored the importance of careful consideration of genetic regulatory elements in these approaches, as promoters typically considered

specific for particular glial subtypes can be expressed more broadly in both neurons and glia (Brandebura et al., 2018; Wan & Corfas, 2017). Similarly, a more detailed understanding of the role of glia on hearing should provide insight into deafness and other disorders that affect the auditory system, including peripheral neuropathies and tinnitus, and is also likely to suggest novel approaches for treatment.

Further comparative investigations of glial interactions within the auditory system through use of high resolution imaging of animal model and human specimens will also be useful to identify any subtle species-specific differences that may underlie functional divergence (Liu, Edin, et al., 2015; Liu et al., 2020). Similarly, proteomic and single cell transcriptional studies of auditory tissue from humans and animals will provide a sharper picture of molecular mechanisms involved in glial function (Lin et al., 2019; Petitpré et al., 2018; Schmitt et al., 2017; Shrestha et al., 2018; Sun et al., 2018). In addition to providing insight into basic processes such as sensory information processing, a more precise understanding of axon–glia interactions in the auditory system is likely to impact development of future regeneration and neural implant treatment strategies for hearing loss.

ACKNOWLEDGMENTS

This work was supported in part by NIH/NIDCD R01 DC018500 and R21 DC017916; and the Lynn and Ruth Townsend Professorship of Communication Disorders (G.C.). We thank Günseli Wallace for critical reading of the manuscript.

CONFLICT OF INTEREST

G.C. is a scientific founder of Decibel Therapeutics, has an equity interest in and has received compensation for consulting. The company was not involved in this study.

ORCID

Gabriel Corfas  <https://orcid.org/0000-0001-5412-9473>

REFERENCES

- Adamson, C. L., Reid, M. A., & Davis, R. L. (2002). Opposite actions of brain-derived neurotrophic factor and neurotrophin-3 on firing features and ion channel composition of murine spiral ganglion neurons. *The Journal of Neuroscience*, 22(4), 1385–1396. <https://doi.org/10.1523/JNEUROSCI.22-04-01385.2002>
- Akil, O., Sun, Y., Vijayakumar, S., Zhang, W., Ku, T., Lee, C.-K., Jones, S., Grabowski, G. A., & Lustig, L. R. (2015). Spiral ganglion degeneration and hearing loss as a consequence of satellite cell death in saposin B-deficient mice. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 35, 3263–3275.
- Almeida, R. G., & Lyons, D. A. (2017). On myelinated axon plasticity and neuronal circuit formation and function. *Journal of Neuroscience*, 37(42), 10023–10034.
- Anniko, M. (1983). Early development and maturation of the spiral ganglion. *Acta Oto-Laryngologica*, 95, 263–276.

- Babola, T. A., Kersbergen, C. J., Wang, H. C., & Bergles, D. E. (2020). Purinergic signaling in cochlear supporting cells reduces hair cell excitability by increasing the extracellular space. *eLife*, *9*, e52160.
- Bagayogo, I. P., & Dreyfus, C. F. (2009). Regulated release of BDNF by cortical oligodendrocytes is mediated through metabotropic glutamate receptors and the PLC pathway. *ASN Neuro*, *1*(1), e00001.
- Bähr, M., Andres, F., Timmerman, V., Nelis, M. E., Broeckhoven, C. V., & Dichgans, J. (1999). Central visual, acoustic, and motor pathway involvement in a Charcot-Marie-Tooth family with an Asn205Ser mutation in the connexin 32 gene. *Journal of Neurology, Neurosurgery & Psychiatry*, *66*(2), 202–206. <http://dx.doi.org/10.1136/jnnp.66.2.202>.
- Bank, L. M., Bianchi, L. M., Ebisu, F., Lerman-Sinkoff, D., Smiley, E. C., Shen, Y.-C., Ramamurthy, P., Thompson, D. L., Roth, T. M., Beck, C. R., Flynn, M., Teller, R. S., Feng, L., Llewellyn, G. N., Holmes, B., Sharples, C., Coutinho-Budd, J., Linn, S. A., Chervenak, A. P., ... Barald, K. F. (2012). Macrophage migration inhibitory factor acts as a neurotrophin in the developing inner ear. *Development*, *139*(24), 4666–4674. <https://doi.org/10.1242/dev.066647>
- Bansal, R., Miyake, H., Nakamura, I., Eto, H., Gotoh, A., Fujisawa, M., Okada, H., Arakawa, S., Kamidono, S., & Hara, I. (2002). Fibroblast growth factors and their receptors in oligodendrocyte development: Implications for demyelination and remyelination. *Developmental Neuroscience*, *24*, 35–46.
- Barbour, B., & Häusser, M. (1997). Intersynaptic diffusion of neurotransmitter. *Trends in Neurosciences*, *20*(9), 377–384.
- Bartheld, C. S., Bahney, J., & Herculano-Houzel, S. (2016). The search for true numbers of neurons and glial cells in the human brain: A review of 150 years of cell counting. *Journal of Comparative Neurology*, *524*(18), 3865–3895. <https://doi.org/10.1002/cne.24040>
- Baydyuk, M., Xu, J., & Wu, L.-G. (2016). The calyx of Held in the auditory system: Structure, function, and development. *Hearing Research*, *338*, 22–31.
- Belin, S., Ornaghi, F., Shackleford, G. G., Wang, J., Scapin, C., Lopez-Anido, C., Silvestri, N., Robertson, N., Williamson, C., Ishii, A., Taveggia, C., Svaren, J., Bansal, R., Schwab, H. M., Nave, K., Fratta, P., D'Antonio, M., Poitelon, Y., Feltri, M. L., & Wrabetz, L. (2018). Neuregulin 1 type III improves peripheral nerve myelination in a mouse model of congenital hypomyelinating neuropathy. *Human Molecular Genetics*, *28*(8), 1260–1273. <https://doi.org/10.1093/hmg/ddy420>.
- Bengtsson, S. L., Nagy, Z., Skare, S., Forsman, L., Forssberg, H., & Ullén, F. (2005). Extensive piano practicing has regionally specific effects on white matter development. *Nature Neuroscience*, *8*(9), 1148–1150. <https://doi.org/10.1038/nn1516>
- Bergamaschi, R., Romani, A., Zappoli, F., Versino, M., & Cosi, V. (1997). MRI and brainstem auditory evoked potential evidence of eighth cranial nerve involvement in multiple sclerosis. *Neurology*, *48*(1), 270–272. <https://doi.org/10.1212/WNL.48.1.270>
- Bergles, D. E., Roberts, J. D. B., Somogyi, P., & Jahr, C. E. (2000). Glutamatergic synapses on oligodendrocyte precursor cells in the hippocampus. *Nature*, *405*(6783), 187–191. <https://doi.org/10.1038/35012083>
- Berret, E., Barron, T., Xu, J., Debner, E., Kim, E. J., & Kim, J. H. (2017). Oligodendroglial excitability mediated by glutamatergic inputs and Nav1.2 activation. *Nature Communications*, *8*(1), 557. <https://doi.org/10.1038/s41467-017-00688-0>.
- Berret, E., Kim, S. E., Lee, S. Y., Kushmerick, C., & Kim, J. H. (2016). Functional and structural properties of ion channels at the nerve terminal depends on compact myelin. *The Journal of Physiology*, *594*(19), 5593–5609. <https://doi.org/10.1113/JP272205>
- Bianchi, L. M., Conover, J. C., Fritsch, B., DeChiara, T., Lindsay, R. M., & Yancopoulos, G. D. (1996). Degeneration of vestibular neurons in late embryogenesis of both heterozygous and homozygous BDNF null mutant mice. *Development*, *122*(6), 1965–1973.
- Birnholtz, J., & Benacerraf, B. (1983). The development of human fetal hearing. *Science*, *222*(4623), 516–518. <https://doi.org/10.1126/science.6623091>
- Boerkoel, C. F., Takashima, H., Garcia, C. A., Olney, R. K., Johnson, J., Berry, K., Russo, P., Kennedy, S., Teebi, A. S., Scavina, M., Williams, L. L., Mancias, P., Butler, I. J., Krajewski, K., Shy, M., & Lupski, J. R. (2002). Charcot-Marie-Tooth disease and related neuropathies: Mutation distribution and genotype-phenotype correlation. *Annals of Neurology*, *51*(2), 190–201. <https://doi.org/10.1002/ana.10089>
- Boettger, T., Hübner, C. A., Maier, H., Rust, M. B., Beck, F. X., & Jentsch, T. J. (2002). Deafness and renal tubular acidosis in mice lacking the K-Cl co-transporter Kcc4. *Nature*, *416*(6883), 874–878. <https://doi.org/10.1038/416874a>
- Bojrab, D., Zhang, B., Jiang, H., Zhang, L., Cohen, D. S., Luo, X., & Hu, Z. (2017). Expression of oligodendrocyte marker during peripheral-central transitional zone formation of the postnatal mouse cochlear nerve. *Otolaryngology-Head and Neck Surgery*, *157*(3), 488–492. <https://doi.org/10.1177/0194599817718806>
- Bortolozzi, M. (2018). What's the function of Connexin 32 in the peripheral nervous system? *Frontiers in Molecular Neuroscience*, *11*, 227. <https://doi.org/10.3389/fnmol.2018.00227>
- Bradbury, E. J., & Burnside, E. R. (2019). Moving beyond the glial scar for spinal cord repair. *Nature Communications*, *10*(1), 3879. <https://doi.org/10.1038/s41467-019-11707-7>.
- Brandebura, A. N., Morehead, M., Heller, D. T., Holcomb, P., Kolson, D. R., Jones, G., Mathers, P. H., & Spirou, G. A. (2018). Glial cell expansion coincides with neural circuit formation in the developing auditory brainstem. *Developmental Neurobiology*, *78*(11), 1097–1116. <https://doi.org/10.1002/dneu.22633>
- Breuskin, I., Bodson, M., Thelen, N., Thiry, M., Borgs, L., Nguyen, L., Stolt, C., Wegner, M., Lefebvre, P. P., & Malgrange, B. (2010). Glial but not neuronal development in the cochleo-vestibular ganglion requires Sox10. *Journal of Neurochemistry*, *114*(6), 1827–1839. <https://doi.org/10.1111/j.1471-4159.2010.06897.x>
- Brown, M. C. (1987). Morphology of labeled efferent fibers in the guinea pig cochlea. *The Journal of Comparative Neurology*, *260*(4), 605–618. <https://doi.org/10.1002/cne.902600412>
- Budak, M., Zochowski, M., Booth, V., Grosh, K., & Corfas, G. (2019). A computational model for hidden hearing loss: Synaptopathy vs. myelin defects. ARO Midwinter Meeting.
- Camarero, G., Avendaño, C., Fernández-Moreno, C., Villar, A., Contreras, J., Pablo, F. D., Pichel, J. G., & Varela-Nieto, I. (2001). Delayed inner ear maturation and neuronal loss in postnatal igf-1-deficient mice. *Journal of Neuroscience*, *21*(19), 7630–7641.
- Campbell, J. P., & Henson, M. M. (1988). Olivocochlear neurons in the brainstem of the mouse. *Hearing Research*, *35*(2–3), 271–274.
- Cant, N. B., & Casseday, J. H. (1986). Projections from the anteroventral cochlear nucleus to the lateral and medial superior olivary nuclei. *The Journal of Comparative Neurology*, *247*(4), 457–476. <https://doi.org/10.1002/cne.902470406>
- Carozzi, V. A., Canta, A., & Chiorazzi, A. (2015). Chemotherapy-induced peripheral neuropathy: What do we know about mechanisms? *Neuroscience Letters*, *596*, 90–107. <https://doi.org/10.1016/j.neulet.2014.10.014>

- Cassinotti, L. R., Ji, L., Cass, N., Desai, A. S., Yuk, M. C., Palermo, A. T., Burns, J. C., & Corfas, G. (2020). *Exploring the impact of peripheral myelin disorders on inner ear structure and function using mouse models of Charcot-Marie-Tooth disease*. ARO Midwinter Meeting.
- Cassinotti, L. R., Kohrman, D. C., Borges, B., Ji, L., Desai, A. S., Yuk, M. C., & Corfas, G. (2020). *Disruption of Schwann cell ErbB receptor signaling results in auditory nerve hypomyelination and hidden hearing loss*. ARO Midwinter Meeting.
- Chan, D. K., & Chang, K. W. (2014). GJB2-associated hearing loss: Systematic review of worldwide prevalence, genotype, and auditory phenotype. *The Laryngoscope*, *124*(2), E34–E53.
- Chen, S., Velardez, M. O., Warot, X., Yu, Z.-X., Miller, S. J., Cros, D., & Corfas, G. (2006). Neuregulin 1-erbB Signaling Is Necessary for Normal Myelination and Sensory Function. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *26*(12), 3079–3086.
- Choi, J. E., Seok, J. M., Ahn, J., Ji, Y. S., Lee, K. M., Hong, S. H., Choi, B.-O., & Moon, I. J. (2018). Hidden hearing loss in patients with Charcot-Marie-Tooth disease type 1A. *Scientific Reports*, *8*, 10335.
- Clause, A., Kim, G., Sonntag, M., Weisz, C. J., Vetter, D. E., Rubsamen, R., & Kandler, K. (2014). The precise temporal pattern of prehearing spontaneous activity is necessary for tonotopic map refinement. *Neuron*, *82*(4), 822–835. <https://doi.org/10.1016/j.neuron.2014.04.001>
- Coate, T. M., Scott, M. K., & Gurjar, M. (2019). Current concepts in cochlear ribbon synapse formation. *Synapse (New York, N. Y.)*, *73*(5), e22087.
- Cope, T. E., Baguley, D. M., & Griffiths, T. D. (2015). The functional anatomy of central auditory processing. *Practical Neurology*, *15*(4), 302–308. <https://doi.org/10.1136/practneurol-2014-001073>
- Coticchia, J. M., Roeder, M. A. D., Zuliani, G. F., Gow, A., & Garbern, J. Y. (2011). Auditory testing profiles of Pelizaeus-Merzbacher disease. *International Journal of Pediatric Otorhinolaryngology Extra*, *6*(1), 23–29. <https://doi.org/10.1016/j.pedex.2010.01.001>
- Coyat, C., Cazevieuille, C., Baudoux, V., Larroze-Chicot, P., Caumes, B., & Gonzalez-Gonzalez, S. (2019). Morphological consequences of acoustic trauma on cochlear hair cells and the auditory nerve. *International Journal of Neuroscience*, *129*, 580–587.
- Cramer, K. S., & Rubel, E. W. (2016). Glial cell contributions to auditory brainstem development. *Frontiers in Neural Circuits*, *10*, 83. <https://doi.org/10.3389/fncir.2016.00083>
- Davis, J. B., & Stroobant, P. (1990). Platelet-derived growth factors and fibroblast growth factors are mitogens for rat Schwann cells. *The Journal of Cell Biology*, *110*, 1353–1360.
- de Biase, L. M. D., Nishiyama, A., & Bergles, D. E. (2010). Excitability and synaptic communication within the oligodendrocyte lineage. *The Journal of Neuroscience*, *30*(10), 3600–3611. <https://doi.org/10.1523/JNEUROSCI.6000-09.2010>
- de Villiers-Sidani, E., Alzghoul, L., Zhou, X., Simpson, K. L., Lin, R. C. S., & Merzenich, M. M. (2010). Recovery of functional and structural age-related changes in the rat primary auditory cortex with operant training. *Proceedings of the National Academy of Sciences*, *107*(31), 13900–13905.
- de Waele, C., Torres, A. C., Josset, P., & Vidal, P. P. (1996). Evidence for reactive astrocytes in rat vestibular and cochlear nuclei following unilateral inner ear lesion. *European Journal of Neuroscience*, *8*(9), 2006–2018. <https://doi.org/10.1111/j.1460-9568.1996.tb01344.x>
- Ding, D. L., McFadden, S. L., & Salvi, R. J. (2002). Calpain immunoreactivity and morphological damage in chinchilla inner ears after carboplatin. *Journal of the Association for Research in Otolaryngology*, *3*, 68–79.
- Dinh, M. L., Koppel, S. J., Korn, M. J., & Cramer, K. S. (2014). Distribution of glial cells in the auditory brainstem: Normal development and effects of unilateral lesion. *Neuroscience*, *278*, 237–252.
- Dodson, H. C., & Mohuiddin, A. (2000). Response of spiral ganglion neurones to cochlear hair cell destruction in the guinea pig. *Journal of Neurocytology*, *29*, 525–537.
- Donegan, M., Kernisant, M., Cua, C., Jasmin, L., & Ohara, P. T. (2013). Satellite glial cell proliferation in the trigeminal ganglia after chronic constriction injury of the infraorbital nerve. *Glia*, *61*, 2000–2008.
- Doty, R. L., Tourbier, I., Davis, S., Rotz, J., Cuzzocreo, J. L., Treem, J., Shephard, N., & Pham, D. L. (2012). Pure-tone auditory thresholds are not chronically elevated in multiple sclerosis. *Behavioral Neuroscience*, *126*(2), 314–324.
- Drulovic, B., Ribaric-Jankes, K., Kostic, V. S., & Sternic, N. (1993). Sudden hearing loss as the initial monosymptom of multiple sclerosis. *Neurology*, *43*(12), 2703–2705.
- Duan, M., Chen, Z., Qiu, J., Ulfendahl, M., Laurell, G., Borg, E., & Ruan, R. (2006). Low-dose, long-term caroverine administration attenuates impulse noise-induced hearing loss in the rat. *Acta Oto-Laryngologica*, *126*(11), 1140–1147.
- Ehret, G. (1979). Quantitative analysis of nerve fibre densities in the cochlea of the house mouse (*Mus musculus*). *Journal of Comparative Neurology*, *183*(1), 73–88.
- Elbaz, B., & Popko, B. (2019). Molecular control of oligodendrocyte development. *Trends in Neurosciences*, *42*(Exp. Neurol. 283 2016), 263–277.
- Elezgarai, I., Bilbao, A., Mateos, J. M., Azkue, J. J., Benitez, R., Osorio, A., Diez, J., Puente, N., Donate-Oliver, F., & Grandes, P. (2001). Group II metabotropic glutamate receptors are differentially expressed in the medial nucleus of the trapezoid body in the developing and adult rat. *Neuroscience*, *104*(2), 487–498.
- Elkon, R., Milon, B., Morrison, L., Shah, M., Vijayakumar, S., Racherla, M., Leitch, C. C., Silipino, L., Hadi, S., Weiss-Gayet, M., Barras, E., Schmid, C. D., Ait-Lounis, A., Barnes, A., Song, Y., Eisenman, D. J., Eliyahu, E., Frolenkov, G. I., Strome, S. E., ... Hertzano R. (2015). RFX transcription factors are essential for hearing in mice. *Nature Communications*, *6*, (1). <http://dx.doi.org/10.1038/ncomms9549>.
- Elson, K., Simmons, A., & Speck, P. (2004). Satellite cell proliferation in murine sensory ganglia in response to scarification of the skin. *Glia*, *45*, 105–109.
- Emmorey, K., Allen, J. S., Bruss, J., Schenker, N., & Damasio, H. (2003). A morphometric analysis of auditory brain regions in congenitally deaf adults. *Proceedings of the National Academy of Sciences*, *100*(17), 10049–10054.
- Erazo-Fischer, E., Striessnig, J., & Taschenberger, H. (2007). The role of physiological afferent nerve activity during in vivo maturation of the calyx of held synapse. *The Journal of Neuroscience*, *27*(7), 1725–1737. <https://doi.org/10.1523/JNEUROSCI.4116-06.2007>
- Ernfors, P., Van De Water, T., Loring, J., & Jaenisch, R. (1995). Complementary roles of BDNF and NT-3 in vestibular and auditory development. *Neuron*, *14*(6), 1153–1164. [https://doi.org/10.1016/0896-6273\(95\)90263-5](https://doi.org/10.1016/0896-6273(95)90263-5)
- Eybalin, M., Norenberg, M. D., & Renard, N. (1996). Glutamine synthetase and glutamate metabolism in the guinea pig cochlea. *Hearing Research*, *101*, 93–101.
- Farah, R., Schmithorst, V. J., Keith, R. W., & Holland, S. K. (2014). Altered white matter microstructure underlies listening difficulties

- in children suspected of auditory processing disorders: A DTI study. *Brain and Behavior*, 4(4), 531–543.
- Fariñas, I., Jones, K. R., Tessarollo, L., Vigers, A. J., Huang, E., Kirstein, M., de Caprona, D. C., Coppola, V., Backus, C., Reichardt, L. F., & Fritsch, B. (2001). Spatial Shaping of Cochlear Innervation by Temporally Regulated Neurotrophin Expression. *Journal of Neuroscience*, 21(16), 6170–6180. <https://doi.org/10.1523/JNEUROSCI.21-16-06170.2001>
- Faulkner, J. R., Herrmann, J. E., Woo, M. J., Tansey, K. E., Doan, N. B., & Sofroniew, M. V. (2004). Reactive astrocytes protect tissue and preserve function after spinal cord injury. *Journal of Neuroscience*, 24(9), 2143–2155. <https://doi.org/10.1523/JNEUROSCI.3547-03.2004>
- Felix, H. (2002). Anatomical differences in the peripheral auditory system of mammals and man. *Advances in Oto-Rhino-Laryngology*, 59, 1–10.
- Fettiplace, R. (2017). Hair cell transduction, tuning, and synaptic transmission in the mammalian cochlea. *Comprehensive Physiology*, 7(4), 1197–1227.
- Fields, R. D. (2008). White matter in learning, cognition and psychiatric disorders. *Trends in Neurosciences*, 31(7), 361–370. <https://doi.org/10.1016/j.tins.2008.04.001>
- Fledrich, R., Stassart, R. M., Klink, A., Rasch, L. M., Prukop, T., Haag, L., Czesnik, D., Kungl, T., Abdelaal, T. A. M., Keric, N., Stadelmann, C., Brück, W., Nave, K.-A., & Sereda, M. W. (2014). Soluble neuregulin-1 modulates disease pathogenesis in rodent models of Charcot-Marie-Tooth disease 1A. *Nature Medicine*, 20(9), 1055–1061.
- Flores-Otero, J., Xue, H. Z., & Davis, R. L. (2007). Reciprocal regulation of presynaptic and postsynaptic proteins in bipolar spiral ganglion neurons by neurotrophins. *Journal of Neuroscience*, 27(51), 14023–14034. <https://doi.org/10.1523/JNEUROSCI.3219-07.2007>
- Ford, M. C., Alexandrova, O., Cossell, L., Stange-Marten, A., Sinclair, J., Kopp-Scheinflug, C., Pecka, M., Attwell, D., & Grothe, B. (2015). Tuning of ranvier node and internode properties in myelinated axons to adjust action potential timing. *Nature Communications*, 6, (1). <http://dx.doi.org/10.1038/ncomms9073>.
- Ford, M. C., Grothe, B., & Klug, A. (2009). Fenestration of the calyx of Held occurs sequentially along the tonotopic axis, is influenced by afferent activity, and facilitates glutamate clearance. *The Journal of Comparative Neurology*, 514(1), 92–106. <https://doi.org/10.1002/cne.21998>
- Foss-Feig, J. H., Schauder, K. B., Key, A. P., Wallace, M. T., & Stone, W. L. (2017). Audition-specific temporal processing deficits associated with language function in children with autism spectrum disorder. *Autism Research*, 10(11), 1845–1856. <https://doi.org/10.1002/aur.1820>
- Foster, A. Y., Bujalka, H., & Emery, B. (2019). Axoglial interactions in myelin plasticity: Evaluating the relationship between neuronal activity and oligodendrocyte dynamics. *Glia*, 67(11), 2038–2049. <https://doi.org/10.1002/glia.23629>
- Fredrich, M., Zeber, A. C., Hildebrandt, H., & Illing, R.-B. (2013). Differential molecular profiles of astrocytes in degeneration and re-innervation after sensory deafferentation of the adult rat cochlear nucleus. *European Journal of Neuroscience*, 38(1), 2041–2056. <https://doi.org/10.1111/ejn.12200>
- Fuchs, P. A., Glowatzki, E., & Moser, T. (2003). The afferent synapse of cochlear hair cells. *Current Opinion in Neurobiology*, 13(4), 452–458. [https://doi.org/10.1016/S0959-4388\(03\)00098-9](https://doi.org/10.1016/S0959-4388(03)00098-9)
- Fuentes-Santamaria, V., Alvarado, J. C., Melgar-Rojas, P., Gabaldon-Ull, M. C., Miller, J. M., & Juiz, J. M. (2017). The role of glia in the peripheral and central auditory system following noise overexposure: Contribution of TNF-alpha and IL-1beta to the pathogenesis of hearing loss. *Frontiers in Neuroanatomy*, 11, 9. <https://www.frontiersin.org/articles/10.3389/fnana.2017.00009/full>.
- Furness, D. N., & Lawton, D. M. (2003). Comparative distribution of glutamate transporters and receptors in relation to afferent innervation density in the mammalian cochlea. *Journal of Neuroscience*, 23(36), 11296–11304. <https://doi.org/10.1523/JNEUROSCI.23-36-11296.2003>
- Furst, M., & Levine, R. A. (2015). Hearing disorders in multiple sclerosis. *Handbook of Clinical Neurology*, 129, 649–665.
- Garbern, J. Y. (2007). Pelizaeus-Merzbacher disease: Genetic and cellular pathogenesis. *Cellular and Molecular Life Sciences*, 64(1), 5–65. <https://doi.org/10.1007/s00018-006-6182-8>
- Garratt, A. N., Voiculescu, O., Topilko, P., Charnay, P., & Birchmeier, C. (2000). A dual role of erbB2 in myelination and in expansion of the Schwann cell precursor pool. *The Journal of Cell Biology*, 148(5), 1035–1046. <https://doi.org/10.1083/jcb.148.5.1035>
- Giuliani, N., Holte, L., Shy, M., & Grider, T. (2019). The audiologic profile of patients with Charcot-Marie Tooth neuropathy can be characterised by both cochlear and neural deficits. *International Journal of Audiology*, 58(12), 1–11. <https://doi.org/10.1080/14992027.2019.1633022>
- Glowatzki, E., Cheng, N., Hiel, H., Yi, E., Tanaka, K., Ellis-Davies, G. C. R., Rothstein, J. D., & Bergles, D. E. (2006). The glutamate-aspartate transporter GLAST mediates glutamate uptake at inner hair cell afferent synapses in the mammalian cochlea. *Journal of Neuroscience*, 26(29), 7659–7664. <https://doi.org/10.1523/JNEUROSCI.1545-06.2006>
- Gold, J. R., & Bajo, V. M. (2014). Insult-induced adaptive plasticity of the auditory system. *Frontiers in Neuroscience*, 8, 110. <https://www.frontiersin.org/articles/10.3389/fnins.2014.00110/full>.
- Goldberg, J. M., & Brown, P. B. (1969). Response of binaural neurons of dog superior olivary complex to dichotic tonal stimuli: Some physiological mechanisms of sound localization. *Journal of Neurophysiology*, 32(4), 613–636. <https://doi.org/10.1152/jn.1969.32.4.613>
- Green, S. H., Bailey, E., Wang, Q., & Davis, R. L. (2012). The Trk A, B, C's of neurotrophins in the cochlea. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology*, 295(11), 1877–1895.
- Grothe, B., & Pecka, M. (2014). The natural history of sound localization in mammals—A story of neuronal inhibition. *Frontiers in Neural Circuits*, 8, 116. <https://www.frontiersin.org/articles/10.3389/fncir.2014.00116/full>.
- Grothe, C., & Ninkhah, G. (2001). The role of basic fibroblast growth factor in peripheral nerve regeneration. *Anatomy and Embryology*, 204, 171–177.
- Hackett, T. A., Guo, Y., Clause, A., Hackett, N. J., Garbett, K., Zhang, P., Polley, D. B., & Mirnics, K. (2015). Transcriptional maturation of the mouse auditory forebrain. *BMC Genomics*, 16(1), 606. <https://bmcbgenomics.biomedcentral.com/articles/10.1186/s12864-015-1709-8>.
- Hakuba, N., Koga, K., Gyo, K., Usami, S.-I., & Tanaka, K. (2000). Exacerbation of noise-induced hearing loss in mice lacking the glutamate transporter GLAST. *Journal of Neuroscience*, 20(23), 8750–8753. <https://doi.org/10.1523/JNEUROSCI.20-23-08750.2000>
- Han, R., Yang, Y. M., Dietrich, J., Luebke, A., Mayer-Pröschel, M., & Noble, M. (2008). Systemic 5-fluorouracil treatment causes a syndrome of delayed myelin destruction in the central nervous system. *Journal of Biology*, 7(4), 12. <https://doi.org/10.1186/jbiol69>.

- Hansen, M. R., Vijapurkar, U., Koland, J. G., & Green, S. H. (2001). Reciprocal signaling between spiral ganglion neurons and Schwann cells involves neuregulin and neurotrophins. *Hearing Research*, *161*, 87–98.
- Harris, K. C., & Dubno, J. R. (2017). Age-related deficits in auditory temporal processing: Unique contributions of neural dyssynchrony and slowed neuronal processing. *Neurobiology of Aging*, *53*, 150–158. <https://doi.org/10.1016/j.neurobiolaging.2017.01.008>.
- Hashimoto, K., Hickman, T. T., Suzuki, J., Ji, L., Kohrman, D. C., Corfas, G., & Liberman, M. C. (2019). Protection from noise-induced cochlear synaptopathy by virally mediated overexpression of NT3. *Scientific Reports*, *9*(1), 15362.
- Hashisaki, G. T., & Rubel, E. W. (1989). Effects of unilateral cochlea removal on anteroventral cochlear nucleus neurons in developing gerbils. *Journal of Comparative Neurology*, *283*(4), 465–473. <https://doi.org/10.1002/cne.902830402>
- Hellmann, M. A., Steiner, I., & Mosberg-Galili, R. (2011). Sudden sensorineural hearing loss in multiple sclerosis: Clinical course and possible pathogenesis. *Acta Neurologica Scandinavica*, *124*(4), 245–249. <https://doi.org/10.1111/j.1600-0404.2010.01463.x>
- Henry, K. R. (1979). Auditory brainstem volume-conducted responses: Origins in the laboratory mouse. *Journal of the American Auditory Society*, *4*(5), 173–178.
- Hribar, M., Suput, D., Carvalho, A. A., Battelino, S., & Vovk, A. (2014). Structural alterations of brain grey and white matter in early deaf adults. *Hearing Research*, *318*, 1–10.
- Hsieh, C. Y., Hong, C. T., & Cramer, K. S. (2007). Deletion of EphA4 enhances deafferentation-induced ipsilateral sprouting in auditory brainstem projections. *Journal of Comparative Neurology*, *504*(5), 508–518. <https://doi.org/10.1002/cne.21465>
- Huang, L. C., Thorne, P. R., Housley, G. D., & Montgomery, J. M. (2007). Spatiotemporal definition of neurite outgrowth, refinement and retraction in the developing mouse cochlea. *Development*, *134*(16), 2925–2933. <https://doi.org/10.1242/dev.001925>
- Ishibashi, T., Dakin, K. A., Stevens, B., Lee, P. R., Kozlov, S. V., Stewart, C. L., & Fields, R. D. (2006). Astrocytes promote myelination in response to electrical impulses. *Neuron*, *49*(6), 823–832. <https://doi.org/10.1016/j.neuron.2006.02.006>
- Jäger, W., Gojny, M., Herrera-Marschitz, M., Brundin, L., Fransson, A., & Canlon, B. (2000). Noise-induced aspartate and glutamate efflux in the guinea pig cochlea and hearing loss. *Experimental Brain Research*, *134*(4), 426–434. <https://doi.org/10.1007/s002210000470>
- Jang, M., Gould, E., Xu, J., Kim, E. J., & Kim, J. H. (2019). Oligodendrocytes regulate presynaptic properties and neurotransmission through BDNF signaling in the mouse brainstem. *eLife*, *8*, e42156.
- Janz, P., & Illing, R.-B. (2014). A role for microglial cells in reshaping neuronal circuitry of the adult rat auditory brainstem after its sensory deafferentation: Microglia in brainstem neuroplasticity. *Journal of Neuroscience Research*, *92*(4), 432–445. <https://doi.org/10.1002/jnr.23334>
- Jeffries, M. A., Urbanek, K., Torres, L., Wendell, S. G., Rubio, M. E., & Fyffe-Maricich, S. L. (2016). ERK1/2 activation in preexisting oligodendrocytes of adult mice drives new myelin synthesis and enhanced CNS function. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *36*(35), 9186–9200. <https://doi.org/10.1523/JNEUROSCI.1444-16.2016>
- Jiang, M., Karasawa, T., & Steyger, P. S. (2017). Aminoglycoside-induced cochleotoxicity: A review. *Frontiers in Cellular Neuroscience*, *11*, 308. <https://doi.org/10.3389/fncel.2017.00308>.
- Jin, Z.-H., Kikuchi, T., Tanaka, K., & Kobayashi, T. (2003). Expression of glutamate transporter GLAST in the developing mouse cochlea. *The Tohoku Journal of Experimental Medicine*, *200*(3), 137–144. <https://doi.org/10.1620/tjem.200.137>
- Johnson, S. L., Ceriani, F., Houston, O., Polishchuk, R., Polishchuk, E., Crispino, G., Zorzi, V., Mammano, F., & Marcotti, W. (2017). Connexin-mediated signaling in nonsensory cells is crucial for the development of sensory inner hair cells in the mouse cochlea. *The Journal of Neuroscience*, *37*(2), 258–268. <https://doi.org/10.1523/JNEUROSCI.2251-16.2016>
- Johnson, S. L., Eckrich, T., Kuhn, S., Zampini, V., Franz, C., Ranatunga, K. M., Roberts, T. P., Masetto, S., Knipper, M., Kros, C. J., & Marcotti, W. (2011). Position-dependent patterning of spontaneous action potentials in immature cochlear inner hair cells. *Nature Neuroscience*, *14*(6), 711–717. <https://doi.org/10.1038/nn.2803>
- Johnson, S. L., Kennedy, H. J., Holley, M. C., Fettiplace, R., & Marcotti, W. (2012). The resting transducer current drives spontaneous activity in prehearing mammalian cochlear inner hair cells. *Journal of Neuroscience*, *32*(31), 10479–10483. <https://doi.org/10.1523/JNEUROSCI.0803-12.2012>
- Johnsson, L.-G. (1974). Sequence of degeneration of Corti's organ and its first-order neurons. *Annals of Otology, Rhinology & Laryngology*, *83*(3), 294–303. <https://doi.org/10.1177/000348947408300303>
- Jones, A., & May, B. J. (2017). Improving the reliability of tinnitus screening in laboratory animals. *Journal of the Association for Research in Otolaryngology*, *18*(1), 183–195. <https://doi.org/10.1007/s10162-016-0597-1>
- Jones, T. A., Leake, P. A., Snyder, R. L., Stakhovskaya, O., & Bonham, B. (2007). Spontaneous discharge patterns in cochlear spiral ganglion cells before the onset of hearing in cats. *Journal of Neurophysiology*, *98*(4), 1898–1908. <https://doi.org/10.1152/jn.00472.2007>
- Kahkonen, S., Yamashita, H., Rytsala, H., Suominen, K., Ahveninen, J., & Isometsa, E. (2007). Dysfunction in early auditory processing in major depressive disorder revealed by combined MEG and EEG. *Journal of Psychiatry and Neuroscience*, *32*(5), 316–322.
- Kaltenbach, J. A., & Afman, C. E. (2000). Hyperactivity in the dorsal cochlear nucleus after intense sound exposure and its resemblance to tone-evoked activity: A physiological model for tinnitus. *Hearing Research*, *140*(1–2), 165–172.
- Kim, D.-J., Park, S.-Y., Kim, J., Lee, D. H., & Park, H.-J. (2009). Alterations of white matter diffusion anisotropy in early deafness. *NeuroReport*, *20*(11), 1032–1036. <https://doi.org/10.1097/WNR.0b013e32832e0cdd>
- Kim, J. H., Renden, R., & von Gersdorff, H. (2013). Dysmyelination of auditory afferent axons increases the jitter of action potential timing during high-frequency firing. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *33*(22), 9402–9407. <https://doi.org/10.1523/JNEUROSCI.3389-12.2013>
- Kim, K. X., & Rutherford, M. A. (2016). Maturation of NaV and KV channel topographies in the auditory nerve spike initiator before and after developmental onset of hearing function. *The Journal of Neuroscience*, *36*(7), 2111–2118.
- Kim, S. E., Turkington, K., Kushmerick, C., & Kim, J. H. (2013). Central dysmyelination reduces the temporal fidelity of synaptic transmission and the reliability of postsynaptic firing during high-frequency stimulation. *Journal of Neurophysiology*, *110*(7), 1621–1630. <https://doi.org/10.1152/jn.00117.2013>
- Kitzes, L. M., Kageyama, G. H., Semple, M. N., & Kil, J. (1995). Development of ectopic projections from the ventral cochlear nucleus to the superior olivary complex induced by neonatal ablation

- of the contralateral cochlea. *Journal of Comparative Neurology*, 353(3), 341–363. <https://doi.org/10.1002/cne.903530303>
- Kleopa, K. A., & Scherer, S. S. (2006). Molecular genetics of X-linked Charcot-Marie-Tooth disease. *NeuroMolecular Medicine*, 8, 107–122.
- Klumpp, R. G., & Eady, H. R. (1956). Some measurements of interaural time difference thresholds. *The Journal of the Acoustical Society of America*, 28(5), 859–860. <https://doi.org/10.1121/1.1908493>
- Kohrman, D. C., Wan, G., Cassinotti, L. R., & Corfas, Gabriel (2020). Hidden hearing loss: A disorder with multiple etiologies and mechanisms. *Cold Spring Harbor Perspectives in Medicine*, 10, (1), a035493. <http://dx.doi.org/10.1101/cshperspect.a035493>.
- Kolson, D. R., Wan, J., Wu, J., Dehoff, M., Brandebura, A. N., Qian, J., Mathers, P. H., & Spirou, G. A. (2015). Temporal patterns of gene expression during calyx of held development. *Developmental Neurobiology*, 76(2), 166–189. <https://doi.org/10.1002/dneu.22306>
- Kopp-Scheinpflug, C., & Tempel, B. L. (2015). Decreased temporal precision of neuronal signaling as a candidate mechanism of auditory processing disorder. *Hearing Research*, 330(Pt B), 213–220.
- Korn, M. J., Koppel, S. J., & Cramer, K. S. (2011). Astrocyte-secreted factors modulate a gradient of primary dendritic arbors in nucleus laminaris of the avian auditory brainstem. *PLoS One*, 6(11), e27383.
- Korn, M. J., Koppel, S. J., Li, L. H., Mehta, D., Mehta, S. B., Seidl, A. H., & Cramer, K. S. (2012). Astrocyte-secreted factors modulate the developmental distribution of inhibitory synapses in nucleus laminaris of the avian auditory brainstem. *The Journal of Comparative Neurology*, 520(6), 1262–1277. <https://doi.org/10.1002/cne.22786>
- Kostandy, B. B. (2012). The role of glutamate in neuronal ischemic injury: The role of spark in fire. *Neurological Sciences*, 33(2), 223–237. <https://doi.org/10.1007/s10072-011-0828-5>
- Kovach, M. J., Campbell, K. C. M., Herman, K., Waggoner, B., Gelber, D., Hughes, L. F., & Kimonis, V. E. (2002). Anticipation in a unique family with Charcot-Marie-Tooth syndrome and deafness: Delineation of the clinical features and review of the literature. *American Journal of Medical Genetics*, 108(4), 295–303. <https://doi.org/10.1002/ajmg.10223>
- Kraus, K. S., & Illing, R.-B. (2004). Superior olivary contributions to auditory system plasticity: Medial but not lateral olivocochlear neurons are the source of cochleotomy-induced GAP-43 expression in the ventral cochlear nucleus. *The Journal of Comparative Neurology*, 475(3), 374–390. <https://doi.org/10.1002/cne.20180>
- Kuan, C.-C., Sano, M., Kaga, K., Kodama, M., & Kodama, K. (2009). Hearing profile and MRI myelination of auditory pathway in Pelizaeus-Merzbacher disease. *Acta Oto-Laryngologica*, 128(5), 539–546. <https://doi.org/10.1080/00016480701258713>
- Land, R., Burghard, A., & Kral, A. (2016). The contribution of inferior colliculus activity to the auditory brainstem response (ABR) in mice. *Hearing Research*, 341, 109–118.
- Lavigne-Rebillard, M., & Pujol, R. (1988). Hair cell innervation in the fetal human cochlea. *Acta Oto-Laryngologica*, 105(5–6), 398–402.
- Leake, P. A., & Hradek, G. T. (1988). Cochlear pathology of long term neomycin induced deafness in cats. *Hearing Research*, 33, 11–33.
- Lieberman, M. C., & Kujawa, S. G. (2017). Cochlear synaptopathy in acquired sensorineural hearing loss: Manifestations and mechanisms. *Hearing Research*, 349, 138–147. <https://doi.org/10.1016/j.heares.2017.01.003>
- Lin, H.-C., Ren, Y., Lysaght, A. C., Kao, S.-Y., & Stankovic, K. M. (2019). Proteome of normal human perilymph and perilymph from people with disabling vertigo. *PLoS One*, 14(6), e0218292.
- Lin, L., Koutnouyan, H., Baird, A., & Ryan, A. F. (1993). Acidic and basic FGF mRNA expression in the adult and developing rat cochlea. *Hearing Research*, 69(1–2), 182–193.
- Lin, S.-C., & Bergles, D. E. (2003). Synaptic signaling between GABAergic interneurons and oligodendrocyte precursor cells in the hippocampus. *Nature Neuroscience*, 7(1), 24–32. <https://doi.org/10.1038/nn1162>
- Lindsey, B. G. (1975). Fine structure and distribution of axon terminals from cochlear nucleus on neurons in the medial superior olivary nucleus of the cat. *The Journal of Comparative Neurology*, 160(1), 81–103. <https://doi.org/10.1002/cne.901600106>
- Liu, C., Glowatzki, E., & Fuchs, P. A. (2015). Unmyelinated type II afferent neurons report cochlear damage. *Proceedings of the National Academy of Sciences*, 112(47), 14723–14727
- Liu, W., Edin, F., Atturo, F., Rieger, G., Löwenheim, H., Senn, P., Blumer, M., Schrott-Fischer, A., Rask-Andersen, H., & Glueckert, R. (2015). The pre- and post-somatic segments of the human type I spiral ganglion neurons—Structural and functional considerations related to cochlear implantation. *Neuroscience*, 284, 470–482.
- Liu, W., Glueckert, R., Schrott-Fischer, A., & Rask-Andersen, H. (2020). Human cochlear microanatomy—An electron microscopy and super-resolution structured illumination study and review. *Hearing, Balance and Communication*, 18(4), 256–269. <https://doi.org/10.1080/21695717.2020.1807259>
- Locher, H., de Groot, J. C. M. J., van Iperen, L., Huisman, M. A., Frijns, J. H. M., & Chuva de Sousa Lopes, S. M. (2014). Distribution and development of peripheral glial cells in the human fetal cochlea. *PLoS One*, 9(1), e88066.
- Lurie, D. I., & Rubel, E. W. (1994). Astrocyte proliferation in the chick auditory brainstem following cochlea removal. *The Journal of Comparative Neurology*, 346(2), 276–288. <https://doi.org/10.1002/cne.903460207>
- Makinodan, M., Rosen, K. M., Ito, S., & Corfas, G. (2012). A critical period for social experience-dependent oligodendrocyte maturation and myelination. *Science*, 337(6100), 1357–1360. <https://doi.org/10.1126/science.1220845>
- Mammano, F. (2018). Inner ear Connexin channels: Roles in development and maintenance of cochlear function. *Cold Spring Harbor Perspectives in Medicine*, 9(7), a033233.
- Mammano, F., & Bortolozzi, M. (2018). Ca²⁺ signaling, apoptosis and autophagy in the developing cochlea: Milestones to hearing acquisition. *Cell Calcium*, 70(J. Acoust. Soc. Am. 87 1990), 117–126.
- Mao, Y., Reiprich, S., Wegner, M., & Fritzsche, B. (2014). Targeted deletion of Sox10 by Wnt1-cre defects neuronal migration and projection in the mouse inner ear. *PLoS One*, 9, e94580.
- Marín, O., & Rubenstein, J. L. R. (2003). Cell migration in the fore-brain. *Annual Review of Neuroscience*, 26(1), 441–483.
- McFadden, S. L., Ding, D., Jiang, H., & Salvi, R. J. (2004). Time course of efferent fiber and spiral ganglion cell degeneration following complete hair cell loss in the chinchilla. *Brain Research*, 997(1), 40–51. <https://doi.org/10.1016/j.brainres.2003.10.031>
- McKay, S. M., & Oleskevich, S. (2007). The role of spontaneous activity in development of the endbulb of Held synapse. *Hearing Research*, 230(1–2), 53–63.
- McLachlan, N. M., Phillips, D. S., Rossell, S. L., & Wilson, S. J. (2013). Auditory processing and hallucinations in schizophrenia. *Schizophrenia Research*, 150(2–3), 380–385.
- Mei, L., & Nave, K.-A. (2014). Neuregulin-ERBB signaling in the nervous system and neuropsychiatric diseases. *Neuron*, 83(1), 27–49. <https://doi.org/10.1016/j.neuron.2014.06.007>

- Meyers, C. A. (2008). How chemotherapy damages the central nervous system. *Journal of Biology*, 7, 11. <https://doi.org/10.1186/jbiol73>.
- Michailov, G. V., Sereda, M. W., Brinkmann, B. G., Fischer, T. M., Haug, B., Birchmeier, C., Role, L., Lai, C., Schwab, M. H., & Nave, K.-A. (2004). Axonal neuregulin-1 regulates myelin sheath thickness. *Science*, 304(5671), 700–703. <https://doi.org/10.1126/science.1095862>
- Mills, A. W. (1960). Lateralization of high-frequency tones. *Journal of the Acoustical Society of America*, 32, 132–134.
- Monje, M. (2018). Myelin plasticity and nervous system function. *Annual Review of Neuroscience*, 41(1), 61–76. <https://doi.org/10.1146/annurev-neuro-080317-061853>
- Moore, D. R., & Kowalchuk, N. E. (1988). Auditory brainstem of the ferret: Effects of unilateral cochlear lesions on cochlear nucleus volume and projections to the inferior colliculus. *Journal of Comparative Neurology*, 272(4), 503–515. <https://doi.org/10.1002/cne.902720405>
- Moore, J. K., & Linthicum, F. H. (2001). Myelination of the human auditory nerve: Different time courses for Schwann cell and glial myelin. *Annals of Otology, Rhinology & Laryngology*, 110, 655–661.
- Moore, J. K., Perazzo, L. M., & Braun, A. (1995). Time course of axonal myelination in the human brainstem auditory pathway. *Hearing Research*, 87(1–2), 21–31.
- Morelli, K. H., Seburn, K. L., Schroeder, D. G., Spaulding, E. L., Dionne, L. A., Cox, G. A., & Burgess, R. W. (2017). Severity of demyelinating and axonal neuropathy mouse models is modified by genes affecting structure and function of peripheral nodes. *Cell Reports*, 18(13), 3178–3191. <https://doi.org/10.1016/j.celrep.2017.03.009>
- Morlet, T., Nagao, K., Bean, S. C., Mora, S. E., Hopkins, S. E., & Hobson, G. M. (2018). Auditory function in Pelizaeus-Merzbacher disease. *Journal of Neurology*, 265(7), 1580–1589. <https://doi.org/10.1007/s00415-018-8884-x>
- Morley, B. J. (1998). ARIA is heavily expressed in rat peripheral auditory and vestibular ganglia. *Molecular Brain Research*, 54(1), 170–174. [https://doi.org/10.1016/S0169-328X\(97\)00355-0](https://doi.org/10.1016/S0169-328X(97)00355-0)
- Morris, J. K., Maklad, A., Hansen, L. A., Feng, F., Sorensen, C., Lee, K. F., Macklin, W. B., & Fritsch, B. (2006). A disorganized innervation of the inner ear persists in the absence of ErbB2. *Brain Research*, 1091, 186–199.
- Moser, T., Grabner, C. P., & Schmitz, F. (2019). Sensory processing at ribbon synapses in the retina and the cochlea. *Physiological Reviews*, 100(1), 103–144. <https://doi.org/10.1152/physrev.00026.2018>
- Mostafapour, S. P., Cochran, S. L., Puerto, N. M. D., & Rubel, E. W. (2000). Patterns of cell death in mouse anteroventral cochlear nucleus neurons after unilateral cochlea removal. *Journal of Comparative Neurology*, 426(4), 561–571. [https://doi.org/10.1002/1096-9861\(20001030\)426:4<561:AID-CNE5>3.0.CO;2-G](https://doi.org/10.1002/1096-9861(20001030)426:4<561:AID-CNE5>3.0.CO;2-G)
- Nascimento, D. S. M., Castro-Lopes, J. M., & Moreira Neto, F. L. (2014). Satellite glial cells surrounding primary afferent neurons are activated and proliferate during monoarthritis in rats: Is there a role for ATF3? *PLoS One*, 9, e108152. <https://doi.org/10.1371/journal.pone.0108152>
- Nelson, K. R., Gilmore, R. L., & Massey, A. (1988). Acoustic nerve conduction abnormalities in Guillain-Barré syndrome. *Neurology*, 38, 1263–1266.
- Nicholson, G., & Corbett, A. (1996). Slowing of central conduction in X-linked Charcot-Marie-Tooth neuropathy shown by brain stem auditory evoked responses. *Journal of Neurology, Neurosurgery & Psychiatry*, 61(1), 43–46. <https://jnp.bmj.com/content/61/1/43>.
- Nimmerjahn, A., & Bergles, D. E. (2015). Large-scale recording of astrocyte activity. *Current Opinion in Neurobiology*, 32, 95–106.
- O'Connor, K. (2012). Auditory processing in autism spectrum disorder: A review. *Neuroscience & Biobehavioral Reviews*, 36(2), 836–854. <https://doi.org/10.1016/j.neubiorev.2011.11.008>
- O'Connor, L. T., Goetz, B. D., Kwicien, J. M., Delaney, K. H., Fletch, A. L., & Duncan, I. D. (1999). Insertion of a retrotransposon in Mbp disrupts mRNA splicing and myelination in a new mutant rat. *The Journal of Neuroscience*, 19(9), 3404–3413.
- Oestreicher, E., Wolfgang, A., & Felix, D. (2002). Neurotransmission of the cochlear inner hair cell synapse-implications for inner ear therapy. *Advances in Oto-Rhino-Laryngology*, 59, 131–139.
- Okano, T., Xuan, S., & Kelley, M. W. (2011). Insulin-like growth factor signaling regulates the timing of sensory cell differentiation in the mouse cochlea. *The Journal of Neuroscience*, 31(49), 18104–18118. <https://doi.org/10.1523/JNEUROSCI.3619-11.2011>
- Panganiban, C. H., Barth, J. L., Darbelli, L., Xing, Y., Zhang, J., Li, H., Noble, K. V., Liu, T., Brown, L. N., Schulte, B. A., Richard, S., & Lang, H. (2018). Noise-induced dysregulation of quaking RNA binding proteins contributes to auditory nerve demyelination and hearing loss. *Journal of Neuroscience*, 38(10), 2551–2568.
- Parks, T. N., & Rubel, E. W. (1975). Organization and development of brain stem auditory nuclei of the chicken: Organization of projections from n. magnocellularis to n. laminaris. *The Journal of Comparative Neurology*, 164(4), 435–448. <https://doi.org/10.1002/cne.901640404>
- Petitpré, C., Wu, H., Sharma, A., Tokarska, A., Fontanet, P., Wang, Y., Helmbacher, F., Yackle, K., Silberberg, G., Hadjab, S., & Lallemand, F. (2018). Neuronal heterogeneity and stereotyped connectivity in the auditory afferent system. *Nature Communications*, 9(1), 3691. <https://doi.org/10.1038/s41467-018-06033-3>.
- Pirvola, U., Cao, Y., Oellig, C., Suoqi, Z., Pettersson, R. F., & Ylikoski, J. (1995). The site of action of neuronal acidic fibroblast growth factor is the organ of Corti of the rat cochlea. *Proceedings of the National Academy of Sciences*, 92(20), 9269–9273.
- Plack, C. J., Barker, D., & Prendergast, G. (2014). Perceptual consequences of “hidden” hearing loss. *Trends in Hearing*, 18, 1–11. <https://doi.org/10.1177/2F2331216514550621>.
- Pujol, R., & Puel, J. L. (1999). Excitotoxicity, synaptic repair, and functional recovery in the mammalian cochlea: A review of recent findings. *Annals of the New York Academy of Sciences*, 884(1), 249–254. <https://doi.org/10.1111/j.1749-6632.1999.tb08646.x>
- Rance, G. (2005). Auditory neuropathy/Dys-synchrony and its perceptual consequences. *Trends in Amplification*, 9(1), 1–43. <https://doi.org/10.1177/108471380500900102>
- Rance, G., Ryan, M. M., Bayliss, K., Gill, K., O'Sullivan, C., & Whitechurch, M. (2012). Auditory function in children with Charcot-Marie-Tooth disease. *Brain*, 135, 1412–1422.
- Rasband, M. N. (2016). Glial contributions to neural function and disease. *Molecular & Cellular Proteomics*, 15(2), 355–361. <https://doi.org/10.1074/mcp.R115.053744>
- Rasband, M. N., Peles, E., Trimmer, J. S., Levinson, S. R., Lux, S. E., & Shrager, P. (1999). Dependence of nodal sodium channel clustering on Paranodal Axoglial contact in the developing CNS. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 19(17), 7516–7528.
- Rattay, F., Lutter, P., & Felix, H. (2001). A model of the electrically excited human cochlear neuron. *Hearing Research*, 153, 43–63.
- Rattay, F., Potrusil, T., Wenger, C., Wise, A. K., Glueckert, R., & Schrott-Fischer, A. (2013). Impact of morphometry, myelination

- and synaptic current strength on spike conduction in human and cat spiral ganglion neurons. *PLoS One*, 8(11), e79256.
- Ray, B., Roy, T. S., Wadhwa, S., & Roy, K. K. (2005). Development of the human fetal cochlear nerve: A morphometric study. *Hearing Research*, 202(1–2), 74–86.
- Renden, R., Taschenberger, H., Puente, N., Rusakov, D. A., Duvoisin, R., Wang, L. Y., Lehre, K. P., & von Gersdorff, H. (2005). Glutamate transporter studies reveal the pruning of metabotropic glutamate receptors and absence of AMPA receptor desensitization at mature calyx of Held synapses. *Journal of Neuroscience*, 25(37), 8482–8497. <https://doi.org/10.1523/JNEUROSCI.1848-05.2005>
- Reyes-Haro, D., Muller, J., Boresch, M., Pivneva, T., Benedetti, B., Scheller, A., Nolte, C., & Kettenmann, H. (2010). Neuron-astrocyte interactions in the medial nucleus of the trapezoid body. *Journal of General Physiology*, 135(6), 583–594. <https://doi.org/10.1085/jgp.200910354>
- Rocha-Muniz, C. N., Befi-Lopes, D. M., & Schochat, E. (2012). Investigation of auditory processing disorder and language impairment using the speech-evoked auditory brainstem response. *Hearing Research*, 294(1–2), 143–152.
- Rocha-Muniz, C. N., Befi-Lopes, D. M., & Schochat, E. (2014). Sensitivity, specificity and efficiency of speech-evoked ABR. *Hearing Research*, 317, 15–22.
- Ropper, A. H., & Chiappa, K. H. (1986). Evoked potentials in Guillain-Barré syndrome. *Neurology*, 36, 587–590.
- Rossi, G., Robecchi, M. G., & Penna, M. (1976). Effects of acoustic trauma on corti's ganglion. *Acta Oto-Laryngologica*, 81, 270–277.
- Rosskothén-Kuhl, N., Hildebrandt, H., Birkenhager, R., & Illing, R. B. (2018). Astrocyte hypertrophy and microglia activation in the rat auditory midbrain is induced by electrical intracochlear stimulation. *Frontiers in Cell Neuroscience*, 12, 43. <https://doi.org/10.3389/fncel.2018.00043>.
- Roy, K., Murtie, J. C., El-Khodori, B. F., Edgar, N., Sardi, S. P., Hooks, B. M., Benoit-Marand, M., Chen, C., Moore, H., O'Donnell, P., Brunner, D., & Corfas, G. (2007). Loss of erbB signaling in oligodendrocytes alters myelin and dopaminergic function, a potential mechanism for neuropsychiatric disorders. *Proceedings of the National Academy of Sciences of the United States of America*, 104(19), 8131–8136.
- Rutherford, M. A., Chapochnikov, N. M., & Moser, T. (2012). Spike encoding of neurotransmitter release timing by spiral ganglion neurons of the cochlea. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 32, 4773–4789.
- Saliu, A., Adise, S., Xian, S., Kudelska, K., & Rodríguez-Contreras, A. (2014). Natural and lesion-induced decrease in cell proliferation in the medial nucleus of the trapezoid body during hearing development. *Journal of Comparative Neurology*, 522(5), 971–985. <https://doi.org/10.1002/cne.23473>
- Salzer, J. L. (2015). Schwann cell myelination. *Cold Spring Harbor Perspectives in Biology*, 7(8), a020529.
- Scapin, C., Ferri, C., Pettinato, E., Zambroni, D., Bianchi, F., Carro, U. D., Belin, S., Caruso, D., Mitro, N., Pellegatta, M., Taveggia, C., Schwab, M. H., Nave, K.-A., Feltri, M. L., Wrabetz, L., & D'Antonio, M. (2018). Enhanced axonal Neuregulin-1 type-III signaling ameliorates neurophysiology and hypomyelination in a Charcot-Marie-Tooth type 1B mouse model. *Human Molecular Genetics*, 28(6), 992–1006. <https://doi.org/10.1093/hmg/ddy411>.
- Schiff, J. A., Cracco, R. Q., & Cracco, J. B. (1985). Brainstem auditory evoked potentials in Guillain-Barré syndrome. *Neurology*, 35, 771–773.
- Schimmang, T., Minichiello, L., Vazquez, E., Jose, I. S., Giraldez, F., Klein, R., & Represa, J. (1995). Developing inner ear sensory neurons require TrkB and TrkC receptors for innervation of their peripheral targets. *Development*, 121(10), 3381–3391.
- Schiweck, J., Eickholt, B. J., & Murk, K. (2018). Important shapeshifter: Mechanisms allowing astrocytes to respond to the changing nervous system during development, injury and disease. *Frontiers in Cellular Neuroscience*, 12, 261. <https://doi.org/10.3389/fncel.2018.00261>.
- Schmitt, H. A., Pich, A., Schröder, A., Scheper, V., Lilli, G., Reuter, G., & Lenarz, T. (2017). Proteome analysis of human perilymph using an intraoperative sampling method. *Journal of Proteome Research*, 16(5), 1911–1923. <https://doi.org/10.1021/acs.jproteome.6b00986>
- Schütz, M., Scimemi, P., Majumder, P., Siaty, R. D. D., Crispino, G., Rodriguez, L., Bortolozzi, M., Santarelli, R., Seydel, A., Sonntag, S., Ingham, N., Steel, K. P., Willecke, K., & Mammano, F. (2010). The human deafness-associated connexin 30 T5M mutation causes mild hearing loss and reduces biochemical coupling among cochlear non-sensory cells in knock-in mice. *Human Molecular Genetics*, 19(24), 4759–4773. <https://doi.org/10.1093/hmg/ddq402>
- Seal, R. P., Akil, O., Yi, E., Weber, C. M., Grant, L., Yoo, J., Clause, A., Kandler, K., Noebels, J. L., Glowatzki, E., Lustig, L. R., & Edwards, R. H. (2008). Sensorineural deafness and seizures in mice lacking vesicular glutamate transporter 3. *Neuron*, 57(2), 263–275.
- Seidl, A. H., & Rubel, E. W. (2016). Systematic and differential myelination of axon collaterals in the mammalian auditory brainstem. *Glia*, 64(4), 487–494. <https://doi.org/10.1002/glia.22941>
- Sekiya, T., Holley, M. C., Hashido, K., Ono, K., Shimomura, K., Horie, R. T., Hamaguchi, K., Yoshida, A., Sakamoto, T., & Ito, J. (2015). Cells transplanted onto the surface of the glial scar reveal hidden potential for functional neural regeneration. *Proceedings of the National Academy of Sciences of the United States of America*, 112(26), E3431–E3440.
- Shore, S. E., Roberts, L. E., & Langguth, B. (2016). Maladaptive plasticity in tinnitus—triggers, mechanisms and treatment. *Nature Reviews Neurology*, 12(3), 150–160.
- Shrestha, B. R., Chia, C., Wu, L., Kujawa, S. G., Liberman, M. C., & Goodrich, L. V. (2018). Sensory neuron diversity in the inner ear is shaped by activity. *Cell*, 174(5), 1229–1246. [e1217](https://doi.org/10.1016/j.cell.2018.04.017).
- Silva, V. A., Gomide, V. C., & Chadi, G. (2005). Fibroblast growth factor-2 immunoreactivity is present in the central and peripheral auditory pathways of adult rats. *Journal of Morphology*, 265(2), 141–151. <https://doi.org/10.1002/jmor.10345>
- Sinclair, J. L., Fischl, M. J., Alexandrova, O., Heß, M., Grothe, B., Leibold, C., & Kopp-Scheinflug, C. (2017). Sound-evoked activity influences myelination of brainstem axons in the trapezoid body. *Journal of Neuroscience*, 37(34), 8239–8255. <https://doi.org/10.1523/JNEUROSCI.3728-16.2017>
- Stange-Marten, A., Nabel, A. L., Sinclair, J. L., Fischl, M., Alexandrova, O., Wohlfrom, H., Kopp-Scheinflug, C., Pecka, M., & Grothe, B. (2017). Input timing for spatial processing is precisely tuned via constant synaptic delays and myelination patterns in the auditory brainstem. *Proceedings of the National Academy of Sciences*, 114(24), E4851–E4858.
- Stankovic, K., Rio, C., Xia, A., Sugawara, M., Adams, J. C., Liberman, M. C., & Corfas, G. (2004). Survival of adult spiral ganglion neurons requires erbB receptor signaling in the inner ear. *Journal of Neuroscience*, 24, 8651–8661.
- Starr, A. (2003). Pathology and physiology of auditory neuropathy with a novel mutation in the MPZ gene (Tyr145->Ser). *Brain*, 126, 1604–1619.

- Stathopoulos, P., Alexopoulos, H., & Dalakas, M. C. (2015). Autoimmune antigenic targets at the node of Ranvier in demyelinating disorders. *Nature Reviews Neurology*, *11*(3), 143–156. <https://doi.org/10.1038/nrneurol.2014.260>
- Steele, C. J., Bailey, J. A., Zatorre, R. J., & Penhune, V. B. (2013). Early musical training and white-matter plasticity in the corpus callosum: Evidence for a sensitive period. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *33*(3), 1282–1290. <https://doi.org/10.1523/JNEUROSCI.3578-12.2013>
- Stolt, C. C., & Michael, W. (2016). Schwann cells and their transcriptional network: Evolution of key regulators of peripheral myelination. *Brain Research*, *1641*(Pt A), 101–110.
- Stotler, W. A. (1953). An experimental study of the cells and connections of the superior olivary complex of the cat. *The Journal of Comparative Neurology*, *98*(3), 401–431. <https://doi.org/10.1002/cne.900980303>
- Su, P., Kuan, C.-C., Kaga, K., Sano, M., & Mima, K. (2008). Myelination progression in language-correlated regions in brain of normal children determined by quantitative MRI assessment. *International Journal of Pediatric Otorhinolaryngology*, *72*(12), 1751–1763. <https://doi.org/10.1016/j.ijporl.2008.05.017>
- Sugawara, M., Corfas, G., & Liberman, M. C. (2005). Influence of supporting cells on neuronal degeneration after hair cell loss. *Journal of the Association for Research in Otolaryngology*, *6*(2), 136–147. <https://doi.org/10.1007/s10162-004-5050-1>
- Sugawara, M., Murtie, J. C., Stankovic, K. M., Liberman, M. C., & Corfas, G. (2007). Dynamic patterns of neurotrophin 3 expression in the postnatal mouse inner ear. *Journal of Comparative Neurology*, *501*(1), 30–37. <https://doi.org/10.1002/cne.21227>
- Sun, S., Babola, T., Pregernig, G., So, K. S., Nguyen, M., Su, S.-S.-M., Palermo, A. T., Bergles, D. E., Burns, J. C., & Müller, U. (2018). Hair cell mechanotransduction regulates spontaneous activity and spiral ganglion subtype specification in the auditory system. *Cell*, *174*(5), 1247–1263. [e1215](https://doi.org/10.1016/j.cell.2018.05.017).
- Susuki, K., Chang, K.-J., Zollinger, D. R., Liu, Y., Ogawa, Y., Eshed-Eisenbach, Y., Dours-Zimmermann, M. T., Osés-Prieto, J. A., Burlingame, A. L., Seidenbecher, C. I., Zimmermann, D. R., Oohashi, T., Peles, E., & Rasband, M. N. (2013). Three mechanisms assemble central nervous system nodes of Ranvier. *Neuron*, *78*(3), 469–482. <https://doi.org/10.1016/j.neuron.2013.03.005>
- Suzuka, Y., & Schuknecht, H. F. (1988). Retrograde cochlear neuronal degeneration in human subjects. *Acta Oto-Laryngologica Supplementum*, *450*, 1–20.
- Suzuki, J., Corfas, G., & Liberman, M. C. (2016). Round-window delivery of neurotrophin 3 regenerates cochlear synapses after acoustic overexposure. *Scientific Reports*, *6*, 24907.
- Tagoe, T., Barker, M., Jones, A., Allcock, N., & Hamann, M. (2014). Auditory nerve perinodal dysmyelination in noise-induced hearing loss. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *34*, 2684–2688.
- Takazawa, T., Ikeda, K., Murata, K., Kawase, Y., Hirayama, T., Ohtsu, M., Harada, H., Totani, T., Sugiyama, K., Kawabe, K., Kano, O., & Iwasaki, Y. (2012). Sudden deafness and facial diplegia in Guillain-Barré Syndrome: Radiological depiction of facial and acoustic nerve lesions. *Internal Medicine*, *51*, 2433–2437.
- Tang, W., Zhang, Y., Chang, Q., Ahmad, S., Dahlke, I., Yi, H., Chen, P., Paul, D. L., & Lin, X. (2006). Connexin29 is highly expressed in cochlear Schwann cells, and it is required for the normal development and function of the auditory nerve of mice. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *26*, 1991–1999.
- Taveggia, C., Thaker, P., Petrylak, A., Caporaso, G. L., Toews, A., Falls, D. L., Einheber, S., & Salzer, J. L. (2007). Type III neuregulin-1 promotes oligodendrocyte myelination. *Glia*, *56*(3), 284–293. <https://doi.org/10.1002/glia.20612>
- Tong, L., Strong, M. K., Kaur, T., Juiz, J. M., Oesterle, E. C., Hume, C., Warchol, M. E., Palmiter, R. D., & Rubel, E. W. (2015). Selective deletion of cochlear hair cells causes rapid age-dependent changes in spiral ganglion and cochlear nucleus neurons. *Journal of Neuroscience*, *35*(20), 7878–7891. <https://doi.org/10.1523/JNEUROSCI.2179-14.2015>
- Travers, B. G., Adluru, N., Ennis, C., Tromp, D. P. M., Destiche, D., Doran, S., Bigler, E. D., Lange, N., Lainhart, J. E., & Alexander, A. L. (2012). Diffusion tensor imaging in autism spectrum disorder: A review. *Autism Research: Official Journal of the International Society for Autism Research*, *5*(5), 289–313. <https://doi.org/10.1002/aur.1243>
- Tritsch, N. X., & Bergles, D. E. (2010). Developmental regulation of spontaneous activity in the mammalian cochlea. *Journal of Neuroscience*, *30*(4), 1539–1550. <https://doi.org/10.1523/JNEUROSCI.3875-09.2010>
- Tritsch, N. X., Yi, E., Gale, J. E., Glowatzki, E., & Bergles, D. E. (2007). The origin of spontaneous activity in the developing auditory system. *Nature*, *450*(7166), 50–55. <https://doi.org/10.1038/nature06233>
- Tritsch, N. X., Zhang, Y.-X., Ellis-Davies, G., & Bergles, D. E. (2010). ATP-induced morphological changes in supporting cells of the developing cochlea. *Purinergic Signalling*, *6*(2), 155–166. <https://doi.org/10.1007/s11302-010-9189-4>
- Trune, D. R. (1982). Influence of neonatal cochlear removal on the development of mouse cochlear nucleus: I. Number, size, and density of its neurons. *Journal of Comparative Neurology*, *209*(4), 409–424. <https://doi.org/10.1002/cne.902090410>
- Tylstedt, S., Kinnefors, A., & Rask-Andersen, H. (1997). Neural interaction in the human spiral ganglion: A TEM study. *Acta Oto-Laryngologica*, *117*, 505–512.
- Tylstedt, S., & Rask-Andersen, H. (2001). A 3-D model of membrane specializations between human auditory spiral ganglion cells. *Journal of Neurocytology*, *30*(6), 465–473.
- Ueda, N., & Kuroiwa, Y. (2008). Sensorineural deafness in Guillain-Barré syndrome. *Brain and Nerve = Shinkei Kenkyu no Shinpo*, *60*, 1181–1186.
- Usami, S.-I., Osen, K. K., Zhang, N., & Ottersen, O. P. (1992). Distribution of glutamate-like and glutamine-like immunoreactivities in the rat organ of Corti: A light microscopic and semiquantitative electron microscopic analysis with a note on the localization of aspartate. *Experimental Brain Research*, *91*(1), 1–11. <https://doi.org/10.1007/BF00230008>
- Uwechue, N. M., Marx, M.-C., Chevy, Q., & Billups, B. (2012). Activation of glutamate transport evokes rapid glutamine release from perisynaptic astrocytes. *The Journal of Physiology*, *590*(10), 2317–2331. <https://doi.org/10.1113/jphysiol.2011.226605>
- van Ruijven, M. W. M., de Groot, J. C. M. J., & Smoorenburg, G. F. (2004). Time sequence of degeneration pattern in the guinea pig cochlea during cisplatin administration. A quantitative histological study. *Hearing Research*, *197*, 44–54.
- Venkatesh, H. S., Johung, T. B., Caretti, V., Noll, A., Tang, Y., Nagaraja, S., Gibson, E. M., Mount, C. W., Polepalli, J., Mitra, S. S., Woo, P. J., Malenka, R. C., Vogel, H., Bredel, M., Mallick, P., & Monje, M. (2015). Neuronal activity promotes glioma growth through neurotrophin-3 secretion. *Cell*, *161*(4), 803–816.

- Verhamme, C., King, R. H. M., ten Asbroek, A. L. M. A., Muddle, J. R., Nourallah, M., Wolterman, R., Baas, F., & van Schaik, I. N. (2011). Myelin and axon pathology in a long-term study of PMP22-overexpressing mice. *Journal of Neuropathology & Experimental Neurology*, *70*(5), 386–398.
- Verkhratsky, A., Nedergaard, M., & Hertz, L. (2015). Why are astrocytes important? *Neurochemical Research*, *40*(2), 389–401. <https://doi.org/10.1007/s11064-014-1403-2>
- Vogler, D. P., Robertson, D., & Mulders, W. H. A. M. (2011). Hyperactivity in the Ventral cochlear nucleus after cochlear trauma. *The Journal of Neuroscience*, *31*(18), 6639–6645. <https://doi.org/10.1523/JNEUROSCI.6538-10.2011>
- Walsh, E., & Romand, R. (1992). Functional development of the cochlea and the cochlear nerve. In *Development of auditory and vestibular systems 2* (pp. 161–210). Elsevier.
- Wan, G., & Corfas, G. (2017). Transient auditory nerve demyelination as a new mechanism for hidden hearing loss. *Nature Communications*, *8*, 14487.
- Wan, G., Corfas, G., & Stone, J. S. (2013). Inner ear supporting cells: Rethinking the silent majority. *Seminars in Cell & Developmental Biology*, *24*(5), 448–459.
- Wan, G., Gómez-Casati, M. E., Gigliello, A. R., Liberman, M. C., & Corfas, G. (2014). Neurotrophin-3 regulates ribbon synapse density in the cochlea and induces synapse regeneration after acoustic trauma. *eLife*, *3*, e03564.
- Wang, H. C., Lin, C. C., Cheung, R., Zhang-Hooks, Y., Agarwal, A., Ellis-Davies, G., Rock, J., & Bergles, D. E. (2015). Spontaneous activity of cochlear hair cells triggered by fluid secretion mechanism in adjacent support cells. *Cell*, *163*(6), 1348–1359. <https://doi.org/10.1016/j.cell.2015.10.070>
- Wang, J., Zhang, B., Jiang, H., Zhang, L., Liu, D., Xiao, X., Ma, H., Luo, X., Bojrab, D., & Hu, Z. (2013). Myelination of the postnatal mouse cochlear nerve at the peripheral-central nervous system transitional zone. *Frontiers in Pediatrics*, *1*, 43. <https://doi.org/10.3389/fped.2013.00043>.
- Wang, S. J., Furusho, M., D'Sa, C., Kuwada, S., Conti, L., Morest, D. K., & Bansal, R. (2009). Inactivation of fibroblast growth factor receptor signaling in myelinating glial cells results in significant loss of adult spiral ganglion neurons accompanied by age-related hearing impairment. *Journal of Neuroscience Research*, *87*(15), 3428–3437.
- Wang, W., Zhang, L. S., Zinsmaier, A. K., Patterson, G., Leptich, E. J., Shoemaker, S. L., Yatskievych, T. A., Gibboni, R., Pace, E., Luo, H., Zhang, J., Yang, S., & Bao, S. (2019). Neuroinflammation mediates noise-induced synaptic imbalance and tinnitus in rodent models. *PLOS Biology*, *17*(6), e3000307.
- Warr, W. B. (1966). Fiber degeneration following lesions in the anterior ventral cochlear nucleus of the cat. *Experimental Neurology*, *14*(4), 453–474. [https://doi.org/10.1016/0014-4886\(66\)90130-0](https://doi.org/10.1016/0014-4886(66)90130-0)
- Wiechers, B., Gestwa, G., Mack, A., Carroll, P., Zenner, H.-P., & Knipper, M. (1999). A changing pattern of brain-derived neurotrophic factor expression correlates with the rearrangement of fibers during cochlear development of rats and mice. *Journal of Neuroscience*, *19*(8), 3033–3042. <https://doi.org/10.1523/JNEUROSCI.19-08-03033.1999>
- Wingard, J. C., & Zhao, H.-B. (2015). Cellular and deafness mechanisms underlying connexin mutation-induced hearing loss—A common hereditary deafness. *Frontiers in Cellular Neuroscience*, *9*, 202. <https://doi.org/10.3389/fncel.2015.00202>.
- Wise, A. K., Pujol, R., Landry, T. G., Fallon, J. B., & Shepherd, R. K. (2017). Structural and ultrastructural changes to Type I spiral ganglion neurons and schwann cells in the deafened guinea pig cochlea. *Journal of the Association for Research in Otolaryngology*, *18*, 751–769.
- Wong, A. W., Xiao, J., Kemper, D., Kilpatrick, T. J., & Murray, S. S. (2013). Oligodendroglial expression of TrkB independently regulates myelination and progenitor cell proliferation. *The Journal of Neuroscience*, *33*(11), 4947–4957. <https://doi.org/10.1523/JNEUROSCI.3990-12.2013>
- Xiao, J., Wong, A. W., Willingham, M. M., van den Buuse, M., Kilpatrick, T. J., & Murray, S. S. (2010). Brain-derived neurotrophic factor promotes central nervous system myelination via a direct effect upon oligodendrocytes. *Neuro-Signals*, *18*(3), 186–202. <https://doi.org/10.1159/000323170>
- Xing, Y., Samuvel, D. J., Stevens, S. M., Dubno, J. R., Schulte, B. A., & Lang, H. (2012). Age-related changes of myelin basic protein in mouse and human auditory nerve. *PLoS One*, *7*, e34500.
- Xu, S.-A., Shepherd, R. K., Chen, Y., & Clark, G. M. (1993). Profound hearing loss in the cat following the single co-administration of kanamycin and ethacrynic acid. *Hearing Research*, *70*(2), 205–215. [https://doi.org/10.1016/0378-5955\(93\)90159-X](https://doi.org/10.1016/0378-5955(93)90159-X)
- Yin, T. C., & Chan, J. C. (1990). Interaural time sensitivity in medial superior olive of cat. *Journal of Neurophysiology*, *64*(2), 465–488. <https://doi.org/10.1152/jn.1990.64.2.465>
- Zdebik, A. A., Wangemann, P., & Jentsch, T. J. (2009). Potassium ion movement in the inner ear: Insights from genetic disease and mouse models. *Physiology*, *24*(5), 307–316. <https://doi.org/10.1152/physiol.00018.2009>
- Zenisek, R., Thaler, N. S., Sutton, G. P., Ringdahl, E. N., Snyder, J. S., & Allen, D. N. (2015). Auditory processing deficits in bipolar disorder with and without a history of psychotic features. *Bipolar Disorders*, *17*(7), 769–780. <https://doi.org/10.1111/bdi.12333>
- Zhou, R., Abbas, P. J., & Assouline, J. G. (1995). Electrically evoked auditory brainstem response in peripherally myelin-deficient mice. *Hearing Research*, *88*, 98–106.
- Zhou, R., Assouline, J. G., Abbas, P. J., Messing, A., & Gantz, B. J. (1995). Anatomical and physiological measures of auditory system in mice with peripheral myelin deficiency. *Hearing Research*, *88*, 87–97.
- Zhou, Z., Liu, Q., & Davis, R. L. (2005). Complex regulation of spiral ganglion neuron firing patterns by neurotrophin-3. *Journal of Neuroscience*, *25*(33), 7558–7566. <https://doi.org/10.1523/JNEUROSCI.1735-05.2005>
- Zilberstein, Y., Liberman, M. C., & Corfas, G. (2012). Inner hair cells are not required for survival of spiral ganglion neurons in the adult cochlea. *Journal of Neuroscience*, *32*(2), 405–410. <https://doi.org/10.1523/JNEUROSCI.4678-11.2012>
- Ziskin, J. L., Nishiyama, A., Rubio, M., Fukaya, M., & Bergles, D. E. (2007). Vesicular release of glutamate from unmyelinated axons in white matter. *Nature Neuroscience*, *10*(3), 321–330. <https://doi.org/10.1038/nn1854>

How to cite this article: Kohrman D, Borges BC, Cassinotti L, Ji L, Corfas G. Axon–glia interactions in the ascending auditory system. *Develop Neurobiol*. 2021;81:546–567. <https://doi.org/10.1002/dneu.22813>