


# Race, Ethnicity, and Goal-Concordance of End-of-Life Palliative Care in Pediatric Oncology

Puja J. Umaretiya, MD <sup>1,2,3,4</sup>; Anran Li, BA<sup>1</sup>; Alana McGovern, BS<sup>1,3</sup>; Clement Ma, PhD<sup>1,3</sup>; Joanne Wolfe, MD, MPH<sup>1,2,3,4</sup>; and Kira Bona, MD, MPH<sup>1,2,3</sup>

**BACKGROUND:** Racial and ethnic minority children with cancer disproportionately receive intensive care at the end of life (EOL). It is not known whether these differences are goal-concordant or disparities. The authors sought to explore patterns of pediatric palliative care (PPC) and health care utilization in pediatric oncology patients receiving subspecialty palliative care at the end-of-life (last 6 months) and to examine goal-concordance of location of death in a subset of these patients. **METHODS:** This was a retrospective cohort study of pediatric oncology patients receiving subspecialty palliative care at a single large tertiary care center who died between January 2013 and March 2017. **RESULTS:** A total of 115 patients including 71 White, non-Hispanic patients and 44 non-White patients (including 12 Black patients and 21 Hispanic patients) were included in the analytic cohort. There were no significant differences in oncologic diagnosis, cause of death, or health care utilization in the last 6 months of life. White and non-White patients had similar PPC utilization including time from initial consult to death and median number of PPC encounters. Non-White patients were significantly more likely to die in the hospital compared to White patients (68% vs 46%,  $P = .03$ ). Analysis of a subcohort with documented preferences ( $n = 45$ ) revealed that 91% of White patients and 93% of non-White patients died in their preferred location of death. **CONCLUSIONS:** Although non-White children with cancer were more likely to die in the hospital, this difference was goal-concordant in our cohort. Subspecialty PPC access may contribute to the achievement of goal-concordant EOL care. *Cancer* 2021;127:3893-3900. © 2021 American Cancer Society.

**KEYWORDS:** disparities, location of death, pediatric oncology, pediatric palliative care, race.

## INTRODUCTION

Subspecialty pediatric palliative care (PPC) aims to improve quality of life and reduce symptom burden in seriously ill children through the provision of goal-concordant care. Children who are racial and ethnic minorities disproportionately receive intensive end-of-life (EOL) care, including hospital or intensive care unit (ICU) level care, active resuscitation, and in-hospital death.<sup>1-6</sup> It is not known whether these observed differences in pediatric patients are goal-concordant or represent disparities in care among racial and ethnic minority patients.

Race-based disparities in quality palliative care at the EOL are well-documented in adult patients. Specifically, adult Black patients are more likely to die in the hospital and receive intensive and life-prolonging care<sup>7-10</sup> that may be discordant with their expressed preferences.<sup>11</sup>

Current quality metrics in palliative care are based on data that adults of all race/ethnicities have a strong preference for care focused on comfort at the end-of-life and death at home,<sup>12,13</sup> and intensive EOL care is associated with poor quality of life and caregiver outcomes.<sup>14,15</sup> Consequently, in-hospital death, peri-death ICU admissions, and short or no hospice enrollment are deemed poor quality care at the end-of-life.<sup>16,17</sup> Given the dearth of preference data for pediatric patients, it is not clear that these quality metrics should be applied to pediatric patients. Ensuring the provision of high quality PPC at end-of-life requires understanding whether racial and ethnic differences in EOL care for children with cancer reflect family preferences or outcome disparities.

One critical and intervenable end-of-life outcome is ensuring that pediatric location of death is consistent with expressed patient and family preferences. Despite identified differences in pediatric location of death by race and ethnicity, data on family preferences among non-White pediatric patients are lacking.<sup>18,19</sup> To address this critical gap, we aimed to explore racial and ethnic differences in patterns of PPC and EOL care utilization over the last 6 months of life in a cohort

**Corresponding Author:** Puja J. Umaretiya, MD, Department of Pediatric Oncology, Dana-Farber Cancer Institute, 450 Brookline Ave, Boston, MA 02215 (puja\_umaretiya@dfci.harvard.edu).

Anran Li's current address: University of Michigan Medical School, Ann Arbor, Michigan

<sup>1</sup>Department of Pediatric Oncology, Dana-Farber Cancer Institute, Boston, Massachusetts; <sup>2</sup>Division of Population Sciences, Dana-Farber Cancer Institute, Boston, Massachusetts; <sup>3</sup>Division of Pediatric Hematology/Oncology, Boston Children's Hospital, Boston, Massachusetts; <sup>4</sup>Department of Psychosocial Oncology and Palliative Care, Dana-Farber Cancer Institute, Boston, Massachusetts

We acknowledge the patients and families who contributed to this study.

**DOI:** 10.1002/cncr.33768, **Received:** December 29, 2020; **Revised:** May 4, 2021; **Accepted:** June 8, 2021, **Published online** July 13, 2021 in Wiley Online Library (wileyonlinelibrary.com)

of pediatric oncology patients receiving palliative care at a large referral center. A key secondary aim was to explore concordance between location of death and expressed family preferences.

## MATERIALS AND METHODS

This was a retrospective cohort study at a single large tertiary pediatric care center. This study was approved by the Dana-Farber Cancer Institute institutional review board.

### **Study Cohort**

The study cohort was derived from the Dana-Farber/Boston Children Hospital (DF/BCH) Pediatric Palliative Care Database that systematically tracks all patients receiving palliative care at DF/BCH. The analytic cohort included all children with available race/ethnicity data who had a primary cancer diagnosis, received both primary oncology and PPC care at DF/BCH during the last 6 months of life, and died between January 1, 2013, and March 15, 2017. At DF/BCH, 80% of pediatric oncology patients who die receive PPC at least 1 month before death. To minimize heterogeneity, foreign-national patients receiving care via embassy pay were excluded ( $n = 7$ ). The final analytic cohort included 115 patients.

### **Data Collection**

Demographic, disease, and care utilization data were extracted from electronic medical records into a REDCap database. Data elements were obtained via electronic data capture and manual abstraction from free text clinical note content. Primary data abstraction was completed by 1 study investigator (P.J.U.). Unclear data element coding was discussed with another investigator (K.B.), and final consensus was reached by both investigators.

### **Exposures**

Racial and/or ethnic minority status was the primary exposure of interest due to previously described racial and ethnic disparities in palliative care. Race/ethnicity was dichotomized as White versus non-White. Exploratory analyses examining socioeconomic status were performed post hoc. Socioeconomic status was characterized at the neighborhood level by linkage of a child's residential zip code at death with US Census data from the 2012-2016 American Community Survey.<sup>20</sup> Zip codes were a priori defined as high-poverty ( $\geq 20\%$  of the population lived below 100% federal poverty level [FPL]) versus low-poverty if ( $< 20\%$  of the population lived below 100% FPL) based on US Census definition.<sup>21</sup>

## **Covariates**

### **Demographics**

Demographic data include age at death, sex (male or female), primary language (English, Spanish, or other), religion (identified religious affiliation or none), and insurance at death (public, private, or both).

### **Disease and treatment variables**

Disease variables included date of first cancer diagnosis, disease type (hematologic malignancy, extracranial solid tumor, brain tumor), number of relapses, number of stem cell transplants, phase 1 or phase 2 clinical trial enrollment (yes or no), and cause of death (progressive disease, treatment-related or other).

### **Care utilization variables**

Hospital utilization variables in the last 6 months of life included number of days in hospital, number of unplanned hospitalizations (defined as admissions not for chemotherapy or planned surgeries), and number of ICU admissions. PPC utilization variables in the last 6 months of life included date, location (inpatient [non-ICU], ICU, outpatient, other), and indication (advance care planning, goals of care, decision-making, psychosocial support, symptom management, logistics/coordination of care, transition to home, parental/sibling bereavement, quality of life, and other) for initial PPC consult, content of initial PPC consult (advance care planning, goals of care, resuscitation status, and symptom burden), total PPC encounters, hospice enrollment (yes or no) and date if applicable, and documented Medical Orders for Life-Sustaining Treatment form (yes or no).

### **EOL care intensity variables**

EOL variables included primary decision-maker (assumed to be parent if not documented and patient  $< 18$  years), date of death, location of death (hospital or home/hospice), mode of death (active resuscitation, discontinuation of life-sustaining therapies, or comfort-focused care), date and content of first documented resuscitation status order, resuscitation status at death (full code or limited resuscitation), medical interventions in last 7 days of life (cardiopulmonary resuscitation [CPR], intubation, intravenous [iv] inotropes).

### **Outcomes**

The primary outcome of interest was EOL patterns of care (including timing of PPC consult, medical interventions in the last 7 days of life, location of death, and resuscitation status at death) stratified by race/ethnicity.

**TABLE 1.** Demographic and Disease Characteristics by Race/Ethnicity

	Total (N = 115), No. (%)	White (N = 71), No. (%)	Non-White (N = 44), No. (%)	P
Age at death, median, (IQR), y	13.5 (6-17.2)	13.6 (6.0-17.4)	13.1 (6.6-17.0)	.61
Female sex	56 (49)	39 (55)	17 (39)	.12
Primary language				<.0001
English	97 (84)	70 (99)	27 (61)	
Spanish	10 (9)	0 (0)	10 (23)	
Other	8 (7)	1 (1)	7 (16)	
Religious affiliation (N = 96)		(N = 58)	(N = 38)	.80
Yes	76 (79)	45 (78)	31 (82)	
No	20 (21)	13 (22)	7 (18)	
Insurance status at death				.005
Public	46 (40)	20 (28)	26 (59)	
Private	31 (27)	23 (32)	8 (18)	
Mixed	38 (33)	28 (39)	10 (23)	
Diagnosis				.78
Hematologic malignancy	25 (22)	14 (20)	11 (25)	
Solid tumor	51 (44)	33 (46)	18 (41)	
Brain tumor	39 (34)	24 (34)	15 (34)	
No. of relapses				.88
0	58 (50)	35 (49)	23 (52)	
1	27 (23)	18 (25)	9 (20)	
2+	30 (26)	18 (25)	12 (27)	
Enrolled in phase 1 or phase 2 clinical trial	28 (24)	17 (24)	11 (25)	>.99
Living in high poverty neighborhood	21 (18)	2 (3)	19 (43)	<.0001
Cause of death				.28
Treatment-related	9	7(10)	2 (5)	
Progressive disease	105	64 (90)	41 (93)	
Other	1	0 (0)	1 (2)	

Abbreviation: IQR, interquartile range.

We additionally explored goal-concordant location of death stratified by race/ethnicity for a subset cohort with available data. Children for whom documented free text-preferred location of death most proximal to date of death matched actual location of death were considered to have experienced goal-concordant location of death. When disagreement was noted in preference for location of death between child and parents (n = 1), child's preference of death was recorded as preferred location of death.

### Statistical Analysis

Patient demographics, disease, and care utilization were summarized for the overall cohort and by race/ethnicity using descriptive statistics including medians, interquartile ranges for continuous measures, frequencies, and proportions for categorical measures. Baseline characteristics were compared between race/ethnicity groups with the  $\chi^2$  and Wilcoxon tests for categorical and continuous variables, respectively. Because of the small sample size, we a priori chose to perform univariate tests of association only for the EOL patterns of care (including timing of PPC consult, medical interventions in the last 7 days of life, location of death, and resuscitation status at death) as the primary outcomes of interest using Wilcoxon rank-sum test for continuous outcomes and Fisher exact test

for categorical outcomes. All analyses were performed using SAS version 9.4 (SAS Institute Inc, Cary, North Carolina). Two-sided *P* values <.05 were considered statistically significant.

## RESULTS

### Study Population Characteristics

A total of 115 patients were included in analysis with a median age at death of 13.5 years (interquartile range [IQR] = 6-17.2 years) (Table 1). Over a third (38%) were non-White, including 12 (10%) Black and 21 (18%) Hispanic (2 patients identified as both). Twenty-one patients (18%) lived in a high-poverty neighborhood, and 46 patients (40%) had public insurance alone. Children living in high-poverty neighborhoods or with public insurance were disproportionately non-White. Disease characteristics did not differ significantly by race/ethnicity. Progression of disease was the primary cause of death in both White (90%) and non-White patients (93%).

### PPC Utilization

Children had a median of 12 total PPC encounters (IQR = 5-23) before death (Table 2). Initial PPC consults occurred a median of 3 months (IQR = 1.2-6.4 months) before death overall, and this did not differ between White

**TABLE 2.** Palliative Care Utilization by Race/Ethnicity

	Total (N = 115), No. (%)	White (N = 71), No. (%)	Non-White (N = 44), No. (%)
Indications for initial PPC consult		(N = 69)	
Decision support	53 (46)	28 (39)	25 (57)
Psychosocial support	52 (46)	34 (49)	18 (41)
Symptom management	64 (57)	38 (54)	26 (59)
Logistics	8 (7)	6 (8)	2 (5)
Parental/sibling bereavement	0 (0)	0 (0)	0 (0)
Goals of care discussed at initial encounter	102 (89)	64 (90)	38 (86)
Advanced care planning discussed at initial encounter	31 (27)	19 (27)	12 (27)
Resuscitation status at initial consult			
Full code	93 (81)	60 (85)	33 (75)
Limited resuscitation	22 (19)	11 (15)	11 (25)
Location of initial consult			
Inpatient, non-ICU	50 (43)	24 (34)	26 (59)
ICU	14 (12)	9 (13)	5 (11)
Outpatient	46 (40)	35 (49)	11 (25)
Other <sup>a</sup>	5 (4)	3 (4)	2 (6)
No. of total PPC encounters, median (IQR)	12 (5-23)	11 (5-23)	13 (7.5-32)

Abbreviations: ICU, intensive care unit; IQR, interquartile range; PPC, pediatric palliative care.

<sup>a</sup>Four initial palliative care consults occurred at home, and 1 occurred at a local rehabilitation center.

**TABLE 3.** Health Care Utilization by Race/Ethnicity

	Total, (N = 115), No. (%)	White (N = 71), No. (%)	Non-White (N = 44), No. (%)
Mode of death			
Active resuscitation	7 (6)	4 (6)	3 (7)
Discontinuation of LST	18 (16)	9 (13)	9 (20)
Comfort care	90 (78)	58 (82)	32 (73)
MOLST form present	59 (52)	38/70 (56)	21 (48)
Hospice involved	57 (50)	40/70 (56)	17 (40)
Days hospitalized in last 6 mo of life, median	22 (9-58)	19.0 (4-53)	36.5 (16.5-86.0)
No. of unplanned admissions, median	2 (1-3)	2 (1-3)	2.5 (1-4)
No. of ICU admissions in last 6 mo			
0	49 (43)	33 (46)	16 (36)
1+	66 (57)	38 (54)	28 (64)
Resuscitation status changed in last 7 d	42 (37)	24 (34)	18 (41)
Radiation received in last 7 d	2 (2)	0 (0)	2 (5)
Chemotherapy received in last 7 d	24 (21)	16 (23)	8 (18)
ICU admission in last 7 d	18 (16)	8 (11)	10 (24)
Length of final admission if death in hospital, median, (IQR), d	11 (5.5-31.5)	11 (6-30)	14 (3-32)

Abbreviations: ICU, intensive care unit; IQR, interquartile range; LST, life-sustaining treatment; MOLST, medical orders for life-sustaining treatment.

and non-White patients (2.6 vs 3.3 months,  $P = .87$ ). Initial PPC consults were more likely to occur in the outpatient setting for White patients (49% White vs 25% non-White). The content of initial PPC encounters did not vary by race/ethnicity and included goals of care discussion for a majority of patients (90% White vs 86% non-White) and advance care planning for a minority of patients (27% for both White and non-White patients). At the time of initial PPC consult, resuscitation status was Full Code for a majority of patients (84% White vs 73% non-White). Patterns of care utilization were also compared across patients living in high- and low-poverty neighborhoods and did not differ significantly from White vs non-White patient comparisons, likely reflective

of the fact that most patients living in high-poverty areas in our cohort were non-White.

#### **Care Utilization in the Last 6 Months of Life**

Patients were hospitalized for a median of 22 days (IQR = 9-58 days), including a median of 2 unplanned admissions in the last 6 months of life (Table 3). Nearly a quarter (24%) of the cohort enrolled on a phase 1 or 2 trial (Table 1). Overall health care utilization was comparable by race/ethnicity.

#### **Patterns of EOL Care**


Location of death differed significantly between White and non-White patients (Table 4). Non-White patients were significantly more likely to die in-hospital compared

**TABLE 4.** EOL Care by Race/Ethnicity

	Total (N = 115), No. (%)	White (N = 71), No. (%)	Non-White (N = 44), No. (%)	P
Location of death (N = 114) <sup>a</sup>				.03
Hospital	62 (54)	32/70 (46)	30 (68)	
Home/hospice	52 (46)	38/70 (54)	14 (32)	
Resuscitation status at death (N = 114) <sup>a</sup>				1.0
Full code	13 (11)	8 (11)	5 (11)	
Limited resuscitation	101 (89)	62 (89)	39 (89)	
Interventions received in the last 7 d	25 (22)	12 (17)	13 (30)	.11
CPR	7 (6)	4 (6)	3 (7)	
Intubation	23 (20)	12 (17)	11 (26)	
Intravenous inotropes	13 (11)	7 (10)	6 (14)	
Time from initial PPC encounter to death, median (IQR), mo	3 (1.2-6.4)	2.6 (1.2-8.1)	3.3 (1.1-6.2)	.87

Abbreviations: CPR, cardiopulmonary resuscitation; EOL, end of life; IQR, interquartile range; PPC, pediatric palliative care.

<sup>a</sup>One patient died in Florida on a Make-A-Wish trip and was excluded from analysis.



**Figure 1.** Proportion of White and non-White patients with goal-concordant location of death.

to White patients (68% vs 46%;  $P = .03$ ). Differences in high-intensity EOL care as measured by utilization of CPR, intubation, or iv inotropes in last 7 days did not reach significance comparing White patients to non-White patients (30% vs 17%;  $P = .11$ ). There was no difference in resuscitation status at time of death by race/ethnicity ( $P = 1.0$ ).

#### **Concordance of Location of Death with Patient/Family Preferences: Subcohort**

Forty-five patients (39%) in the analytic cohort had documented preference for location of death available for analysis, with similar proportions of non-White patients (14 of 44, 32%) and White patients (31 of 71, 44%) as in the overall cohort. The subcohort did not differ significantly from the overall cohort in terms of key sociodemographic covariates including language, insurance status, and proportion living in high-poverty neighborhoods. Similar to the overall cohort, non-White patients in this subcohort remained more likely to die in the hospital

than White patients (64% vs 29%). Location of death was concordant with expressed preference in a majority of both non-White and White patients (93% vs 91%, respectively) (Fig. 1). Among non-White patients, 9 of 9 in-hospital deaths (100%) were goal-concordant, whereas 4 of 5 home deaths (80%) were goal-concordant. Similarly, among White patients, 21 of 22 home deaths (95%) were goal-concordant and 8 of 9 in-hospital deaths (89%) were goal-concordant. In this subcohort, public insurance, high-poverty zip code, and non-English primary language were also associated with higher proportions of in-hospital death (58%, 70%, and 80%, respectively); all of these in-hospital deaths were goal-concordant.

#### **DISCUSSION**

Non-White pediatric oncology patients receiving subspecialty PPC were significantly more likely than White patients to die in-hospital (68% vs 46%;

$P = .03$ ), similar to previously published data.<sup>1-4,6</sup> This race/ethnicity-associated difference appears to be goal-concordant based on subcohort analysis of documented family preferences. Specifically, among those with documented preferences, 91% of White patients and 93% of non-White patients died in their preferred location of death. Although fewer non-White patients in our cohort preferred to die at home, all of those who expressed this preference successfully achieved goal-concordant location of death. We identified no differences between White and non-White patients with regard to PPC patterns of care including frequency of PPC encounters or content of initial PPC consultation, and this finding is consistent with prior studies.<sup>22,23</sup> Uniform PPC access in our cohort may have contributed to the fact that there were also no differences between non-White and White patients in intensity of EOL care or resuscitation status in the last 7 days of life. It may have also led to goal-concordant locations of death for the patients in our subcohort.

These data contribute substantially to the existing data regarding family and patient preferences for location of death in children that include largely White populations. Prior studies identified death at home as preferred among adolescents<sup>18</sup> and parents of children with cancer.<sup>19</sup> However, a majority of adolescents and parents were White. Our data suggest that pediatric race-associated differences in location of death may, in fact, be goal-concordant and thus reflective of different preferences, not inequitable care. The implications of this for PPC research and care delivery are significant. Research to date has relied on adult metrics to interpret pediatric data, including a cohort study of pediatric oncology patients across California that reported disparities in the intensity of EOL care among White and non-White patients.<sup>2</sup> Although non-White patients were more likely to die in the hospital in this cohort, other measures of intensity did not vary significantly between White and non-White patients. Interpreted in the context of our concordance data, these results may, in fact, reflect differential preferences rather than disparities. Importantly, these data suggest that the application of adult quality metrics to children with cancer may be inappropriate,<sup>24</sup> and recent efforts to define high quality pediatric EOL care highlight the importance of death in a preferred location, rather than in the hospital.<sup>25</sup>

There are several reasons that non-White pediatric patients and their families may prefer to die in the hospital. First, race is a social construct<sup>26</sup> and may serve as a proxy

for multiple other factors (culture, religion, language, and perceived trustworthiness of the health care system) that shape family preferences. These factors may differ between racial and ethnic groups (eg, Black vs Hispanic). For example, a recent study suggests that Spanish-speaking bereaved parents preferred death at home,<sup>27</sup> although preferences in hindsight during bereavement may differ from those before the death of a child. Notably, the parents in that study additionally expressed that home death was preferable only if there were adequate resources to support their child at home. Our sample size limits our ability to analyze differences between these groups. Second, observed differences may reflect the downstream effects of structural racism leading to ongoing economic disadvantage for some racial and ethnic groups. Non-White patients in our cohort were more likely to live in high-poverty neighborhoods. Families living in high-poverty neighborhoods are more likely to be low-income and experience material hardship—eg, difficulty putting food on the table, keeping the heat or electricity on, or unstable housing.<sup>29,30</sup> They may also have less access to hospice or home services. It is possible that observed racial/ethnic differences in preferred location of death reflect resource and access needs that parents may perceive as barriers to a quality death at home. Finally, although the timing and frequency of PPC was similar between White and non-White patients, we were unable to assess the quality of communication, as well as implicit and explicit biases, in these encounters. It has previously been demonstrated that providers believe that Black and Hispanic parents are less interested in prognostic communication, although a majority of Black and Hispanic families desire detailed prognostic information.<sup>28</sup> Differences in communication may contribute to the finding that although the frequency of EOL discussions between Black and White patients is similar, Black patients have less terminal illness awareness.<sup>11</sup> The next steps to providing equitable, goal-concordant care at the EOL will involve understanding the primary drivers of these preferences. Although our data suggest that public insurance, high-poverty neighborhood, and non-English primary language were also associated with a preference for hospital death, focused research efforts to explore the perspectives of non-White racial and ethnic groups are necessary to learn how these factors and their intersectionality influence parent preferences.

Our study had several limitations. Our cohort derives from a single large, tertiary care center, which may limit generalizability of these findings. The retrospective nature of this study limits a broader understanding of family and patient preferences that were abstracted from the medical record. Provider documentation may

not clearly reflect all discussions relevant to the full scope of patient and family preferences. Additionally, concordance data were available only for a subset of our analytic cohort and may represent a biased sample of families particularly skilled at advocating for their preferences at EOL. Compared to location of death for the entire cohort, a smaller percentage of White patients in the concordance subcohort died in the hospital (46% vs 29%) whereas a similar number of non-White patients died in the hospital (68% vs 64%). This said, all patients in this cohort had active PPC involvement that may mitigate this potential bias by offering patients the opportunity to discuss hopes and worries in the context of advanced illness.

Among a cohort of pediatric oncology patients receiving subspecialty PPC, race and ethnicity-associated differences in location of death were concordant with family preferences. As such, in-hospital death in our pediatric cohort appears to reflect quality—not poor—PPC care delivery. Further investigation exploring the factors that inform racial/ethnic differences in family goals for EOL care are clearly needed to ensure the provision of equitable and goal-concordant PPC.

## FUNDING SUPPORT

Kira Bona is supported by the National Cancer Institute at the National Institutes of Health (K07CA211847).

## CONFLICTS OF INTEREST DISCLOSURES

Clement Ma reports participating on the Data Safety Monitoring Boards of 2 trials. The remaining authors made no disclosures.

## AUTHOR CONTRIBUTIONS

**Puja J. Umaretiya:** Conceptualization, methodology, investigation, data curation, formal analysis, writing—original draft, and writing—review and editing. **Anran Li:** Formal analysis and writing—review and editing. **Alana McGovern:** Formal analysis and writing—review and editing. **Clement Ma:** Formal analysis and writing—review and editing. **Joanne Wolfe:** Conception, methodology, and writing—review and editing. **Kira Bona:** Conceptualization, methodology, investigation, data curation, formal analysis, supervision, and writing—review and editing.

## REFERENCES

- Cawkwell PB, Gardner SL, Weitzman M. Persistent racial and ethnic differences in location of death for children with cancer. *Pediatr Blood Cancer*. 2015;62:1403-1408.
- Johnston EE, Alvarez E, Saynina O, Sanders L, Bhatia S, Chamberlain LJ. Disparities in the intensity of end-of-life care for children with cancer. *Pediatrics*. 2017;140:e20170671.
- Shah A, Diggins N, Stiller C, Murphy D, Passmore J, Murphy MF. Place of death and hospital care for children who died of cancer in England, 1999-2006. *Eur J Cancer*. 2011;47:2175-2181.
- Kaye EC, DeMarsh S, Gushue CA, et al. Predictors of location of death for children with cancer enrolled on a palliative care service. *Oncologist*. 2018;23:1525-1532.
- Kaye EC, Gushue CA, DeMarsh S, et al. Impact of race and ethnicity on end-of-life experiences for children with cancer. *Am J Hosp Palliat Care*. 2019;36:767-774.
- Rajeshuni N, Johnston EE, Saynina O, Sanders LM, Chamberlain LJ. Disparities in location of death of adolescents and young adults with cancer: a longitudinal, population study in California. *Cancer*. 2017;123:4178-4184.
- Mitchell SL, Teno JM, Roy J, Kabumoto G, Mor V. Clinical and organizational factors associated with feeding tube use among nursing home residents with advanced cognitive impairment. *JAMA*. 2003;290:73-80.
- Goodlin SJ, Zhong Z, Lynn J, et al. Factors associated with use of cardiopulmonary resuscitation in seriously ill hospitalized adults. *JAMA*. 1999;282:2333-2339.
- Virnig BA, Kind S, McBean M, Fisher E. Geographic variation in hospice use prior to death. *J Am Geriatr Soc*. 2000;48:1117-1125.
- Hanchate A, Kronman AC, Young-Xu Y, Ash AS, Emanuel E. Racial and ethnic differences in end-of-life costs: why do minorities cost more than Whites? *Arch Intern Med*. 2009;169:493-501.
- Mack JW, Paulk ME, Viswanath K, Prigerson HG. Racial disparities in the outcomes of communication on medical care received near death. *Arch Intern Med*. 2010;170:1533-1540.
- Barnato AE, Anthony DL, Skinner J, Gallagher PM, Fisher ES. Racial and ethnic differences in preferences for end-of-life treatment. *J Gen Intern Med*. 2009;24:695-701.
- Wright AA, Keating NL, Ayanian JZ, et al. Family perspectives on aggressive cancer care near the end of life. *JAMA*. 2016;315:284-292.
- Wright AA, Zhang B, Ray A, et al. Associations between end-of-life discussions, patient mental health, medical care near death, and caregiver bereavement adjustment. *JAMA*. 2008;300:1665-1673.
- Wright AA, Keating NL, Balboni TA, Matulonis UA, Block SD, Prigerson HG. Place of death: correlations with quality of life of patients with cancer and predictors of bereaved caregivers' mental health. *J Clin Oncol*. 2010;28:4457-4464.
- Palliative and End-of-Life Care 2015-2016. National Quality Forum. Accessed August 12, 2020. [https://www.qualityforum.org/Publications/2016/12/Palliative\\_and\\_End-of-Life\\_Care\\_2015-2016.aspx](https://www.qualityforum.org/Publications/2016/12/Palliative_and_End-of-Life_Care_2015-2016.aspx)
- Earle CC, Park ER, Lai B, Weeks JC, Ayanian JZ, Block S. Identifying potential indicators of the quality of end-of-life cancer care from administrative data. *J Clin Oncol*. 2003;21:1133-1138.
- Jacobs S, Perez J, Cheng YI, Sill A, Wang J, Lyon ME. Adolescent end of life preferences and congruence with their parents' preferences: results of a survey of adolescents with cancer. *Pediatr Blood Cancer*. 2015;62:710-714.
- Kassam A, Skiadareis J, Alexander S, Wolfe J. Parent and clinician preferences for location of end-of-life care: home, hospital or freestanding hospice? *Pediatr Blood Cancer*. 2014;61:859-864.
- American Community Survey: data tables and tools. US Census Bureau. Accessed August 12, 2020. <https://www.census.gov/acs/www/data/data-tables-and-tools/>
- Krieger N, Chen JT, Waterman PD, Rehkopf DH, Subramanian SV. Race/ethnicity, gender, and monitoring socioeconomic gradients in health: a comparison of area-based socioeconomic measures—the public health disparities geocoding project. *Am J Public Health*. 2003;93:1655-1671.
- Kaye EC, Jerkins J, Gushue CA, et al. Predictors of late palliative care referral in children with cancer. *J Pain Symptom Manage*. 2018;55:1550-1556.
- Baker JN, Rai S, Liu W, et al. Race does not influence do-not-resuscitate status or the number or timing of end-of-life care discussions at a pediatric oncology referral center. *J Palliat Med*. 2009;12:71-76.
- Johnston EE, Rosenberg AR, Kamal AH. Pediatric-specific end-of-life care quality measures: an unmet need of a vulnerable population. *J Oncol Pract*. 2017;13:e874-e880.
- Johnston EE, Martinez I, Wolfe J, Asch SM. Quality measures for end-of-life care for children with cancer: a modified Delphi approach. *Cancer*. Published March 30, 2021. doi:10.1002/cncr.33546
- Williams DR, Sternthal M. Understanding racial-ethnic disparities in health: sociological contributions. *J Health Soc Behav*. 2010;51(suppl):S15-S27.

27. Johnston EE, Molina J, Martinez I, et al. Bereaved parents' views on end-of-life care for children with cancer: quality marker implications. *Cancer*. 2020;126:3352-3359.
28. Ilowite MF, Cronin AM, Kang TI, Mack JW. Disparities in prognosis communication among parents of children with cancer: the impact of race and ethnicity. *Cancer*. 2017;123:3995-4003.
29. Sandel M, Sheward R, Ettinger de Cuba S, et al. Unstable housing and caregiver and child health in renter families. *Pediatrics*. 2018;141:e20172199.
30. Frank DA, Casey PH, Black MM, et al. Cumulative hardship and wellness of low-income, young children: multisite surveillance study. *Pediatrics*. 2010;125:e1115-e1123.