

Quigg Kellen Haley (Orcid ID: 0000-0003-4365-2298)

**Title:** Transcutaneous CO2 Monitoring as Indication for Inpatient NIV Initiation in Patients with ALS

**Author List:** Kellen H. Quigg, MD<sup>1</sup>, Matthew W. Wilson, MD<sup>2</sup>, Philip J. Choi, MD<sup>2</sup>

1. Department of Neurology, University of Michigan, Ann Arbor, MI
2. Department of Internal Medicine, Division of Pulmonary and Critical Care Medicine, University of Michigan, Ann Arbor, MI

**Corresponding Author Information:**

Philip J. Choi, MD

Division of Pulmonary and Critical Care Medicine

3916 Taubman Center/SPC 5360

1500 E. Medical Center Dr

Ann Arbor, MI 48109

[Pchoi@med.umich.edu](mailto:Pchoi@med.umich.edu)

**Conflict of interest:**

Kellen Quigg: none

Matthew Wilson: none

Philip Choi: none

**Funding information:** none

**Prior abstract presentation:** American Thoracic Society 2020 Virtual Meeting; June 24, 2020

**Key Words:** amyotrophic lateral sclerosis, hypercarbia, non-invasive ventilation, respiratory failure, transcutaneous CO2

**Word Count Manuscript:** 1376

**Word Count Abstract:** 244

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/mus.27457](https://doi.org/10.1002/mus.27457)

This article is protected by copyright. All rights reserved.

## **Abstract**

**Introduction/Aims:** Amyotrophic lateral sclerosis (ALS) is characterized by profound muscle weakness, including diaphragmatic weakness resulting in hypercapnic respiratory failure. While non-invasive ventilation (NIV) is usually initiated in the home, patients presenting with hypercapnic respiratory failure may be at high risk of adverse outcomes with delays in treatment. We aim to describe the clinical utility of transcutaneous CO<sub>2</sub> (TCO<sub>2</sub>) to assess the need for inpatient initiation of NIV.

**Methods:** Eight patients from the University of Michigan Pranger ALS clinic were directly admitted to the hospital for urgent initiation of NIV between May 2020-May 2021. A retrospective review of electronic medical records, including pre-hospital pulmonary function assessments, hospitalization blood gases and NIV use metrics was performed.

**Results:** All eight patients had symptoms of respiratory insufficiency at time of admission, though not all patients had forced vital capacity (FVC) measurements that would identify need for NIV. All patients had measured TCO<sub>2</sub> > 45 mmHg. Seven of eight patients had worsening hypercapnia after admission, indicating advanced respiratory failure. All patients were titrated to tolerance of continuous nocturnal NIV while in the hospital, with an average length of stay of 6.5 days (range 3-8). All patients demonstrated compliance with NIV, >4hours, at post-hospital follow up.

**Discussion:** Many current ambulatory measurements underestimate, or incompletely evaluate, respiratory dysfunction, and arterial blood gases are not typically readily available. Outpatient TCO<sub>2</sub> measurements can serve as a useful screening tool to identify ALS patients who would benefit from inpatient initiation and titration of NIV.

# **Transcutaneous CO<sub>2</sub> Monitoring as Indication for Inpatient NIV Initiation in Patients with ALS**

## **Introduction**

Amyotrophic lateral sclerosis (ALS) is characterized by profound muscle weakness, including diaphragmatic weakness resulting in hypercapnic respiratory failure(1). Forced vital capacity (FVC) and maximum inspiratory pressure (MIP) are commonly used as surrogates for respiratory function, with an FVC <50% of predicted values typically indicative of significant respiratory muscle involvement(2). Additionally, these tests generally underestimate respiratory dysfunction, are not designed to approximate alveolar hypoventilation leading to hypercapnic respiratory failure and become unreliable with increasing neuromuscular incoordination and worsening bulbar weakness. The ALS Functional Rating Scale Revised (ALSF<sub>RS</sub>-R) contains questions addressing respiratory function(3), but alone may not accurately predict impending respiratory failure.

The mainstay of respiratory support in patients with ALS is non-invasive ventilation (NIV)(4). While most patients can safely initiate NIV at home, there are no clear or reliable criteria that would identify those patients at high risk of imminent respiratory collapse in need of inpatient admission. Transcutaneous CO<sub>2</sub> (TCO<sub>2</sub>) monitoring is a non-invasive, reliable means of measuring PaCO<sub>2</sub> in an ambulatory setting(5, 6). It is a well-validated proxy for PaCO<sub>2</sub>, typically underestimating measurements by 5-8mmHg(5). We aim to describe how TCO<sub>2</sub> monitoring can be a more useful surrogate for hypercapnia to identify patients with ALS who may benefit from hospital admission for immediate initiation and titration of NIV.

## **Methods**

We identified eight patients that presented to the University of Michigan Pranger ALS clinic from May 2020-May 2021 who were directly admitted for initiation or titration of NIV based on elevated TCO<sub>2</sub> levels. Decisions to admit patients were made based on elevated TCO<sub>2</sub> (TCO<sub>2</sub> >45mmHg), declining FVC measurements, and clinical symptoms concerning for impending respiratory failure in the judgement of the Pulmonology/Critical Care attending evaluating the patient in clinic. Data was collected via retrospective review of the medical records. All baseline clinic measurements were obtained either on day of admission or within 5 days prior. Pulmonary function measurements were obtained in clinic by a trained pulmonary function technician. All patients underwent three reproducible spirometry tests, with appropriateness of the patient effort indicated by the technician. The best of the three tests was chosen for interpretation by a pulmonary physician. Percent predicted values were based on sex, age, height, and race(7). Clinic TCO<sub>2</sub> was obtained using a Digital Monitoring System (SenTec AG, Switzerland) (**Supplementary Figure 1**). ALSFRS-R questionnaires were administered by trained providers as part of routine ALS multidisciplinary care. Inpatient NIV settings were titrated to patient tolerance for continuous nocturnal ventilation with additional adjustments for daytime support as needed with nasal or mouthpiece interface. Readiness for discharge was determined by tolerance of NIV plus stable or improving PaCO<sub>2</sub> prior to discharge. Discharge blood gases were obtained in-hospital within 24 hours of discharge. Post-hospitalization TCO<sub>2</sub> was obtained at the next in-person visit, typically within 30 days. NIV usage was determined based on data downloads using remote monitoring systems. Average daily use was recorded from these downloads. This study was approved by the Institutional Review Board.

## Results

Key demographic information and clinical measurements are presented in **Table 1**. Pre-hospitalization, inpatient and post-discharge respiratory measurements are presented in **Table 2**.

All 8 patients presented to clinic with symptoms or signs of respiratory insufficiency, such as dyspnea on exertion or apparent accessory muscle use. Seven patients were in clinic for their first multidisciplinary clinic visit after receiving the diagnosis of ALS. Patient 5 had previously been diagnosed with sleep apnea and had been started on nocturnal continuous positive airway pressure (CPAP) with 2L continuous nasal canula bleed through. Patient 7 was an established patient and had been using NIV for an average of 8-10 hours/night for at least 3 months prior to admission from clinic. All other patients were naïve to supplementary respiratory support. All patients had some bulbar dysfunction as captured by the ALSFRS-R. Only two patients had bulbar subscores <9. One patient had documented cognitive impairment.

All patients were directly admitted to a moderate care pulmonary service at University of Michigan Health, comprised of pulmonary/critical care physicians and respiratory therapists trained in NIV management. Average length of hospital stay was 6.5 days (range 3-8). Seven patients had worsening hypercapnia early in their admission, signifying advanced respiratory failure. All patients were titrated to tolerance of continuous nocturnal NIV while in the hospital. Seven patients required additional daytime ventilatory support via mouthpiece ventilation (MPV). At follow up, all patients were using NIV an average of >4 hours/day. Five patients reported at least 20 hours of NIV use per day. All patients, other than Patient 5, who had concomitant chronic obstructive pulmonary disease (COPD), had an improvement in measured pCO<sub>2</sub> prior to discharge or at follow up visits. Patient 6 had a stable TCO<sub>2</sub> at follow up 30 days after hospitalization, but at return visit 3 months later had a measured TCO<sub>2</sub> of 38 mmHg.

## Discussion

Most patients with ALS are initiated on NIV in the outpatient setting, prior to developing hypercapnia, but those who present with signs and symptoms of advanced respiratory failure may require inpatient admission for more rapid titration. There are no clear objective criteria based on pulmonary function test values or physical examination that would identify patients who need to be admitted. Importantly in our currently described cohort of patients, ambulatory measurements such as FVC, SpO<sub>2</sub>, and the respiratory symptom portion of the ALSFRS-R did not seem to provide consistent and persuasive evidence of impending respiratory failure, with two patients admitted with FVC > 50% predicted. This implies that effort-dependent measurements and subjective symptom reports alone are not sufficient to capture the complexities or provide sensitive markers of alveolar hypoventilation leading to hypercapnic respiratory failure.

Based on our experience with these 8 patients, TCO<sub>2</sub> measurement is a reliable, noninvasive means by which to identify patients with significant hypoventilation. Based on in-hospital PaCO<sub>2</sub> measurements, the patients identified in our case series could have progressed to fulminant respiratory failure requiring intubation and possibly death with any delay in starting NIV therapy. TCO<sub>2</sub> measurements in clinic provided important information that led to a decision to admit the patient for monitored titration of NIV, as the presence of hypercapnia indicated the patient's inability to maintain adequate minute ventilation without mechanical respiratory support. By admitting these patients, we were able to not only expeditiously initiate NIV, but also increase the probability of NIV tolerance and adherence in the immediate period after being seen in clinic, which has been shown to improve survival in ALS(8). European evaluations have shown improved survival with rapid titration of NIV settings, particularly in patients without

severe bulbar involvement(9). When comparing planned outpatient versus inpatient titration, waiting time for initiation and adverse events (death and admission for respiratory failure) were higher in the outpatient group and survival was extended for those admitted; however, this was a very small group of patients (n=12)(10).

There are several limitations to this case series. Foremost is the small sample size and retrospective nature of a case series. Additionally, all patients were seen in an ALS Center of Excellence with the presence of a pulmonary critical care physician trained in the management of neuromuscular disease. These patients were admitted to a pulmonary moderate care unit that frequently initiates NIV in patients with various conditions. These resources may not be available in smaller centers. Because outpatient initiation of NIV for ALS patients is primarily based on pulmonary function testing as opposed to PaCO<sub>2</sub>, it is unclear what the natural course of home NIV initiation for hypercapnic patients would be. While prospective controlled studies comparing home versus inpatient initiation of NIV for hypercapnic ALS patients would be ideal, the feasibility of these studies would be challenging given the rapidly progressive nature of ALS. Future studies looking at the incidence of hypercapnia at the first visit to an ALS Center would be useful.

In summary, although NIV is usually initiated in the home, patients presenting with hypercapnic respiratory failure may be at high risk of adverse outcomes with delays in treatment. Outpatient TCO<sub>2</sub> measurements can serve as a useful adjunctive measurement to identify ALS patients who would benefit from inpatient initiation and titration of NIV and avoid unwanted morbidity with prolonged outpatient titration, or more acute decompensation resulting in unwanted intubation or death. Further studies assessing the survival benefit of ambulatory TCO<sub>2</sub> monitoring and rapid initiation of NIV to tolerance with correction of CO<sub>2</sub> are also warranted.

**Abbreviations:**

ALS: amyotrophic lateral sclerosis

FVC: forced vital capacity

MIP: maximum inspiratory pressure

ALSFRS-R: ALS Functional Rating Scale-Revised

NIV: non-invasive ventilation

TCO<sub>2</sub>: transcutaneous CO<sub>2</sub>

CPAP: continuous positive airway pressure

COPD: chronic obstructive pulmonary disease

MPV: mouthpiece ventilation

**References**

1. L. I. Bruijn, T. M. Miller, D. W. Cleveland, Unraveling the mechanisms involved in motor neuron degeneration in ALS. *Annu Rev Neurosci* **27**, 723-749 (2004).
2. N. Lechtzin, J. Rothstein, L. Clawson, G. B. Diette, C. M. Wiener, Amyotrophic lateral sclerosis: evaluation and treatment of respiratory impairment. *Amyotroph Lateral Scler Other Motor Neuron Disord* **3**, 5-13 (2002).
3. J. M. Cedarbaum *et al.*, The ALSFRS-R: a revised ALS functional rating scale that incorporates assessments of respiratory function. BDNF ALS Study Group (Phase III). *J Neurol Sci* **169**, 13-21 (1999).
4. S. Niedermeyer, M. Murn, P. J. Choi, Respiratory Failure in Amyotrophic Lateral Sclerosis. *Chest* **155**, 401-408 (2019).
5. S. Aarrestad *et al.*, Validity of transcutaneous PCO<sub>2</sub> in monitoring chronic hypoventilation treated with non-invasive ventilation. *Respir Med* **112**, 112-118 (2016).

6. M. K. Rafiq *et al.*, Using transcutaneous carbon dioxide monitor (TOSCA 500) to detect respiratory failure in patients with amyotrophic lateral sclerosis: a validation study. *Amyotroph Lateral Scler* **13**, 528-532 (2012).
7. J. F. Morris, A. Koski, L. C. Johnson, Spirometric standards for healthy nonsmoking adults. *Am Rev Respir Dis* **103**, 57-67 (1971).
8. J. Ackrivo, J. Y. Hsu, J. Hansen-Flaschen, L. Elman, S. M. Kawut, Noninvasive Ventilation Use Is Associated with Better Survival in Amyotrophic Lateral Sclerosis. *Ann Am Thorac Soc* **18**, 486-494 (2021).
9. S. C. Bourke *et al.*, Effects of non-invasive ventilation on survival and quality of life in patients with amyotrophic lateral sclerosis: a randomised controlled trial. *Lancet Neurol* **5**, 140-147 (2006).
10. D. O'Brien *et al.*, The optimisation of noninvasive ventilation in amyotrophic lateral sclerosis: a systematic review. *Eur Respir J* **54**, (2019).

**Table 1: Demographics and Baseline Clinical Features of Patients Directly Admitted for Inpatient Initiation of Non-Invasive Ventilation**

Patient	Demographics			ALSFRS-R Scores				
	Age (years)	Gender (Male/Female)	BMI (kg/m <sup>2</sup> )	Total Score (0-48)	Bulbar Sub-score (0-12)	Dyspnea (0-4)	Orthopnea (0-4)	Respiratory Insufficiency (0-4)
1	73	F	25.5	29	9	2	1	4
2	54	M	28.9	35	11	2	3	4
3	65	M	21.7	38	8	3	4	4
4	62	F	39.2	NA	NA	NA	NA	NA
5	75	F	18.8	37	9	2	4	4 †
6	56	M	19.1	29	10	3	0	3
7	70	F	21.6	35	10	4	4	4
8	69	F	26.17	23	5	1	2	4

ALSFRS-R = ALS Functional Rating Scale-Revised; BMI = body mass index; NA = Not assessed. Obtaining outpatient clinical measurements were sometimes limited due to the COVID-19 pandemic.

†: Patient 5 had previously been using nocturnal CPAP machine at home

**Table 2: Respiratory Measurements of Patients Directly Admitted for Inpatient Initiation of Non-Invasive Ventilation**

Patient	Pre-Admission Clinic Measurements				Inpatient Serum Blood Gas Measurements				Post-Hospitalization Clinic Measurements	
	TCO2 (mmHg)	MIP (cmH20)	FVC (% predicted)	SpO2 (%)	Admission pH	Admission pCO2 (mmHg)	Discharge pH	Discharge pCO2 (mmHg)	TCO2 (mmHg)	NIV (hours)
1	62	-25	NA	91	7.35	84	7.4	75	45	>2
2	64	-45	66	88	7.38	66	NA	NA	43	>2
3	56	-18	46	96	7.33	72	7.41	59	47	>2
4	58	-29	39	87	7.38	66	7.4	58	46	>2
5	48	-28	61	93*	7.47‡	51	7.42	55	48	14
6	52	NA	NA	96	7.41‡	56	7.41	51	53	24
7	56	-19	44	94	7.44	54	7.5‡	37	43	14
8	59	-14	47	94	7.38‡	62‡	7.39‡	56	50	7-14

TCO2 = transcutaneous CO2; MIP = mean inspiratory pressure; FVC = forced vital capacity;

SpO2 = oxygen saturation by pulse oximetry; NIV = non-invasive ventilation; NA = not assessed.

\*: Obtained while on baseline 2L nasal cannula oxygen support

‡: Value obtained from venous blood gas, rather than arterial

Author Manuscript