

PROFESSOR JUNG-WOOK KIM (Orcid ID : 0000-0002-9399-2197)

Article type : Letter to the Editor

## Novel Homozygous *KREMEN1* Mutation causes Ectodermal Dysplasia

Yejin Lee<sup>1</sup>, Hong Zhang<sup>2</sup>, Figen Seymen<sup>3</sup>, Mine Koruyucu<sup>3</sup>, Yelda Kasimoglu<sup>3</sup>,  
Zang Hee Lee<sup>4</sup>, Jan C.-C. Hu<sup>2</sup>, James P. Simmer<sup>2</sup>, Jung-Wook Kim<sup>1,5</sup>

<sup>1</sup>Department of Pediatric Dentistry

School of Dentistry & DRI, Seoul National University, Seoul, Korea.

<sup>2</sup>Department of Biologic and Materials Sciences, School of Dentistry, University of  
Michigan, Ann Arbor, MI, USA.

<sup>3</sup>Department of Pedodontics, Faculty of Dentistry  
Istanbul University, Istanbul, Turkey.

<sup>4</sup>Department of Cell and Developmental Biology  
School of Dentistry & DRI, Seoul National University, Seoul, Korea.

<sup>5</sup>Department of Molecular Genetics  
School of Dentistry & DRI, Seoul National University, Seoul, Korea.

1) **Short Title:** *KREMEN1* Mutation

2) **Key Words:** Syndromic; Oligodontia; *KREMEN1*; Mutation; Ectodermal  
Dysplasia

3) **Date of submission:** 3-May-2021

**\*\*Correspondence to:**

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ODI.13921](https://doi.org/10.1111/ODI.13921)

This article is protected by copyright. All rights reserved

Jung-Wook Kim, D.D.S., M.S.D, Ph.D.

Department of Molecular Genetics

Department of Pediatric Dentistry

School of Dentistry & Dental Research Institute, Seoul National University

101 Daehak-ro, Jongno-gu, Seoul, 03080, Republic of Korea.

Tel: +82-2-2072-2639, Fax: +82-2-744-3599

E-mail: pedoman@snu.ac.kr

Tooth agenesis is one of the most common dental genetic disorders. Single or combination of multiple genetic factors are believed to be involved in the molecular pathogenesis in addition to the various harmful environmental factors (Thesleff, 2003). Hypodontia or oligodontia can occur as an isolated form or syndromic phenotype (Song, Bae, & Kim, 2020; Yang, Liang, Yue, & Bian, 2020). Because the tooth organ is ectodermally derived, the most frequent manifestation of tooth agenesis is as a syndromic phenotype of ectodermal dysplasia (ED). ED is a collection of genetic disorders affecting more than two tissues with ectodermal origin (Wright et al., 2019) and the current search in Online Mendelian Inheritance in Man database resulted in 233 EDs of which 65 EDs have tooth agenesis.

EDA/NF- $\kappa$ B and WNT/ $\beta$ -catenin pathways are important signaling pathways involved in tooth development (Yu, Wong, Han, & Cai, 2018). Mutations in genetic factors involved in this pathway, such as *WNT10A*, *WNT10B*, *LRP6* and *AXIN2*, have been identified and characterized. Recently, a homozygous recessive mutation in the gene encoding kringle containing transmembrane protein 1 (*KREMEN1*) has been determined to be responsible for ED with tooth agenesis in an extended Palestinian family (Issa et al., 2016). Clinical features were reported as oligodontia affecting both the primary and permanent teeth, abnormal hair distribution and brittle scalp hair, thin eyebrows and eyelashes, soft and glossy facial skin, and some dysmorphic features. Subsequently, two homozygous mutations of *KREMEN1* in Turkish families (Dinckan et al., 2018) and compound heterozygous mutations in a Thai

family (Intarak et al., 2018) have been reported to be involved with oligodontia with ED phenotype (Table S1).

In this study, we identified a novel homozygous mutation (c.97+2T>A) in a consanguineous Turkish family by whole-exome sequencing (Figure S1). The mutation destroyed the splicing donor site of intron 1 and would lead to a frameshift and a premature stop codon (Figure 1). This mutation would result in a lack of functional KREMEN1 due to the nonsense-mediated mRNA decay. The affected daughter was first evaluated at age 26-month old with multiple missing teeth. At age 8, it was apparent that she has curly fluffy hair, thin eyebrows and eyelashes, mildly dry skin and perioral pigmentation, but no dysmorphic facial features. The parents and the first unaffected child in this study were carriers but have no missing tooth or other features related to ED.

Kremen has a context-dependent biphasic Wnt signaling activity: Kremen potentiates Wnt/ $\beta$ -catenin signaling by maintaining LRP5/6 at the plasma membrane in the absence of Dickkopf1 (Dkk1), however, Kremen increases Dkk1-mediated Wnt inhibition in the presence of Dkk1 (Cselenyi & Lee, 2008). The mutation identified in this study would result in a lack of functional KREMEN1 at the plasma membrane. This report will not only expand the mutational spectrum of rare *KREMEN1* mutations but also provide further evidence to support the idea of *KREMEN1* as a candidate for oligodontia with mild to moderate ED symptoms.

## ACKNOWLEDGEMENTS

We thank the participants in this study for their cooperation. This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (NRF-2018R1A5A2024418 and NRF-2020R1A2C2100543) and National Institute of Dental and Craniofacial Institute (DE015846). The authors declare no potential conflicts of interest with respect to the authorship and/or publication of this article.

## CONFLICT OF INTEREST

The authors declare no conflict of interests.

## AUTHOR CONTRIBUTIONS

Y. Lee and H. Zhang conducted experiments and data analyses, critically revised manuscript; M. Koruyucu, Y. Kasimoglu and Z.H. Lee contributed to sample collection, clinical evaluation, data analysis and interpretation, critically revised manuscript; F. Seymen, J. Hu, J. Simmer, and J.-W. Kim, contributed to conception, design, data acquisition, analysis, and interpretation, drafted manuscript, critically revised the manuscript. All authors gave final approval and agree to be accountable for all aspects of the work.

## REFERENCES


- Cselenyi, C. S., & Lee, E. (2008). Context-dependent activation or inhibition of Wnt-beta-catenin signaling by Kremen. *Sci Signal*, *1*(8), pe10. doi:10.1126/stke.18pe10
- Dinckan, N., Du, R., Petty, L. E., Coban-Akdemir, Z., Jhangiani, S. N., Paine, I., . . . Letra, A. (2018). Whole-Exome Sequencing Identifies Novel Variants for Tooth Agenesis. *J Dent Res*, *97*(1), 49-59. doi:10.1177/0022034517724149
- Intarak, N., Theerapanon, T., Srijunbarl, A., Suphapeetiporn, K., Pornaveetus, T., & Shotelersuk, V. (2018). Novel compound heterozygous mutations in KREMEN1 confirm it as a disease gene for ectodermal dysplasia. *Br J Dermatol*, *179*(3), 758-760. doi:10.1111/bjd.16541
- Issa, Y. A., Kamal, L., Rayyan, A. A., Dweik, D., Pierce, S., Lee, M. K., . . . Kanaan, M. (2016). Mutation of KREMEN1, a modulator of Wnt signaling, is responsible for ectodermal dysplasia including oligodontia in Palestinian families. *Eur J Hum Genet*, *24*(10), 1430-1435. doi:10.1038/ejhg.2016.29
- Song, J. S., Bae, M., & Kim, J. W. (2020). Novel TSPEAR mutations in non-syndromic oligodontia. *Oral Dis*, *26*(4), 847-849. doi:10.1111/odi.13316
- Thesleff, I. (2003). Epithelial-mesenchymal signalling regulating tooth morphogenesis. *J Cell Sci*, *116*(Pt 9), 1647-1648.
- Wright, J. T., Fete, M., Schneider, H., Zinser, M., Koster, M. I., Clarke, A. J., . . . Morasso, M. I. (2019). Ectodermal dysplasias: Classification and organization by phenotype, genotype and molecular pathway. *Am J Med Genet A*, *179*(3), 442-447. doi:10.1002/ajmg.a.61045
- Yang, L., Liang, J., Yue, H., & Bian, Z. (2020). Two novel mutations in MSX1 causing oligodontia. *PLoS One*, *15*(1), e0227287. doi:10.1371/journal.pone.0227287

Yu, M., Wong, S. W., Han, D., & Cai, T. (2018). Genetic analysis: Wnt and other pathways in nonsyndromic tooth agenesis. *Oral Dis*. doi:10.1111/odi.12931

## FIGURE LEGENDS

**Figure 1** Clinical photos of proband, gene diagram and mutations of *KREMEN1*, a minigene splicing assay and an illustration of *KREMEN1* function. (a) Facial photo of the proband at age 8 years. Curly fluffy hair, thin eyebrows and eyelashes, mildly dry skin, and slight perioral pigmentation, but no other dysmorphic facial features can be seen. (b) Hands of the proband at age 8 years. No abnormality can be seen. (c) Gene diagram of *KREMEN1*. Gene structure (NM\_032045.5) shows 10 exons. White boxes indicate UTRs and grey boxes indicate coding exons. Exon numbers are shown above the boxes and previously reported mutations are shown. The mutation (c.97+2T>A) identified in this study is shown below the diagram. Nucleotide sequences with capital letter in bold are exon 1 and sequences with lower cases are intronic sequence. Amino acids encoded are indicated with black underlines in the exonic sequence. Mutated nucleotide is shown in red color. (d) In vitro splicing assay with a minigene showed a 251 bp normal splicing product (red arrow) only in the wild-type (W) not in the mutant vector (M). Black arrow indicates a band with vector sequence. Intron 1 retention due to the disruption of the splicing donor site would result in the premature translation termination in the intron 1 [p.(Glu33Glyfs\*316)]. However, mutant mRNA would be degraded by nonsense-mediated mRNA decay, resulting in a lack of functional *KREMEN1* (Appendix materials and methods). (e) *KREMEN1* binding to LRP5/6 potentiates WNT signaling at the plasma membrane in the absence of DKK1 (left illustration). However, *KREMEN1* increases DKK1-mediated WNT signaling inhibition by promoting the endocytosis of the *KREMEN1*-DKK1-LRP5/6 complex (right illustration).


(a)




(b)




(c)



(d)



(e)



This article is protected by copyright. All rights reserved