

Bertha Madeline (Orcid ID: 0000-0001-6361-3515)
Mellinger Jessica (Orcid ID: 0000-0001-7364-5035)

Trends in the Inpatient Burden of Alcohol Related Liver Disease Among Women Hospitalized in the United States

Madeline Bertha MD MSc¹, Kerby Shedden PhD², Jessica Mellinger MD MSc³

¹ Department of Internal Medicine, Northwestern University Hospital, Chicago IL; ² Department of Statistics, University of Michigan, Ann Arbor, MI; ³ Department of Internal Medicine, Michigan Medicine, Ann Arbor, MI.

Conflicts of Interest:

MB: None

KS: None

JLM: None

Grant Support: Dr. Mellinger is supported by an NIAAA K23 Career Development Award (AA026333).

Word Count: 1875

Corresponding Author:

Madeline Bertha
1340 N Astor Street Apt 2501
Chicago IL, 60610
Madeline.Bertha@gmail.com
9177339926

Abbreviations:

AC	alcohol-related cirrhosis
AH	alcoholic hepatitis
ALD	alcohol-related liver disease
AUD	alcohol use disorder
NIS	National Inpatient Sample

Short Title: Trends in ALD Among Hospitalized Women

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/liv.15277](https://doi.org/10.1111/liv.15277)

This article is protected by copyright. All rights reserved.

ABSTRACT:

Alcohol-related liver disease (ALD) is the leading indication for liver transplantation in the United States, but disparities for women with ALD exist. We sought to characterize trends in ALD hospitalizations and mortality among women. Using the National Inpatient Sample, we evaluated ALD and non-ALD discharges from 2003-2017. Multivariable logistic regression was used to evaluate mortality. ALD increased more rapidly among women, with alcohol hepatitis (AH) rising the most. When stratified by age and sex, changes in alcohol related cirrhosis (AC) and AH were greater in women in nearly all age groups. Similar increases were present when stratified by race, notably for Native American and Asian women. AH mortality increased in women in almost all age groups. While ALD remains predominantly male, discharges and mortality have disproportionately increased among women, particularly in young women and Native Americans. These findings shed light on populations in need of intensive public health interventions.

Key Words:

Alcohol Related Liver Disease

Alcohol Related Hepatitis

National Inpatient Sample

Hospitalized Women

INTRODUCTION: Alcohol-related liver disease (ALD) represents a spectrum disease ranging from fatty liver to advanced cirrhosis. ALD is now the most common indication for liver transplant in the United States (US) and represents an important public health problem.¹ It is increasingly well recognized that ALD disproportionately affects women because of increased physiologic vulnerability to alcohol's toxic effects as well as to disproportionate increases in rates of heavy alcohol use amongst women.²⁻⁴ Despite well-documented rising rates of ALD mortality among US women⁵, fewer studies have evaluated the intersection between sex, race, and age on ALD epidemiology and inpatient burden. Our goal was to specifically characterize the burden of ALD hospitalizations and mortality among women in the US as well as to define trends in ALD discharges and mortality for intersections of sex, age, and race.

METHODS: This study evaluated data from the National Inpatient Sample (NIS), the largest all-payer inpatient care database in the US. Using NIS "core" and "severity" data files for years 2003 through 2017, we defined three discharge subpopulations based on ICD-9 and ICD-10 code algorithms for AC, alcoholic hepatitis (AH) and non-alcohol-related cirrhosis, as previously published⁶. For our primary analysis we defined total ALD discharges as a combination of AC and AH discharges. We then analyzed subpopulations of AC alone and AH alone. Weighted and unweighted counts and marginal rates of discharges and in-hospital deaths were calculated for each disease subpopulation and year and reported per 100,000 discharges or deaths per year. Weighted trends in discharges and mortality were calculated for each disease group of interest against year and also reported as discharges or deaths per 100,000 discharges or deaths per year. Logistic regression, controlling for age, insurance status, income, race/ethnicity, year and chronic liver disease complications, was used to identify associations with mortality during hospital discharges for each population. We also calculated the marginal odds ratios between each explanatory factor in the logistic regression models and in-hospital mortality. Statistical analysis was performed using R and Python. IRB approval was waived for this study.

RESULTS: There were 382.2 discharges per 100,000 total all-cause discharges in 2003 and 358.0 discharges for every 100,000 discharges in 2017. Of all-cause discharges, 225.3 and 201.9/100,000 were women, respectively. Female ALD discharges made up 6.4 and 12.0/100,000 of the discharges in 2003 and 2017, respectively (**Table 1**). While ALD discharges increased among both sexes, they increased to a greater extent in women compared to men, (87.5% vs 57.8%). AH discharges rose most with a relative increase of 125.8% (16.2/100,000 in

2003 to 36.5/100,000 in 2017) for women compared to 89.4% for men. When stratified by sex and age (**Figure 1**), the relative changes in AC and AH discharges were greater in women compared to men across all age groups, except for AC discharges among men ≥ 65 . Most notably, women <34 had a 205.8% relative increase in total ALD discharges compared to 104.5% among men. In particular, women <34 had an increase in AC discharges of 145.6% (vs 40.9% among men) and AH discharges of 264.6% (vs 167.2% among men). When stratified by race, ALD discharges were greater for women among all race/ethnicities, but particularly for Native American women with an increase of 1,014% (341 in 2013 to 3,800 in 2017) and Asian women with an increase of 234%, driven primarily by AC discharge increases (323 in 2013 to 1,080 in 2017) (**Figure 1**).

There were 8.5 deaths for every 100,000 deaths (8.5/100,000) in 2003 and 7.0 deaths for every 100,000 deaths in 2017 (4.3 and 3.3/100,000 deaths among women) (**Table 1**). Total female and male ALD deaths decreased (-1.85% and -15.1%), driven by a decrease in AC deaths. AH deaths increased for both women and men (45.6% and 39.5%, respectively) (**Table 1**). When stratified by age (**Figure 1**), AH mortality increased among all age groups and sex, except women aged 35-44. AH mortality increases among women exceeded men for ages ≥ 65 (112.1% vs 8.2%) and 45-54 (26.9% vs 5.9%). In contrast, AC mortality decreased for all women except the ages of <34 and 55-64 (increase of 37.9% and 46.1%, respectively). When stratified by race/ethnicity, ALD mortality increased notably among the Native American populations, especially for women (594%) compared to men (318%). These rates were similar when categorized by AH and AC mortality (**Figure 1**). In multivariate analysis, stratified by AC vs AH, there was no significant difference in odds of death due to AH between sex ($p=0.52$), but there was an 8% decrease in odds of death due to AC among women compared to men ($p<0.01$).

DISCUSSION: This study of the US inpatient population showed an increasing burden of ALD discharges and mortality for young women of all races/ethnicities. While ALD remains a predominantly male disease, women are catching up, as ALD discharges increased more rapidly among women compared to men. Furthermore, despite an overall decrease in AC mortality, AH mortality has been increasing in women. Significant increases in alcohol use among women may be driving these findings.⁴ Women are more susceptible to ALD at lower levels of consumption with more rapid development of liver disease complications and death.³ These increases in ALD discharges and AH mortality among women, if persistent, will result in

increased AC deaths, suggestions that have been corroborated by projections of future ALD burden.⁷ Taken together, all of these findings may explain why the gap between death rates due to ALD among women and men continues to narrow.⁸

Similar to prior reports⁵, we found that AC discharges and mortality among women <34 progressed at an unprecedented rate. We noted both a marked increase in AC discharges and mortality among young women compared to men, a trend likely fed by high rates of alcohol misuse and AUD amongst high school and college age women^{4, 9, 10} However, despite an overall decrease in ALD mortality among older women, AH specific mortality in women >65 had a relative percentage increase of 112%, emphasizing that future public health interventions should not be limited to young women.

In our subgroup analysis, ALD discharges were greater for women among all race/ethnicities (excepting AH for Asian/Pacific Islander women). Our study not only supports previously published results that identify Hispanics as a population at substantially increased risk of developing ALD compared to non-Hispanic Whites and African Americans but sheds light on other at-risk race/ethnicities.⁸ In particular, we noted a dramatic relative percentage increase in both AH and AC related discharges and mortality among Native American women, both independently and compared to Native American men. While previous studies have highlighted AC and AUD as a leading cause of mortality among Native Americans, we are one of the first to highlight the increased burden of ALD among Native American women.^{11, 12}

Our study has limitations that are inherent to the use of a large claims database. Reliance of ICD9/10 codes for identification of patients with ALD is subject to both misclassification bias and reporting error. Additionally, the NIS reports data per hospitalization and not individual, which does not allow for patient-level longitudinal analysis but does allow for more detailed analysis of inpatient burden. Our analysis is subject to confounding variables that we are unable to account for. However, our findings are in agreement with previously published epidemiological trends for AUD and ALD. The observational nature of this data is hypothesis generating and does not allow for causal inference.

In conclusion, with ALD and AUD rates climbing rapidly, a public health emergency is underway for women in the US. As the number of women with ALD continues to increase, primary

preventive, gender-tailored interventions are urgently needed. Our results shed new light on subpopulations of women in need of intensive public health interventions.

References

1. Lee BP, Vittinghoff E, Dodge JL, et al. National Trends and Long-term Outcomes of Liver Transplant for Alcohol-Associated Liver Disease in the United States. *JAMA Intern Med* 2019;179:340-348.
2. Guy J, Peters MG. Liver disease in women: the influence of gender on epidemiology, natural history, and patient outcomes. *Gastroenterol Hepatol (N Y)* 2013;9:633-9.
3. Szabo G. Women and alcoholic liver disease - warning of a silent danger. *Nat Rev Gastroenterol Hepatol* 2018;15:253-254.
4. Grant BF, Chou SP, Saha TD, et al. Prevalence of 12-Month Alcohol Use, High-Risk Drinking, and DSM-IV Alcohol Use Disorder in the United States, 2001-2002 to 2012-2013: Results From the National Epidemiologic Survey on Alcohol and Related Conditions. *JAMA Psychiatry* 2017;74:911-923.
5. Tapper EB, Parikh ND. Mortality due to cirrhosis and liver cancer in the United States, 1999-2016: observational study. *BMJ* 2018;362:k2817.
6. Mellinger JL, Shedden K, Winder GS, et al. The high burden of alcoholic cirrhosis in privately insured persons in the United States. *Hepatology* 2018;68:872-882.
7. Julien J, Ayer T, Bethea ED, et al. Projected prevalence and mortality associated with alcohol-related liver disease in the USA, 2019-40: a modelling study. *Lancet Public Health* 2020;5:e316-e323.
8. Yoon YH, Chen CM, Slater ME, et al. Trends in Premature Deaths From Alcoholic Liver Disease in the U.S., 1999-2018. *Am J Prev Med* 2020;59:469-480.
9. Jones CM, Clayton HB, Deputy NP, et al. Prescription Opioid Misuse and Use of Alcohol and Other Substances Among High School Students - Youth Risk Behavior Survey, United States, 2019. *MMWR Suppl* 2020;69:38-46.
10. McCance-Katz EF. The Substance Abuse and Mental Health Services Administration (SAMHSA): New Directions. *Psychiatr Serv* 2018;69:1046-1048.
11. Sancar F, Abbasi J, Bucher K. Mortality Among American Indians and Alaska Natives. *JAMA* 2018;319:112.
12. Moon AM, Yang JY, Barritt ASt, et al. Rising Mortality From Alcohol-Associated Liver Disease in the United States in the 21st Century. *Am J Gastroenterol* 2020;115:79-87.

Table 1: Weighted Trends in Alcohol-Related liver Disease Discharges and Deaths Over Time









Discharges	2003		2017		Absolute Change	Relative Change
	Total	%	Total	%		
TOTAL	38,220,591	100%	35,798,453		-2,422,138	-6.34%
ALD	239,807	0.63%	392,980	1.10%	153,173	63.87%
AC	181,676	0.48%	281,490	0.79%	99,814	54.94%
AH	58,131	0.15%	111,490	0.31%	53,359	91.79%
Non-AC	190,090	0.50%	335,375	0.94%	145,285	76.43%
Gender						
<i>Female</i>	22,533,664	100%	20,186,250		-2,347,414	-10.42%
ALD	63,973	0.28%	119,950	0.59%	55,977	87.5%
AC	47,796	0.21%	83,420	0.41%	35,624	74.5%
AH	16,177	0.07%	36,350	0.18%	20,353	125.8%
Non-AC	94,147	0.42%	158,230	0.78%	64,083	68.1%
<i>Male</i>	15,686,927	100%	15,612,203		-74,724	-0.48%
ALD	175,834	1.12%	277,530	1.78%	101,696	57.8%
AC	133,880	0.85%	198,070	1.27%	64,190	47.9%
AH	41,954	0.27%	79,460	0.51%	37,506	89.4%
Non-AC	95,943	0.61%	177,145	1.13%	81,202	84.6%
Deaths						
TOTAL	849,267		698,690		-150,577	-17.7%
ALD	20,394	2.4%	17,430	2.49%	-2,964	-14.5%
AC	18,201	2.14%	14,330	2.05%	-3,871	-21.3%
AH	2193	0.26%	3100	0.44%	907	41.4%
Non-AC	15,335	1.81%	12,295	1.76%	-3,040	-19.8%
Gender						
<i>Female</i>	431,985		331,730		-100,255	-23.2%
ALD	5364	1.24%	5265	1.59%	-99	-1.9%
AC	4684	1.08%	4275	1.29%	-409	-8.7%
AH	680	0.16%	990	0.30%	310	45.6%
Non-AC	6866	1.59%	5550	1.67%	-1316	-19.2%
<i>Male</i>	417,282		366,960		-50,322	-12.1%
ALD	15,030	3.6%	12,760	3.48%	-2270	-15.1%
AC	13,517	3.24%	10,650	2.9%	-2867	-21.2%
AH	1513	0.36%	2110	0.57%	597	39.5%
Non-AC	8470	2.03%	6745	1.84%	-1725	20.4%

*ALD: alcohol-related liver disease; AC: alcohol-related cirrhosis; AH: alcoholic hepatitis

Figure 1: Relative Percentage Change (2003-2017) in Alcohol-Related Cirrhosis and Alcoholic Hepatitis Discharges and Mortality Stratified by Gender, Age, and Race/Ethnicity

Author Manuscript

Author Manuscript



LIV_15277_Figure 1.tif