

Can PATH Study susceptibility measures predict e-cigarette and cigarette use among American youth one year later?

Ruoyan Sun, PhD*, David Mendez, PhD**, and Kenneth E. Warner, PhD**

*University of Alabama at Birmingham, School of Public Health, Department of Health Care Organization and Policy, Birmingham, AL 35294.

**University of Michigan, School of Public Health, Department of Health Management and Policy, Ann Arbor, MI 48109.

Corresponding: Ruoyan Sun, PhD

University of Alabama at Birmingham
Health Care Organization and Policy
310C Ryals Public Health Building
1665 University Boulevard
Birmingham, AL 35294
(205)934-9387 || rysun@uab.edu

Running Head: PATH susceptibility predicts subsequent use

Conflict of Interest Declaration: None

Word Count: 2000

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/add.15808](https://doi.org/10.1111/add.15808)

ABSTRACT

Aims: To investigate whether e-cigarette and cigarette susceptibility predict e-cigarette and cigarette use among American youth 1 year later.

Design and Setting: Longitudinal data from the Population Assessment of Tobacco and Health (PATH) Study – a four-stage, stratified probability cohort study of youth (12-17 years old) sampled from the U.S. civilian, non-institutionalized population. Multivariable logistic regression was used to estimate the association between initial product-specific susceptibility and subsequent cigarette smoking and e-cigarette use while controlling for sociodemographic characteristics, exposure to nicotine users, and behavioral risk factors.

Participants: The sample included 8,841 adolescent never nicotine users at initial survey who participated in both wave 4 (2016-2017) and wave 4.5 (2017-2018) of PATH.

Measurements: We measured cigarette and e-cigarette susceptibility (defined as a lack of a firm commitment to not use cigarettes or e-cigarettes) among never nicotine users at baseline (wave 4) as well as cigarette and e-cigarette use at 12-month follow-up (wave 4.5).

Findings: Youth e-cigarette susceptibility was statistically significantly ($p < 0.05$) associated with e-cigarette use 1 year later, for both past 12-month (adjusted odds ratio [aOR] = 2.99, 95% confidence interval [CI], 2.29-3.90) and past 30-day e-cigarette use (aOR = 2.73, 95% CI, 1.78-4.16), but not with cigarette smoking (aOR = 1.05; 95% CI, 0.64-1.73 for past 12-month smoking and aOR = 0.65; 95% CI, 0.29-1.45 for past 30-day smoking). Smoking susceptibility predicted subsequent smoking in the past 12 months (aOR = 1.82, 95% CI, 1.09-3.03) and past 30 days (aOR = 3.32, 95% CI (1.33-8.29), but not e-cigarette use in the past 12 months (aOR = 0.96, 95% CI, 0.77-1.19) or past 30 days (aOR = 1.11, 95% CI, 0.82-1.51).

Conclusion: E-cigarette and cigarette susceptibility measures appear to predict product-specific use among youth 1 year later.

INTRODUCTION

In the last two decades, cigarette smoking has declined steadily among U.S. adolescents [1,2]. In 2020, 1.6% of middle school and 4.6% of high school students reported past-30-day use of cigarettes, compared with 12.8% and 34.8% in 1999 [3-5]. While cigarette use has dropped, use of electronic- or e-cigarettes has increased substantially among adolescents [6,7]. Since 2014, e-cigarettes have become the most commonly used nicotine-containing product among American middle and high school students [8,9]. From 2017 to 2018, current use (past 30-day use) of e-cigarettes increased from 11.7% to 20.8% among high school students and from 3.3% to 4.9% among middle school students [3]. The rates grew even higher in 2019, with 27.5% of high school and 10.5% of middle school students having used e-cigarettes in the past 30 days [10]. E-cigarette use fell significantly in 2020 to 19.6% of high school and 4.7% of middle school students [11]. Rates as high as those in the US are less common in other countries. For example, the prevalence of current e-cigarette use among 11-18 year olds in the UK was 4.1% in 2021 [12].

The concept of cigarette susceptibility, defined as the lack of a firm commitment to not use cigarettes, was introduced in 1995 [13]. Research has validated susceptibility as an independent predictor of smoking onset among adolescents [14-17]. Lately, studies examining e-cigarette susceptibility with the same construct have emerged. Using both cross-sectional [18,19] and longitudinal data [20-22], these studies have reported a significant positive association between e-cigarette susceptibility and subsequent use. However, little is known about the predictability of product-specific susceptibility with subsequent use of that nicotine-containing product, controlling for susceptibility to other products. Only one study [20] included both cigarette and e-cigarette susceptibilities. It found that cigarette susceptibility predicted e-cigarette use, but e-cigarette susceptibility did not predict future smoking.

Our study, investigating product-specific susceptibility and subsequent product use, examines if e-cigarette and cigarette susceptibility predict e-cigarette and cigarette use among American youth one year later. Using longitudinal data from the Population Assessment of Tobacco and Health (PATH) Study, we assess whether product-specific susceptibility is a significant independent predictor of initiation, controlling for sociodemographic characteristics, exposure to nicotine users, and behavioral risk factors. We include alcohol and marijuana use among our behavioral risk factors.

METHODS

PATH is a nationally representative, longitudinal cohort study of adults and youth in the U.S. Computer-assisted personal interviewing is used to collect self-reported data on nicotine product use and related health behaviors. A full description of the PATH design and methods is available [23]. Our sample included youth (12-17) who participated in both wave 4 (2016-2017) and wave 4.5 (2017-2018) of PATH. The weighted wave 4.5 response rate for the wave 1 cohort (2013-2014) was 74.6% and the weighted wave 4.5 response rate for the wave 4 cohort was 89.1%. We examined a subsample of youth who had never used any nicotine-containing product by wave 4 ($n = 8,841$), including life-time use of cigarettes, e-cigarettes, cigars, pipe, hookah, snus, smokeless tobacco, bidis, kreteks, and dissolvable tobacco.

Measures

The principal independent variables of interest are cigarette and e-cigarette susceptibility. Other covariates include sociodemographic characteristics, exposure to nicotine users, and behavioral risk factors of adolescent never nicotine users in wave 4. The dependent variables, smoking and e-cigarette use behaviors, are evaluated by respondents' reports in wave 4.5 of whether they had smoked or used e-cigarettes at all during the past 12 months (any use) and during the past 30 days (current use).

Susceptibility

In PATH, participants were asked four survey questions on potential use of cigarettes and e-cigarettes: “Have you ever been curious about using [product]?”, “Do you think that you will try a [product] soon?”, “If one of your best friends were to offer you a [product], would you [use] it?”, and “Do you think you will use a [product] in the next year?” Based on previous literature [13,14], we constructed a binary variable of susceptibility. Participants answering “not at all curious” to the first question and “definitely not” to the last three were considered not susceptible to cigarettes or e-cigarettes; the rest were regarded as susceptible.

Sociodemographic Characteristics

Sociodemographic variables included age (12-14 vs 15-17), sex (male vs female), race/ethnicity (non-Hispanic white, non-Hispanic black, Hispanic, and non-Hispanic other), highest parental education (high school/GED or less, some college, college or above), household income (<\$50,000, \$50,000-\$100,000, >\$100,000), and school grades (higher than mostly B's vs not).

Exposure to Nicotine Users

Family nicotine use (vs none) was evaluated by asking if anyone living with the respondent uses any nicotine-containing products(s). Youth exposure to secondhand smoke (vs no) was determined by asking respondents if they had been around others who were smoking, including at home, in a car, at school, or outdoors, for at least a total of an hour during the past seven days. Best friends' nicotine use (0 vs 1) is scored 1 if participants answered a positive number to any

of the questions asking “How many of your best friends use [product]?” The nicotine-containing products include cigarettes, e-cigarettes, cigarillos, snus and smokeless tobacco.

Behavioral Risk Factors

Behavioral risk factors included any past 12-month use (vs none) of alcohol and marijuana.

Analysis

All of the analyses were conducted using Stata version 16 (StataCorp LLC, College Station, TX).

We performed a complete case analysis via multivariable logistic regressions to examine the association between product-specific susceptibility measures at baseline (wave 4) and product consumption a year later (wave 4.5), controlling for sociodemographic characteristics, exposure to nicotine users, and behavioral risk factors. E-cigarette susceptibility has the most missing values, at 8.5%. Logistic regressions on missing values imply they are missing at random. We incorporated sample weights via the *svy* command in Stata to produce nationally representative results. The analysis was not pre-registered, and the results should be considered exploratory.

We also calculated the Variance Inflation Factor (VIF) to assess potential multicollinearity among our control variables. The VIF score is 1.35, implying limited multicollinearity. Pearson’s chi-squared test were performed in Table 1 to compare distributions of categorical variables.

RESULTS

Table 1 shows strong univariate associations between cigarette and e-cigarette susceptibility questions and subsequent smoking and e-cigarette use behaviors. Considering the binary measure of susceptibility (last four rows of the table), 18.5% of participants who were susceptible to e-cigarettes in PATH wave 4 used e-cigarettes in the subsequent 12 months while only 4.6% who were not susceptible did so ($p<0.001$). A higher percentage of those who were susceptible to e-cigarettes used e-cigarettes in the past 30 days as well (8.4% vs 1.9%, $p<0.001$). We report similar results for cigarette smoking. Compared with non-susceptible adolescents, a higher proportion of adolescents susceptible to cigarettes reported smoking later, both in the 12 months following wave 4 (3.6% vs 1.1%, $p<0.001$) and in the past 30 days (1.5% vs 0.4%, $p<0.001$).

Table 2 presents the relationships between other covariates and susceptibility measures. Among adolescent never nicotine users, higher proportions of the older group (15-17) are susceptible to e-cigarettes ($p<0.001$), based on Pearson's chi-squared test. We also report that higher proportions of youth are susceptible to e-cigarettes if they have family members who use nicotine ($p<0.001$), are exposed to secondhand smoke ($p<0.001$), have best friend(s) who used nicotine-containing products ($p<0.001$), or have past 12-month marijuana ($p<0.001$) and alcohol ($p<0.001$) use. Similarly, higher proportions of never nicotine users are susceptible to cigarettes if they are older ($p<0.001$), of the female sex ($p=0.018$), have family members who use nicotine products ($p<0.001$), are exposed to secondhand smoke ($p<0.001$), have best friend(s) who used

nicotine-containing products ($p < 0.001$), or have past 12-month marijuana ($p < 0.001$) and alcohol ($p < 0.001$) use. Multivariable logistic regressions reveal that significant factors affecting e-cigarette and cigarette susceptibility include race, family nicotine product use, secondhand smoke, best friends' nicotine product use, marijuana and alcohol consumption.

Table 3 reports weighted associations in adjusted odds ratios (aORs) of binary susceptibility measures with subsequent e-cigarette and cigarette use, controlling for our list of covariates. The aORs were 2.99 (95% CI, 2.29-3.90) between e-cigarette susceptibility and any use of e-cigarettes in the next 12 months, and 2.73 (95% CI, 1.78-4.16) for subsequent current (past 30-day) e-cigarette use, while the aORs for cigarette susceptibility were not significant. The aORs for cigarette susceptibility and subsequent cigarette smoking were 1.82 (95% CI, 1.09-3.03) for past 12-month smoking and 3.32 (95% CI, 1.33-8.29) for current smoking. E-cigarette susceptibility was not statistically significantly associated with subsequent cigarette use.

For sensitivity analysis, we tested 1) the weighted association between continuous susceptibility measures with subsequent product use, reported in supplementary Table S1; and 2) the unweighted association between the binary susceptibility measures and subsequent product use, reported in Table S2. The results in both cases are similar to our main findings.

DISCUSSION

This study examined e-cigarette and cigarette susceptibility among adolescent never nicotine users as predictors of e-cigarette and cigarette use within the next year. Our results agree with previous literature that susceptibility measures are significant independent predictors of future e-cigarette and cigarette use (any use and current use). E-cigarettes susceptibility is associated with subsequent any use and current use of e-cigarettes. Cigarette susceptibility is associated with any use and current use of cigarettes one year later. Our study contributes to the literature by examining more recent data (2016-2018) and controlling for important risky behaviors among never nicotine users, including marijuana and alcohol use.

In addition, by including both e-cigarette and cigarette susceptibility in our models, we investigated the predictability of product-specific susceptibility measures. We found that e-cigarette susceptibility was significantly associated with future e-cigarette use behaviors, but not with cigarette smoking behaviors. Similarly, smoking susceptibility was a significant predictor for subsequent cigarette smoking but not for e-cigarette use. Especially given all of our control variables, these findings imply that product-specific susceptibility is an independent risk factor for the future use of that specific product only. As such, we conclude that future studies of the association between e-cigarette use and smoking should include a cigarette-specific susceptibility measure as an independent risk factor.

Limitations

Our analysis examined, and hence only applies to, how susceptibility measures predict youth never nicotine users' subsequent product use. Other studies have found that the predictability of susceptibility measures varies by study sample. For example, the association of susceptibility and product initiation is stronger among never nicotine users [24,25], compared to those who have used other nicotine products. More research is needed to further examine how study sample influences susceptibility's association with product use.

We were unable to include in our study one risk factor included in many other studies [25-28]: a psychosocial measure of risk-taking propensity. PATH includes questions on sensation seeking but poses them only to new participants in each wave. As such we could not get a wave 4 sensation seeking measure for all respondents. Given our inclusion of other psychoactive substance use, however, this omission would be unlikely to affect our conclusion.

Another limitation is the reliability and validity of the self-reported data from PATH.

Tourangeau et al. [29] reinterviewed a subset of adults and youth from wave 4 of PATH and then followed self-report responses with saliva samples. They concluded that the self-report data on nicotine/tobacco use are both reliable and valid. More generally, Bachman et al. [30] reviewed a substantial body of evidence regarding multiple surveys of youth self-reporting of drug use, including nicotine/tobacco use, and found that the evidence supports the validity of the self-reported data.

We excluded from our analysis individuals who aged out of the wave 4.5 follow-up adolescent survey by having turned 18. PATH wave 4.5 is a special survey of 12-17 year-olds only. We believe that these 18 year-olds should not be substantially different from respondents who finished both surveys, thus making it unlikely that their exclusion would change our results.

Conclusion

E-cigarette and cigarette susceptibility measures are statistically significantly associated with e-cigarette and cigarette use a year later, respectively. These susceptibility measures predict product-specific use, where e-cigarette susceptibility predicts future e-cigarette use but not cigarette use and cigarette susceptibility predicts future cigarette smoking but not e-cigarette use. When available, as in PATH, susceptibility questions should be included in prospective studies examining the determinants of youth substance use.

References

1. Meza R, Jimenez-Mendoza E, Levy DT. Trends in tobacco use among adolescents by grade, sex, and race, 1991-2019. *JAMA Netw Open*. 2020;3(12):e2027465.
2. Dutra LM, Glantz SA. E-cigarettes and national adolescent cigarette use: 2004–2014. *Pediatrics*. 2017;139(2):e20162450.
3. Gentzke AS, Creamer M, Cullen KA, et al. Vital signs: tobacco product use among middle and high school students—United States, 2011–2018. *MMWR Morb Mortal Wkly Rep*. 2019;68:157-64.

4. Wang TW, Gentzke AS, Creamer MR et al. Tobacco product use and associated factors among middle and high school students—United States, 2019. *MMWR Morb Mortal Wkly Rep.* 2019;68(12):1-22.
5. Centers for Disease Control and Prevention. Tobacco use among middle and high school students—United States, 1999. *JAMA.* 2000;283(9):1134–6.
6. Miech R, Johnston L, O’Malley PM, Bachman JG, Patrick ME. Trends in adolescent vaping, 2017–2019. *N Engl J Med.* 2019;381(15):1490-1.
7. Chadi N, Hadland SE, Harris SK. Understanding the implications of the “vaping epidemic” among adolescents and young adults: A call for action. *Subst Abus.* 2019;40(1):7-10.
8. Centers for Disease Control and Prevention. About electronic cigarettes (e-cigarettes). 2020. https://www.cdc.gov/tobacco/basic_information/e-cigarettes/about-e-cigarettes.html (Accessed October 19, 2021).
9. US Department of Health and Human Services. E-cigarette use among youth and young adults. Atlanta, GA: US Department of Health and Human Services, CDC; 2016.
10. U.S. Food & Drug Administration. Youth tobacco use: results from the National Youth Tobacco Survey. 2020. <https://www.fda.gov/tobacco-products/youth-and-tobacco/youth-tobacco-use-results-national-youth-tobacco-survey> (Accessed October 19, 2021).
11. Wang TW, Neff LJ, Park-Lee E, Ren C, Cullen KA, King BA. E-cigarette use among middle and high school students—United States, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(37):1310-2.
12. Action on Smoking and Health (ASH). Use of e-cigarettes among young people in Great Britain. 2021. <https://ash.org.uk/wp-content/uploads/2021/07/Use-of-e-cigarettes-among-young-people-in-Great-Britain-2021.pdf> (Accessed October 19, 2021).
13. Pierce JP, Farkas AJ, Evans N, Gilpin E. An improved surveillance measure for adolescent smoking? *Tob Control.* 1995;4(suppl 1):S47–S56.
14. Pierce JP, Choi WS, Gilpin EA, Farkas AJ, Merritt RK. Validation of susceptibility as a predictor of which adolescents take up smoking in the United States. *Health Psychol.* 1996; 15(5):355-61.

15. Jackson C, Dickinson D. Cigarette consumption during childhood and persistence of smoking through adolescence. *Arch Pediatr Adolesc Med.* 2004;158(11): 1050-6.
16. Spelman AR, Spitz MR, Kelder SH, Prokhorov AV, Bondy ML, Frankowski RF, et al. Cognitive susceptibility to smoking: Two paths to experimenting among Mexican origin youth. *Cancer Epidemiol Biomarkers Prev.* 2009;18(12):3459-67.
17. Barrington-Trimis JL, Bae D, Schiff S, Davis J, Unger JB, Leventhal AM. Characterizing the predictive validity of measures of susceptibility to future use of combustible, vaporized and edible cannabis products in adolescent never-users. *Addiction.* 2020;115(12):2339-48.
18. Saddleson ML, Kozlowski LT, Giovino GA, Hawk LW, Murphy JM, MacLean MG, et al. Risky behaviors, e-cigarette use and susceptibility of use among college students. *Drug Alcohol Depend.* 2015;149:25-30.
19. Kwon E, Seo DC, Lin HC, Chen Z. Predictors of youth e-cigarette use susceptibility in a US nationally representative sample. *Addict Behav.* 2018;82:79-85.
20. Nicksic NE, Barnes AJ. Is susceptibility to e-cigarettes among youth associated with tobacco and other substance use behaviors one year later? Results from the PATH study. *Prev Med.* 2019;121:109-14.
21. Bold KW, Kong G, Cavallo DA, Camenga DR, Krishnan-Sarin S. E-cigarette susceptibility as a predictor of youth initiation of e-cigarettes. *Nicotine Tob Res.* 2017;20(1):140-4.
22. Cole AG, Kennedy RD, Chaurasia A, Leatherdale ST. Exploring the predictive validity of the susceptibility to smoking construct for tobacco cigarettes, alternative tobacco products, and e-cigarettes. *Nicotine Tob Res.* 2019;21(3):323-30.
23. Hyland A, Ambrose BK, Conway KP, et al. Design and methods of the Population Assessment of Tobacco and Health (PATH) Study. *Tob Control.* 2017;26(4):371-8.
24. Barrington-Trimis JL, Leventhal AM, Alonzo TA, et al. Performance of cigarette susceptibility index among e-cigarette and hookah users. *Drug Alcohol Depend.* 2018;183:43-50.
25. Barrington-Trimis JL, Liu F, Unger JB, et al. Evaluating the predictive value of measures of susceptibility to tobacco and alternative tobacco products. *Addict Behav.* 2019;96:50-5

26. Lozano P, Barrientos-Gutierrez I, Arillo-Santillan E, Morello P, Mejia R, Sargent JD, et al. A longitudinal study of electronic cigarette use and onset of conventional cigarette smoking and marijuana use among Mexican adolescents. *Drug Alcohol Depend.* 2017;180:427-30.
27. Primack BA, Soneji S, Stoolmiller M, Fine MJ, Sargent JD. Progression to traditional cigarette smoking after electronic cigarette use among US adolescents and young adults. *JAMA Pediatr.* 2015;169(11):1018-23.
28. Pierce JP, Chen R, Leas EC, White MM, Kealey S, Stone MD, Benmarhnia T, Trinidad DR, Strong DR, Messer K. Use of E-cigarettes and Other Tobacco Products and Progression to Daily Cigarette Smoking. *Pediatrics.* 2021;147(2):e2020025122.
29. Tourangeau R, Yan T, Sun H, Hyland A, Stanton CA. Population Assessment of Tobacco and Health (PATH) reliability and validity study: selected reliability and validity estimates. *Tob control.* 2019;28(6): 663-8.
30. Bachman JG, Johnston LD, O'Malley PM, Schulenberg JE, Miech RA. The Monitoring The Future Project after four decades: design and procedures. Monitoring the Future Occasional Paper Series, Paper #82. Ann Arbor, MI: University of Michigan, Institute for Social Research, 2015. <http://www.monitoringthefuture.org/pubs/occpapers/mtf-occ82.pdf> (Accessed October 19, 2021).

Table 1. Wave 4 never nicotine product users' weighted rates of use of e-cigarettes and cigarettes in Wave 4.5 by product-specific susceptibility measures, among US youth in the PATH study (N = 8,841).

| Susceptibility Responses, Wave 4 | Product Use ^a , Wave 4.5 | | | |
|---|---|--------------------------|------------------------------|-------------------------|
| | Electronic Cigarettes N (weighted %) | | Cigarettes N (weighted %) | |
| | Any Use (n = 628) | Current Use (n = 277) | Any Use (n = 151) | Current Use (n = 60) |
| Ever been curious about using [product]? | | | | |
| Yes | 279 (19.3) | 123 (8.6) | 66 (4.1) | 27 (1.7) |
| No | 314 (5.5) | 150 (2.4) | 85 (1.2) | 33 (0.4) |
| p-value | <0.001 ^b | <0.001 | <0.001 | <0.001 |
| Think you will try a [product] soon? | | | | |
| Yes | 183 (25.9) | 89 (12.6) | 35 (5.2) | 15 (2.4) |
| No | 437 (6.3) | 184 (2.6) | 116 (1.5) | 45 (0.5) |
| p-value | <0.001 | <0.001 | <0.001 | <0.001 |
| Would you use a [product] if one of your best friends offered you one? | | | | |
| Yes | 261 (22.6) | 135 (11.8) | 47 (4.6) | 20 (2.2) |
| No | 357 (5.6) | 138 (2.1) | 104 (1.4) | 40 (0.5) |
| p-value | <0.001 | <0.001 | <0.001 | <0.001 |
| Think you will use a [product] in the next year? | | | | |
| Yes | 225 (25.8) | 117 (13.4) | 44 (6.11) | 17 (2.46) |
| No | 395 (5.9) | 156 (2.3) | 107 (1.35) | 43 (0.52) |
| p-value | <0.001 | <0.001 | <0.001 | <0.001 |
| Binary measure: susceptible to any of the four questions | | | | |
| Yes | 360 (18.5) | 164 (8.4) | 80 (3.6) | 33 (1.5) |
| No | 260 (4.6) | 109 (1.9) | 71 (1.1) | 27 (0.4) |
| p-value | <0.001 | <0.001 | <0.001 | <0.001 |

Notes.

^a Any use: past 12-month use. Current use: past 30-day use.

^b Pearson's chi-squared test of independence was performed to assess the association.

Table 2. Descriptive statistics of never nicotine product users among US youth in the PATH study, by product susceptibility (N = 8,841).

| | Electronic Cigarettes | | p-value ^a | Cigarettes | | p-value |
|--------------------------------------|---|---|----------------------|---|---|--------------|
| | Susceptible (n = 2059, 25.3%) Weighted % (95% CI) | Unsusceptible (n = 6032, 74.7%) Weighted % (95% CI) | | Susceptible (n = 2252, 25.4%) Weighted % (95% CI) | Unsusceptible (n = 6575, 74.6%) Weighted % (95% CI) | |
| <i>Age</i> | | | 0.309 | | | 0.972 |
| 12-14 | 22.8 (21.5,24.2) | 77.2 (75.8,78.6) | | 23.8 (22.4,25.2) | 76.3 (74.8,77.6) | |
| 15-17 | 30.1 (28.1,32.1) | 70.0 (67.9,72.0) | | 28.8 (27.2,30.3) | 71.3 (69.7,72.8) | |
| <i>Sex</i> | | | 0.249 | | | 0.123 |
| Male | 25.3 (23.6,27.0) | 74.8 (73.0,76.4) | | 24.3 (23.0,25.8) | 75.7 (74.3,77.0) | |
| Female | 25.4 (23.8,27.0) | 74.6 (73.0,76.0) | | 26.5 (25.1,27.9) | 73.5 (72.1,74.9) | |
| <i>Race/Ethnicity</i> | | | <0.001 | | | <0.001 |
| NH White | 23.9 (22.3,25.6) | 76.1 (74.4,77.7) | | 24.5 (23.0,26.1) | 75.5 (73.9,77.0) | |
| NH Black | 27.8 (24.7,31.2) | 72.2 (68.8,75.3) | | 25.5 (22.9,28.4) | 74.5 (71.6,77.1) | |
| Hispanic | 27.0 (24.8,29.3) | 73.0 (70.7,75.2) | | 26.9 (24.8,29.1) | 73.1 (70.9,75.2) | |
| Other | 25.8 (22.3,29.6) | 74.2 (70.5,77.7) | | 26.1 (23.2,29.3) | 73.9 (70.7,76.8) | |
| <i>Parents education</i> | | | 0.069 | | | 0.250 |
| High school/GED or less | 24.6 (22.9,26.5) | 75.4 (73.6,77.1) | | 25.1 (23.3,27.0) | 74.9 (73.0,76.7) | |
| Some college | 25.5 (23.7,27.4) | 74.5 (72.7,76.3) | | 25.6 (23.8,27.4) | 74.4 (72.6,76.2) | |
| College or higher | 25.3 (24.1,26.7) | 74.7 (73.4,76.9) | | 25.4 (23.6,27.3) | 74.6 (72.7,76.4) | |
| <i>Household income</i> | | | 0.745 | | | 0.438 |
| < 50k | 25.6 (23.8,27.4) | 74.4 (72.6,76.2) | | 25.9 (24.3,27.7) | 74.1 (72.3,75.7) | |
| 50k to 100k | 26.2 (23.8,28.8) | 73.8 (71.2,76.2) | | 24.9 (22.4,27.4) | 75.2 (72.6,77.6) | |
| > 100k | 25.4 (23.2,27.8) | 74.6 (72.2,76.8) | | 26.1 (24.2,28.2) | 73.9 (71.8,75.8) | |
| <i>Grades higher than mostly B's</i> | | | 0.211 | | | 0.132 |
| Yes | 24.9 (23.4,26.5) | 75.1 (73.5,76.6) | | 24.8 (23.5,26.1) | 75.2 (73.9,76.6) | |
| No | 26.8 (24.6,29.1) | 73.2 (70.9,75.4) | | 27.2 (25.0,29.6) | 72.8 (70.4,75.0) | |
| <i>Family nicotine use</i> | | | 0.013 | | | 0.006 |
| Yes | 30.9 (28.7,33.2) | 69.1 (66.8,71.3) | | 31.4 (29.4,33.4) | 68.6 (66.6,70.6) | |
| No | 23.2 (21.8,24.6) | 76.8 (75.4,78.2) | | 23.3 (22.0,24.5) | 76.8 (75.5,78.0) | |

| | | | | | | |
|------------------------------------|-------------------------|-------------------------|------------------|-------------------------|-------------------------|------------------|
| <i>Secondhand smoke</i> | | | <0.001 | | | <0.001 |
| Yes | 34.3 (31.9,36.7) | 65.7 (63.3,68.1) | | 34.5 (32.3,36.7) | 65.5 (63.3,67.7) | |
| No | 22.2 (20.8,23.7) | 77.8 (76.3,79.2) | | 22.4 (21.1,23.8) | 77.6 (76.3,78.9) | |
| <i>Best friends' nicotine use</i> | | | <0.001 | | | <0.001 |
| Yes | 47.9 (45.3,50.5) | 52.1 (49.6,54.7) | | 41.9 (39.4,44.4) | 58.1 (55.6,60.6) | |
| No | 19.2 (18.1,20.5) | 80.8 (79.6,81.9) | | 21.2 (20.2,22.3) | 78.8 (77.7,79.8) | |
| <i>Past 12-month marijuana use</i> | | | 0.031 | | | 0.808 |
| Yes | 54.9 (46.2,63.4) | 45.1 (36.6,53.8) | | 40.5 (31.2,50.5) | 59.5 (49.5,68.8) | |
| No | 24.5 (23.3,25.8) | 75.5 (74.2,76.7) | | 24.9 (23.8,26.0) | 75.1 (74.0,76.2) | |
| <i>Past 12-month alcohol use</i> | | | <0.001 | | | <0.001 |
| Yes | 47.9 (45.0,50.9) | 52.1 (49.1,55.0) | | 45.2 (42.4,48.0) | 54.8 (52.0,57.6) | |
| No | 20.5 (19.3,21.8) | 79.5 (78.2,80.7) | | 21.6 (20.5,22.7) | 78.4 (77.3,79.6) | |

Notes.

^a Multivariable logistic regression was performed with sample weights, in which electronic cigarette or cigarette susceptibility (yes vs no) was the outcome variable, and covariates included age, sex, race/ethnicity, parents' education, household income, school grades, family tobacco use, secondhand smoke, best friends' tobacco use, and past 12-month use of marijuana and alcohol.

Bolded figures are statistically significant at $p < 0.05$.

Table 3. Weighted association of binary susceptibility measures with subsequent e-cigarette and cigarette use among US youth in the PATH study (N = 8,841).

| Wave 4 variables | Wave 4.5 outcomes ^a | | | | | | | |
|--|-----------------------------------|------------------|-----------------------------------|------------------|-----------------------------------|--------------|-----------------------------------|--------------|
| | Electronic Cigarettes | | | | Cigarettes | | | |
| | Any Use aOR (95% CI) | p-value | Current Use aOR (95% CI) | p-value | Any Use aOR (95% CI) | p-value | Current Use aOR (95% CI) | p-value |
| Susceptibility Measure | | | | | | | | |
| Susceptible to e-cigarettes (vs no) | 2.99 (2.29-3.90) | <0.001 | 2.73 (1.78-4.16) | <0.001 | 1.05 (0.64-1.73) | 0.839 | 0.65 (0.29-1.45) | 0.291 |
| Susceptible to cigarettes (vs no) | 0.96 (0.77-1.19) | 0.707 | 1.11 (0.82-1.51) | 0.500 | 1.82 (1.09-3.03) | 0.022 | 3.32 (1.33-8.29) | 0.011 |
| Covariates | | | | | | | | |
| Age (12-14 vs 15-17) | 1.29 (1.04-1.62) | 0.024 | 1.19 (0.88-1.60) | 0.248 | 1.40 (0.87-2.24) | 0.164 | 1.90 (0.87-4.15) | 0.105 |
| Female (vs male) | 0.99 (0.79-1.25) | 0.959 | 1.27 (0.96-1.69) | 0.089 | 0.97 (0.64-1.49) | 0.905 | 0.98 (0.51-1.86) | 0.940 |
| Race/Ethnicity (Ref: NH white) | | | | | | | | |
| NH black | 0.34 (0.22-0.51) | <0.001 | 0.27 (0.13-0.56) | 0.001 | 0.27 (0.09-0.83) | 0.023 | 0.18 (0.04-0.89) | 0.035 |
| Hispanic | 0.75 (0.56-1.02) | 0.067 | 0.75 (0.52-1.09) | 0.130 | 0.76 (0.45-1.30) | 0.317 | 0.57 (0.23-1.41) | 0.223 |
| Parents education (Ref: High school/GED or less) | | | | | | | | |
| Some college | 1.07 (0.79-1.46) | 0.656 | 1.04 (0.65-1.68) | 0.860 | 0.96 (0.51-1.80) | 0.906 | 1.57 (0.69-3.59) | 0.283 |
| College or higher | 1.50 (1.09-2.07) | 0.014 | 1.31 (0.83-2.06) | 0.240 | 0.92 (0.48-1.76) | 0.802 | 0.88 (0.33-2.33) | 0.796 |
| Household income (Ref: < 50k) | | | | | | | | |
| 50k to 100k | 1.12 (0.82-1.54) | 0.471 | 1.35 (0.87-2.10) | 0.182 | 1.04 (0.59-1.83) | 0.885 | 1.10 (0.54-2.27) | 0.786 |
| > 100k | 1.32 (0.99-1.76) | 0.057 | 1.70 (1.06-2.72) | 0.027 | 0.83 (0.45-1.54) | 0.558 | 0.67 (0.24-1.87) | 0.437 |

| | | | | | | | | |
|---------------------------------------|-----------------------------------|------------------|-----------------------------------|------------------|------------------------------------|------------------|-----------------------------------|--------------|
| Grades higher than mostly B's (vs no) | 0.70 (0.54-0.91) | 0.008 | 0.69 (0.45-1.05) | 0.085 | 0.46 (0.29-0.75) | 0.002 | 0.32 (0.15-0.68) | 0.003 |
| Family nicotine use (vs no) | 1.72 (1.39-2.13) | <0.001 | 1.32 (0.93-1.87) | 0.122 | 1.75 (1.13-2.70) | 0.012 | 1.89 (0.92-3.90) | 0.083 |
| Secondhand smoke (vs no) | 1.00 (0.79,1.28) | 0.977 | 1.08 (0.75-1.56) | 0.674 | 1.41 (0.81-2.45) | 0.219 | 1.17 (0.52-2.63) | 0.695 |
| Best friends' nicotine use (vs no) | 2.51 (1.94-3.25) | <0.001 | 1.99 (1.39-2.83) | <0.001 | 2.43 (1.57-3.78) | <0.001 | 3.27 (1.56-6.88) | 0.002 |
| Past 12-month alcohol use (vs no) | 2.43 (1.98-2.98) | <0.001 | 2.35 (1.70-3.25) | <0.001 | 1.60 (0.93-2.73) | 0.083 | 1.38 (0.61-3.13) | 0.442 |
| Past 12-month marijuana use (vs no) | 1.59 (0.87-2.89) | 0.128 | 2.23 (1.10-4.52) | 0.027 | 4.93 (2.16-11.21) | <0.001 | 3.23 (0.94-11.19) | 0.063 |

Notes.

^a Any use: past 12-month use. Current use: past 30-day use.

Bolded figures are statistically significant at $p < 0.05$.