

Editorial for “Aortic pulse wave velocity evaluated by 4D flow MRI across the adult lifespan”

Alexander Fyrdahl, PhD¹

¹Department of Radiology, University of Michigan, Ann Arbor, MI, USA

Corresponding author:
Alexander Fyrdahl, PhD
fyrdahl@med.umich.edu

Pulse wave velocity (PWV) is an established measure of arterial stiffness and is a strong predictor of cardiovascular events and all-cause mortality (1). PWV represents the rate at which the pressure wave caused by the cardiac pumping travels through the arteries. As the arterial walls become stiffer, either due to advanced age or other conditions, the rate at which the pressure wave travels through the arteries increases.

Conventionally, PWV is determined by measuring the time delay of the arrival of the pulse wave at the carotid artery and the femoral artery (2) using tonometry, oscillometry, or Doppler ultrasound. In the field of magnetic resonance imaging, estimation of PWV has been performed by imaging the velocity of flow in the aorta, either using a two-dimensional time-resolved phase-contrast sequence with through-plane velocity encoding and multiple slices (3), or using a three-dimensional time-resolved phase-contrast (4D flow) sequence with multiple velocity encoding directions (4,5). From these measurements, PWV can be calculated based on the distance and time delay of the flow profile in two or more imaging planes.

Establishing normative values for 4D-flow-derived PWV is important for future studies., however, such normative values are currently missing in the literature. In this issue of JMRI, the article “Aortic pulse wave velocity evaluated by 4D flow MRI across the adult lifespan” by Jarvis et. al. (6) used a 4D flow method to investigate differences in PWV in an age- and sex-stratified cohort of 100 healthy adults to better understand how PWV is impacted by normal aging. The evaluation also considered the correlation between PWV and cardiac functional parameters such as end-diastolic volume, stroke volume, myocardial mass, and mean aortic blood flow velocity.

As expected, the authors found that PWV increased with age, in concordance with previous work (7). The increase was found to be approximately 1 m/s per decade. Group-wise analysis based on stage of adulthood showed significant differences in PWV between groups. PWV was shown to correlate with other age-dependent factors such as decreased cardiac function and reduced aortic flow velocity, however, no difference in PWV due to aging was found between men and women. Previous studies have shown varying results with respect to sex, with some studies finding that the rate of PWV increases more with advancing age in men than in women (8), whereas others did not find sex-related differences in PWV (9). Another important aspect of this study was that it evaluated the repeatability of 4D-flow-derived PWV in a subset of the cohort. An intraclass correlation coefficient of 0.75 was found, indicating good test-retest repeatability. A limitation discussed by the authors is that the temporal resolution used in the study was lower than what was used in previous studies of 4D-flow-derived PWV. However, no formal consensus on suitable scan parameters for the determination of PWV has been established so far. Overall, this is an important study, and even though it may not be sufficient to establish normative values for 4D-flow derived PWV

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/jmri.28050](https://doi.org/10.1002/jmri.28050)

on its own, the work by Jarvis et. al. successfully demonstrates the utility of 4D-flow-derived PWV for studying the impact of aging on the aorta.

1. Vlachopoulos C, Aznaouridis K, Stefanadis C. Prediction of Cardiovascular Events and All-Cause Mortality With Arterial Stiffness. A Systematic Review and Meta-Analysis. *J. Am. Coll. Cardiol.* 2010;55:1318–1327.
2. Van Bortel LM, Laurent S, Boutouyrie P, et al. Expert consensus document on the measurement of aortic stiffness in daily practice using carotid-femoral pulse wave velocity. *J. Hypertens.* 2012;30:445–448.
3. Boese JM, Bock M, Schoenberg SO, Schad LR. Estimation of aortic compliance using magnetic resonance pulse wave velocity measurement. *Phys. Med. Biol.* 2000;45:1703–1713.
4. Markl M, Wallis W, Brendecke S, Simon J, Frydrychowicz A, Harloff A. Estimation of global aortic pulse wave velocity by flow-sensitive 4D MRI. *Magn. Reson. Med.* 2010;63:1575–1582.
5. Wentland AL, Wieben O, François CJ, Boncyk C, Munoz Del Rio A, Johnson KM, Grist TM, Frydrychowicz A. Aortic pulse wave velocity measurements with undersampled 4D flow-sensitive MRI: Comparison with 2D and algorithm determination. *J. Magn. Reson. Imaging* 2013;37:853–859.
6. [Placeholder reference] Aortic pulse wave velocity evaluated by 4D flow MRI across the adult lifespan. *J. Magn. Reson. Imaging* 2021;XX:YY–ZZ
7. Dyverfeldt P, Ebbers T, Länne T. Pulse wave velocity with 4D flow MRI: Systematic differences and age-related regional vascular stiffness. *Magn. Reson. Imaging* 2014;32:1266–1271.
8. AlGhatrif M, Strait JB, Morrell CH, Canepa M, Wright J, Elango P, Scuteri A, Najjar SS, Ferrucci L, Lakatta EG. Longitudinal trajectories of arterial stiffness and the role of blood pressure: The Baltimore longitudinal study of aging. *Hypertension* 2013;62:934–941.
9. Łoboz-Rudnicka M, Jaroch J, Kruszyńska E, Bociąga Z, Rzyckowska B, Dudek K, Szuba A, Łoboz-Grudzień K. Gender-related differences in the progression of carotid stiffness with age and in the influence of risk factors on carotid stiffness. *Clin. Interv. Aging* 2018;13:1183–1191.

Editorial for “Aortic pulse wave velocity evaluated by 4D flow
MRI across the adult lifespan”

Alexander Fyrdahl, PhD¹

¹Department of Radiology, University of Michigan, Ann Arbor, MI, USA

Corresponding author:
Alexander Fyrdahl, PhD
fyrdahl@med.umich.edu