

## EPIDEMIOLOGY

# Higher Household Air Pollution Levels Correlate with Poorer Cognitive Function in the Longitudinal Aging Study in India (LASI)

Sara Adar<sup>1</sup> | Jennifer D'Souza<sup>2</sup> | Gavin Shaddick<sup>3</sup> | Kenneth M Langa<sup>1,4</sup> | Alden L Gross<sup>5</sup> | Marco Angrisani<sup>6</sup> | Pranali Khobragade<sup>6</sup> | Joyita Banerjee<sup>7</sup> | Aparajita Chattopadhyay<sup>8</sup> | A. B. Dey<sup>7</sup> | T.V. Sekher<sup>8</sup> | David Bloom<sup>9</sup> | Jinkook Lee<sup>6</sup>

<sup>1</sup>University of Michigan, Ann Arbor, MI, USA

<sup>2</sup>University of Michigan School of Public Health, Ann Arbor, MI, USA

<sup>3</sup>University of Exeter, Exeter, United Kingdom

<sup>4</sup>Veterans Affairs Center for Clinical Management Research, Ann Arbor, MI, USA

<sup>5</sup>Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

<sup>6</sup>University of Southern California, Los Angeles, CA, USA

<sup>7</sup>All India Institute of Medical Sciences, New Delhi, India

<sup>8</sup>International Institute for Population Sciences, Mumbai, India

<sup>9</sup>Harvard T.H. Chan School of Public Health, Boston, MA, USA

**Correspondence**

Sara Adar, University of Michigan, Ann Arbor, MI, USA.

Email: [sadar@umich.edu](mailto:sadar@umich.edu)

**Abstract**

**Background:** Indians have some of the highest fine particulate matter (PM<sub>2.5</sub>) exposures in the world, not only due to high ambient air levels but also because of household combustion of biomass fuel. To understand the burdens of these exposures for older adults, we evaluated associations of indoor PM<sub>2.5</sub> with cognitive function in India.

**Method:** We used data from the 2017-2019 baseline survey of the nationally-representative Longitudinal Aging Study in India (LASI) to derive a general cognition score for each participant. For indoor PM<sub>2.5</sub> levels, we used an India-specific prediction model informed by participant cooking and housing characteristics. We estimated associations between indoor PM<sub>2.5</sub> with cognitive function using linear mixed-effects models after adjustment for age, sex, individual and community-level socioeconomic variables, region, clustering by village, and sample weights.

**Result:** We observed a population of 61,708 participants aged 45 and over. Many of these individuals (48%) used highly polluting fuel, resulting in average indoor PM<sub>2.5</sub> concentrations of 205 ± 110 µg/m<sup>3</sup>. After adjustment for individual and area-level confounders, we found that indoor PM<sub>2.5</sub> was associated with poorer cognitive function. For each 100 µg/m<sup>3</sup> of indoor PM<sub>2.5</sub>, we observed -0.027 (95% CI: -0.033, -0.02) lower cognition scores. These associations were strongest among women as well as among adults less than age 65 years. When scaling these associations to the observed interquartile range (150 µg/m<sup>3</sup>), household exposures resulted in cognitive differences comparable to those observed between people 2 years apart.

**Conclusion:** Among older adults living in India, household PM<sub>2.5</sub> is associated with poorer cognition at older ages even after adjustment for individual sociodemographic characteristics and regional trends.