

Note

A Remark about Permutations

WALTER FEIT, ROGER LYNDON, AND LEONARD L. SCOTT*

*Department of Mathematics, Yale University, New Haven, Connecticut 06520;
Department of Mathematics, University of Michigan, Ann Arbor, Michigan 48104;
and Department of Mathematics, University of Virginia,
Charlottesville, Virginia 22903*

Communicated by N. S. Mendelsohn

Received April 1, 1974

In [2] Ree proved a theorem about permutation groups by making use of a formula for the genus of Riemann surfaces. The purpose of this note is to provide a direct proof of Ree's theorem.

Let Ω be a finite set of n letters. For a permutation π on Ω define $v(\pi) = \sum_{i=1}^k (l_i - 1)$ where π is a product of disjoint cycles of lengths l_1, \dots, l_k .

LEMMA 1. *Let T be a minimal set of transpositions generating a transitive group on Ω . Then*

- (a) $|T| = n - 1$,
- (b) *the product in any order of all elements of T is an n -cycle.*

Proof. Define a graph G_T on Ω by joining α to β whenever $(\alpha\beta) \in T$. The fact that T generates a transitive group means that G_T is connected. Moreover G_T is a tree because of the minimality of T . Now (a) is immediate.

Let $\tau_1, \dots, \tau_{n-1}$ be the elements of T . We will show by induction on $|\Omega|$ that $\tau_1 \cdots \tau_{n-1}$ is an n -cycle. Of course we may rearrange the order of $\tau_1, \dots, \tau_{n-1}$ by any cyclic permutation, so we may assume $\tau_1 = (\alpha\beta)$ where α is joined only to β in G_T . Thus $G_{T - \{\tau_1\}}$ is a tree on $\Omega - \{\alpha\}$. By induction $\tau_2 \cdots \tau_{n-1}$ is an $(n - 1)$ -cycle $(\beta\gamma \cdots)$. So $\tau_1\tau_2 \cdots \tau_{n-1} = (\beta\alpha\gamma \cdots)$ is an n -cycle as required.

* This work was partially supported by the N.S.F., and represents independent work of the second author, and the first and third author in collaboration.

REE'S THEOREM. Let π_1, \dots, π_m be permutations on Ω such that $\pi_1 \cdots \pi_m = 1$. Then

$$v(\pi_1) + \cdots + v(\pi_m) \geq 2(n - s),$$

where s is the number of orbits of the group generated by π_1, \dots, π_m .

Proof. As Ree observes it clearly suffices to settle the case $s = 1$. We may also assume each π_i is a cycle, for if $\pi = \rho_1 \cdots \rho_k$ is a product of disjoint cycles, we have $v(\pi) = v(\rho_1) + \cdots + v(\rho_k)$. Finally if $\rho = (\alpha_1 \cdots \alpha_k)$ is a k -cycle, then $\rho = \tau_1 \cdots \tau_{k-1}$ where $\tau_i = (\alpha_1 \alpha_{i+1})$, and $v(\rho) = k - 1 = v(\tau_1) + \cdots + v(\tau_{k-1})$. So we may even assume that each π_i is a transposition.

Select from π_1, \dots, π_m a minimal subset $\pi_{i_1}, \dots, \pi_{i_r}$ which generates a group transitive on Ω . By Lemma 1 $r = n - 1$ and $\pi_{i_1} \cdots \pi_{i_r}$ is an n -cycle. From the equation $\pi_1 \cdots \pi_m = 1$ we obtain, by moving $\pi_{i_1}, \dots, \pi_{i_{n-1}}$ to the left, an equation $\pi_{i_1} \cdots \pi_{i_{n-1}} \pi_{n'} \cdots \pi_{m'} = 1$ where the π_j 's are conjugates of those π_i 's not among $\pi_{i_1}, \dots, \pi_{i_{n-1}}$.

Since $\pi_{i_1} \cdots \pi_{i_{n-1}}$ is an n -cycle, its inverse $\pi_{n'} \cdots \pi_{m'}$ is also an n -cycle. Thus $\pi_{n'}, \dots, \pi_{m'}$ generate a transitive group and so $m - n + 1 \geq n - 1$ by part (a) of Lemma 1. Hence $v(\pi_1) + \cdots + v(\pi_m) = m \geq 2(n - 1)$ and the proof is complete.

The function $v(\pi)$ has arisen previously in connection with Riemann surfaces. See for instance [1, Definition 3].

The following lemma provides alternative descriptions of $v(\pi)$.

Let $l(\pi)$ be the smallest integer l such that π is a product of l transpositions.

Let $t(\pi)$ be the multiplicity of 1 as a characteristic value of π considered as a permutation matrix.

LEMMA 2. $l(\pi) = v(\pi) = n - t(\pi)$.

Proof. Clearly $v(\pi) + t(\pi) = n$. Since a k -cycle is a product of $k - 1$ transpositions we see that $l(\pi) \leq v(\pi)$. Each transposition has the characteristic value 1 with multiplicity $n - 1$. Thus $t(\pi) \geq n - l(\pi)$. Therefore,

$$l(\pi) \leq v(\pi) = n - t(\pi) \leq l(\pi),$$

as required.

REFERENCES

1. M. FRIED, On a Conjecture of Schur, *Mich. Math. J.* 17 (1970), 41-55.
2. R. REE, A Theorem on Permutations, *J. Combinatorial Theory* 10 (1971), 174-175.