

## ON SYMMETRIC AND SKEW-SYMMETRIC DETERMINANTAL VARIETIES

JOE HARRIS† and LORING W. TU‡

(Received 26 June 1981)

### INTRODUCTION

LET  $E$  and  $F$  be differentiable vector bundles over a compact oriented manifold  $M$  and  $f: E \rightarrow F$  a bundle map between them. The locus of points in  $M$  where the rank of  $f$  does not exceed a given integer  $r$  is called the *degeneracy locus* of rank  $r$  of  $f$ , denoted  $M_r$ . As such a locus is defined locally by the vanishing of all the  $(r + 1)$  by  $(r + 1)$  minors of the matrix of  $f$ , in the algebraic category it is called a *determinantal variety*. In [17] Thom observed that the cohomology class of the degeneracy locus of a general map  $f: E \rightarrow F$  should be a polynomial in the Chern classes of  $E$  and  $F$ . This polynomial was found by Porteous in [16]:

$$[M_r] = (-1)^{(m-r)(n-r)} \Delta_{m-r, n-r} \left( \frac{c_i(E)}{c_i(F)} \right)$$

where if  $a(t) = \sum_{-\infty}^{\infty} a_k t^k$  is a formal series, then by definition

$$\Delta_{p,q}(a) = \det \begin{pmatrix} a_p & a_{p+1} & \cdots & a_{p+q-1} \\ a_{p-1} & a_p & & \vdots \\ a_{p-q+1} & \cdots & \cdots & a_p \end{pmatrix}.$$

The purpose of this article is to derive the analogous formulas for symmetric and skew-symmetric maps; these are expressed in Theorems 1 and 8. By a “squaring principle”, these results can be extended to a twisted bundle map  $f: E \rightarrow E^* \otimes L$ , where  $L$  is a line bundle (see Theorem 10). As applications of these formulas, we compute the degree of a determinantal variety, symmetric or not, and derive some numerical criteria for a variation of Hodge structure to be nontrivial.

We are indebted to the referee for detailed criticisms and to James Damon for pointing out an alternative method of deriving the degeneracy locus formulas (Theorems 1 and 8).

### §1. THE DEGENERACY LOCUS FORMULAS

#### (a) *Symmetric and skew-symmetric maps*

Let  $V$  be a vector space and  $V^*$  its dual vector space. A linear map  $f: V \rightarrow V^*$  is said to be *symmetric* if  $(f(x), y) = (f(y), x)$  for all  $x, y$  in  $V$ , where  $(\cdot, \cdot)$  is the dual pairing between  $V^*$  and  $V$ . Equivalently,  $f$  is symmetric if  $f = {}^t f$ . Relative to a basis for  $V$  and its dual basis

†Supported in part by NSF Grant MCS78-04008.

‡Supported in part by NSF Grant MSC80-02272.

Part of the work was done during the Montreal NATO Conference in August 1980; both authors wish to acknowledge the hospitality of the Séminaire de Mathématiques Supérieures at the University of Montreal.



polynomial  $P_r(c_1(S^*), \dots, c_k(S^*))$  gives the locus where a section of  $\text{Sym}^2 S^*$  has rank at most  $r$ , then its pullback

$$j^*P_r(c_1(S^*), \dots, c_n(S^*)) = P_r(c_1(E^*), \dots, c_n(E^*))$$

gives the degeneracy locus  $M_r$ .

One way of getting a section of  $\text{Sym}^2 S^*$  over  $G(n, m)$  is to take a smooth quadric in  $\mathbb{C}^m$  and restrict it to each  $n$ -plane in  $\mathbb{C}^m$ . The problem may now be stated in purely geometric terms:

*Given a smooth quadric  $Q$  in  $\mathbb{C}^m$ , let  $M_r$  be the locus of  $n$ -planes  $\Lambda$  in  $\mathbb{C}^m$  such that the restriction of  $Q$  to  $\Lambda$  has rank at most  $r$ . Find the cohomology class of  $M_r$  in  $H^*(G(n, m))$ .*

It turns out to be simpler, in the matter of indices, to consider the locus  $S_r = M_{n-r}$ ; we will show

$$(3) \quad S_r \sim 2^r \sigma_{r, r-1, \dots, 1}$$

in the homology of  $G(n, m)$ . We start by proving the

**PROPOSITION (4):** *If  $Q: W \rightarrow W^*$  is a quadric and  $V \subset W$  a linear subspace of codimension  $k$  in  $W$ , then*

$$\text{rank } Q|_V \geq \text{rank } Q - 2k$$

*Proof.* Note that if  $i$  is the inclusion of  $V$  in  $W$ , then the quadric  $Q|_V$  corresponds to the map

$$\tilde{Q}: V \xrightarrow{i} W \xrightarrow{Q} W^* \xrightarrow{i^*} V^*;$$

since composing with  $i$  or  $i^*$  can drop the rank of a map by at most  $k$ , the proposition follows.

Now, to compute the class of  $S_r$  in general, we intersect it with the Schubert cycles of the complementary codimension, i.e. we compute

$$S_r \cdot \sigma_b, \quad \text{where } \sum b_i = n(m-n) - \binom{r+1}{2}.$$

Now  $S_r \cdot \sigma_b$  is nonempty if and only if there are  $n$ -planes  $\Lambda$  in  $S_r$  such that

$$\dim(\Lambda \cap V_{m-n+i-b_i}) \geq i \quad \text{for } i = 1, \dots, n,$$

where the  $V$ 's form a generic flag of the indicated dimensions in  $\mathbb{C}^m$ . Let  $Q_i$  be the section  $Q \cap V_{m-n+i-b_i}$ . It is a smooth quadric of rank  $m-n+i-b_i$ ; furthermore, because  $\Lambda \cap Q$  has rank at most  $n-r$  by the definition of  $\Lambda$ ,  $(\Lambda \cap V_{m-n+i-b_i}) \cap Q_i$  also has rank at most  $n-r$ . By Proposition (4),

$$n-r \geq (m-n+i-b_i) - 2(m-n-b_i).$$

Therefore,

$$b_i \leq m-r-i.$$

By the definition of the Schubert cycles,  $b_i \leq m - n$ . Hence,

$$(1.5) \quad \begin{aligned} b_i &\leq m - n && \text{for } 1 \leq i \leq n - r, \\ b_i &\leq m - r - i && \text{for } n - r + 1 \leq i \leq n. \end{aligned}$$

Since by assumption

$$\sum_i b_i = n(m - n) - \binom{r + 1}{2},$$

equality must hold in (1.5). In other words,  $S_r \cdot \sigma_b \neq 0$  only if

$$b = m - n, \dots, m - n, m - n - 1, \dots, m - n - r.$$

This proves that  $S_r$  is a single-condition Schubert cycle:

$$S_r \sim a\sigma_{r,r-1,\dots,1} \text{ for some constant } a.$$

To determine  $a$  we have

$$a = S_r \cdot \underbrace{\sigma_{m-n,\dots,m-n,m-n-1,\dots,m-n-r}}_{n-r}$$

So  $a$  is the number of  $\Lambda$  in  $S_r$  satisfying the Schubert conditions

$$\dim(\Lambda \cap V_{n-r+2i}) \geq n - r + i, \quad i = 0, \dots, r.$$

and this number is readily computed: letting  $\Lambda_{n-r+i} = \Lambda \cap V_{n-r+2i}$ , we see that  $\Lambda_{n-r} = V_{n-r}$ , while for  $i = 1, \dots, r$ ,  $\Lambda_{n-r+i}$  must be spanned by  $\Lambda_{n-r+i-1}$  and either one of the two null-vectors of the quadratic form induced by  $Q$  in the 2-dimensional vector space

$$(V_{n-r+2i} \cap \Lambda_{n-r+i-1}^\perp) / \Lambda_{n-r+i-1}.$$

Thus  $a = 2^r$ , and Formula (3) is established.

To express this Schubert cycle in terms of the Chern classes of the dual of the universal subbundle  $S^*$  we apply Giambelli's formula (Griffiths and Harris[11], p. 205). Under the natural isomorphism

$$\begin{array}{ccc} S^* & & Q \\ \downarrow & & \downarrow \\ G(n, V) & \longrightarrow & G(m - n, V^*), \quad m = \dim V, \end{array}$$

the Schubert cycle  $\sigma_{r,r-1,\dots,1}$  in  $G(n, V)$  corresponds to  $\sigma_{r,r-1,\dots,1}$  in  $G(m - n, V^*)$ . Giambelli's formula gives

$$\sigma_{r,r-1,\dots,1} = \det \begin{pmatrix} \sigma_r & \sigma_{r+1} & \cdots \\ \sigma_{r-2} & \sigma_{r-1} & \cdots \\ & & \sigma_1 \end{pmatrix} \text{ in } G(m - n, V)$$

which corresponds to

$$\sigma_{r,r-1,\dots,1} = \det \begin{pmatrix} c_r & c_{r+1} & \cdots & & \\ c_{r-2} & c_{r-1} & & & \\ & & \cdots & & \\ & & & & c_1 \end{pmatrix}, \quad \text{where } c_i = c_i(S^*) \text{ in } G(n, V).$$

This completes the proof of Theorem 1.

*Remark 6.* Two cases of this formula, rank at most  $k - 1$  and rank 0, may be proved directly. Since  $M_{k-1}$  is the locus where  $\det f = 0$  and

$$\text{def } f: \det E \longrightarrow \det E^*$$

is a section of the line bundle  $(\det E^*) \otimes (\det E^*)$ ,

$$[M_{k-1}] = c_1((\det E^*)^{\otimes 2}) = 2c_1(E^*).$$

For  $M_0$  we note that it is the zero locus of a section of  $\text{Sym}^2 E^*$ , and is represented by the top Chern class  $c\binom{k+1}{2}(\text{Sym}^2 E^*)$ . A direct, though somewhat tricky, algebraic computation yields

$$c\binom{k+1}{2}(\text{Sym}^2 E^*) = 2^k \begin{vmatrix} c_k & c_{k+1} & \cdots & & \\ c_{k-2} & c_{k-1} & & & \\ & & \ddots & & \\ & & & & c_1 \end{vmatrix}, \quad \text{where } c_i = c_i(E^*).$$

*Remark 7.* The method of reduction to the Grassmannian in the proof of Theorem 2 is not practical for a Porteous locus, since if two bundles  $E$  and  $F$  are involved, it would be necessary to embed  $M$  into a product of Grassmannians  $G(m, N) \times G(n, N')$  and it is not easy to carry out explicitly the Schubert calculus on such a product.

*Remark.* A different approach to the proof of Theorem 1 is to mimic the standard construction used in deriving Porteous' formula: that is, we pass to the Grassmannian bundle  $G = G(n - r, E)$ . If  $Q$  is the universal quotient bundle on  $G$ , then we have a natural inclusion

$$\text{Sym}^2 Q^* \hookrightarrow \text{Sym}^2 \pi^* E^*;$$

let  $\Sigma$  denote the quotient bundle and  $\tilde{f}$  the image in  $\Sigma$  of the section  $\pi^* f$  of  $\text{Sym}^2 \pi^* E^*$ . Then  $M_r$ —the locus of  $x \in M$  such that  $f_x$  is the pullback to  $E_x$  of a quadric on some  $r$ -dimensional quotient of  $E_x$ —is just the image in  $M$  of the zero locus of  $\tilde{f}$ . The class of  $M_r$  is then just the Gysin image  $\pi_* e(\Sigma)$  of the Euler class of  $\Sigma$ . James Damon has shown us, via the referee, how this approach can be successfully carried out.

**SKEW-SYMMETRIC MAPS**

Again let  $E$  be a smooth complex vector bundle of rank  $n$  over a manifold  $M$ . A bundle map  $f: E \rightarrow E^*$  is *skew-symmetric* if  $f = -f'$ . Such a map may be viewed as a global section

of the exterior bundle  $\Lambda^2 E^*$  whose fiber at each point  $x$  in  $M$  consists of all the skew forms on the fiber  $E_x$ . Locally  $f$  is represented by a skew-symmetric matrix of functions. We now let  $M_r$  be the locus of points  $x$  in  $M$  where  $f_x$  has rank at most  $r$  and ask for the cohomology class of  $M_r$ . The result is

**THEOREM 8.** *Let  $E$  be a smooth complex vector bundle of rank  $n$  over a compact oriented manifold  $M$  and  $f: E \rightarrow E^*$  a general skew-symmetric bundle map. Then for any even  $r$ , the cohomology class of the degeneracy locus  $M_r$  is*

$$\begin{vmatrix} c_{n-r-1} & c_{n-r} & \cdots & & & \\ & c_{n-r-3} & c_{n-r-2} & & & \\ & & & \ddots & & \\ & & & & & c_1 \end{vmatrix},$$

where  $c_i = c_i(E^*)$ .

*Proof.* The proof here is exactly analogous to the symmetric case. First, we may reduce the problem to the following:

Given a nondegenerate skew form  $Q$  on  $\mathbb{C}^m$ , let  $S_r$  be the locus of  $n$ -planes  $\Lambda$  in  $\mathbb{C}^m$  such that  $Q|_\Lambda$  has rank at most  $n - r$ . Find the cohomology class of  $S_r$  in  $H^*(g(n, m))$ .

Secondly, since Proposition 4 applies just as well to skew forms as to quadrics, we see that

$$\begin{aligned} S_r &\sim a \cdot \sigma_{r-1, r-2, \dots, 1} \\ a &= S_r \cdot \sigma_{m-n, \dots, m-n, m-n-1, \dots, m-n-r+1} \\ &= \# \{ \Lambda \in S_r; \dim \Lambda \cap V_{n-r+2i-1} \geq n-r+i, \quad i = 1, \dots, r \}. \end{aligned}$$

Finally, the number  $a$  is easy to compute: if  $\Lambda$  satisfies these conditions and  $\Lambda_i = \Lambda \cap V_{n-r+2i-1}$ , we must have

$$\begin{aligned} \Lambda_1 &= V_{n-r+1} \\ \Lambda_2 &= \text{span of } \Lambda_1 \cap V_{n-r+3} \text{ and } \Lambda_1 \\ \Lambda_3 &= \text{span of } \Lambda_2 \cap V_{n-r+5} \text{ and } \Lambda_2 \\ &\vdots \end{aligned}$$

and finally

$$\Lambda = \Lambda_r = \text{span of } \Lambda_{r-1} \cap V_{n+r-1} \text{ and } \Lambda_{r-1}.$$

Thus,  $a = 1$ .

$$S_r \sim \sigma_{r-1, r-2, \dots, 1}$$

and Theorem 8 follows from Giambelli's formula as before.

**(b) Twisted symmetric and skew-symmetric maps (the squaring principle)**

Let  $E$  be a complex vector bundle and  $L$  a complex line bundle over  $M$ . A bundle map  $f: E \rightarrow E^* \otimes L$  is locally given by a square matrix of functions;  $f$  is said to be *symmetric*

or *skew-symmetric* according as the local matrix is symmetric or skew-symmetric around each point. Equivalently, if  $1_L: L \rightarrow L$  is the identity map and  $f^*: E \otimes L^* \rightarrow E^*$  the dual of  $f$ , then  $f$  is *symmetric* if  $f = f^* \otimes 1_L$ , and *skew-symmetric* if  $f = -f^* \otimes 1_L$ . Here we identify  $E \otimes L^* \otimes L$  with  $E$  by the natural pairing on  $L^*$  and  $L$ :

$$\langle , \rangle: E \otimes L^* \otimes L \xrightarrow{\sim} E.$$

Given such a twisted map  $f$ , one may again ask for the cohomology class of its degeneracy locus.

In case the line bundle  $L$  is a square, say  $L = K^2$ , the answer can be easily computed from Theorem 1 for

$$(\text{Sym}^2 E^*) \otimes L = \text{Sym}^2 (E^* \otimes K).$$

We claim that even when  $L$  is not the square of a line bundle, the same formula holds. To justify this, we will mimic the idea of the splitting principle and establish the following "squaring principle".

**PROPOSITION 9.** (The squaring principle). *Given a line bundle  $L$  over a manifold  $M$ , there exist a manifold  $X$ , a map  $\sigma: X \rightarrow M$  and a line bundle  $K$  on  $X$  such that*

- (a)  $\sigma^*L \cong K \otimes K$
- (b) *the induced map in cohomology  $\sigma^*: H^*(M, \mathbb{Z}) \rightarrow H^*(X, \mathbb{Z})$  is injective.*

This proposition says that we may compute the class of the degeneracy locus of a twisted symmetric bundle map  $f: K \rightarrow E^* \otimes L$  by positing the existence of a line bundle  $K = \sqrt{L}$  with  $K^{\otimes 2} \cong L$  and applying Theorem 2 to  $E \otimes K$ ; since the formula involves only even multiples of  $c_1(K)$  and we have  $2c_1(K) = c_1(L)$ , the answer will in the end involve only the Chern classes of  $E$  and  $L$ . The skew-symmetric case being completely analogous, we have the

**THEOREM 10.** (The degeneracy locus formula for twisted symmetric maps). *Let  $E$  be a rank  $k$  complex vector bundle and  $L$  a complex line bundle over  $M$ .*

- (a) *If  $f: E \rightarrow E^* \otimes L$  is a general symmetric map, then the cohomology class of  $M_f$  is the same as in Theorem 1, but with  $c_i = c_i(E^* \otimes \sqrt{L})$ ;*
- (b) *If  $f: E \rightarrow E^* \otimes L$  is a general skew-symmetric map, then the cohomology class of  $M_f$  is the same as in Theorem 8, but with  $c_i = c_i(E^* \otimes \sqrt{L})$ .*

*Proof of Proposition 9.* Let  $f_L: M \rightarrow \mathbb{C}P^\infty$  be the classifying map of the line bundle  $L$ , and  $g: \mathbb{C}P^\infty \rightarrow \mathbb{C}P^\infty$  the classifying map of  $S^{\otimes 2}$ , where  $S$  is the universal subbundle over  $\mathbb{C}P^\infty$ . Define  $X$  to be the fiber product of  $f_L$  and  $g$ . Then we have the following commutative diagram

$$\begin{array}{ccc}
 & & S^{\otimes 2} \\
 & & \swarrow \\
 X & \xrightarrow{h} & \mathbb{C}P^\infty \\
 \downarrow \sigma & & \downarrow g \\
 M & \xrightarrow{f_L} & \mathbb{C}P^\infty \\
 & & \swarrow S
 \end{array}$$

and

$$\sigma^*L = \sigma^*f_L^*S = h^*g^*S = h^*(S^{\otimes 2}) = (h^*S)^{\otimes 2}.$$

Hence  $X$  satisfies Property (a) of the proposition.

To show the injectivity of  $\sigma^*: H^*(M) \rightarrow H^*(X)$ , we first note that  $g^*: H^*(\mathbb{C}P^\infty) \rightarrow H^*(\mathbb{C}P^\infty)$  is injective, because  $H^*(\mathbb{C}P^\infty) = \mathbb{Z}[a]$  where  $a = c_1(S)$ , and

$$g^*(a) = g^*(c_1(S)) = c_1(g^*(S)) = c_1(S^{\otimes 2}) = 2a.$$


The injectivity of  $\sigma$  now follows from the following topology lemma.

LEMMA 11. *Let  $X$  be the fiber product of  $A \rightarrow C$  and  $B \rightarrow C$ .*

$$\begin{array}{ccc} X & \longrightarrow & B \\ f \downarrow & & \downarrow g \\ A & \longrightarrow & C \end{array}$$

Assume  $C$  to be simply connected. If  $g^*: H^*(C) \rightarrow H^*(B)$  is injective, then so is  $f^*: H^*(A) \rightarrow H^*(X)$ .

*Proof.* Recall from homotopy theory that every map is a fibering up to homotopy equivalence. So we may view  $g: B \rightarrow C$  as a fibering, say with fiber  $F$ , and  $f: X \rightarrow A$  has the same fiber. Since  $C$  is simply connected, the  $E^2$  term of the spectral sequence of the fibering  $g$  is the group  $H^*(C, H^*(F))$ . Because  $g^*$  is injective, the differentials which hit



the bottom row must be all zero. By the commutativity of the two spectral sequence diagrams, the same is true of the differentials of  $f: X \rightarrow A$ . Hence  $f^*: H^*(A) \rightarrow H^*(X)$  is injective.

Q.E.D.

(c) **Application: The degrees of determinantal varieties**

As an application of the degeneracy locus formulas we will now compute the degrees of determinantal varieties in a projective space.

PROPOSITION 12. (a) *Let  $V$  be the space of all  $m \times n$  matrices and  $V_k$  those of rank at most  $k$ . Then in the projective space  $\mathbb{P}(V)$ ,*

$$\deg \mathbb{P}(V_k) = \prod_{\alpha=0}^{m-k-1} \frac{\binom{n+\alpha}{m-1-\alpha}}{\binom{n-k+\alpha}{m-k-\alpha}}$$

(b) *Let  $W$  be the space of all  $n \times n$  symmetric matrices and  $S_r$  those of corank at least  $r$ . Then in  $\mathbb{P}(W)$*

$$\deg \mathbb{P}(S_r) = \prod_{\alpha=0}^{r-1} \frac{\binom{n+\alpha}{r-\alpha}}{\binom{2\alpha+1}{\alpha}}$$

(c) If in (b) the symmetric matrices are replaced by the skew symmetric matrices and  $n - r$  is even, then

$$\deg P(S_r) = \frac{1}{2^{r-1}} \prod_{\alpha=0}^{r-2} \frac{\binom{n+\alpha}{r-1-\alpha}}{\binom{2\alpha+1}{\alpha}}.$$

*Remark.* In (b),  $\mathbb{P}(W)_r = \mathbb{P}\binom{n+1}{2}^{-1}$  is the space of all quadrics in  $\mathbb{P}^{n-1}$  and  $\mathbb{P}(S_r)$  is the subvariety of quadrics of corank of at least  $n$ .

Because the computation of all three parts are very similar, we will carry out only the symmetric case.

Let  $(X_{ij})_{1 \leq i, j \leq n}$ ,  $X_{ij} = X_{ji}$ , be the homogeneous coordinates on  $\mathbb{P}(W)$ . Then  $\mathbb{P}(S_r)$  is the locus where  $\text{rank}(X_{ij}) \leq n - r$ . Since  $(X_{ij})$  may be regarded as a section of  $\text{Hom}^s(\mathcal{O}^{\oplus n}, \mathcal{O}^{\oplus n} \otimes \mathcal{O}(1))$  over  $\mathbb{P}(W)$ ,  $\mathbb{P}(S_r)$  is the degeneracy locus of corank  $r$  of this bundle.

If we write formally

$$\text{Sym Hom}(\mathcal{O}^{\oplus n}, \mathcal{O}^{\oplus n} \otimes \mathcal{O}(1)) = \text{Sym}^2\left(\mathcal{O}^{\oplus n} \otimes \mathcal{O}\left(\frac{1}{2}\right)\right) = \text{Sym}^2\left(\mathcal{O}\left(\frac{1}{2}\right)^{\oplus n}\right),$$

then by Theorem 1.10, the cohomology class of  $\mathbb{P}(S_r)$  is given by

$$2^r \begin{vmatrix} c_r & c_{r+1} & & & \\ c_{r-2} & c_{r-1} & & & \\ & & \ddots & & \\ & & & & c_1 \end{vmatrix}$$

where

$$c_i = c_i\left(\mathcal{O}\left(\frac{1}{2}\right)^{\oplus n}\right).$$

Since

$$c\left(\mathcal{O}\left(\frac{1}{2}\right)^{\oplus n}\right) = \left(1 + \frac{1}{2}\omega\right)^n = \sum_{i=0}^n \binom{n}{i} \frac{\omega^i}{2^i},$$

the degree of  $\mathbb{P}(S_r)$  is

$$(13) \quad \frac{1}{2 \binom{r}{2}} \begin{vmatrix} \binom{n}{r} & \binom{n}{r+1} & & & \\ \binom{n}{r-2} & \binom{n}{r-1} & & & \\ \vdots & & \ddots & & \\ & & & & \binom{n}{1} \end{vmatrix}$$

To complete the proof of the proposition, then, we have to evaluate this determinant: we do this in slightly greater generality in the following lemma, whose proof may be found in [1]:

LEMMA 14. For any  $n, r$  and  $a_1, \dots, a_r$  such that  $n \geq a_i + r - 1$  for all  $i$ ,

$$\det \begin{pmatrix} \binom{n}{a_1} & \binom{n}{a_1-1} & \cdots & \binom{n}{a_1-r+1} \\ \vdots & & & \vdots \\ \binom{n}{a_r} & \binom{n}{a_r-1} & \cdots & \binom{n}{a_r-r+1} \end{pmatrix} = \prod_{z=1}^r (n+r-1-a_z)! (a_z)! \prod_{1 \leq i < j \leq r} (a_j - a_i)$$

Applying this to the matrix in (13) with rows and columns reversed, we take

$$a_i = 2i - 1$$

and compute

$$\prod_{1 \leq i < j \leq r} (a_j - a_i) = \prod_{1 \leq i < j \leq r} (2j - 2i) = 2^{\binom{r}{2}} \prod_{i=1}^{r-1} i!;$$

we conclude that

$$\begin{aligned} \deg \mathbb{P}(S_r) &= \frac{1}{2^{\binom{r}{2}}} \det \begin{pmatrix} \binom{n}{r} & \cdots & \binom{n}{2r-1} \\ \binom{n}{r-2} & & \\ & \ddots & \\ & & \binom{n}{1} \end{pmatrix} \\ &= \prod_{\alpha=1}^r \frac{(\alpha-1)!(n+\alpha-1)!}{(2\alpha-1)!(n+r-2\alpha)!} \\ &= \prod_{\alpha=0}^{r-1} \frac{\alpha!(n+\alpha)!}{(2\alpha+1)!(n-r+2\alpha)!} \\ &= \prod_{\alpha=0}^{r-1} \frac{\binom{n+\alpha}{r-\alpha}}{\binom{2\alpha+1}{\alpha}}. \end{aligned}$$

Note that if  $r = n - 1$ , then

$$\begin{aligned} \deg \mathbb{P}(S_r) &= \frac{0!1! \cdots (n-2)!n!(n+1)! \cdots (2n-2)!}{1!3! \cdots (2n-3)!1!3! \cdots (2n-3)!} \\ &= \frac{1}{(n-1)!} \frac{1!2!3! \cdots (2n-2)!}{(1!3! \cdots (2n-3)!)^2} \\ &= \frac{1}{(n-1)!} \frac{2!4!6! \cdots (2n-2)!}{1!3!5! \cdots (2n-3)!} \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{(n-1)!} \cdot 2 \cdot 4 \cdot 6 \dots \cdot (2n-2) \\
 &= 2^{n-1},
 \end{aligned}$$

as expected (the variety  $\mathbb{P}(S_{n-1})$  of double hyperplanes is just the quadric Veronese embedding of  $(\mathbb{P}^{n-1})^*$  in  $\mathbb{P}^{\binom{n+1}{2}-1}$ ); while if  $r = 1$  we have

$$\text{deg } \mathbb{P}(S_1) = n$$

as expected. If  $r = 2$ ,

$$\text{deg } \mathbb{P}(S_2) = \frac{(n+1)!}{3!(n-2)!} = \binom{n+1}{3},$$

and if  $r = 3$ ,

$$\text{deg } \mathbb{P}(S_3) = \frac{n}{3} \binom{n+2}{5}.$$

*Remark.* The computation of Proposition 12 in fact proves something slightly more general; namely, if  $L$  is a complex line bundle over a compact oriented manifold  $M$  and  $(s_{ij})$  a transversal  $n \times n$  symmetric matrix of global sections of  $L$ , then the cohomology class of the locus where this matrix has corank at least  $r$  is

$$\left( \prod_{\alpha=0}^{r-1} \binom{n+\alpha}{r-\alpha} / \binom{2\alpha+1}{\alpha} \right) c_1(L)^{\binom{r+1}{2}}.$$

Analogous formulas of course hold for a general matrix or a skew-symmetric matrix of sections of a line bundle.

**APPLICATIONS TO VARIATION OF HODGE STRUCTURE**

Let  $\pi: X \rightarrow B$  be a family of curves of genus  $g$  over a compact base space, and  $H^{1,0}$  and  $H^{0,1}$  the associated Hodge bundles. In this section we will deduce, from the symmetric degeneracy locus formula, inequalities among the Chern numbers of the base space and the Hodge bundles when the variation of Hodge structure is nontrivial. For the definitions of these terms, see for instance [9] or [18].

Let  $S$  be the universal subbundle and  $Q$  the universal quotient bundle on the Grassmannian  $G(k, n)$ , and let  $\phi: B \rightarrow G(g, 2g)/\Gamma$  be the period map of the family of curves above. In this context the pullback to  $B$  of the universal subbundle over  $G(g, 2g)$  is the Hodge bundle  $H^{1,0}$  and the pullback to  $B$  of the universal quotient bundle is  $\cup H^1(X, \mathbb{C})/H^{1,0}(X) = H^{0,1}$ . So the tangent bundle to  $G(g, 2g)/\Gamma$  pulls back to  $\text{Hom}(H^{1,0}, H^{0,1})$  and the differential of the period map gives a bundle map

$$\phi_* : TB \rightarrow \text{Hom}(H^{1,0}, H^{0,1}).$$

It is well known that the period map may be given locally by  $g \times 2g$  matrices of the form  $(I, Z(t))$ ,  $t \in B$ , where the matrix  $Z(t)$  is *holomorphic* as a function of  $t$  and is *symmetric* by the Hodge–Riemann bilinear relations (Griffiths[10], pp. 613 and 616). So if  $v$  is a tangent vector to  $B$  at  $t$ , then  $\phi_*(v)$  is a symmetric map from  $H_t^{1,0}$  to  $H_t^{0,1}$ , i.e.

$$\phi_* : TB \rightarrow \text{Sym}^2(H^{1,0})^*.$$

Write  $E$  for the bundle  $H^{1,0}$  as well as its pullback to  $TB$ . Then the differential of the period map may be viewed as a section of  $\text{Sym}^2 E^*$  over  $TB$ . Since  $\phi_*(\lambda v) = \lambda \phi_*(v)$  for  $\lambda$  a complex number and  $v$  a tangent vector to  $B$ , if we projectivize the tangent bundle  $TB$ ,  $\phi_*$  gives rise to a section of  $\text{Sym}^2 E^* \otimes \mathcal{O}(1)$  over  $P = \mathbb{P}(TB)$ , where  $\mathcal{O}(1)$  is the universal line bundle over  $P$  which restricts to the hyperplane bundle on each fiber.

We will study several cases separately.

(a) *The base  $B$  is a curve and there are no singular fibers*

In this case the line bundle  $\det(\text{Sym}^2 E^* \otimes \mathcal{O}(1))$  has a holomorphic section  $\det(\phi_*)$  over  $P$ . Assume that the variation of Hodge structure is not constant. Then it follows from the local Torelli theorem that the differential of the period map  $\phi_*$  has maximal rank at some point, i.e.  $\det(\phi_*)$  is not identically zero. Therefore,

$$c_1(\det(\text{Sym}^2 E^* \otimes \mathcal{O}(1))) \geq 0.$$

Noting that  $c_1(\text{Sym}^2 E^*) = (g + 1)c_1(E^*)$ , this simplifies to

$$(15) \quad g(g(B) - 1) \geq c_1(E).$$

By a result of Griffiths [9, Corollary 7.10, p. 147] when the base is a complete curve and there are no singular fibers,  $c_1(E) \geq 0$ , with strict inequality if the variation of Hodge structure is nontrivial. Combined with (15), this gives

**PROPOSITION 15.1.** *A family of smooth curves over  $\mathbb{P}^1$  or an elliptic curve necessarily has trivial variation of Hodge structure.*

*Remark.* 15.2. The twisted degeneracy locus formula (Theorem 10) gives exactly the same inequality as (15), as follows. If the variation of Hodge structure over  $B$  is not trivial, then by the local Torelli theorem the period map  $\phi$  has maximal rank at some point of  $B$ . Hence the degeneracy locus  $M_{g-1}$  is not all of  $B$ . By Theorem 10,

$$[M_{g-1}] = -2c_1\left(E \otimes \mathcal{O}\left(-\frac{1}{2}\right)\right) = -2(c_1(E) - g(g(B) - 1)).$$

Because this is an effective divisor,

$$g(g(B) - 1) - c_1(E) \geq 0.$$

Note that here because the tangent bundle  $TB$  is a line bundle,  $P = \mathbb{P}(TB) = B$  and  $\mathcal{O}(-1)$  is the tangent bundle  $TB$ .

*Remark* 15.3. Proposition 15.1 also follows from the fact that the classifying space  $G(g, 2g)/\Gamma$  is a bounded domain in some  $\mathbb{C}^n$ . If the base is  $\mathbb{P}^1$ , the period map gives a holomorphic map from  $\mathbb{P}^1$  into  $\mathbb{C}^n$  and so must be a constant. If the base is a torus, the period map can be lifted to a holomorphic map  $\tilde{\phi}$  on the universal covering  $\mathbb{C}$  of the torus

$$\begin{array}{ccc} \mathbb{C} & \xrightarrow{\tilde{\phi}} & G(g, 2g)/\Gamma \subset \mathbb{C}^n \\ \downarrow & \searrow \phi & \\ B & \longrightarrow & \end{array}$$

Since there are no bounded entire holomorphic functions on  $\mathbb{C}$  except the constants,  $\tilde{\phi}$  and therefore  $\phi$  also are constant.

*Remark 15.4.* To underscore the fact that the inequality (15) holds only if the variation of Hodge structure is nontrivial, consider a product family of smooth curves over  $\mathbb{P}^1$ . Inequality (15) gives  $-g \geq 0$ , which is clearly false. The argument breaks down because in this case  $\det(\phi_*)$  is identically zero, so that the degeneracy locus  $M_{g-1}$  does not have the expected dimension.

(b) **The base  $B$  is a curve and there are singular curves**

Let  $D$  be the divisor on  $B$  parametrizing the singular fibers. The period map  $\phi$  is defined only over  $B - D$ . Now one version of the theorem on regular singular point says that the differential of the period map has at most simple poles over  $D$ . Therefore  $\phi_*$  is a section of  $\text{Sym}^2 E^* \otimes \mathcal{O}(D) \otimes \mathcal{O}(1)$  over  $P = \mathbb{P}(TB)$ , where we write  $\mathcal{O}(D)$  for the pullback of the line bundle to  $P$ . Applying the twisted degeneracy locus formula (Theorem 10), we get

$$g(g(B) - 1) + \frac{\deg D}{2} \geq c_1(E). \quad (16)$$

If the nontriviality of the variation of Hodge structure is measured by the degree of the Hodge bundle  $H^{1,0}$ , then (16) says that *the more nontrivial the variation of Hodge structure, either the larger the genus or the base curve or the greater the number of singular fibers.*

(c) **The base  $B$  is a surface and there are no singular fibers**

The differential of the period map  $\phi_*$  is a section of  $\text{Sym}^2 E^* \otimes \mathcal{O}(1)$  over  $P = \mathbb{P}(TB)$ , which is a compact threefold. If it is a general section, then the degeneracy locus formula gives an effective cycle of codimension 3 on  $P$ ,

$$M_{g-2} = -4(c_1c_2 - c_3) \geq 0 \quad (17)$$

where

$$c_i = c_i\left(E \otimes \mathcal{O}\left(\frac{1}{2}\right)\right).$$

Recall that if  $E$  is a rank  $g$  vector bundle and  $L$  is a line bundle, then the Chern classes of the tensor product  $E \otimes L$  are given by

$$c_r(E \otimes L) = \sum_{i=0}^r \binom{g-i}{r-i} c_i(E) c_1^{r-i}(L).$$

Recall also that if  $x = c_1(\mathcal{O}(1))$  is the hyperplane class on  $P$  and  $\pi: P \rightarrow B$  is the projection map, then the Gysin map  $\pi_*$  is given by

$$\pi_* x = 1,$$

$$\pi_* x^2 = c_1(-B) = -c_1(B),$$

$$\pi_* x^3 = c_2(-B) = c_1^2(B) - c_2(B),$$

and so on. Using these facts, (1.20) may be simplified to

$$2((g-1)c_1^2(E) + 2c_2(E)) + (g+1)(g-1)c_1(E)c_1(B) + \binom{g+1}{3}(c_1^2(B) - c_2(B)) \geq 0.$$

This formula holds whenever the degeneracy locus of the differential of the period map, considered as a section of  $\text{Sym}^2 E^*$  over  $TB$ , has the expected codimension 3.

## REFERENCES

1. E. ARBARELLO, M. CORNALBA, P. GRIFFITHS and J. HARRIS: *Geometry of Algebraic Curves*, to appear.
2. W. BARTH: Counting singularities of quadratic forms on vector bundles. In *Vector Bundles and Differential Equations. Proc. of the Nice Conf.*, 12–17 June 1979 (Edited by André Hirschowitz. Birkhäuser, Boston, 1–19).
3. J. BOARDMAN: Singularities of differentiable maps. *Publ. Math. I.H.E.S.* **33** (1967), 21–57.
4. J. DAMON: The Gysin homomorphism for flag bundles. *Am. J. Math.* **95** (1973), 643–659.
5. J. DAMON: The Gysin homomorphism for flag bundles: applications. *Am. J. Math.* **96** (1974), 248–260.
6. W. FULTON and R. LAZARSFELD: On the connectedness of degeneracy loci and special divisors. *Acta Math.* **146** (1981), 271–283.
7. D. GIESEKER: Stable curves and special divisors, I, preprint.
8. M. GOLUBITSKY and V. GUILLEMIN: *Stable Mappings and Their Singularities*, Graduate Text in Math. 14. Springer-Verlag, New York (1973).
9. P. GRIFFITHS: Periods of integrals on algebraic manifolds III. *Publ. Math. I.H.E.S.* **38** (1970), 125–180.
10. P. GRIFFITHS: Complex analysis and algebraic geometry. *Bull. Am. Math. Soc.* **1** (1979), 595–626.
11. P. GRIFFITHS and J. HARRIS: *Principles of Algebraic Geometry*. Wiley, New York (1978).
12. G. KEMPF: Schubert methods with an application to algebraic curves. *Publication of the Mathematisch Centrum*. Amsterdam (1971).
13. S. KLEIMAN: The enumerative theory of singularities. In *Real and Complex Singularities (Proc. 9th Nordic Summer School/NAVF Sympos. Math., Oslo)* (1976) (Edited by P. Holm), pp. 297–396. Sijthoff and Noordhoff, Alphen aanden Rijn (1977).
14. S. KLEIMAN and D. LAKSOV: Another proof of the existence of special divisors. *Acta Math.* **132** (1974), 163–176.
15. H. I. LEVINE: Singularities of Differentiable Maps, based on lectures by R. Thom, *Proc. of Liverpool Singularities Symposium I*, Lecture Notes in Math. 192. Springer-Verlag, New York (1971), 1–89.
16. I. R. PORTEOUS: Simple singularities of maps, *Proc. Liverpool Singularities Symposium I*, Lecture Notes in Math. 192 (Edited by C. T. C. Wall), Springer-Verlag, New York (1971), 286–307.
17. R. THOM: Les ensembles singuliers d'une application différentiable et leur propriétés homologiques. Séminaire de Topologie de Strasbourg, Dec. 1957.
18. L. TU: Variation of Hodge structure and the local Torelli problem, *Thesis*, Harvard University (1979). *Note.* It is mentioned in Barth[2] that the formulas for the cohomology class of a symmetric determinantal variety has also appeared in the following paper:
19. T. JOSEFIK, A. LASCoux and P. PRAGACZ: Classes of determinantal varieties associated with symmetric and antisymmetric matrices. *Izvestia* **45** (1981), 662–673.

*Department of Mathematics*  
*Brown University*  
*Providence, RI 02912, U.S.A.*

*Department of Mathematics*  
*University of Michigan*  
*Ann Arbor, MI 48109, U.S.A.*

*School of Mathematics*  
*Institute for Advanced Study*  
*Princeton, NJ 08540, U.S.A.*