

## Growth and nitrogen accretion of dinitrogen-fixing *Alnus glutinosa* (L.) Gaertn. under elevated carbon dioxide

Christoph S. Vogel<sup>1</sup>, Peter S. Curtis<sup>1</sup> & Richard B. Thomas<sup>2\*</sup>

<sup>1</sup>Department of Plant Biology, The Ohio State University, 1735 Neil Avenue, Columbus, OH 43210, USA and The University of Michigan Biological Station, Pellston, MI 49769, USA; <sup>2</sup>Department of Botany, Duke University, Durham, NC 27708, USA (\*Present address: Department of Biology, University of West Virginia, Morgantown WV 26506, USA)

Received 23 March 1996; accepted in revised form 26 November 1996

**Key words:** Black alder, Carbon Dioxide Enrichment, Delta <sup>15</sup>N Analysis, Nitrogen Fixation, Root Nodules

### Abstract

Short-term studies of tree growth at elevated CO<sub>2</sub> suggest that forest productivity may increase as atmospheric CO<sub>2</sub> concentrations rise, although low soil N availability may limit the magnitude of this response. There have been few studies of growth and N<sub>2</sub> fixation by symbiotic N<sub>2</sub>-fixing woody species under elevated CO<sub>2</sub> and the N inputs these plants could provide to forest ecosystems in the future. We investigated the effect of twice ambient CO<sub>2</sub> on growth, tissue N accretion, and N<sub>2</sub> fixation of nodulated *Alnus glutinosa* (L.) Gaertn. grown under low soil N conditions for 160 d. Root, nodule, stem, and leaf dry weight (DW) and N accretion increased significantly in response to elevated CO<sub>2</sub>. Whole-plant biomass and N accretion increased 54% and 40%, respectively. Delta-<sup>15</sup>N analysis of leaf tissue indicated that plants from both treatments derived similar proportions of their total N from symbiotic fixation suggesting that elevated CO<sub>2</sub> grown plants fixed approximately 40% more N than did ambient CO<sub>2</sub> grown plants. Leaves from both CO<sub>2</sub> treatments showed similar relative declines in leaf N content prior to autumnal leaf abscission, but total N in leaf litter increased 24% in elevated compared to ambient CO<sub>2</sub> grown plants. These results suggest that with rising atmospheric CO<sub>2</sub> N<sub>2</sub>-fixing woody species will accumulate greater amounts of biomass N through N<sub>2</sub> fixation and may enhance soil N levels by increased litter N inputs.

### Introduction

As atmospheric CO<sub>2</sub> levels rise, growth of many woody plant species is expected to increase (Bazzaz et al. 1990; Ceulemans & Mousseau 1994). Low soil N availability, however, may limit the magnitude of both the photosynthetic (Tissue et al. 1993; Curtis et al. 1995) and the overall growth response (Pregitzer et al. 1995; Wong et al. 1992) of trees to CO<sub>2</sub> enrichment. Negative acclimation of photosynthesis is commonly observed in trees grown at elevated CO<sub>2</sub> (Gunderson & Wullschlegler 1994) and may be due to the decline in leaf N content that often occurs at elevated CO<sub>2</sub>, especially under conditions of low soil N availability (Curtis et al. 1995; Norby et al. 1986a). Linked soil-plant process models also indicate that the stimulation of growth by elevated CO<sub>2</sub> in N limited forest ecosystems will be

considerably less than what has been observed in short term studies unless nutrient supply increases to match carbon supply (Comins 1994, Comins & McMurtrie 1993; Kirschbaum et al. 1994; Pastor and Post 1988).

The close coupling between carbon assimilation and symbiotic N<sub>2</sub> fixation suggests that N<sub>2</sub> fixation should increase as atmospheric CO<sub>2</sub> concentrations rise (Sinclair 1992; Stulen and den Hertog 1993). Indeed, increased N<sub>2</sub> fixation with CO<sub>2</sub> enrichment has been well documented in annual legume species (Phillips et al. 1976; Ryle et al. 1992). This could have important consequences for forested ecosystems if increased N<sub>2</sub> fixation in woody species leads to greater soil N availability and an amelioration of nutrient limitations to CO<sub>2</sub> responses among associated non-fixing species. Results from a number of studies suggest that N<sub>2</sub>-fixing woody plants have the capacity to

increase both C assimilation and total N<sub>2</sub> fixation as atmospheric CO<sub>2</sub> levels rise. Arnone & Gordon (1990) and Thomas et al. (1991) reported greater whole plant N<sub>2</sub> fixation and total plant N accretion in several N<sub>2</sub>-fixing woody species grown at elevated compared to ambient CO<sub>2</sub>. Norby (1987) observed increased whole plant N<sub>2</sub> fixation as well as an increase in the proportion of nodule mass to whole-plant mass in high CO<sub>2</sub> grown *Alnus glutinosa*. Also working with *A. glutinosa*, Vogel & Curtis (1995) found increased leaf N content and no downregulation of photosynthesis in plants grown under elevated CO<sub>2</sub> and low N soil conditions.

In this study we examined the relationship between growth at elevated CO<sub>2</sub> and N accretion (g N organ<sup>-1</sup>) in plant fractions of N<sub>2</sub>-fixing *A. glutinosa* growing on a low N soil. Actinorhizal (*Frankia* nodulated) symbiotic N<sub>2</sub>-fixing species can contribute significant amounts of fixed N<sub>2</sub> to temperate forest ecosystems (Dawson 1983), leading to increased growth of co-occurring tree species (Cote & Camire 1984; Dawson 1986). For example, actinorhizal *A. glutinosa* and *Elaeagnus umbellata* Thunb. increased total and/or available soil N concentrations in mixed plantings of *Juglans nigra* L. compared to control plots containing *J. nigra* only (Friedrich & Dawson 1984; Paschke et al. 1989). Increased N<sub>2</sub> fixation under elevated CO<sub>2</sub>, resulting in greater N levels in above- and belowground tissues, could therefore further increase soil availability of N for use by associated plants. We hypothesized that CO<sub>2</sub> enriched *A. glutinosa* would show increased growth, especially in nodule biomass and leaf area, leading to increased N<sub>2</sub> fixation and a concomitant increase in whole-plant N accretion.

## Materials and methods

In mid-May, 1993, ten open bottom root boxes (0.6 × 0.6 × 0.6 m) were placed into the soil at the University of Michigan Biological Station, in northern Lower Michigan, USA (45°34' N latitude, 84°40' W longitude). The lower half of each box was filled with the Rubicon sand (164 μg N g<sup>-1</sup>) that underlay the site and the upper half was filled with a mixture of 20% locally derived Kalkaska series topsoil and 80% Rubicon sand (202 μg N g<sup>-1</sup>). The soil mixture had a PO<sub>4</sub><sup>-3</sup> concentration of 110 μg PO<sub>4</sub><sup>-3</sup> g<sup>-1</sup> as determined by an acid extraction procedure.

Five, three-month old *Alnus glutinosa* (L.) Gaertn. (black alder) seedlings were transplanted into each of the root boxes on Julian Day (JD) 149 (May 29,

1993). The plants were grown from seed in sterilized peat:perlite:vermiculite potting medium under greenhouse conditions of ambient CO<sub>2</sub>. At the time of transplanting, seedlings had an average leaf area (LA) and stem height of 126 cm<sup>2</sup> and 16.3 cm, respectively, and nodulation was not apparent. Three days after transplanting, seedlings were inoculated with a homogenized suspension of locally collected *Alnus rugosa* (DuRoi) Sprengel nodules (0.075 M KPO<sub>4</sub>, pH 7.0).

Immediately after transplanting, small open top chambers (0.7 × 0.7 × 1.0 m tall, Curtis & Teeri, 1992) were placed over the plants in each of the ten root boxes and CO<sub>2</sub> partial pressure elevated in five of the chambers. The experiment was arranged in a randomized block design with two CO<sub>2</sub> treatments (elevated and ambient) and five blocks (replicates). Carbon dioxide dispensing procedure and monitoring of temperature and CO<sub>2</sub> inside the chambers was as described previously (Vogel & Curtis, 1995). Daytime CO<sub>2</sub> partial pressure in the ambient and elevated CO<sub>2</sub> treatments were 34.9 ± 0.02 Pa and 69.5 ± 0.36 Pa (mean ± SE), respectively. Daytime temperature averaged 2.6 °C higher inside than outside chambers and there was no significant temperature difference between ambient and elevated CO<sub>2</sub> treatments.

Stem height and LA of each plant was measured non-destructively at the time of planting and on 5 other dates during the growing season. Individual leaf areas (ILA) of leaves were calculated using leaf length (*l*, cm) and width (*w*, cm) and the equation ILA = 0.762(*lw*) - 2.52, *r*<sup>2</sup> = 0.99, *n* = 46, derived from the destructive harvest of a separate set of plants. Stem elongation and leaf growth had ceased before JD 293, the final date stem height and LA measurements were taken.

Leaf litter was collected daily as leaves abscised during late October and early November. Complete above- and below-ground harvest took place on JD 308 (November 4). At the time of harvest, 20% (%DW) of the leaves had not naturally abscised and were removed from the stems, dried, weighed, and added to previously collected leaf litter for bulk leaf mass and total N analysis. The rooting volume was excavated by hand and sieved to remove excess soil and the root fraction was stored frozen. Frozen roots and root nodules were washed free of soil and hand-separated. All organ fractions were dried to constant weight at 65 °C immediately after harvesting or washing to determine dry weight. Daily, whole plant net assimilation rate (NAR) was calculated as: NAR = (g whole-plant DW)/(m<sup>2</sup> LA \* *D*) where LA = maximal leaf area (JD 293) and *D* = duration of the experiment in days (160 d).

Seasonal changes in leaf N content ( $\text{g N m}^{-2}$ ) were measured by sampling leaf tissue from the youngest mature leaf (most recent fully expanded leaf, generally the 7th leaf from the apex) six times from mid-summer through leaf abscission. The 7th leaf from the apex continued to be sampled for leaf N after leaf initiation had ceased in autumn. Leaf tissue was frozen on dry ice immediately after sampling and subsequently freeze dried. Freeze dried leaf and oven dried bulk leaf, stem, root and nodule samples were ground to a powder and analyzed for total N and C concentration with an elemental analyzer (Carlo Erba, Milan, Italy).

Delta  $^{15}\text{N}$  of dried bulk leaf tissue from the harvest were compared to that of the soil in which the plants were grown in order to assess the source of nitrogen accumulating in plant tissue. Natural abundance of  $^{15}\text{N}$  for each sample was measured using a SIRA Series II stable isotope ratio mass spectrometer (VG ISOGAS, Middlewich, UK). Following combustion of the samples in an elemental analyzer (Carlo Erba NA 1500, Milan, Italy) the gaseous emissions were passed to the mass spectrometer using helium as the carrier (Thomas et al. 1991). Isotopic composition ( $\delta^{15}\text{N}$ ) was expressed as units of abundance of  $^{15}\text{N}$  per mil ( $\text{‰}$ ), where  $\delta^{15}\text{N} = (R_{\text{sample}}/R_{\text{standard}} - 1) \times 1000$  and  $R = ^{15}\text{N}/^{14}\text{N}$  with atmospheric  $\text{N}_2$  as the standard.


Statistical analyses were performed using analysis of variance (ANOVA) for randomized block design where the unit of replication was a single chamber. Seasonal LA and stem height and final harvest growth, tissue N, and  $\delta^{15}\text{N}$  data were expressed on a per plant basis from pooled chamber data (5 plants chamber $^{-1}$ ). Seasonal leaf N data were collected by sampling an individual plant from each chamber on each date.

## Results

Growth at elevated  $\text{CO}_2$  resulted in significantly greater leaf area plant $^{-1}$  (LA) and greater stem height compared to growth at ambient  $\text{CO}_2$  (Table 1). These differences were evident 52 d after  $\text{CO}_2$  treatment had begun and were sustained throughout the growing season (Figure 1). Treatment effects on final LA were due to an increase in both the total number of leaves plant $^{-1}$  (LN) and the size of individual leaves in elevated compared to ambient grown plants (Table 1). Specific leaf area (SLA) was significantly reduced with  $\text{CO}_2$  enrichment (Table 1).

**Table 1.** Leaf area plant $^{-1}$  (LA), stem height, mean area leaf $^{-1}$  (MAL), number of leaves plant $^{-1}$  (LN), and specific leaf area (SLA) of *A. glutinosa* seedlings grown at elevated (70 Pa) and ambient (35 Pa)  $\text{CO}_2$  on JD 293 (post bud-set). Specific leaf area measurements were from the 7th leaf from the apex. Mean  $\pm$  (SE),  $n = 5$  (5 individuals pooled within chambers). \* =  $P \leq 0.05$ , \*\* =  $P \leq 0.01$ , \*\*\* =  $P \leq 0.001$

	Elevated	Ambient	$\Delta$ %
LA ( $\text{cm}^2$ )	2521 (58.5)	2140 (28.8)	+18 ***
Stem Ht (cm)	111.6 (5.28)	92.9 (2.96)	+20 *
MAL ( $\text{cm}^2$ )	97.8 (1.33)	87.8 (0.58)	+11 ***
LN	25.8 (0.29)	24.4 (0.21)	+ 6 **
SLA ( $\text{m}^2/\text{kg}$ )	13.8 (0.35)	16.6 (0.57)	-17 **



**Figure 1.** Leaf area plant $^{-1}$  and stem height plant $^{-1}$  of *A. glutinosa* seedlings grown at elevated  $\text{CO}_2$  (70 Pa, ●) or ambient  $\text{CO}_2$  (35 Pa, ○). Final measurements were taken after bud set when leaf area was maximal. Carbon dioxide treatments began on JD 149. Vertical bars represent one SE of mean ( $n = 5$  with 5 individuals pooled within chambers).

Total above-ground biomass increased 50% under elevated compared to ambient  $\text{CO}_2$ , with stem and leaf biomass increases of 58% and 36%, respectively (Table 2). Below-ground growth was also stimulated, with significant  $\text{CO}_2$  effects on both total root and nodule biomass. There was no difference between  $\text{CO}_2$  treatments in fine root (<0.5 mm diameter) biomass, or in root:shoot ratio at the final harvest (Table 2). The relatively greater  $\text{CO}_2$  effect on whole plant DW (+54%) compared to total LA (+18%) was reflected in

Table 2. Above and below ground biomass fractions, total biomass (g DW plant<sup>-1</sup>), root: shoot ratio, and daily net assimilation rate (NAR) of *A. glutinosa* seedlings grown at elevated (70 Pa) and ambient (35 Pa) CO<sub>2</sub> for 160 d. Plants were harvested on November 4, 1993 after 80% of the leaves had naturally abscised. NAR based on maximal leaf area measured on JD 293. Mean ± (SE), *n* = 5 (5 individuals pooled within chambers), except for fine roots where *n* = 3. \*\* = *P* ≤ 0.01, \*\*\* = *P* ≤ 0.001, ns = not significant at *P* ≤ 0.05.

	Elevated	Ambient	Δ %
Above ground DW (g)			
Leaf	15.9 (0.35)	11.7 (0.67)	+36 ***
Stem	35.7 (2.68)	22.6 (0.89)	+58 **
Total	51.6 (2.92)	34.3 (1.55)	+50 ***
Below ground DW (g)			
Fine Roots	6.1 (1.27)	4.8 (0.76)	+27 ns
All Roots	57.5 (1.82)	36.4 (1.11)	+58 ***
Nodule	1.6 (0.05)	0.9 (0.56)	+78 ***
Total	59.1 (1.82)	37.3 (1.16)	+58 ***
Whole plant DW (g)	110.7 (4.38)	71.7 (2.10)	+54 ***
Root: shoot	1.15 (0.05)	1.09 (0.04)	+ 5 ns
NAR (g m <sup>-2</sup> d <sup>-1</sup> )	2.74 (0.07)	2.09 (0.04)	+31 ***

the significant increase in daily NAR with CO<sub>2</sub> enrichment (Table 2). The ratio of nodule DW to maximal LA (g nodule DW m<sup>-2</sup> LA) also increased significantly (*P* = 0.01) in elevated (6.24 ± 0.26, mean ± SE) compared to ambient (4.32 ± 0.53) CO<sub>2</sub> grown plants.

Seasonal changes in leaf N content differed between plants in the two CO<sub>2</sub> treatments (Figure 2). Leaves from ambient grown plants had an essentially constant N content between JD 208 and 264 while those from elevated CO<sub>2</sub> plants showed a 50% increase during the same period. Apparent retranslocation of leaf N, calculated as the difference in N content of abscised leaves and leaves with maximal N content (Figure 2) was 30% higher at elevated compared to ambient CO<sub>2</sub>, although this effect was only marginally significant (*P* = 0.15). The *relative* reduction in leaf N content, however, was not different between elevated (-42%) and ambient (-39%) CO<sub>2</sub> grown plants. While maximum leaf N content was significantly greater at high CO<sub>2</sub>, there was no significant CO<sub>2</sub> effect on N content of abscised leaf number 7 or of the bulk leaf fraction (Figure 2).

Nitrogen accretion increased in both above- and belowground tissues with CO<sub>2</sub> enrichment, ranging

from a 24% increase in leaves to a 74% increase in nodules (Table 3). Tissue N concentration (%DW) and C:N ratios were either not significantly changed (stem, roots and nodules) or changed only slightly (leaves). The relatively greater increase in whole-plant biomass (+58%) compared to N accretion (+40%) at elevated CO<sub>2</sub> resulted in a 9% reduction in whole-plant N concentration and a 10% increase in C:N ratio in elevated compared to ambient CO<sub>2</sub> grown plants (Table 3). Leaves from both CO<sub>2</sub> treatments were depleted in <sup>15</sup>N relative to the atmosphere resulting in negative δ<sup>15</sup>N values (Table 3) while a pooled sample of the low N soil in which the plants were grown was enriched in <sup>15</sup>N (δ<sup>15</sup>N<sub>soil</sub> = 2.6 ‰/‰). While we can not determine the proportion of tissue N that was fixed by the plants to that attained from the soil, these data suggest that most of the N accumulated in the leaves of our plants could be attributed to N<sub>2</sub> fixation. In addition, the similar δ<sup>15</sup>N values for ambient and elevated CO<sub>2</sub> grown black alder leaves suggest that similar proportions of N were fixed by plants from each CO<sub>2</sub> treatment.

Table 3. Leaf, stem, root, nodule, and whole plant N accretion, concentration, C:N ratio, and leaf  $\delta^{15}\text{N}$  ( $^{\circ}/_{\infty}$ ) of *A. glutinosa* seedlings grown at elevated (70 Pa) and ambient (35 Pa)  $\text{CO}_2$  for 160 d. Plants were harvested on November 4, 1993 after 80% of the leaves had naturally abscised. Mean  $\pm$  (SE),  $n = 5$  (5 individuals pooled within chambers). \* =  $P \leq 0.05$ , \*\* =  $P \leq 0.01$ , \*\*\* =  $P \leq 0.001$ , ns = not significant at  $P \leq 0.05$ .

	Elevated	Ambient	$\Delta$ %
N accretion ( $\text{g organ}^{-1}$ )			
Leaf	0.301 (0.011)	0.243 (0.015)	+24 *
Stem	0.445 (0.017)	0.315 (0.017)	+41 ***
Root	0.721 (0.049)	0.491 (0.022)	+47 **
Nodule	0.033 (0.003)	0.019 (0.003)	+74 ***
Whole Plant	1.500 (0.069)	1.068 (0.047)	+40 ***
N concentration (% DW)			
Leaf	1.89 (0.042)	2.08 (0.051)	- 9 **
Stem	1.26 (0.067)	1.39 (0.029)	- 9 ns
Root	1.25 (0.051)	1.35 (0.030)	- 7 ns
Nodule	2.10 (0.145)	2.04 (0.100)	+ 3 ns
Whole Plant	1.35 (0.013)	1.49 (0.024)	- 9 ***
C:N ratio			
Leaf	25.9 (0.35)	24.0 (0.57)	+ 8 **
Stem	40.6 (1.93)	36.7 (0.75)	+11 ns
Root	38.2 (1.72)	35.0 (0.78)	+ 9 ns
Nodule	24.5 (1.54)	23.6 (0.71)	+ 4 ns
Whole Plant	36.0 (0.47)	32.8 (0.58)	+10 **
Leaf $\delta^{15}\text{N}$ ( $^{\circ}/_{\infty}$ )	-1.6 (0.05)	-1.46 (0.07)	- 9 ns

## Discussion

Increased growth and whole-plant N accretion in  $\text{CO}_2$  enriched *A. glutinosa* grown in a low N soil was consistent with the positive  $\text{CO}_2$  effects reported for other  $\text{N}_2$ -fixing woody species (Arnone & Gordon 1990; Norby 1987; Thomas et al. 1991). In our study, the significant increase in whole-plant LA, combined with a decline in SLA, lead to a large increase in leaf litter mass at elevated compared to ambient  $\text{CO}_2$ . This increase in litter mass, coupled with a small reduction in litter N concentration, resulted in a 24% increase in total N returned to the soil with  $\text{CO}_2$  enrichment. The significant increase in leaf litter N accretion and only modest increase in litter C:N ratio (+8%) at high  $\text{CO}_2$  suggests there will be little  $\text{CO}_2$  effect on litter decomposition rates (Cotrufo and Ineson 1996; Couteaux et al. 1991), and supports the suggestion of Gifford

(1992) that rising atmospheric  $\text{CO}_2$  may lead to greater symbiotic N inputs to terrestrial ecosystems.

Ours is the first report of leaf N content in naturally abscised leaves of a high  $\text{CO}_2$  grown,  $\text{N}_2$ -fixing tree and only limited data exist for litter N content in other woody species under elevated  $\text{CO}_2$  conditions. Norby et al. (1986b) reported a 23% reduction in N concentration, and a 30% increase in C:N ratio, in elevated  $\text{CO}_2$  grown *Quercus alba* L. leaf litter, while Couteaux et al. (1991) found a 48% decline in *Castanea sativa* Mill litter N concentration and a 85% increase in C:N ratio due to elevated  $\text{CO}_2$ . Diaz et al. (1993) found that on a productive soil foliar N concentrations of herbaceous species declined while soil microbial C and N increased in response to a doubling of ambient  $\text{CO}_2$  concentration. The authors suggested that increased C assimilation at high  $\text{CO}_2$  caused an increase in soil C, thus stimulating growth of microbial populations that




Figure 2. Seasonal leaf N content of the 7th leaf from the apex of *A. glutinosa* seedlings grown at elevated CO<sub>2</sub> (70 Pa, ●) or ambient CO<sub>2</sub> (35 Pa, ○) and whole canopy leaf N content after leaf abscission of plants grown at elevated (■) or ambient CO<sub>2</sub> (□). Final measurement of 7th leaf N content was of recently abscised leaves. All measurements were from leaves that had developed after CO<sub>2</sub> treatment had begun. Vertical bars represent one SE of mean ( $n = 4,5$ ). \* =  $P \leq 0.05$ .

consequently sequestered more soil N. Increased leaf litter C:N ratios at elevated CO<sub>2</sub> could also result in net immobilization of soil N. However, the contrast in the response of litter C:N ratio to elevated CO<sub>2</sub> observed for non-fixing species vs. that of *A. glutinosa* suggests that soil N availability in systems containing N<sub>2</sub> fixers would not decline, but rather would increase owing to the substantially greater litter N content of N<sub>2</sub> fixers at elevated CO<sub>2</sub>.

To date, little work has been done on the retranslocation efficiency of N in leaves grown at elevated CO<sub>2</sub> and any effect that elevated CO<sub>2</sub> may have on this process could influence N return to forested ecosystems. Under ambient CO<sub>2</sub> conditions temperate woody deciduous plants typically resorb 40–80% of their leaf N prior to leaf abscission (Chapin & Kedrowski 1983). Dinitrogen-fixing species generally show somewhat lower net resorption, with 20–50% of leaf N being translocated to perennial tissues (Cote et al. 1989; Vogel & Dawson 1993). The apparent retranslocation of 39% to 42% of leaf N content we observed in *A. glutinosa* is within this range and suggests that elevated CO<sub>2</sub> will have little effect on the dynamics of autumnal leaf N retranslocation.

The negative leaf  $\delta^{15}\text{N}$  in both ambient and elevated CO<sub>2</sub> grown plants from our study compared with the positive soil  $\delta^{15}\text{N}$  indicated a significant proportion of leaf N was symbiotically fixed (Focht 1987; Shearer et al. 1983), as was expected given the low soil N status and the abundant root nodules we observed. There was, however, no CO<sub>2</sub> effect on the proportion of total N derived from N<sub>2</sub> fixation, as leaf  $\delta^{15}\text{N}$  did not differ among treatments. Although we cannot determine the actual proportion of leaf N that was derived from symbiotic N<sub>2</sub> fixation, our data do rule out any greater or lesser reliance on nodule derived N at high CO<sub>2</sub>.

An increase in whole-plant N content of 40% at high CO<sub>2</sub> combined with similar leaf  $\delta^{15}\text{N}$  between CO<sub>2</sub> treatments suggests that high CO<sub>2</sub> grown *A. glutinosa* had an approximately 40% greater whole-plant nitrogenase activity. Increased whole-plant nitrogenase activity has been found in other N<sub>2</sub>-fixing woody plants grown under elevated CO<sub>2</sub> (Arnone & Gordon 1990; Norby 1987; Thomas et al. 1991). Greater whole-plant nitrogenase activity at elevated compared to ambient CO<sub>2</sub> was likely due both to increased nodule biomass and increased specific nitrogenase activity (Vogel and Curtis 1995). Furthermore, higher instantaneous (Vogel & Curtis 1995) and daily NAR under elevated CO<sub>2</sub> resulted in a significant increase in the ratio of nodule biomass to LA consistent with the close coupling of net CO<sub>2</sub> assimilation and nitrogenase activity in *A. glutinosa* (Gordon & Wheeler 1978; Dawson & Gordon 1979).

Ceulemans & Mousseau (1994), in their review of CO<sub>2</sub> responses in 64 woody species, concluded that root:shoot ratio typically increased at high CO<sub>2</sub>, with more pronounced responses under conditions of low N availability. We found that root biomass in elevated CO<sub>2</sub> grown *A. glutinosa* increased to a similar extent as did shoot biomass, resulting in no CO<sub>2</sub> effect on root:shoot ratio. Norby (1987) also observed either no change or a decline in root weight ratio of N<sub>2</sub>-fixing *Robinia pseudoacacia* L., *Elaeagnus angustifolia* L., and *A. glutinosa* in response to elevated CO<sub>2</sub>. The difference in allocational responses to CO<sub>2</sub> among N<sub>2</sub>-fixing and non-fixing plants may be due to the high metabolic cost of N<sub>2</sub> fixation. Root nodules are strong sinks for photosynthate (Tjepkema 1985) and the large increase in nodule biomass, producing comparatively more available N, due to high CO<sub>2</sub> observed in *A. glutinosa* and other N<sub>2</sub>-fixing trees (Norby 1987; Thomas et al. 1991) could account for a significant consumption of photosynthate that might otherwise be available for root growth. Increased N availability due to greater

whole-plant nitrogenase activity at high CO<sub>2</sub> may also have influenced allocation of C to roots. Sellstedt et al. (1986) showed that root:shoot ratio increased, whole plant biomass decreased, and whole plant N content decreased in *Alnus incana* (L.) Moench inoculated with a *Frankia* strain inefficient in N<sub>2</sub> fixation compared to plants inoculated with a more efficient strain.

It is important to note, however, that a lack of a CO<sub>2</sub> effect on root biomass recovered at the final harvest does not remove the possibility of CO<sub>2</sub> effects on root turnover during the course of the experiment. For example, Pregitzer et al. (1995) found no difference between high and low soil N grown *Populus x euramericana* (Dode) Guinier in harvested fine root biomass, but using minirhizotron observations they found significant treatment effects on fine root production and mortality over the course of the growing season. The ability of nodulated *Alnus* species to increase soil N levels and the higher total N content of elevated compared to ambient CO<sub>2</sub> grown alder root tissues suggest that root turnover could play an important role in soil N accretion as atmospheric CO<sub>2</sub> levels rise. Clearly, more intensive observations of root growth dynamics will be necessary before we can determine the magnitude of CO<sub>2</sub> effects on belowground C and N inputs in this system.

Our results suggest that future increases in atmospheric CO<sub>2</sub> will lead to greater whole plant nitrogenase activity in woody N<sub>2</sub>-fixing species and that soil N inputs will increase due to greater leaf litter N content. Given the similarities in litter C:N ratio between elevated and ambient CO<sub>2</sub> treatments, there would be an increase in N mineralization rate that could sustain increased growth of associated non-fixing trees under the predicted higher CO<sub>2</sub> atmosphere.

### Acknowledgements

We would like to thank Mark Kubiske, Tony Sutterly and Richard Spray for construction and maintenance support, and James Teeri for his continued support of global change research. Thanks are also extended to Kurt Pregitzer for C and N analyses and Larry Giles and the Duke University Phytotron mass spectrometer facility for  $\delta^{15}\text{N}$  analyses. This research was supported by funds from the U.S. Department of Energy (National Institute for Global Environmental Change), the University of Michigan Biological Station, and NRI Competitive Grants Program/USDA (92-37100-7535; Plant Response to the Environment).

### References

- Arnore, J. A. & Gordon, J. C. 1990. Effect of nodulation, nitrogen fixation and CO<sub>2</sub> enrichment on the physiology, growth and dry mass allocation of seedlings of *Alnus rubra* Bong. *New Phytologist* 116: 55–66.
- Bazzaz, F. A., Coleman, J. S. & Morse, S. R. 1990. Growth responses of seven major co-occurring tree species of the northeastern United States to elevated CO<sub>2</sub>. *Can. J. Forest Res.* 20: 1479–1484.
- Ceulemans, R. & Mousseau, M. 1994. Effects of elevated atmospheric CO<sub>2</sub> on woody plants. *New Phytol.* 127: 425–446.
- Chapin, F. S. & Kedrowski, R. A. 1983. Seasonal changes in nitrogen and phosphorus fractions and autumn retranslocation in evergreen and deciduous taiga trees. *Ecology* 64: 376–391.
- Comins, H. N. 1994. Equilibrium analysis of integrated plant-soil models for prediction of the nutrient limited growth response to CO<sub>2</sub> enrichment. *J. Theor. Biol.* 71: 369–385.
- Comins, H. N. & McMurtrie, R. E. 1993. Long-term response of nutrient-limited forests to CO<sub>2</sub> enrichment; Equilibrium behavior of plant-soil models. *Ecol. Appl.* 3: 666–681.
- Cote, B. & Camire, C. 1984. Growth, nitrogen accumulation, and symbiotic dinitrogen fixation in pure and mixed plantings of hybrid poplar and black alder. *Plant Soil* 78: 209–220.
- Cote, B. Vogel, C. S. & Dawson, J. O. 1989. Autumnal changes in tissue nitrogen of autumn olive, black alder and eastern cottonwood. *Plant Soil* 118: 23–32.
- Cotrufo, M. F. & Ineson, P. 1996. Elevated CO<sub>2</sub> reduces field decomposition rates of *Betula pendula* (Roth.) leaf litter. *Oecologia* 106: 525–530.
- Couteaux, M. M., Mousseau, M., Celerier, M. L. & Bottner, P. 1991. Increased atmospheric CO<sub>2</sub> and litter quality: decomposition of sweet chestnut leaf litter with animal food webs of different complexities. *Oikos* 61: 54–64.
- Curtis, P. S. & Teeri, J. A. 1992. Seasonal responses of leaf gas exchange to elevated carbon dioxide in *Populus grandidentata*. *Can. J. Forest Res.* 22: 1320–1325.
- Curtis, P. S., Vogel, C. S., Pregitzer, K. S., Zak, D. R. & Teeri, J. A. 1995. Interacting effects of soil fertility and atmospheric CO<sub>2</sub> on leaf area growth and carbon gain physiology in *Populus x euramericana* (Dode) Guinier. *New Phytol.* 129: 253–263.
- Dawson, J. O. 1983. Dinitrogen fixation in forest ecosystems. *Can. J. Forest Res.* 29: 979–992.
- Dawson, J. O. 1986. Actinorhizal plants: Their use in forestry and agriculture. *Outlook Agric.* 15: 202–208.
- Dawson, J. O. & Gordon, J. C. 1979. Nitrogen fixation in relation to photosynthesis in *Alnus glutinosa*. *Bot. Gazette* 140 (suppl): s70–s75.
- Diaz, S., Grime, J. P., Harris, J. & McPherson, E. 1993. Evidence of a feedback mechanism limiting plant response to elevated carbon dioxide. *Nature* 364: 616–617.
- Focht, D. D. 1987. Measurement of biological nitrogen fixation by <sup>15</sup>N techniques. pp. 257–288. In: Elkan, G. H. (ed), *Symbiotic nitrogen fixation technology*. Marcel Dekker, Inc., New York.
- Friedrich, J. M. & Dawson, J. O. 1984. Soil nitrogen concentration and *Juglans nigra* growth in mixed plots with nitrogen-fixing *Alnus*, *Elaeagnus*, *Lespedeza*, and *Robinia* species. *Can. J. Forest Res.* 14: 864–868.
- Gifford, R. M. 1992. Implications of the globally increasing atmospheric CO<sub>2</sub> concentration and temperature for the Australian terrestrial carbon budget: integration using a simple model. *Australian J. Bot.* 40: 527–543.

- Gordon, J. C. & Wheeler, C. T. 1978. Whole-plant studies on photosynthesis and acetylene reduction in *Alnus glutinosa*. *New Phytol.* 80: 179–186.
- Gunderson, C. A. & Wullschleger, S. D. 1994. Photosynthetic acclimation in trees to rising atmospheric CO<sub>2</sub>: A broader perspective. *Photosynthesis Res.* 39: 369–388.
- Kirschbaum, M. U. F., King, D. A., Comins, H. N., McMurtrie, R. E., Medlyn, B. E., Pongracic, S., Murty, D., Keith, H., Raison, R. J., Khanna, P. K. & Sheriff, D. W. 1994. Modelling forest response to increasing CO<sub>2</sub> concentration under nutrient-limited conditions. *Plant Cell Environ.* 17: 1081–1099.
- Norby, R. J. 1987. Nodulation and nitrogenase activity in nitrogen-fixing woody plants stimulated by CO<sub>2</sub> enrichment of the atmosphere. *Physiol. Plantarum* 71: 77–82.
- Norby, R. J., O'Neill, E. G. & Luxmoore, R. J. 1986a. Effects of atmospheric CO<sub>2</sub> enrichment on the growth and mineral nutrition of *Quercus alba* seedlings in nutrient-poor soil. *Plant Physiol.* 82: 83–89.
- Norby, R. J., Pastor, J. & Melillo, J. M. 1986b. Carbon-nitrogen interactions in CO<sub>2</sub>-enriched white oak: physiological and long-term perspectives. *Tree Physiol.* 2: 233–241.
- Paschke, M. W., Dawson, J. O. & David, M. B. 1989. Soil nitrogen mineralization in plantations of *Juglans nigra* interplanted with actinorhizal *Elaeagnus umbellata* or *Alnus glutinosa*. *Plant Soil* 118: 33–42.
- Pastor, J. & Post, W. M. 1988. Response of northern forests to CO<sub>2</sub>-induced climate change. *Nature* 334: 55–58.
- Phillips, D. A., Newell, K. D., Hassell, S. A. & Felling, C. E. 1976. The effect of CO<sub>2</sub> enrichment on root nodule development and symbiotic N<sub>2</sub> reduction in *Pisum sativum* L. *Amer. J. Bot.* 63: 356–362.
- Pregitzer, K. S., Zak, D. R., Curtis, P. S., Kubiske, M. E., Teeri, J. A. & Vogel, C. S. 1995. Atmospheric CO<sub>2</sub>, soil nitrogen, and fine root turnover. *New Phytol.* 129: 579–585.
- Ryle, G. J. A., Powell, C. E. & Davidson, I. A. 1992. Growth of white clover, dependent on N<sub>2</sub> fixation, in elevated CO<sub>2</sub> and temperature. *Ann. Bot.* 70: 213–220.
- Sellstedt, A., Huss-Danell, K. & Ahlqvist A. 1986. Nitrogen fixation and biomass production in symbiosis between *Alnus incana* and *Frankia* strains with different hydrogen metabolism. *Physiol. Plantarum* 66: 99–107.
- Shearer, G., Kohl, D. H., Virginia, R. A., Bryan, B. A., Skeeters, J. L., Nilsen, E. T., Sharifi, M. R. & Rundel P. W. 1983. Estimates of N<sub>2</sub>-fixation from variation in the natural abundance of <sup>15</sup>N in Sonoran Desert ecosystems. *Oecologia* 56: 365–373.
- Sinclair, T. R. 1992. Mineral nutrition and plant growth response to climate change. *J. Exper. Bot.* 43: 1141–1146.
- Stulen, I. & den Hertog, J. 1993. Root growth and functioning under atmospheric CO<sub>2</sub> enrichment. *Vegetatio* 104/105: 99–115.
- Thomas, R. B., Richter, D. D., Ye, H., Heine, P. R. & Strain, B. R. 1991. Nitrogen dynamics and growth of seedlings of an N-fixing tree (*Gliricidia sepium* (Jacq) Walp.) exposed to elevated atmospheric carbon dioxide. *Oecologia* 88: 415–421.
- Tissue, D. T., Thomas, R. B. & Strain, B. R. 1993. Long-term effects of elevated CO<sub>2</sub> and nutrients on photosynthesis and rubisco in loblolly pine seedlings. *Plant, Cell Environ.* 16: 859–865.
- Tjepkema, J. D. 1985. Utilization of photosynthate for nitrogen fixation in seedlings of *Myrica gale* and *Alnus rubra*. Pp. 183–192. In: Ludden, P. W. & Burris, J. E. (eds), *Nitrogen Fixation and CO<sub>2</sub> metabolism*. Elsevier, New York.
- Vogel, C. S. & Curtis, P. S. 1995. Leaf gas exchange and nitrogen dynamics of N<sub>2</sub>-fixing, field grown *Alnus glutinosa* under elevated atmospheric CO<sub>2</sub>. *Global Change Biol.* 1: 55–56.
- Vogel, C. S. & Dawson, J. O. 1993. Changes in tissue nitrogen and phosphorus and foliar free amino acids in autumn olive, black locust, American sycamore, and honey locust during autumn. *Can. J. Forest Res.* 23: 665–672.
- Wong, S. C., Kriedemann, P. E. & Farquhar, G. D. 1992. CO<sub>2</sub> × nitrogen interaction on seedling growth of four species of Eucalypt. *Austr. J. Bot.* 40: 457–472.