

Detection of Mammalian microRNA Expression by In Situ Hybridization With RNA Oligonucleotides

Monika Deo,¹ Jenn-Yah Yu,^{1,2} Kwan-Ho Chung,^{1,2} Melissa Tippens,¹ and David L. Turner^{1-3*}

Erratum accepted 29 December 2006

The original article to which this Erratum refers was published in *Developmental Dynamics* 235:2538–2548

In the original published version of this article, an erroneous concentration was published in the Experimental Procedures section. The original article included “100 mg/ml heparin” to describe the hybridization buffer, but it should correctly read “100 µg/ml heparin.” The corrected sentence is published in full below. The authors regret this error.

“Hybridization buffer: 50% formamide, 0.3 mg/ml yeast RNA (ICN), 100 µg/ml heparin, 1× Denhardt’s solution (0.2% BSA, 0.2% polyvinylpyrrolidone, 0.2% Ficoll 400), 0.1% Tween 20, 0.1% CHAPS, 5 mM EDTA, and 0.3 nmol/ml 12-mer random sequence DNA oligonucleotides.”

¹Molecular and Behavioral Neuroscience Institute, University of Michigan, Ann Arbor, Michigan

²Program in Neuroscience, University of Michigan, Ann Arbor, Michigan

³Department of Biological Chemistry, University of Michigan, Ann Arbor, Michigan

*Correspondence to: David L. Turner, Molecular and Behavioral Neuroscience Institute, University of Michigan, 5062 BSRB, 109 Zina Pitcher Place, Ann Arbor, MI 48109-2200. E-mail: dlturner@umich.edu

DOI 10.1002/dvdy.21079

Published online 26 January 2007 in Wiley InterScience (www.interscience.wiley.com).