

**MECHANISMS OF CORTICOTROPIN-RELEASING FACTOR (CRF) GENE  
REGULATION IN THE FROG *XENOPUS LAEVIS***

**by**

**Meng Yao**

A dissertation submitted in partial fulfillment  
of the requirements for the degree of  
Doctor of Philosophy  
(Molecular Cellular and Developmental Biology)  
in The University of Michigan  
2007

Doctoral Committee:

Professor Robert J. Denver, Chair  
Professor Pamela A. Raymond  
Professor Audrey F. Seasholtz  
Associate Professor Cunming Duan

© Meng Yao  
All rights Reserved  
2007

**To my mother and father.**

## ACKNOWLEDGMENTS

It is my great pleasure to acknowledge the people without whom I could never have completed my dissertation.

First I would like to thank my advisor Bob Denver. He has guided me every step of the way of my graduate study. His broad span of knowledge in the biological field and critical thinking of every scientific question have always impressed me and will continue to inspire me in my future career.

I thank my committee members, Dr. Cunming Duan, Dr. Pamela Raymond, Dr. Audrey Seasholtz, and Dr. Gisela Wilson for their insight and inspiration. I have learned tremendously from their rich experiences in different biological disciplines, and their input has been of great help for the development of my dissertation research.

I have had the great pleasure of working with very kind and talented people in the laboratory. I especially want to thank Nicole Westphal, Graham Boorse, Erica Crespi, and Fang Hu for their contribution to my dissertation project.

I am very grateful for the funding sources that have supported me through my graduate study. I have received funding from the National Science Foundation, Rackham Graduate School, and Department of Molecular Cellular and Developmental Biology.

Professor Pan at the Peking University has also been an immeasurable help to me, and I have always looked up to him during my growth as a person and scientist. I am extremely thankful to my dear friend Yuliang, who has endured me and wholeheartedly supported me through the final stage of my graduate life. I would also like to thank my friend Paul, who has helped me in many ways through the years and made my life enjoyable.

My very special thanks go to whom I owe everything I am today, my mother and father. This work would have been impossible without their unconditional love and support through all these years. I also would like to thank my grandmother who has taught me so much and inspired my interests in biology at a very early age.

## TABLE OF CONTENTS

<b>DEDICATION.....</b>	<b>ii</b>
<b>ACKNOWLEDGMENTS.....</b>	<b>iii</b>
<b>LIST OF TABLES.....</b>	<b>vii</b>
<b>LIST OF FIGURES.....</b>	<b>viii</b>
<b>LIST OF APPENDICES.....</b>	<b>xi</b>
<b>CHAPTER</b>	
<b>1 INTRODUCTION.....</b>	<b>1</b>
Overview.....	1
Introduction to the neuroendocrine stress system.....	2
Physiological regulation of CRF gene expression in the CNS.....	5
Molecular mechanisms of transcriptional regulation of CRF genes.....	13
Organization of the dissertation.....	28
Acknowledgements.....	29
References.....	34
<b>2 DISTRIBUTION AND ACUTE STRESSOR-INDUCED ACTIVATION OF CORTICOTROPIN-RELEASING FACTOR NEURONS IN THE CNS OF <i>XENOPUS LAEVIS</i>.....</b>	<b>54</b>
Abstract.....	54
Introduction.....	56
Materials and Methods.....	59
Results.....	65
Discussion.....	69
Acknowledgements.....	76
References.....	87

**3 STRUCTURAL AND FUNCTIONAL CONSERVATION OF VERTEBRATE CORTICOTROPIN-RELEASING FACTOR GENES: EVIDENCE FOR A CRITICAL ROLE FOR A CONSERVED CYCLIC AMP RESPONSE ELEMENT.....95**

Abstract.....95  
Introduction.....96  
Materials and Methods.....99  
Results.....111  
Discussion.....122  
Acknowledgements.....129  
References.....140

**4 DISTRIBUTION AND CORTICOSTEROID REGULATION OF GLUCOCORTICOID RECEPTOR IN THE CENTRAL NERVOUS SYSTEM OF *XENOPUS LAEVIS*.....146**

Abstract.....146  
Introduction.....147  
Materials and Methods.....152  
Results.....157  
Discussion.....164  
Acknowledgements.....173  
References.....185

**5 DIFFERENTIAL REGULATION OF CORTICOTROPIN-RELEASING FACTOR (CRF) GENE EXPRESSION BY GLUCOCORTICOIDS IN THE BRAIN OF *XENOPUS LAEVIS*.....193**

Abstract.....193  
Introduction.....194  
Materials and Methods.....197  
Results.....204  
Discussion.....210  
Acknowledgements.....218  
References.....230

**6 CONCLUSIONS.....235**

Stress activation of the CRF genes.....236  
Feedback regulation of CRF expression by glucocorticoids.....238  
Conserved regulatory elements in the CRF promoters.....240  
Conserved functions of the limbic system and implications for evolution of the vertebrate brain.....242

Future directions.....244  
Concluding remarks.....249  
References.....251

**APPENDICES.....255**

## LIST OF TABLES

<b>Table 1.1.</b> Repressor element-1/neuron-restrictive silencing element (RE-1/NRSE) sequences found within the introns of CRF genes of human, mouse, rat, sheep, chicken, and <i>Xenopus laevis</i> (CRFa).....	30
<b>Table 2.1.</b> Abbreviations.....	77
<b>Table 3.1.</b> Sequences of oligonucleotide probes used in EMSA. ....	130
<b>Table 4.1.</b> Abbreviations.....	174
<b>Table 5.1.</b> Sequences of primers and probes used in RT-qPCR. ....	219
<b>Table 5.2.</b> Sequences of oligonucleotide probes used in EMSA.....	220

## LIST OF FIGURES

<b>Figure 1.1.</b> Alignment of vertebrate CRF genes.....	31
<b>Figure 1.2.</b> ClustalW alignment of the proximal promoter regions of human (h), chimp (pt), sheep (sh), mouse (m), rat (r), and <i>Xenopus laevis</i> (x; genes a and b) CRF genes.....	32
<b>Figure 1.3.</b> Schematic representation of the major pathways involved in the regulation of CRF gene transcription at the proximal promoter.....	33
<b>Figure 2.1.</b> Photomicrographs of transverse sections through the brain of a juvenile <i>X. laevis</i> at the level of the median eminence showing the specificity of the affinity-purified anti-xCRF IgG. ....	78
<b>Figure 2.2.</b> Schematic coronal illustration of CRF-ir distribution in forebrain regions of juvenile <i>X. laevis</i> .....	79
<b>Figure 2.3.</b> Schematic coronal illustration of CRF-ir distribution in hindbrain and rostral spinal cord regions of juvenile <i>X. laevis</i> .....	80
<b>Figure 2.4.</b> Photomicrographs of transverse sections through the forebrain and part of the midbrain of juvenile <i>X. laevis</i> showing the distribution of CRF-ir positive somata and fibers. ....	81
<b>Figure 2.5.</b> Photomicrographs of transverse sections through part of the midbrain and the hindbrain of juvenile <i>X. laevis</i> showing the distribution of CRF-ir positive somata and fibers. ....	82
<b>Figure 2.6.</b> Plasma corticosterone concentrations of <i>X. laevis</i> juveniles exposed to shaking stress for various times.....	83
<b>Figure 2.7.</b> Effects of shaking stress on CRF-ir in the anterior preoptic area (POA), medial amygdala (MeA), and bed nucleus of the stria terminalis (BNST) of juvenile <i>X. laevis</i> .....	84
<b>Figure 2.8.</b> Changes in Fos-LI in the anterior preoptic area (POA) in control (A) and shaking-stressed (B) juvenile <i>X. laevis</i> .....	85
<b>Figure 2.9.</b> Colocalization of CRF-ir and Fos-LI in the anterior preoptic area (POA).....	86

<b>Figure 3.1.</b> ClustalW alignment of 5' promoter regions of human (h), chimp (pt), sheep (sh), mouse (m), rat (r), and <i>Xenopus</i> CRF (x) genes.....	131
<b>Figure 3.2.</b> Effects of shaking stressor on CRF expression in the anterior preoptic area (POA) of juvenile <i>X. laevis</i> . .....	132
<b>Figure 3.3.</b> Effects of shaking stressor on phosphorylated CREB immunoreactivity (pCREB-ir) in the anterior preoptic area (POA) of juvenile <i>X. laevis</i> .....	133
<b>Figure 3.4.</b> Regulation of the frog CRF promoters by activation of the PKA/CREB pathway in transiently transfected PC-12 cells.....	135
<b>Figure 3.5.</b> Effects of exchanging the CREs present in the xCRFa and xCRFb genes on promoter activity in transiently transfected PC-12 and XLT-15 cells.....	136
<b>Figure 3.6.</b> Binding of CREB to the CREs in the CRF promoters as analyzed by electrophoretic mobility shift assay (EMSA).....	137
<b>Figure 3.7.</b> Association of CREB or acetylated histone H4 with the CRE region of the CRFb proximal promoter as analyzed by chromatin immunoprecipitation (ChIP) assay.....	138
<b>Figure 3.8.</b> Effects of injection of ovine CRF (oCRF) or shaking stressor on the activity of CRF promoter constructs transfected into <i>X. laevis</i> brain.....	139
<b>Figure 4.1.</b> . The affinity purified GR IgG recognizes <i>in vitro</i> synthesized <i>Xenopus laevis</i> GR.....	175
<b>Figure 4.2.</b> The affinity-purified GR IgG recognizes a protein that shows nuclear translocation in response to dexamethasone treatment in A6 cells.....	176
<b>Figure 4.3.</b> Photomicrographs of transverse sections through the anterior preoptic area (POA) of the brain of a juvenile <i>Xenopus laevis</i> showing the specificity of the affinity-purified anti-GR IgG.....	177
<b>Figure 4.4.</b> Schematic coronal illustration of glucocorticoid receptor (GR)-immunoreactivity (ir) distribution in the brain of juvenile <i>Xenopus laevis</i> .....	178
<b>Figure 4.5.</b> Photomicrographs of transverse sections through the forebrain and part of the midbrain of juvenile <i>Xenopus laevis</i> showing the distribution of glucocorticoid receptor (GR)-immunoreactive (ir) cells.....	179
<b>Figure 4.6.</b> Photomicrographs of transverse sections through the midbrain and the hindbrain of juvenile <i>Xenopus laevis</i> showing the distribution of glucocorticoid receptor (GR)-immunoreactive (ir) cells.....	180
<b>Figure 4.7.</b> Plasma corticosterone concentrations of <i>Xenopus laevis</i> juveniles following various treatments.....	181

<b>Figure 4.8.</b> Effects of corticosterone manipulation on glucocorticoid receptor (GR)-immunoreactivity (ir) in the anterior preoptic area (POA) of juvenile <i>Xenopus laevis</i> .....	182
<b>Figure 4.9.</b> Effects of corticosterone manipulation on glucocorticoid receptor (GR)-immunoreactivity (ir) in (A) the medial pallium (mp), (B) the medial amygdala (MeA), and (C) the bed nucleus of stria terminalis (BNST) of juvenile <i>Xenopus laevis</i> .....	183
<b>Figure 4.10.</b> Effects of corticosterone manipulation on glucocorticoid receptor (GR)-immunoreactivity (ir) in the rostral portion of anterior pituitary of juvenile <i>Xenopus laevis</i> .....	184
<b>Figure 5.1.</b> Effects of corticosterone manipulation on CRF immunoreactivity (ir) in the anterior preoptic area (POA) of juvenile <i>Xenopus laevis</i> . ....	221
<b>Figure 5.2.</b> Effects of corticosterone manipulation of CRF immunoreactivity (ir) in the (A) the anterior preoptic area (POA), (B) the medial amygdala (MeA), and (C) the bed nucleus of stria terminalis (BNST) revealed by densitometric analyses.....	222
<b>Figure 5.3.</b> Colocalization of CRF immunoreactivity (ir) and GR immunoreactivity (ir) in (A) the anterior preoptic area (POA), and (B) the medial pallium (mp).....	223
<b>Figure 5.4.</b> Analysis of (A) xCRFb hnRNA, (B) xCRFa hnRNA, and (C) CRF mRNA in the anterior preoptic area (POA)/hypothalamus of juvenile <i>Xenopus laevis</i> by reverse transcriptase quantitative polymerase chain reaction (RT-qPCR).....	224
<b>Figure 5.5.</b> Effects of treatment with glucocorticoids on the activity of the frog CRF promoters in transiently transfected PC-12 cells.....	226
<b>Figure 5.6.</b> Effects of i.c.v. injection of corticosterone (CORT) on the activity of CRFb promoter construct transfected into tadpole brain by electroporation mediated (EM) gene transfer. ....	227
<b>Figure 5.7.</b> Binding of recombinant <i>X. laevis</i> GR (rxGR) to the GRE sites in the CRF promoters analyzed by electrophoretic mobility shift assay (EMSA).....	229

## LIST OF APPENDICES

### Appendix

- A. Oligonucleotides used for reverse transcriptase-PCR detection of primary xCRFa and xCRFb transcripts.....255
- B. Oligonucleotides and fluorescent-labeled probes (Taqman) used for reverse transcriptase quantitative PCR (RTqPCR) for xCRFa hnRNA, xCRFb hnRNA, and xCRF mRNA (recognize transcripts of both a and b genes).....256
- C. Oligonucleotides and fluorescent-labeled probes (Taqman) used in ChIP assays for quantitative PCR analysis of xCRFb core promoter (CRFb CRE) and xCRF coding region (CRF CDS; recognize both a and b genes)..... 257
- D. Time course of fold-increase of pGL3-xCRFb576 and pGL3-xCRFb[CREmut] by forskolin in transiently transfected PC-12 cells.....258
- E. Effects of exchanging the CREs present in the xCRFa and xCRFb genes on the fold-increase of the promoter activity induced by forskolin (Fsk; 25  $\mu$ M) in transiently transfected PC-12 cells.....259
- F. Schematic coronal illustration of mineralocorticoid receptor (MR)-immunoreactivity (ir) distribution in the brain of juvenile *Xenopus laevis*...260
- G. Schematic coronal illustration of urocortin 1 (UCN1)-immunoreactivity (ir) distribution in the brain of juvenile *Xenopus laevis*.....262
- H. Schematic coronal illustration of urocortin 3 (UCN3)-immunoreactivity (ir) distribution in the brain of juvenile *Xenopus laevis*.....265