

EVIDENCE-BASED REVIEW OF TREATMENT OPTIONS FOR PATIENTS WITH GLOTTIC CANCER

Dana M. Hartl, MD, PhD,^{1,2} Alfio Ferlito, MD, DLO, DPath, FRCSEd *ad hominem*, FRCS (Eng, Glasg, Ir) *ad eundem*, FDSRCS *ad eundem*, FHKCORL, FRCPath, FASCP, IFCAP,³ Daniel F. Brasnu, MD,⁴ Johannes A. Langendijk, MD, PhD,⁵ Alessandra Rinaldo, MD, FRCSEd *ad hominem*, FRCS (Eng, Ir) *ad eundem*, FRCSEd Glasg,³ Carl E. Silver, MD,⁶ Gregory T. Wolf, MD⁷

¹Department of Otolaryngology–Head and Neck Surgery, Institut Gustave Roussy, Villejuif, France. E-mail: dana.hartl@igr.fr

²Laboratoire de Phonétique et de Phonologie, Sorbonne Nouvelle, Paris, France

³Department of Surgical Sciences, ENT Clinic, University of Udine, Udine, Italy

⁴Department of Otolaryngology–Head and Neck Surgery, and Voice, Biomaterials and Head and Neck Oncology Research Laboratory, University Paris V, European Hospital Georges Pompidou, Paris, France

⁵Department of Radiation Oncology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

⁶Departments of Surgery and Otolaryngology–Head and Neck Surgery, Albert Einstein College of Medicine, Montefiore Medical Center, Bronx, New York

⁷Department of Otolaryngology–Head and Neck Surgery, University of Michigan, Ann Arbor, Michigan

Accepted 21 May 2010

Published online 6 January 2011 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/hed.21528

Abstract: Evidence-based medicine integrates the best available data in decision making, with the goal of minimizing physicians' and patients' subjectivity. In 2006, the American Society of Clinical Oncology edited clinical practice guidelines for the use of larynx preservation strategies. The objective of this review was to evaluate the current levels of evidence for glottic squamous cell carcinoma. Current guidelines for early stage glottic cancer are based on low-level evidence. Conservation surgery (open or transoral) and radiation therapy are all valid options for T1 and selected T2 lesions. For advanced lesions, surgery and combined chemotherapy and radiation are options. High-level evidence favors combined chemotherapy and radiation therapy or altered fractionation radiation therapy as nonsurgical strategies for organ preservation, compared with radiation therapy alone. The optimal combination of chemotherapy, targeted therapy, and radiation therapy remains to be demonstrated, however, and for high-volume tumors, total laryngectomy may still be warranted. © 2011 Wiley Periodicals, Inc. *Head Neck* 33: 1638–1648, 2011

Keywords: glottic cancer; surgery; transoral laser; radiotherapy; chemotherapy

Evidence-based medicine is the “conscientious, explicit and judicious use of current best evidence in making decisions about the care of individual patients,” and “integrating experience with the best available data in decision making.”¹ The role of clinical research is to allow us to rationalize and refine

our practices for the benefit of our patients: “Experience is good but wisdom is better. On what should we rely for treatment decisions—anecdotes, opinions, and persuasion, or more objective data, systematic reviews, and thoughtful interpretation of available clinical reports? Use of the best information, thus minimizing bias and opinion, is the key element that separates ethical medical practitioners from quacks and charlatans.”²

But why do we need evidence-based medicine? As technology advances, there is an increasing number of therapeutic options at our disposal. As globalization advances, we see the spectrum of the variations in clinical practice expand. Modern medicine includes the patient in therapeutic decision making, with informed consent obtained based on objective information and sound communication with healthcare providers. Finally, as knowledge in decision psychology progresses, it has become increasingly apparent that physicians are not as objective as they think they are when making decisions for patient treatment.¹

Evidence in medicine can be classified as “high-level evidence,” evidence with high statistical power and a low risk of bias, and “low-level evidence” with low statistical power (or none at all) and a high risk of bias. Between the highest and lowest levels, groups of experts throughout the world have developed various classification schemes, to compare different medical studies according to their methodology, statistical power, and risk of bias, generally with 4 or 5 levels.

In 2006, the American Society of Clinical Oncology edited clinical practice guidelines for the use of larynx-

Correspondence to: D. M. Hartl

© 2011 Wiley Periodicals, Inc.

preservation strategies, based on a systematic review of the literature.³ The objective of this review was to evaluate the level of current evidence concerning these treatment options for glottic squamous cell carcinoma. For the purposes of this review, we have chosen a 5-level classification to rate levels of evidence⁴: level I, large randomized trials or meta-analyses of randomized trials; level II, small randomized trials; level III, nonrandomized studies with contemporaneous controls; level IV, nonrandomized studies with historical controls; level V, case studies with no controls.

QUESTIONS

1. What is the evidence in favor of any 1 particular type of treatment for early-stage glottic tumors (T1a)⁵ without anterior commissure (AC) involvement?

High rates of local control and laryngeal function preservation have been shown for patients with early glottic tumors using radiation therapy (RT), transoral laser resection, or open partial laryngeal surgery. The reported rates of local control with conservation laryngeal surgery for T1 glottic carcinoma—transoral laser resection or traditional open partial laryngectomies—range from 85% to 100%,^{6–46} and reported local control with RT alone ranges from 84% to 95% (Table 1).^{47–66} These data are based on nonrandomized studies with historical or contemporaneous controls (levels III and IV). A systematic review in the Cochrane Database of Systematic Reviews found only 1 prospective randomized trial comparing RT and surgery.⁶⁷ That study was fraught with methodological flaws, however, leading the authors of the review to conclude that “there is currently insufficient evidence to guide management decisions on the most effective treatment.” To date, no well-conducted prospective randomized study has been performed to provide level I or level II evidence in favor of any 1 of these different types of approaches. The best level evidence to date, comparing surgery (open or laser) to RT for T1a glottic carcinoma, shows no final local control or survival difference among these different treatments (Table 2),^{46,50,68} although 1 study⁴⁶ (101 patients, level 3 evidence) found a significant improvement in final laryngeal preservation for patients initially treated surgically.

The current evidence regarding exclusive chemotherapy for treating glottic tumors is based on retrospective and prospective nonrandomized studies (level IV) and shows a rate of local control with chemotherapy alone to be in the range of 8% to 12% of all patients initially treated with induction chemotherapy for T1 to T3 glottic tumors.⁶⁹ Thus, in a comparison of treatment modalities, the (low-level) evidence suggests that surgery and RT provide higher initial local control rates than those of exclusive chemotherapy.

The majority of the evidence (level III) shows that RT and transoral laser resection provide comparable

Table 1. Retrospective studies (level IV evidence) of initial local control rates for T1–T2 glottic carcinoma.

T classification	Open cordectomy ^{12–14}	Frontal anterior laryngectomy ^{12,15,16}	Hemi-laryngectomy ^{11,17–23}	Frontal anterior laryngectomy ^{16,24,25}	SCPL ^{26–33}	Laser ^{6–11,34–46}	RT ^{47–66}
T1a	90% to 100%	93% to 96%	91% to 100%	100%	98% to 100%	85% to 100%	84% to 95%
T1a-b and T2 AC	—	—	86% to 98%	—	70% to 95%	—	T1: 43% to 91% T1T2*: 58% to 74%
T2	—	92%	69% to 78%	92% to 94%	88% to 96%	66% to 100%	50% to 85%

Abbreviations: AC, anterior commissure; SCPL, supracricoid partial laryngectomy; RT, radiation therapy.
*Mixed cohorts.

Table 2. Highest-level evidence for treatment options for T1/T1a laryngeal carcinoma.

Reference	No. of patients	Methodology	Group	Outcome	Evidence level
Gourin et al ⁶⁸	89	Retrospective nonrandomized cohort, T1 all laryngeal sites	RT vs surgery	Survival: no difference	III
Jones et al ⁵⁰	364	Retrospective nonrandomized cohort, T1 glottic and supraglottic	RT vs surgery (laser or open resection)	Initial local control: no difference	III
Stoeckli et al ⁴⁶	101	Retrospective nonrandomized cohort, T1 glottic tumors	RT vs laser	Initial local control, survival: no difference Final laryngeal preservation: initial surgery better than RT	III

Abbreviation: RT, radiation therapy.

results in terms of voice outcomes and voice-related quality of life.^{70–75} However, 1 study found a better voice handicap score after laser surgery versus RT,⁷⁶ and 2 other studies found that voice after RT was better than that after laser surgery^{77,78} (all level III studies). Voice results depend on the depth of the laser resection, but level IV evidence shows that voice can return to normal for a number of patients (45% of patients in the study by Vilaseca et al⁷⁹). This evidence reflects the fact that not all tumors classified as T1a are the same. They differ in depth of invasion of the thyroarytenoid muscle. Depth of invasion may constitute a bias in retrospective studies and a major bias when comparing retrospective studies from different centers. Level III evidence shows that the post-operative morbidity is lower with transoral laser resection than that with open conservation surgery, and that, despite comparable oncologic outcomes, transoral resection has for the most part replaced open surgery for tumors classified as T1a not involving the AC.⁸⁰

What criteria then can be used to choose from among these 2 treatment options if there is no strong evidence in favor of 1 treatment over the other in terms of local control and voice-related quality of life? Cost is increasingly being integrated into medical decision making and expert-driven recommendations. Four level III studies (prospective non-randomized) have shown that RT for T1 glottic carcinoma is significantly more costly than transoral laser resection, even though the calculation of cost varies among countries.^{70–72,81} Technical aspects of both treatments are also a consideration. Transoral laser resection may not be possible because of patient morphology or comorbidities, or may not be possible because of the absence of a laser or a surgeon experienced in the technique. Alternatively, the availability and logistics of 6 to 7 weeks of RT and the need for careful cancer follow-up also become decision-making factors. Finally, after a full discussion and complete patient information, physician and patient preferences also may be taken into account.

2. What is the evidence for treatment of T1a or T1b glottic tumors involving the AC?

Cancers involving the AC pose a particular anatomical problem for tumor spread because of the proximity of the thyroid cartilage to the mucosa, the absence of perichondrium or conus elasticus, the absence of anatomic barriers to tumor spread toward the petiole of the epiglottis or to the cricothyroid membrane, and the early ossification of the cartilage at this level, which lowers its resistance to invasion by the tumor. Deep extension at this level may have no effect on vocal fold mobility, leading to a risk of understaging.^{82,83} The small size of the region and the early ossification of the cartilage contribute to the difficulties in radiological staging and diagnosis of cartilage invasion. CT and MRI criteria to optimize the diagnosis of thyroid cartilage invasion are still being assessed, and under-staging or overstaging still occur in 25% to 50% of cases.^{83–86} Finally, adequate exposure of the AC for transoral laser resection is challenging in some patients.⁸⁷

As Bradley et al⁸³ clearly show in their comprehensive review, AC involvement for T1 tumors can mean many things. T1a tumors involving 1 cord up to the ipsilateral half of the AC are different from superficial T1b tumors involving both cords and the AC and from T1b tumors arising at the AC itself, with or without infiltration, ulceration, or spread to the vocal folds. It is also clear from this review that there has been no study directly comparing treatment modalities, such as RT, open surgery, or transoral laser surgery, for comparable tumors involving the AC, taking into account the differences in subtypes of AC involvement, which do not appear in the TNM classification. Scarring of the AC after either surgery or RT generally results in poor voice quality.

Furthermore, the existing literature includes heterogeneous populations of tumors, mixing the results of superficial and infiltrating tumors, and tumors classified T1a, T1b, and T2.⁸³

The bulk of the literature concerning AC lesions aims at determining whether AC involvement is a risk factor for reduced local control. Most of these studies include T1a, T1b, and T2 tumors involving the AC, so that individualizing information regarding T1 tumors with AC involvement is difficult. These

retrospective studies provide conflicting level III and level IV evidence (Tables 1 and 3), but it would seem that at least certain subtypes of tumors involving the AC (infiltrating tumors arising at the AC itself) are of worse prognosis than tumors that do not involve the AC.⁸³ The literature, however, does not provide high-level evidence as to the optimum treatment for these lesions. Rates of local control and laryngeal preservation ranging from 75% to 98% have been reported from retrospective studies of T1 tumors involving the AC using RT,^{47,48,83} open surgery,^{12,88} and transoral laser resection.^{34–39}

A recent retrospective study by Sachse et al⁸⁹ (level III, Table 3) found no difference between open surgery and transoral laser resection for T1 or T2 tumors involving the AC, but showed that local control was significantly decreased if the AC was involved, compared with tumors not involving the AC. Bron et al⁷ and Zohar et al⁹⁰ both found that surgery provided better initial local control than RT. Rucci et al⁹¹ retrospectively compared their surgical and radiotherapy cohorts of tumors with AC involvement and found that surgery as first-line treatment provided significantly better local control (86% vs 74%), but that for “pure AC cancers” RT provided better initial local control, although salvage surgery was less effective after RT. This study illustrates the heterogeneity of the subgroups of tumors involving the AC, the bias inherent in retrospective studies, and the limitations of comparing these studies.

In light of the above-cited evidence (levels III–IV) there is no evidence favoring surgery over RT for T1 tumors involving the AC. For laryngeal tumors in general, Silver et al⁷⁹ recently analyzed the decline in open surgery. As a result of the morbidity of open surgery compared with that of transoral laser resection or RT, these latter techniques have largely supplanted open conservation surgery for early stage laryngeal cancers. For early-stage tumors, open surgery is currently mostly reserved for recurrences.^{80,92} In some cases, however, open surgery should still be considered an option as first-line treatment, in accord with the experience and technical possibilities of the surgeon and the radiation oncologist.⁹³ Other factors that can influence this decision include the ease of operative exposure and sufficient possibilities for close patient follow-up.

3. What treatment should be favored for higher classification glottic tumors (T2) with or without impaired vocal fold mobility?

Comparing retrospective case series with historical controls (level IV evidence), the initial local control for T2 tumors with normal vocal fold mobility treated with RT is comparable to the results of open surgery and transoral laser resection, ranging from 84% to 95% (Table 1).^{8,9,19,40,41,47,48} Level III studies do not provide compelling evidence in favor of surgery versus RT (Table 4).^{33,46,50,68} Level III evidence favors

Table 3. Highest-level evidence for treatment options for T1–T2 glottic carcinoma involving the anterior commissure.

Reference	No. of patients	Methodology	Group	Outcome	Evidence level
Sachse et al ⁸⁹	119	Retrospective nonrandomized cohort	Open surgery vs laser	Local control: no difference	III
Bron et al ⁷	156	Retrospective nonrandomized cohort	Surgery (laser or open) vs RT	Initial local control and final laryngeal preservation: surgery better than RT	III
Zohar et al ⁹⁰	67	Retrospective nonrandomized cohort	Open surgery vs RT	Initial local control: open surgery better than RT	III
Rucci et al ⁹¹	182	Retrospective nonrandomized cohort	Open surgery vs RT	Initial local control: RT better than surgery Final local control (after surgical salvage): initial surgery better than initial RT	III

Abbreviation: RT, radiation therapy.

Reference	No. of patients	Methodology	Group	Outcome	Evidence level
Gourin et al ⁶⁸	98	Retrospective nonrandomized cohort, T2 all laryngeal sites	RT vs surgery	Survival: no difference	III
Jones et al ⁵⁰	124	Retrospective nonrandomized cohort	RT vs surgery (laser or open resection)	Initial local control: no difference	III
Stoeckli et al ⁴⁶	39	Retrospective nonrandomized cohort	RT vs laser	Initial local control and final laryngeal preservation: surgery better than RT	III
Marandas et al ³³	66	Retrospective nonrandomized cohort, T2 with impaired mobility	RT vs open surgery	Initial local control surgery 88%, RT 79% (no statistical analysis)	III

Abbreviation: RT, radiation therapy.

supracricoid partial laryngectomy as opposed to a vertical laryngectomy to optimize local control if open surgery is chosen, however.¹⁹

For the subgroup of T2 tumors with impaired vocal fold mobility, local control with RT alone falls to 76% (at best).^{47,48,94-96} Similarly, Peretti et al^{40,41} found a significant decrease in local control with transoral laser resection for T2 tumors with deep extension involving the paraglottic space or extending to the roof of the ventricle and the false vocal fold (17%), compared with the entire group of tumors classified as T2 (77%). This evidence (level III) emphasizes, yet again, the bias inherent in comparing T2 tumors since not all T2 tumors are similar, and they do not have the same prognosis.

Retrospective studies (level IV) of supracricoid partial laryngectomy for patients with T2 tumors with impaired vocal fold mobility show a high rate of local control (88% to 94%), albeit at the cost of removing three fourths of the larynx and a large portion of the structures not involved by the tumor.^{33,97,98} Thus, for T2 tumors with impaired mobility, level IV evidence seems to favor open conservation surgery over RT or transoral laser resection when local control is the endpoint. However, voice function, the risk of persistent aspiration, and the higher risk in elderly patients are other factors to integrate into decision making.

There is no evidence directly comparing conservation surgery (open or transoral) with chemoradiation or other nonsurgical organ preservation protocols (RT with cetuximab or altered fractionation schedules) for T2 tumors. Outcomes of prospective randomized studies (level I evidence) for stage III and stage IV tumors (including those classified as T2N+) show better results for primary chemoradiation with cisplatin versus RT alone or versus induction chemotherapy followed by RT,⁹⁹ but also for RT plus cetuximab versus RT alone,¹⁰⁰ and for altered-fractionation schedules versus RT alone.¹⁰¹ When conservation surgery is not possible for a T2 tumor, attributed to tumor extensions or lack of local surgical possibilities, high-level evidence would favor chemoradiation, RT plus cetuximab, or altered-fractionation RT over RT alone for organ preservation.

For T2 tumors with impaired vocal cord mobility amenable to conservation surgery, there is only low-level evidence to guide treatment decisions. As stated in the American Society of Clinical Oncology's guidelines, tumor evaluation, patient selection, and "local expertise" are necessary but provide only subjective elements for decision making. The relative morbidity of each approach must be weighed, but has never been directly compared. Open conservation surgery exposes patients to permanent dysphonia and often some degree of dysphagia or aspiration, and requires an experienced surgical team. Nonsurgical organ preservation protocols incorporating chemotherapy

carry hematologic, renal, and/or cutaneous toxicities, and RT exposes patients to late complications such as laryngeal edema and fibrosis, which can lead to dysphagia, dysphonia, and/or dyspnea.^{102,103}

4. Is there evidence favoring a particular organ preservation strategy for higher-classification (T3–T4) tumors arising at the glottic level?

There are no studies directly comparing organ-preservation surgery with nonsurgical organ-preservation protocols for advanced-stage laryngeal tumors in a prospective manner with comparable patient groups. Given the evidence (level IV) showing excellent local control, functional organ preservation, and survival for T3 laryngeal carcinoma using supracricoid partial laryngectomy,^{27,104} this approach remains an option for eligible patients, providing that the surgical expertise is available. Supracricoid partial laryngectomy remains an alternative to total laryngectomy (TL) or nonsurgical organ preservation for selected patients with tumors classified as T3 and T4a.

Survival and local control by TL followed by RT have not been bettered by the use of nonsurgical organ-preservation strategies. In the only prospective randomized study comparing TL and RT with nonsurgical organ preservation, the Department of Veterans Affairs (VA) Laryngeal Cancer Study,¹⁰⁵ disease-specific survival was shown to be the same in the 2 groups (2-year overall survival 68% in both groups, level I evidence). A further prospective randomized trial, the Radiation Therapy Oncology Group (RTOG) 91-11 Study,⁹⁹ comparing 3 arms—concurrent chemoradiation (CRT), induction chemotherapy, and RT (the VA protocol) and RT alone—showed that overall survival did not differ among the 3 groups (2-year survival of 74%, 76%, and 75%, respectively). Disease-free survival was significantly better in the 2 arms with chemotherapy, however (61% and 52% vs 44% for RT alone). Organ preservation was better for the CRT arm (88% vs 75% vs 70% at 2 years). Local control and survival have also been shown to be higher with accelerated and hyperfractionated radiotherapy compared with conventional RT (level I evidence).¹⁰¹ RT combined with cetuximab has been shown to improve survival for patients with advanced head and neck tumors (level I), although this improvement was not significant for the subset of patients with advanced laryngeal cancer.¹⁰⁰ Finally, the addition of docetaxel to induction chemotherapy improves oncological results and organ preservation in advanced-stage tumors compared with cisplatin and fluorouracil alone followed by chemoradiation (level I).^{106,107} A meta-analysis of large randomized trials (level I evidence)^{108,109} has shown improved survival for CRT using cisplatin, compared with sequential treatments or RT alone, but the role of newer therapies (taxanes and targeted therapies) remains to be examined.

Lower-level evidence, however, challenges the notion that nonsurgical organ preservation strategies

provide survival rates comparable to the “gold standard” of total laryngectomy followed by RT (Table 5).^{68,99–101,105–107,109–111} An analysis of the National Cancer Data Base (NCDB) information on laryngeal cancer led Hoffman et al¹¹⁰ to demonstrate that, contrary to most other cancers, 5-year overall survival for laryngeal cancer had decreased significantly from the period of 1985 to 2001, from 68.1% to 64.7%. This decrease paralleled a decline in patients’ socioeconomic status over that period. A subgroup analysis, however, showed that this decrease was significant for T3N0 supraglottic cancer, but not for T4N0 or T3N+ supraglottic tumors, and that the decrease for glottic tumors overall was significant only when comparing 1985–1987 with 1994–1996. Survival after chemoradiation was not significantly different from survival after surgery with RT for T3N0 laryngeal or T3N0 glottic tumors, but was worse after RT alone.^{110,112,113} A smaller database study came to similar conclusions for stage III laryngeal tumors, showing no difference in survival between CRT and TL.¹¹¹ This study, however, showed significantly better survival after TL compared with CRT or RT for stage IV tumors.¹¹¹ This study also showed a significant association between survival and race, socioeconomic status, and access to healthcare. Finally, a recent large retrospective case series demonstrated improved survival for T4 tumors treated surgically compared with CRT or RT.⁶⁸

Furthermore, despite the impeccable methodology and high statistical power of these randomized controlled trials, there still remains a certain risk in applying the results of these studies to all patients with stage III and stage IV tumors.¹¹⁴ A close look at the patients included in the RTOG 91-11 study, for example, reveals that approximately two thirds of the tumors were supraglottic tumors, 40% of the patients had mobile vocal folds, only 10% of the tumors were classified as T4, and the performance status was >80% for approximately 95% of the patients.⁹⁹ This study specifically excluded “large volume” T4 disease, defined as “tumor penetrating through the cartilage or extending more than 1 cm into the base of the tongue.”^{99,112} The clinical guidelines, or “standard of care” determined by the American Society of Clinical Oncology in 2006³ state that “evidence supports the use of larynx-preservation approaches for appropriately selected patients without a compromise in survival; however no larynx-preservation approach offers a survival advantage compared with TL.” The optimization of nonsurgical organ preservation (as with surgical organ preservation strategies) still relies on adequate patient selection. For appropriately selected patients, the options of CRT, induction chemotherapy with taxanes, RT and cetuximab, and altered fractionation RT all provide a high level of organ preservation without compromising survival, compared with RT.

Table 5. Highest-level evidence for treatment options for advanced stage (III–IV) laryngeal carcinoma.

Reference	No. of patients	Methodology	Group	Outcome	Evidence level
Wolf et al ¹⁰⁵	332	Prospective randomized, larynx	TL + RT vs Induction chemo* + RT in responders	Overall survival: no difference; 64% larynx preservation	I
Forastiere et al ⁹⁹	547	Prospective randomized, larynx	CRT vs chemo* + RT in responders vs RT	Overall survival: no difference. Disease-free survival: better if CRT or chemo. Larynx preservation best for CRT (88%)	I
Bonner et al ¹⁰⁰	424	Prospective randomized, all head and neck sites (1/4 larynx)	RT + cetuximab vs RT	Overall survival and locoregional control: RT + cetuximab better	I
Posner et al ¹⁰⁶	501	Prospective randomized, all head and neck sites (18% larynx)	Induction TPF vs chemo* both followed by CRT	Overall survival and locoregional control: TPF better	I
Pointreau et al ¹⁰⁷	213	Prospective randomized, larynx + hypopharynx	Induction TPF vs chemo* both followed by RT	Organ preservation: TPF better (70% vs 58%)	I
Bourhis et al ¹⁰¹	6,515	Meta-analysis prospective randomized trials (head and neck, all sites)	Hyperfractionated or accelerated RT vs RT	Overall survival and locoregional control: altered fractionation better	I
Pignon et al ¹⁰⁹	17,346	Meta-analysis prospective randomized trials (head and neck, all sites)	CRT vs sequential chemotherapy and RT vs RT	Overall survival better if chemotherapy; CRT better than other regimens	I
Hoffman et al ¹¹⁰	158,426	Retrospective database (larynx all sites)	Treatments 1985–2001 (surgery vs RT vs CRT)	Overall survival: —decrease for T3N0 supraglottic but not for T4N0 or T3N+; —no significant decrease for T3N0 glottic and no difference between CRT and TL + RT (66%), but RT alone worse (48%)	III
Chen et al ¹¹¹	7,019	Retrospective database (larynx all sites)	TL vs RT vs CRT or sequential chemotherapy (chemo-RT)	Overall survival: stage III: TL and chemo-RT better than RT; stage IV: TL better than chemo-RT or RT	III
Gourin et al ⁶⁸	451	Retrospective database (larynx all sites)	Surgery vs RT vs CRT	Overall survival: stage III: no difference between surgery and CRT/RT but CRT better than RT; stage IV: surgery better than CRT or RT	III

Abbreviations: TL, total laryngectomy; RT, radiation therapy; Chemo*, chemotherapy with cisplatin and 5-fluorouracil; CRT, concurrent chemoradiation with cisplatin; TPF, chemotherapy with docetaxel, cisplatin, and 5-fluorouracil.

There is currently no evidence directly comparing these different nonsurgical organ-preservation protocols (chemoradiation, RT plus cetuximab, hyperfractionated or accelerated RT, or induction chemotherapy with taxanes) among each other, and only the VA study directly compared their protocol with TL plus RT, still considered as the “gold standard” in terms of local control and survival for locally advanced laryngeal carcinoma. Thus, for nonsurgical organ preservation, in light of these more recent treatment options, the optimal strategy is still unclear, and for “large volume” laryngeal carcinoma with widespread cartilage or tongue base infiltration, the current trend still tends to favor TL followed by RT.^{112,113,115,116}

DISCUSSION

The aim of this review was to show that the levels of evidence for treatment selection remain very low. The American Society of Clinical Oncology published clinical practice guidelines in 2006, with recommendations based on current evidence. These guidelines admit that the recommendations for T1T2 lesions of the glottis are based on “comparison of outcomes from case series/prospective single-arm studies” (level III). The recommendations reflect this lack of high-level evidence. RT, open surgery, or transoral laser resection are all options for patients with tumors classified as T1T2 in these guidelines. The authors do make clear that multimodality therapy is not recommended for patients with these early-stage lesions, not because evidence shows a lack of advantage, but rather, it seems, from a risk-benefit analysis, where the morbidity of multimodality treatment probably outweighs the oncological benefit for these patients whose prognosis remains excellent with single-modality therapy. Although treatment selection may be single modality, all patients benefit from an expert multidisciplinary evaluation in which all treatment options and functional issues are evaluated and discussed.

A major advantage of the guidelines is to emphasize organ-preservation strategies, whether surgical or nonsurgical, and to emphasize the need for a multidisciplinary teamwork approach for each patient. The tracheostoma and loss of a physiological voice after TL decrease quality of life.¹¹⁷⁻¹¹⁹ If similar oncological outcomes, without other functional morbidity, can be obtained with alternatives to TL, they are to be preferred, whether with organ-preservation surgery or nonsurgical organ-preservation strategies.

Future prospective randomized studies with adequate statistical power should aim to answer the questions not addressed in the current literature. It has become apparent that all T1a tumors are not equal, in terms of depth of vocalis muscle invasion. One question is: for T1a tumors with comparable depth of vocal fold invasion, which treatment provides the best voice quality and voice-related quality of life?

It has become evident that not all tumors involving the AC are of worse prognosis, but that some subtypes are and may require more aggressive treatment. For each subgroup of T1 tumors involving the AC (T1a, T1b superficial and bicordal, or infiltrating T1b arising at the AC), which treatment provides the best local control? Which treatment provides the best ultimate laryngeal preservation? Which treatment provides the best voice outcome? From studies comparing different subtypes of tumors classified as T2, it is apparent that not all T2 tumors are the same. For superficial tumors classified as T2, how do different treatment modalities compare? For T2 tumors with impaired mobility amenable to conservation surgery, how does organ-preservation surgery compare with nonsurgical strategies in terms of local control, final organ-preservation, and quality of life? Finally, among the nonsurgical organ-preservation strategies for advanced laryngeal cancer now at our disposal, how do they compare in terms of local control, final organ preservation, and acute and late toxicities and when is a TL still preferable?

Prospective randomized trials with comparable cohorts are necessary to improve our levels of evidence for treatment decision making. Trials are currently under way concerning nonsurgical organ-preservation strategies for patients with advanced laryngeal cancer. These prospective multicenter trials providing a high level of evidence often involve support from the private sector, but even these trials may be fraught with bias in patient selection and evaluation of endpoints,¹¹⁶ arising from the subjective nature of clinical tumor staging/classification (evaluation of tumor extensions and vocal fold mobility). Information from radiological studies is not always pertinent in the initial evaluation or evaluation of response to treatment for laryngeal cancer because of the complexities and subjectivity of the interpretation of laryngeal imaging. Finally, results of complex clinical trials may not be reproducible in the general setting, as a result of epidemiological factors, and patient self-selection for clinical trials that may constitute a bias toward better outcomes, in and of itself.¹²⁰ Because of the heterogeneity observed when comparing past clinical trials, new recommendations¹¹⁵ attempt to redefine inclusion criteria, pertinent stratification variables, and acceptable endpoints for future larynx-preservation clinical trials. Developing clinical trials involving surgery meets with similar difficulties in selecting patients, in eliminating bias in patient selection and surgical expertise in different centers, and in financing studies involving surgery. Academic support for this type of prospective study is necessary. Finally, outcomes in laryngeal cancer are in part related to socioeconomic inequalities, or the “deprivation gap” described by Rachet et al¹²¹ that healthcare systems and governments are currently striving to remedy.

The future holds even more questions. Molecular profiling for treatment selection is still in its infancy for laryngeal cancer, but research in this domain will

give way to new therapeutic paradigms. Robotic surgery, future miniaturization of robotic instruments, image-guided surgery, and other technological advances may improve tumor visualization and resection, and enlarge the indications for transoral minimally invasive surgery. Finally, and in the very near future, better patient selection based on tumor biology, new targeted treatments, and new combined modality therapies will be offering even more options for organ preservation.

CONCLUSIONS

Current treatment guidelines for early-stage glottic cancer are based on low-level evidence. Conservation surgery (open or transoral laser resection) and RT are all still valid options for treating T1 and selected T2 glottic lesions. Subjective selection criteria are still the basis for treatment choice for these early-stage lesions. For advanced lesions not amenable to conservation surgery, high-level evidence favors combined-modality therapy, chemotherapy and RT, or altered fractionation RT, as nonsurgical strategies for organ preservation, compared with RT alone. The optimal combination of chemotherapy, targeted therapy, and RT remains to be demonstrated. Finally, for large volume, infiltrating lesions, TL followed by RT still has its place in our armamentarium.

Acknowledgments. This review was written by members and invitees of the International Head and Neck Scientific Group (www.IHNSG.com).

REFERENCES

1. Elstein AS. On the origins and development of evidence-based medicine and medical decision making. *Inflamm Res* 2004;53 (Suppl 2):S184–S189.
2. Blakley BW, Magit AE. Letter to the Editor. Response to: Hornibrook J. The role of tonsillectomy in reducing recurrent pharyngitis: a systematic review. *Otolaryngol Head Neck Surg* 2009; 141:155–156.
3. Pfister DG, Laurie SA, Weinstein GS, et al. American Society of Clinical Oncology clinical practice guidelines for the use of larynx-preservation strategies in the treatment of laryngeal cancer. *J Clin Oncol* 2006;24:3693–3704.
4. Sackett DL. Rules of evidence and clinical recommendations on the use of antithrombotic agents. *Chest* 1989; 95 (Suppl 2):S2–S4.
5. Sobin LH, Wittekind C. UICC International Union Against Cancer. TNM classification of malignant tumors, 6th edition. New York: Wiley-Liss; 2002. pp 36–42.
6. Hartl DM, Brasnu DF, Fried MP. Conservation surgery for glottic cancer. In: Fried MP, Ferlito A, editors. *The larynx*, 3rd edition. San Diego, CA: Plural Publishing; 2009. pp 515–543.
7. Bron LP, Soldati D, Zouhair A, et al. Treatment of early stage squamous-cell carcinoma of the glottic larynx: endoscopic surgery or cricohyoidoepiglottopexy versus radiotherapy. *Head Neck* 2001;23:823–829.
8. Ambrosch P. The role of laser microsurgery in the treatment of laryngeal cancer. *Curr Opin Otolaryngol Head Neck Surg* 2007;15:82–88.
9. Peretti G, Piazza C, Cocco D, et al. Transoral CO₂ laser treatment for T_{1a}-T₃ glottic cancer: The University of Brescia experience on 595 patients. *Head Neck* 2009; Nov 9 [Epub ahead of print].

10. Motta G, Esposito E, Motta S, Tartaro G, Testa D. CO₂ laser surgery in the treatment of glottic cancer. *Head Neck* 2005;27:566–574.
11. Gallo A, de Vincentiis M, Mancio V, Simonelli M, Fiorella ML, Shah JP. CO₂ laser cordectomy for early-stage glottic carcinoma: a long-term follow-up of 156 cases. *Laryngoscope* 2009;112:370–374.
12. Laccourreye O, Weinstein G, Brasnu D, Trotoux J, Laccourreye H. Vertical partial laryngectomy: a critical analysis of local recurrence. *Ann Otol Rhinol Laryngol* 1991;100:68–71.
13. Neel HB, Devine KS, Desanto LW. Laryngofissure and cordectomy for early cordal carcinoma: outcome in 182 patients. *Otolaryngol Head Neck Surg* 1980;88:79–84.
14. Thomas JV, Olsen KS, Neel HB, DeSanto LW, Suman VJ. Early glottic carcinoma treated with open laryngeal procedures. *Arch Otolaryngol Head Neck Surg* 1994;120:264–268.
15. Brumund KT, Guitierrez-Fonseca R, Garcia D, Babin E, Hans S, Laccourreye O. Frontolateral vertical partial laryngectomy without tracheotomy for invasive squamous cell carcinoma of the true vocal cord: a 25-year experience. *Ann Otol Rhinol Laryngol* 2005;114:314–322.
16. Giovanni A, Guelfucci B, Gras R, Yu P, Zanaret M. Partial frontolateral laryngectomy with epiglottic reconstruction for management of early-stage glottic carcinoma. *Laryngoscope* 2001;111:663–668.
17. Soo KC, Shah JP, Gopinath KS, Gerold FP, Jacques DP, Strong EW. Analysis of prognostic variables and results after supraglottic partial laryngectomy. *Am J Surg* 1988;156:301–305.
18. Leroux-Robert J. A statistical study of 620 laryngeal carcinomas of the glottic region personally operated upon more than five years ago. *Laryngoscope* 1975;85:1440–1444.
19. Laccourreye O, Laccourreye L, Garcia D, Gutierrez-Fonseca R, Brasnu D, Weinstein G. Vertical partial laryngectomy versus supracricoid partial laryngectomy for selected carcinomas of the true vocal cord classified as T2N0. *Ann Otol Rhinol Laryngol* 2000;109:965–971.
20. Bailey BJ, Calcaterra TC. Vertical subtotal laryngectomy and laryngoplasty. Review of experience. *Arch Otolaryngol* 1971;93:232–237.
21. Biller HF, Lawson W. Partial laryngectomy for cancer with marked limitation or fixation of the vocal cord. *Laryngoscope* 1986;96:61–64.
22. Daniilidis J, Nicolaou A, Fountzilias G, Sombolos K. Vertical partial laryngectomy: our results after treating 81 cases of T2 and T3 laryngeal carcinomas. *J Laryngol Otol* 1992;106:349–352.
23. Ogura JH, Sessions DG, Spector GJ. Analysis of surgical therapy for epidermoid carcinoma of the laryngeal glottis. *Laryngoscope* 1975;85:1522–1530.
24. Tucker HM, Benninger MS, Roberts JK, Wood BJ, Levine HL. Near-total laryngectomy with epiglottic reconstruction: long-term results. *Arch Otolaryngol Head Neck Surg* 1989; 115:1341–1344.
25. Mallet Y, Chevalier D, Darras JA, Wiel E, Desaulty A. Near total laryngectomy with epiglottic reconstruction. Our experience of 65 cases. *Eur Arch Otorhinolaryngol* 2001;258:488–491.
26. Brasnu DF. Supracricoid partial laryngectomy with cricohyoidopexy in the management of laryngeal carcinoma. *World J Surg* 2003;27:817–823.
27. Dufour X, Hans S, de Monès E, Brasnu D, Ménard M, Laccourreye O. Local control after supracricoid partial laryngectomy for “advanced” endolaryngeal squamous cell carcinoma classified as T3. *Arch Otolaryngol Head Neck Surg* 2004;130:1092–1099.
28. Laccourreye O, Brasnu D, Merite-Drancy A, et al. Cricohyoidopexy in selected infrahyoid epiglottic carcinomas presenting with pathological preepiglottic space invasion. *Ann Otolaryngol Head Neck Surg* 1993;119:881–886.
29. Alajmo E, Fini-Storchi O, Agostini V, Polli G. Conservation surgery for cancer of the larynx in the elderly. *Laryngoscope* 1985;95:203–205.
30. Maurice N, Crampette L, Mondain M, Guerrier B. Subtotal laryngectomy with cricohyoidopexy: carcinologic results and early functional follow-up. A propos of 49 cases. *Ann Otolaryngol Chir Cervicofac* 1994;111:435–442.
31. Piquet JJ, Chevalier D. Subtotal laryngectomy with cricohyoido-epiglottopexy for the treatment of extended glottic carcinomas. *Am J Surg* 1991;162:357–361.
32. Laccourreye O, Weinstein GS, Brasnu D, et al. A clinical trial of continuous cisplatin-fluorouracil induction chemotherapy and supracricoid partial laryngectomy for glottic carcinoma classified as T2. *Cancer* 1994;74:2781–2790.

33. Marandas P, Hartl DM, Charffedine I, Le Ridant AM, Schwaab G, Luboinski B. T2 laryngeal carcinoma with impaired mobility: subtypes with therapeutic implications. *Eur Arch Otorhinolaryngol* 2002;259:87–90.
34. Peretti G, Nicolai P, Piazza C, Redaelli de Zinis LO, Valentini S, Antonelli AR. Oncological results of endoscopic resections of Tis and T1 glottic carcinomas by carbon dioxide laser. *Ann Otol Rhinol Laryngol* 2001;110:820–826.
35. Hartl DM, de Monès E, Hans S, Janot F, Brasnu D. Treatment of early-stage glottic cancer by transoral laser resection. *Ann Otol Rhinol Laryngol* 2007;116:832–836.
36. Pearson BW, Salassa JR. Transoral laser microresection for cancer of the larynx involving the anterior commissure. *Laryngoscope* 2003;113:1104–1112.
37. Steiner W, Ambrosch P, Rödel RMW, Kron M. Impact of anterior commissure involvement of local control of early glottic carcinoma treated by laser microresection. *Laryngoscope* 2004;114:1485–1491.
38. Davis RK, Hadley K, Smith ME. Endoscopic vertical partial laryngectomy. *Laryngoscope* 2004;114:236–240.
39. Chone CT, Yonehara E, Martins JEF, Altemani A, Crespo AN. Importance of anterior commissure in recurrence of early glottic cancer after laser endoscopic resection. *Arch Otolaryngol Head Neck Surg* 2007;133:882–887.
40. Peretti G, Piazza C, Mensi MC, Magnoni L, Bolzoni A. Endoscopic treatment of cT2 glottic carcinoma: prognostic impact of different pT subcategories. *Ann Otol Rhinol Laryngol* 2005;114:579–586.
41. Peretti G, Piazza C, Bolzoni A, et al. Analysis of recurrences in 322 Tis, T1, or T2 glottic carcinomas treated by carbon dioxide laser. *Ann Otol Rhinol Laryngol* 2004;113:853–858.
42. Sigston E, de Monès E, Babin E, et al. Early-stage glottic cancer. Oncological results and margins in laser cordectomy. *Arch Otolaryngol Head Neck Surg* 2006;132:147–152.
43. Davis RK. Endoscopic surgical management of early glottic carcinoma. *Otolaryngol Clin North Am* 1997;30:79–86.
44. Eckel HE, Thumfart WF. Laser surgery for the treatment of larynx carcinomas: indications, techniques and preliminary results. *Ann Otol Rhinol Laryngol* 1992;101:113–118.
45. Steiner W, Vogt P, Ambrosch P, Kron M. Transoral carbon dioxide laser microsurgery for recurrent glottic carcinoma after radiotherapy. *Head Neck* 2004;26:477–484.
46. Stoeckli SJ, Schnieper I, Huguenin P, Schmid S. Early glottic carcinoma: treatment according to patient's preference? *Head Neck* 2003;25:1051–1056.
47. Mendenhall WM, Amdur RJ, Morris CG, Hinerman RW. T1–T2N0 squamous cell carcinoma of the glottic larynx treated with radiation therapy. *J Clin Oncol* 2001;19:4029–4036.
48. Smee RI, Meagher NS, Williams JR, Broadley K, Bridger GP. Role of radiotherapy in early glottic carcinoma. *Head Neck* 2009;32:850–859.
49. Cellai E, Frata P, Magrini SM, et al. Radical radiotherapy for early glottic cancer: results in a series of 1087 patients from two Italian radiation oncology centers. I. The case of T1N0 disease. *Int J Radiat Oncol Biol Phys* 2005;63:1378–1386.
50. Jones DA, Mendenhall CM, Kirwan J, et al. Radiation therapy for management of T1-T2 glottic cancer at a private practice. *Am J Clin Oncol* 2010; Jan 4. [Epub ahead of print].
51. Le QT, Fu KK, Kröll S, et al. Influence of time and fractionation on local control of T1–T2 glottic carcinoma. *Int J Radiat Oncol Biol Phys* 1997;39:1–2.
52. Johansen LV, Overgaard J, Hjelan-Hansen N, et al. Primary radiotherapy for T1 squamous cell carcinoma of the larynx: analysis of 478 patients treated from 1963 to 1985. *Int J Radiat Oncol Biol Phys* 1990;18:1307–1313.
53. Wang CC, Efid JT. Does prolonged treatment course adversely affect local control of carcinoma of the larynx? *Int J Radiat Oncol Biol Phys* 1994;29:657–660.
54. Mittal B, Rao DV, Marks JE, et al. Role of radiation in the management of early vocal carcinoma. *Int J Radiat Oncol Biol Phys* 1983;9:997–1002.
55. Harwood AR. Cancer of the larynx. The Toronto experience. *J Otolaryngol Suppl* 1982;11:3–21.
56. Fernberg JO, Ringborg U, Silversward C, et al. Radiation therapy in early glottic cancer. Analysis of 177 consecutive cases. *Acta Otolaryngol* 1989;108:478–481.
57. Sjögren EV, Wiggenraad RG, Le Cessie S, Snijder S, Pomp J, de Jong RJ. Outcome of radiotherapy in T1 glottic carcinoma: a population-based study. *Eur Arch Otorhinolaryngol* 2009;266:735–744.
58. Dinshaw KA, Sharma V, Agarwal JP, Ghoshi S, Havaldar R. Radiation therapy in T1–T2 glottic carcinoma: influence of various treatment parameters on local control/complications. *Int J Radiat Oncol Biol Phys* 2000;48:723–735.
59. Groome PA, O'Sullivan B, Mackillop WJ, et al. Compromised local control due to treatment interruptions and late treatment breaks in early glottic cancer: population-based outcomes study supporting need for intensified treatment schedules. *Int J Radiat Oncol Biol Phys* 2006;64:1002–1012.
60. Jin J, Liao ZM, Gao L, Huang XD, Xu GZ. Analysis of prognostic factors for T1N0M0 glottic cancer treated with definitive radiotherapy alone: experience of the cancer hospital of Peking Union Medical College and the Chinese Academy of Medical Sciences. *Int J Radiat Oncol Biol Phys* 2002;54:471–478.
61. Howell-Burke D, Peters LJ, Goepfert H, et al. T2 glottic carcinoma: recurrence, salvage, and survival after definitive radiotherapy. *Arch Otolaryngol Head Neck Surg* 1990;116:830–835.
62. Karim ABMF, Kralendonk JH, Yap LY, et al. Heterogeneity of stage II glottic carcinoma and its therapeutic implications. *Int J Radiat Oncol Biol Phys* 1987;13:313–317.
63. Kaplan MJ, Johns ME, McLean WC, et al. Stage II glottic carcinoma: prognostic factors and management. *Laryngoscope* 1983;93:725–728.
64. Mantravadi RVP, Liebner EJ, Haas RE, Skolnik EM, Appelbaum EL. Cancer of the larynx: prognostic factors in radiation therapy. *Radiology* 1983;149:311–314.
65. Harwood AR, DeBoer G. Prognostic factors in T2 glottic cancer. *Cancer* 1980;45:991–995.
66. Hendrickson FR. Radiation therapy treatment of larynx cancers. *Cancer* 1985;55:2058–2061.
67. Dey P, Arnold D, Wight R, McKenzie K, Kelley CG, Wilson J. Radiotherapy versus open surgery versus endolaryngeal surgery (with or without laser) for early laryngeal squamous cell cancer. *Cochrane Database Syst Rev* 2002;2:CD002027.
68. Gourin CG, Conger BT, Sheils C, Bilodeau PA, Coleman TA, Porubsky ES. The effect of treatment on survival in patients with advanced laryngeal carcinoma. *Laryngoscope* 2009;119:1312–1317.
69. Hartl DM, Brasnu DF. Chemotherapy alone for glottic carcinoma: a need for higher-level evidence. *Ann Otol Rhinol Laryngol* 2009;118:543–545.
70. Smith JC, Johnson JT, Cognetti DM, et al. Quality of life, functional outcome, and costs of early glottic cancer. *Laryngoscope* 2003;113:68–76.
71. Goor KM, Peeters AJ, Mahieu HF, et al. Cordectomy by CO₂ laser or radiotherapy for small T1a glottic carcinomas: costs, local control, survival, quality of life, and voice quality. *Head Neck* 2007;29:128–136.
72. Cragle SP, Brandenburg JH. Laser cordectomy or radiotherapy: cure rates, communication, and cost. *Otolaryngol Head Neck Surg* 1993;108:648–654.
73. Sjögren EV, van Rossum MA, Langeveld TMP, et al. Voice outcome in T1a midcord glottic carcinoma. Laser surgery vs radiotherapy. *Arch Otolaryngol Head Neck Surg* 2008;134:965–972.
74. Cohen SM, Garrett CG, Dupont WD, Ossoff RH, Courey MS. Voice-related quality of life in T1 glottic cancer: irradiation versus endoscopic excision. *Ann Otol Rhinol Laryngol* 2006;115:581–586.
75. Wedman J, Heimdal JH, Elstad I, Olofsson J. Voice results in patients with T1a glottic cancer treated by radiotherapy or endoscopic measures. *Eur Arch Otorhinolaryngol* 2002;259:547–550.
76. Peeters AJ, van Gogh CD, Goor KM, Verdonck de Leeuw IM, Langendijk JA, Mahieu HF. Health status and voice outcome after treatment for T1a glottic carcinoma. *Eur Arch Otorhinolaryngol* 2004;261:534–540.
77. Krenqli M, Policarpo M, Manfreda I, et al. Voice quality after treatment for T1a glottic carcinoma-radiotherapy versus laser cordectomy. *Acta Oncol* 2004;43:284–289.
78. Rydell R, Schalen L, Fex S, Elnor A. Voice evaluation before and after laser excision vs. radiotherapy of T1a glottic carcinoma. *Acta Otolaryngol* 1995;115:560–565.
79. Vilaseca I, Huerta P, Blanch JL, Fernández-Planas AM, Jiménez C, Bernal-Sprekelsen M. Voice quality after CO₂ laser cordectomy—what can we really expect? *Head Neck* 2008;30:43–49.
80. Silver CE, Beitler JJ, Shaha AR, Rinaldo A, Ferlito A. Current trends in initial management of laryngeal cancer: the declining use of open surgery. *Eur Arch Otorhinolaryngol* 2009;266:1333–1352.
81. Brandenburg JH. Laser cordectomy versus radiotherapy: an objective cost analysis. *Ann Otol Rhinol Laryngol* 2001;110:312–318.

82. Nakayama M, Brandeburg JH. Clinical underestimation of laryngeal cancer: Predictive indicators. *Arch Otolaryngol Head Neck Surg* 1993;119:950–957.
83. Bradley PJ, Rinaldo A, Suárez C, et al. Primary treatment of the anterior vocal commissure squamous carcinoma. *Eur Arch Otorhinolaryngol* 2006;263:879–888.
84. Fernandes R, Gopalan P, Spyridakou C, Joseph G, Kumar M. Predictive indicators for thyroid cartilage involvement in carcinoma of the larynx seen on spiral computed tomography scans. *J Laryngol Otol* 2006;120:857–860.
85. Becker M. Neoplastic invasion of laryngeal cartilage: radiologic diagnosis and therapeutic implications. *Eur J Radiol* 2000;33:216–229.
86. Becker M, Zbären P, Casselman JW, Kohler R, Dulguerov P, Becker CD. Neoplastic invasion of laryngeal cartilage: reassessment of criteria for diagnosis at MR imaging. *Radiology* 2008;249:551–559.
87. Steiner W, Ambrosch P. Endoscopic laser surgery of the upper aerodigestive tract. New York: Thieme; 2000.
88. Laccourreye O, Muscatello L, Laccourreye L, Naudo P, Brasnu D, Weinstein G. Supracricoid partial laryngectomy with cricohyoidoepiglottopexy for “early” glottic carcinoma classified as T1-T2N0 invading the anterior commissure. *Am J Otolaryngol* 1997;18:385–390.
89. Sachse F, Stool W, Rudack C. Evaluation of treatment results with regard to initial anterior commissure involvement in early glottic carcinoma treated by external partial surgery or transoral laser microresection. *Head Neck* 2009;31:531–537.
90. Zohar Y, Rahima M, Shvili Y, Talmi YP, Lurie H. The controversial treatment of anterior commissure carcinoma of the larynx. *Laryngoscope* 1992;102:69–72.
91. Rucci L, Gallo O, Fini-Storchi O. Glottic cancer involving anterior commissure: surgery vs radiotherapy. *Head Neck* 1991;13:403–410.
92. Mendenhall WM, Werning JW, Hinerman RW, Amdur R, Vilaret DB. Management of T1–T2 glottic carcinomas. *Cancer* 2004;100:1786–1792.
93. Ferlito A, Silver CE, Howard DJ, Laccourreye O, Rinaldo A, Owen R. The role of partial laryngeal resection in current management of laryngeal cancer: a collective review. *Acta Otolaryngol* 2000;120:456–465.
94. Frata P, Cellai E, Magrini SM, et al. Radical radiotherapy for early glottic cancer: results in a series of 1087 patients from two Italian radiation oncology centers. II. The case of T2N0 disease. *Int J Radiat Oncol Biol Phys* 2005;63:1387–1394.
95. Burke LS, Greven KM, McGuirt WT, Case D, Hoen HM, Raben M. Definitive radiotherapy for early glottic carcinoma: prognostic factors and implications for treatment. *Int J Radiat Oncol Biol Phys* 1997;38:1001–1006.
96. Fein DA, Mendenhall WM, Parsons JT, Million RR. T1–T2 squamous cell carcinoma of the glottic larynx treated with radiotherapy: a multivariate analysis of variables potentially influencing local control. *Int J Radiat Oncol Biol Phys* 1993;25:605–611.
97. Chevalier D, Laccourreye O, Brasnu D, Laccourreye H, Piquet JJ. Cricohyoidoepiglottopexy for glottic carcinoma with fixation or impaired motion of the true vocal cord: 5-year oncologic results with 112 patients. *Ann Otol Rhinol Laryngol* 1997;106:364–369.
98. Laccourreye O, Diaz EM Jr, Bassot V, Muscatello L, Garcia D, Brasnu D. A multimodal strategy for the treatment of patients with T2 invasive squamous cell carcinoma of the glottis. *Cancer* 1999;85:40–46.
99. Forastiere AA, Goepfert H, Maor M, et al. Concurrent chemotherapy and radiotherapy for organ preservation in advanced laryngeal cancer. *N Engl J Med* 2003;349:2091–2098.
100. Bonner JA, Harari PM, Giralt J, et al. Radiotherapy plus cetuximab for locoregionally advanced head and neck cancer: 5-year survival data from a phase 3 randomised trial, and relation between cetuximab-induced rash and survival. *Lancet Oncol* 2010;11:21–28.
101. Bourhis J, Overgaard J, Audry H, et al. Hyperfractionated or accelerated radiotherapy in head and neck cancer: a meta-analysis. *Lancet* 2006;368:843–854.
102. Matchay M, Moughan J, Trotti A, et al. Factors associated with severe late toxicity after concurrent chemoradiation for locally advanced head and neck cancer: an RTOG analysis. *J Clin Oncol* 2008;26: 3582–3589.
103. Givens DJ, Karnell LH, Gupta AK, et al. Adverse events associated with concurrent chemoradiation therapy in patients with head and neck cancer. *Arch Otolaryngol Head Neck Surg* 2009;135:1209–1217.
104. Laccourreye O, Salzer SJ, Brasnu D, Shen W, Laccourreye H, Weinstein GS. Glottic carcinoma with a fixed true vocal cord: outcomes after neoadjuvant chemotherapy and supracricoid partial laryngectomy with cricohyoidoepiglottopexy. *Otolaryngol Head Neck Surg* 1996;114:400–406.
105. Wolf GT, Hong WK, Gross Fisher S, et al. Induction chemotherapy plus radiation compared with surgery plus radiation in patients with advanced laryngeal cancer (The Department of Veterans Affairs [VA] Laryngeal Cancer Study Group). *N Engl J Med* 1991;324:1685–1690.
106. Posner MR, Hershock DM, Blajman CR, et al. Cisplatin and fluorouracil alone or with docetaxel in head and neck cancer. *N Engl J Med* 2007;357:1705–1715.
107. Pointreau Y, Garaud P, Chapet S, et al. Randomized trial of induction chemotherapy with cisplatin and 5-fluorouracil with or without docetaxel for larynx preservation. *J Natl Cancer Inst* 2009;101:498–506.
108. Pignon JP, Bourhis J, Domenge C, Designé L. Chemotherapy added to locoregional treatment for head and neck squamous-cell carcinoma: three meta-analyses of updated individual data. MACH-NC Collaborative Group. Meta-Analysis of Chemotherapy on Head and Neck Cancer. *Lancet* 2000;255:949–955.
109. Pignon JP, le Maître A, Maillard E, Bourhis J, for the MACH-NC Collaborative Group. Meta-analysis of chemotherapy in head and neck cancer (MACH-NC): an update on 93 randomised trials and 17,346 patients. *Radiation Oncol* 2009;92:4–14.
110. Hoffman HT, Porter K, Karnell LH, et al. Laryngeal cancer in the United States: changes in demographics, patterns of care, and survival. *Laryngoscope* 2006;116(Suppl 111):1–13.
111. Chen AY, Halpern M. Factors predictive of survival in advanced laryngeal cancer. *Arch Otolaryngol Head Neck Surg* 2007;133:1270–1276.
112. Forastiere AA. Larynx preservation and survival trends: should there be concern? *Head Neck* 2010;32:14–17.
113. Wolf GT. Reexamining the treatment of advanced laryngeal cancer: the VA laryngeal cancer study revisited. *Head Neck* 2010;32:7–14.
114. Olsen KD. Reexamining the treatment of advanced laryngeal cancer. *Head Neck* 2010;32:1–7.
115. Lefebvre JL, Ang KK, for the Larynx Preservation Consensus Panel. Larynx preservation clinical trial design: key issues and recommendations—a consensus panel summary. *Head Neck* 2009;31:429–441.
116. Forastiere AA, Trotti AM. Searching for less toxic larynx preservation: a need for common definitions and metrics. *J Natl Cancer Inst* 2009;101:129–131.
117. DeSanto LW, Olsen KD, Perry WC, Rohe DE, Keith RL. Quality of life after surgical treatment of cancer of the larynx. *Ann Otol Rhinol Laryngol* 1995;104: 763–769.
118. Weinstein GS, El-Sawy MM, Ruiz C, et al. Laryngeal preservation with supracricoid partial laryngectomy results in improved quality of life when compared with total laryngectomy. *Laryngoscope* 2001;111:191–199.
119. Fung K, Lyden TH, Lee J, et al. Voice and swallowing outcomes of an organ-preservation trial for advanced laryngeal cancer. *Int J Radiat Oncol Biol Phys* 2005;63:1395–1399.
120. Clark AL, Lammiman MJ, Goode K, Cleland JG. Is taking part in clinical trials good for your health? A cohort study. *Eur J Heart Fail* 2009;11:1078–1083.
121. Rachet B, Quinn MJ, Cooper N, Coleman MP. Survival from cancer of the larynx in England and Wales up to 2001. *Br J Cancer* 2008;99:S35–S37.