

Mode Competition in Relativistic Magnetrons and Injection Locking in KW Magnetrons*

V.B. Neculaes, R.M. Gilgenbach, M. Lopez, Y.Y. Lau, M. Jones, W. White, P. Pengvanich, M.D. Johnston, T. Strickler, T.A. Spencer¹, J. Luginsland², M. Haworth¹, K. Cartwright¹, P. Mardahl, ¹ T. Murphy, ¹ D. Price³

*Intense Energy Beam Interaction Laboratory
Nuclear Engineering and Radiological Sciences Dept.
University of Michigan, Ann Arbor, MI 48109 USA*

¹ *Air Force Research Lab, Kirtland AFB, NM*

² *SAIC*

³ *Titan Corp.*

Abstract. Both relativistic and nonrelativistic magnetrons are under experimental and theoretical investigation at U of M. Relativistic (Titan-6-vane) magnetron experiments (300-400 kV, 1-10 kA, 0.5 microsecond) investigate mode control with various output coupling geometries. Mode competition between the pi mode and the 2/3 pi mode has been characterized for two-versus-three output extractors for comparison with particle in cell simulations. Phase measurements and time-frequency-analysis are performed for mode identification. Peak microwave output power on the order 0.5 GW has been measured, assuming equal output from 3 waveguides.

Nonrelativistic (4 kV, <1A, kW microwave power) magnetron experiments are performed on commercial oven magnetrons for an in-depth investigation of crossed-field injection-locking and noise. Injection-locking is demonstrated by utilizing an oven magnetron as a reflection amplifier. Noise generation is explored as a function of injected signal and cathode conditions.

*Research supported by the AFOSR, AFRL, and the DUST (S&T) under the Innovative Microwave Vacuum Electronics MURI Program Managed by the Air Force Office of Scientific Research, and Northrop Grumman Corp.