

Supplementary online data


Bimolecular fluorescence complementation analysis of eukaryotic fusion products

Ho-Pi Lin*, Claudius Vincenz†, Kevin W. Eliceiri‡, Tom K. Kerppola† and Brenda M. Ogle*†§¹

*Department of Biomedical Engineering, University of Wisconsin-Madison, Madison, WI 53706, U.S.A., †Howard Hughes Medical Institute and Department of Biological Chemistry, University of Michigan-Ann Arbor, Ann Arbor, MI 48109, U.S.A., ‡Laboratory for Optical and Computational Instrumentation, University of Wisconsin-Madison, WI 53706, U.S.A., and §The Material Science Program, University of Wisconsin-Madison, Madison, WI 53706, U.S.A.

Figure S1 | Growth rate of COS-1 cells seeded at high density

COS-1 cells used for fusion experiments were seeded at high-density (~95%) confluency. At this confluency, the cells are in plateau phase and so undergo minimal net growth during the period of time typically monitored for cell fusion (i.e. 1–4 days after seeding). Shown here is a typical growth curve (from three separate experiments) for COS-1 cells at the confluency used for cell fusion experiments, although BiFC transduction and fusion were not induced in this case. Arrow indicates typical time point for induction of fusion (d1) after mixing of BiFC transfected populations (d0).



Received 16 March 2010/18 June 2010; accepted 1 July 2010

Published as Immediate Publication 1 July 2010, doi:10.1042/BC20100033

¹To whom correspondence should be addressed (email ogle@wisc.edu).