

Table S2. Estimated Cumulative Percent and 95% Confidence Interval of 1320 Women with at least 1, 2, or 3 Occurrences of Potentially Abnormal Uterine Bleeding³⁻⁶, The Study of Women’s Health Across the Nation (SWAN) Menstrual Calendar Substudy

Bleeding Characteristic	At Least 1 % (95% CI)^a	At Least 2 % (95% CI)^a	At Least 3 % (95% CI)^a
Menses Duration >8 days^b			
All Stages ^c	96.0 (90.1, 99.9)	88.2 (66.5, 98.5)	87.1 (76.3, 94.7)
During Early Transition ^d	89.9 (83.8, 94.4)	85.2 (78.7, 92.9)	73.9 (58.9, 86.9)
During Late Transition ^e	90.4 (78.8, 97.1)	88.4 (63.9, 99.0)	86.3 (69.1, 98.6)
Menses Duration ≥10 days^{f,g}			
All Stages ^c	91.0 (83.0, 96.2)	87.5 (64.9, 98.3)	77.7 (56.7, 93.2)
During Early Transition ^d	84.4 (76.2, 91.0)	69.8 (57.1, 81.7)	51.6 (39.4, 65.2)
During Late Transition ^e	92.3 (76.2, 99.1)	73.1 (51.8, 90.6)	54.6 (33.9, 77.9)
Menses Duration ≥15 days^h			
All Stages ^c	58.5 (52.1, 65.0)	40.3 (30.0, 52.6)	29.8 (17.4, 48.1)
During Early Transition ^d	57.9 (44.7, 71.8)	35.7 (24.3, 50.4)	23.5 (14.1, 37.6)
During Late Transition ^e	50.2 (31.6, 50.2)	41.6 (22.2, 68.3)	22.8 (10.9, 44.1)
Menses includes ≥ 6 days of spotting^g			
All Stages ^c	90.4 (83.0, 95.5)	87.0 (63.9, 98.3)	66.8 (55.2, 78.0)
During Early Transition ^d	87.3 (78.4, 93.8)	69.0 (58.1, 79.2)	51.2 (40.7, 61.2)
During Late Transition ^e	90.5 (71.7, 98.7)	89.0 (64.3, 99.1)	75.2 (49.6, 94.2)
Menses includes ≥3 heavy days of bleeding^g			
All Stages ^c	70.6 (56.6, 83.3)	44.6(37.6, 52.2)	34.5 (30.2, 39.2)
During Early Transition ^d	60.1 (48.1, 72.3)	53.7 (38.7, 70.3)	30.7 (25.3, 40.1)
During Late Transition ^e	54.5 (39.5, 70.8)	50.7 (26.6, 80.0)	34.7 (16.5, 63.7)

^a Cumulative percent and 95% confidence interval calculated by Kaplan-Meier Product Limit to adjust for right-censoring

^b Prolonged menses defined by FIGO^{3,6}

^c 1320 women were observed during the study

^d 963 had the start of the early transition stage observed

^e 815 women had the start of the late transition observed.

^f Prolonged menses defined by Belsey 1997⁷

^g Cutoff defined by approximate upper 5% of menstrual cycles in SWAN menstrual calendar substudy

^h Prolonged menses defined by WHO⁸