



RESEARCH ARTICLE

10.1002/2015JD024631

Key Points:

- CH₄ emissions from the Bakken region of North Dakota quantified
- First emission estimate using in situ CH₄ measurements
- CH₄ sources dominated by oil- and gas-related activities

Correspondence to:

J. Peischl,
jeff.peischl@noaa.gov

Citation:

Peischl, J., et al. (2016), Quantifying atmospheric methane emissions from oil and natural gas production in the Bakken shale region of North Dakota, *J. Geophys. Res. Atmos.*, 121, 6101–6111, doi:10.1002/2015JD024631.

Received 9 DEC 2015

Accepted 22 APR 2016

Accepted article online 11 MAY 2016

Published online 25 MAY 2016

Quantifying atmospheric methane emissions from oil and natural gas production in the Bakken shale region of North Dakota

J. Peischl^{1,2}, A. Karion^{1,3,4}, C. Sweeney^{1,3}, E. A. Kort⁵, M. L. Smith⁵, A. R. Brandt⁶, T. Yeskoo⁷, K. C. Aikin^{1,2}, S. A. Conley⁸, A. Gvakharia⁵, M. Trainer², S. Wolter^{1,3}, and T. B. Ryerson²

¹Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, Colorado, USA, ²Chemical Sciences Division, NOAA Earth System Research Laboratory, Boulder, Colorado, USA, ³Global Monitoring Division, NOAA Earth System Research Laboratory, Boulder, Colorado, USA, ⁴Now at the National Institute of Standards and Technology, Gaithersburg, Maryland, USA, ⁵Department of Climate and Space Sciences and Engineering, University of Michigan, Ann Arbor, Michigan, USA, ⁶Department of Energy Resources Engineering, Stanford University, Stanford, California, USA, ⁷Department of Civil and Environmental Engineering, Stanford University, Stanford, California, USA, ⁸Department of Land, Air, and Water Resources, University of California, Davis, California, USA

Abstract We present in situ airborne measurements of methane (CH₄) and ethane (C₂H₆) taken aboard a NOAA DHC-6 Twin Otter research aircraft in May 2014 over the Williston Basin in northwestern North Dakota, a region of rapidly growing oil and natural gas production. The Williston Basin is best known for the Bakken shale formation, from which a significant increase in oil and gas extraction has occurred since 2009. We derive a CH₄ emission rate from this region using airborne data by calculating the CH₄ enhancement flux through the planetary boundary layer downwind of the region. We calculate CH₄ emissions of (36 ± 13), (27 ± 13), (27 ± 12), (27 ± 12), and (25 ± 10) × 10³ kg/h from five transects on 3 days in May 2014 downwind of the Bakken shale region of North Dakota. The average emission, (28 ± 5) × 10³ kg/h, extrapolates to 0.25 ± 0.05 Tg/yr, which is significantly lower than a previous estimate of CH₄ emissions from northwestern North Dakota and southeastern Saskatchewan using satellite remote sensing data. We attribute the majority of CH₄ emissions in the region to oil and gas operations in the Bakken based on the similarity between atmospheric C₂H₆ to CH₄ enhancement ratios and the composition of raw natural gas withdrawn from the region.

1. Introduction

From 1981 to 2003, North Dakota accounted for less than 1.5% of United States (U.S.) crude oil production (U.S. Energy Information Administration (EIA), www.eia.gov). However, the onset of directional drilling and hydraulic fracturing has led to a large increase in oil and natural gas extraction from shale formations in the U.S. in recent years. One such shale formation, the Bakken, is located in the Williston Basin of northwestern North Dakota. As a result, North Dakota accounted for 12.5% of the U.S. crude oil production in May 2014 (EIA), due mainly to oil extraction from the Bakken shale formation (Figure 1), surpassing the oil production from all U.S. regions except Texas and federal Gulf of Mexico (EIA).

In addition to crude oil, natural gas is also extracted from the Bakken shale formation. The region was responsible for 1.4% of U.S. natural gas production in May 2014 (EIA). Natural gas is a mixture primarily composed of methane (CH₄), with ethane (C₂H₆) typically the second largest component. Emissions of natural gas to the atmosphere may occur during various stages of oil and natural gas extraction and processing. Consequently, the U.S. Environmental Protection Agency (EPA) greenhouse gas (GHG) inventory estimates that CH₄ emissions to the atmosphere from the oil and gas industry account for 29% of 2013 U.S. anthropogenic CH₄ emissions (EPA 430-R-15-004, <http://www3.epa.gov/climatechange/Downloads/ghgemissions/US-GHG-Inventory-2015-Main-Text.pdf>). Two recent studies have shown measurement-based estimates of CH₄ emissions from the oil and gas industry are in agreement with the EPA GHG inventory for large gas-producing regions [Peischl et al., 2015; Karion et al., 2015]. However, other studies have shown that GHG inventories underestimate CH₄ emissions from other oil and natural gas-producing regions [Pétron et al., 2012; Peischl et al., 2013; Karion et al., 2013; Pétron et al., 2014; Brandt et al., 2014]. Therefore, studies that estimate CH₄ emissions from the oil and gas industry are necessary to constrain regional and national GHG emissions inventories and ultimately inform decisions based on the climate impacts of U.S. fuel choices.

©2016. The Authors.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.




Figure 1. Monthly average daily oil and natural gas production from the Bakken shale play through May 2014. Source: U.S. Energy Information Administration (downloaded October 2015).

Schneising et al. [2014] reported an increase in CH_4 emissions from the Bakken region of northwestern North Dakota and southeastern Saskatchewan of $0.99 \pm 0.65 \text{ Tg CH}_4/\text{yr}$ between the years 2006–2008 and 2009–2011 based on measurements from the Scanning Imaging Absorption spectrometer for Atmospheric Chartography (SCIAMACHY) instrument on board the Envisat satellite. *Schneising et al.* [2014] further estimated an “energy content leak rate” for the Bakken, i.e., the energy content of the natural gas emitted to the atmosphere

divided by the sum of the energy content of the oil and natural gas extracted from the ground, of $10.1\% \pm 7.3\%$ by comparing the increase in CH_4 emissions with the increase in oil and natural gas production from this region over the same time period. We interpret this as a loss of $44 \pm 32\%$ of the natural gas withdrawn, as discussed in section 5.5. Several studies using in situ airborne measurements and the mass balance method have recently quantified CH_4 emissions to the atmosphere from oil and natural gas-producing regions such as the Uinta Basin of Utah [Karion *et al.*, 2013], the Denver-Julesburg Basin of northeastern Colorado [Pétron *et al.*, 2014], the Haynesville region of northwestern Louisiana and eastern Texas, the Arkoma Basin of Arkansas, the Marcellus region of northeastern Pennsylvania [Peischl *et al.*, 2015], and the Barnett region of central Texas [Karion *et al.*, 2015]. These aircraft studies report a variety of atmospheric CH_4 emission rates, ranging from $(14\text{--}80) \times 10^3 \text{ kg CH}_4/\text{h}$, with rates of natural gas losses to the atmosphere ranging from 0.18% to 11.7% of the natural gas extracted from the ground from these regions. The variability in emissions and loss rates from these regions likely stems from a variety of factors, including whether oil is extracted in addition to natural gas, the amount of aboveground fossil fuel processing that occurs, the well count, the fossil fuel production volume, the age of the infrastructure, and the type of activity occurring on days when the aircraft studies occur. For comparison, the average hourly emission increase found by *Schneising et al.* [2014] was $(110 \pm 70) \times 10^3 \text{ kg CH}_4/\text{h}$ over the years 2006–2008 to 2009–2011. This CH_4 emission rate is the largest emission reported from all oil and gas-producing regions studied to date, which warrants further examination. Here we derive atmospheric CH_4 emission rates from the Bakken region of North Dakota using in situ airborne data and the aircraft mass balance technique and compare to the estimate by *Schneising et al.* [2014].

We estimate the total CH_4 emitted to the atmosphere from the Bakken region of North Dakota using measurements taken aboard a chemically instrumented National Oceanic and Atmospheric Administration (NOAA) DHC-6 Twin Otter aircraft in May 2014 during the Twin Otter Projects Defining Oil/gas Well emissionNs (TOPDOWN) field project, based out of Minot, ND (Figure 2). The NOAA Twin Otter flew on 12 days in the Bakken region, with transects performed upwind, over, and downwind of this region. The Twin Otter typically flew at $\sim 65 \text{ m/s}$, and full field transects normally took 45 min. A flight was typically 3 to 3.5 h in duration. On six of the flight days, the Twin Otter refueled and took off again for a second flight. Figure 2 shows all the flight tracks of the NOAA Twin Otter around the Bakken region in May 2014. Using the mass balance technique, we determine the CH_4 emissions from this region on 3 days with steady winds. We then attribute the CH_4 emissions in the region to oil and natural gas extraction operations based on the similarity of atmospheric C_2H_6 to CH_4 enhancement ratios during the mass balance transects with the abundances of C_2H_6 and CH_4 in natural gas extracted from the Bakken region. Finally, we estimate the natural gas loss rate to the atmosphere from oil and gas operations in the Bakken region by converting the CH_4 emission to a natural gas emission and dividing by the amount of natural gas extracted in the region in May 2014.

2. Instrumentation

CH_4 , carbon dioxide (CO_2), carbon monoxide (CO), and water vapor (H_2O) were measured by wavelength-scanned cavity ringdown spectrometry using a Picarro G2401-m [Karion *et al.*, 2015]. CH_4 was measured once every 2 s




Figure 2. Map of the Bakken region. The NOAA Twin Otter flight tracks are shown with thin black lines. Active oil and gas wells in North Dakota (ND), Montana (MT), Saskatchewan (SK), and Manitoba (MB) are shown as light blue circles. Natural gas was sampled from 710 wells, colored mustard yellow, and analyzed for CH₄, C₂H₆, and other trace gas abundance [Brandt *et al.*, 2015]. Point sources from the 2013 EPA GHG Reporting Program inventory are shown as open dark blue circles sized by inventory CH₄ emissions. Urban area outlines, including Minot and Williston, ND, are colored orange. The approximate extent of the Williston Basin is shaded gray in the overview map.

with an estimated accuracy of ± 1.4 ppb (all uncertainties herein are 1σ); the precision of this measurement was ± 0.2 ppb. All CH₄ measurements herein are reported as dry air mole fractions expressed as nanomole/mole or parts per billion. C₂H₆ was measured by tunable infrared laser direct absorption spectroscopy using an Aerodyne mini ethane analyzer [Smith *et al.*, 2015]. C₂H₆ was measured every second with an estimated accuracy of $\pm 7\%$; 1 s precision was $\pm (0.02 \text{ to } 0.21)$ ppb, depending on flight conditions. Additionally, black carbon was measured using a single particle soot photometer [Schwarz *et al.*, 2015], ozone was measured using a 2B Technologies analyzer, and flasks of air samples were later analyzed for hydrocarbons and other trace gases. Two differential GPS antennae were mounted above the fuselage of the Twin Otter and provided aircraft heading, altitude, latitude, longitude, ground speed, and course over ground. Ambient temperature was measured with a Rosemount deiced total temperature sensor, model number 102CP2AF, calibrated before and after the field project, with a precision of $\pm 0.2^\circ\text{C}$ and an estimated accuracy of $\pm 1.0^\circ\text{C}$. Meteorological and navigational measurements were made once per second by various sensors aboard the NOAA Twin Otter. Potential temperature, θ , is calculated using the following equation:

$$\theta = T \times \left(\frac{1000}{P} \right)^{0.286} \quad (1)$$

where T is ambient temperature in units of Kelvin and P is ambient pressure in units of millibar. The ambient wind speed and wind direction were derived from the true air speed, heading, speed over ground, and course over ground measurements aboard the Twin Otter using the technique of Conley *et al.* [2014]. We estimate uncertainties in the wind speed of ± 1 m/s and wind direction of $\pm 6^\circ$ during typical level flight conditions, based on an analysis of the wind calibration flight pattern data [Conley *et al.*, 2014].

3. Other Data

The following inventories are used in the current analysis: the EPA 2013 GHG Inventory (publication EPA 430-R-15-004, downloaded September 2015, <http://www3.epa.gov/climatechange/Downloads/ghgemissions/US-GHG-Inventory-2015-Main-Text.pdf>), and the EPA GHG Reporting Program (GHGRP) inventory

(downloaded November 2015, <http://www2.epa.gov/ghgreporting/subpart-w-reported-data>). The GHGRP data include the latitude, longitude, and estimated CH₄ emission of known point sources (Figure 2). Some latitude-longitude coordinates in the 2013 EPA GHGRP inventory data set were modified to match the actual locations determined using satellite imagery.

Natural gas sample analysis data are used from the work of *Brandt et al.* [2015] by way of the North Dakota Industrial Commission, Department of Mineral Resources, Oil and Gas Division (<https://www.dmr.nd.gov/oilgas/>). Samples were taken from 710 wells across the Bakken region (Figure 2) and analyzed for the molar composition of C₁-C₆ alkanes, CO₂, and other trace gases. North Dakota well locations were downloaded from the same website (accessed September 2014). North Dakota well production data are from the North Dakota State Industrial Commission May 2014 Oil and Gas Production Report (<https://www.dmr.nd.gov/oilgas/mprindex.asp>). Bakken production data were obtained from the EIA Web site (<http://www.eia.gov/petroleum/drilling/>, downloaded October 2015).

Terrain height below the Twin Otter aircraft is estimated using a digital elevation map (DEM) from the U.S. Geological Survey (GTOPO30, 1996, <https://lta.cr.usgs.gov/GTOPO30>). The DEM has a resolution of 30 arc sec (~900 m) and is based on World Geodetic System 1984 data. Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model back trajectories [*Stein et al.*, 2015; <https://ready.arl.noaa.gov/HYSPLIT.php>] are used to evaluate the evolution of wind fields over time and are run with the following settings: Meteorology: North American Mesoscale Forecast System, 12 km; Vertical Motion: Model vertical velocity (which uses the vertical velocity field from the meteorological data). County-level livestock populations were downloaded from the U.S. Department of Agriculture (USDA) National Agriculture Statistics Service (NASS) (<http://www.nass.usda.gov>). We assume an average emission of 67.9 kg CH₄/yr per head from enteric fermentation (EPA GHG inventory) and no emission of C₂H₆. Bison populations are estimated from a North Dakota Department of Agriculture website (<http://www.nd.gov/aitc/agfacts/Livestock/>). We assume that bison populations scale with cattle populations and assume an average emission of 72 kg CH₄/yr per head [*Kelliher and Clark*, 2010]. County-level manure emissions estimates were downloaded from the National Renewable Energy Laboratory (<http://maps.nrel.gov/biomass>) and were based on 2002 USDA data. In cases where the NOAA Twin Otter flight track transects a county, an apportionment based on geographical area is used to determine livestock and manure emissions within the study region.

4. Mass Balance Technique

The technique used to estimate CH₄ fluxes is the mass balance approach [*White et al.*, 1976]. This technique has been used extensively to estimate CH₄ emissions from oil and gas-producing regions [*Karion et al.*, 2013; *Pétron et al.*, 2014; *Peischl et al.*, 2015; *Karion et al.*, 2015]. Further, the mass balance technique was used to estimate C₂H₆ emissions from an oil and gas-producing region [*Smith et al.*, 2015] and black carbon and C₂H₆ emissions from the Bakken region during the 2014 TOPDOWN project [*Schwarz et al.*, 2015; *Kort et al.*, 2016].

The mass balance technique is used as follows. We calculate the horizontal flux of a species *X* through a vertical plane defined by an aircraft transect:

$$\text{flux} = v \cos(\alpha) \int_{z_0}^{z_1} \int_{-y}^y (X - X_{bg}) dy dz \quad (2)$$

where $v \cos(\alpha)$ is the magnitude of the component of the wind velocity normal to the flight track averaged over the course of a transect, z_0 is the terrain height in meters above sea level (masl), z_1 is the adjusted mixing height in meters above sea level, $(X - X_{bg})$ is the CH₄ enhancement above the background, and y is the crosswind distance [*White et al.*, 1976]. We define the background CH₄ mixing ratios as those found immediately outside the plume downwind of the Bakken region. This is discussed in more detail, below. We define the adjusted mixing height in meters above ground level (magl) as described by *Peischl et al.* [2015]: $z_1 = (3z_{PBL} + z_e)/4$, where z_{PBL} is the well-mixed planetary boundary layer (PBL) depth, and z_e is the entrainment height or the altitude at which mixing from the PBL has reached estimated free tropospheric CH₄ mixing ratios. Below, we compare this method of determining the adjusted mixing height with that of *Schwarz et al.* [2015]. All other calculations are similar to those reported by *Peischl et al.* [2015].




Figure 3. Map showing the five transects downwind of the central Bakken shale formation from which CH_4 fluxes were calculated. North Dakota natural gas production data are summed into a $\sim 5.6 \times 5.6$ km grid. Flight tracks are colored and sized by CH_4 mixing ratio for the transects shown in Figure 4. The arrow indicates the average wind speed and direction during the transects, where the length of the arrow is proportional to the wind speed. (a) The 13 May transect at approximately 350 m agl. (b) The 14 May northeast-to-southwest transect at approximately 350 m agl and southwest-to-northeast transect at approximately 650 m agl. (c) The 22 May east-to-west transect at approximately 500 m agl. (d) The 22 May west-to-east transect at approximately 1000 m agl.

5. Results and Discussion

5.1. CH_4 Emissions to the Atmosphere From the Bakken Study Region

The NOAA Twin Otter aircraft flew three flights dedicated to determining the CH_4 mass balance from the Bakken region on 13, 14, and 22 May (Figure 3). Eight additional flight days were dedicated to characterizing emissions from CH_4 point sources, flaring, and general field emissions. One flight day, 21 May, was dedicated to mass balance, but because the wind speeds were light (~ 2 – 3 m/s), we cannot verify that they were steady over a full traverse of the Bakken region, and we therefore discard this day from our analysis. For the three mass balance flights used here, the winds measured aboard the Twin Otter were steady throughout the Bakken region, in agreement with HYSPLIT back trajectories.

The 13, 14, and 22 May flight tracks sampled downwind of the majority of the highest-producing wells in the Bakken region (Figure 3). The flight tracks in Figure 3 are colored by the CH_4 mixing ratio measured within the PBL along the transects used in the mass balance analysis.

Figure 4 shows the time series of CH_4 and aircraft altitude for the downwind transects highlighted in Figure 3. Also plotted in Figure 4 is the background CH_4 mixing ratio used in equation (2). We use the C_2H_6 measurement,




Figure 4. Time series of CH_4 measurements (black) from the five transects downwind of the Bakken shale region shown in Figure 3. The blue trace shows the Twin Otter altitude above ground level. The red trace represents the background CH_4 used for the flux calculations; the dashed red traces represent the estimated $\pm 1\sigma$ uncertainty of the background. The shaded areas show the CH_4 enhancement above the background along the transects.

which typically has a greater plume signal to background variability ratio than the CH_4 measurement, to help define the plume background, especially on days such as 22 May, when the CH_4 plume enhancements were only a factor of 2–4 greater than the CH_4 background variability. The CH_4 background is further selected to be consistent with the C_2H_6 data, such that the ratio of their derived fluxes is equal to the molar enhancement ratio of C_2H_6 to CH_4 observed on the downwind transects.

Examples of the mixing height determination are shown with three vertical profiles in Figure 5. Figure 5 (left) shows the vertical profile immediately before the transect shown in Figure 4a; Figure 5 (middle) shows the vertical profile immediately before the second transect shown in Figure 4b; Figure 5 (right) shows the vertical profile immediately after the first transect shown in Figure 4c. We calculate the adjusted mixing height, z_1 , following the method of Peischl *et al.* [2015] to account for any transport of CH_4 through the top of the PBL. We increase the effective PBL depth by one quarter the distance between the well-mixed PBL, defined by the dashed lines in Figure 5, and entrainment height, z_e , defined by the dotted lines in Figure 5, and




Figure 5. One vertical profile is plotted from each of the three flight days used for the mass balance. Dashed horizontal black lines indicate the well-mixed PBL depth. Dotted horizontal black lines indicate the entrainment height. The yellow circles indicate the average adjusted mixing heights used for the mass balance calculations. The orange diamonds indicate the average adjusted mixing heights ($h \times \beta$) used by Schwarz *et al.* [2015]. Solid vertical gray lines indicate the CH_4 background; dashed vertical gray lines represent the $\pm 1\sigma$ uncertainty of the background.

include an uncertainty equal to this same distance. This adjusted mixing height and uncertainty encompasses a range of possible sources for the CH_4 enhancement above the free tropospheric levels immediately above the PBL: z_{PBL} if the CH_4 enhancements above the PBL are solely due to preexisting layers aloft, and halfway between z_{PBL} and z_e if all CH_4 enhancements above the PBL are due to detrainment from the PBL. The resulting adjusted mixing heights and their uncertainties from each vertical profile during a flight are fit with a line. We use the value of this fit at the time of the transect, along with the corresponding 1σ confidence interval of the fit, to determine z_1 for equation (2), and plot the resulting adjusted mixing heights as filled yellow circles in Figure 5. For comparison, the mixed height determined by Schwarz *et al.* [2015] using a different method are shown with orange diamonds. For each mass balance transect in this analysis, the mixed height determinations are well within the uncertainties of the two different estimation techniques.

Table 1 lists the parameters and their uncertainties used in equation (2) to calculate the CH_4 flux for each transect. The largest uncertainties for the 13 May flight were from the uncertainty of the background CH_4 value ($\pm 25\%$), the variability of the wind speed ($\pm 20\%$), and the uncertainty of z_1 ($\pm 18\%$). The largest uncertainties for the 14 May flight were from the variability of the wind speed ($\pm 34\%$), the uncertainty of the background CH_4 value ($\pm 30\%$), and the uncertainty of z_1 ($\pm 21\%$). Finally, the largest uncertainties for the 22 May flight were from the uncertainty of the background CH_4 value ($\pm 27\%$), the variability of the wind speed ($\pm 24\%$), and the uncertainty of z_1 ($\pm 21\%$). The CH_4 fluxes calculated from the five downwind transects are (36 ± 13) , (27 ± 13) , (27 ± 12) , (27 ± 12) , and $(25 \pm 10) \times 10^3$ kg/h. The two flux determinations from the 14 and 22 May flights did not depend on transect altitude, indicating the emissions were mixed vertically within the PBL by the time they were sampled by the Twin Otter. The $1/\sigma^2$ -weighted average of the five CH_4

Table 1. Data Used in Equation (2) From Five Transects on the 13, 14, and 22 May 2014 Flights and the Resulting CH₄ Fluxes

Transect	Terrain Height (masl)	z ₁ , (masl)	Wind Direction (°)	Wind Speed (m/s)	CH ₄ Flux (10 ²⁶ molecule/s)	CH ₄ Flux (10 ³ kg/h)
13 May	680 ± 60	2810 ± 265	312 ± 9	13.3 ± 2.7	3.8 ± 1.4	36 ± 13
14 May, #1	695 ± 85	2545 ± 395	339 ± 14	7.7 ± 2.6	2.8 ± 1.4	27 ± 13
14 May, #2	705 ± 80	2655 ± 245	339 ± 8	8.0 ± 2.3	2.8 ± 1.3	27 ± 12
22 May, #1	655 ± 45	2340 ± 355	177 ± 10	7.5 ± 1.8	2.8 ± 1.2	27 ± 12
22 May, #2	660 ± 45	2320 ± 300	180 ± 9	7.4 ± 1.7	2.7 ± 1.1	25 ± 10

emissions estimates for the Bakken region is $28 (\pm 5) \times 10^3$ kg CH₄/h (Table 2), suggesting relatively small day-to-day variability in the total emissions rate for this short study. We therefore assume that this emission rate represents the average May 2014 CH₄ emission from the Bakken region.

5.2. Oil and Natural Gas Production in the Bakken Study Region

Monthly oil and natural gas production from the Bakken region is estimated for May 2014 using both EIA and North Dakota state data. The EIA reports gross gas withdrawal volumes in thousands of cubic feet (MCF) at 60° F and 6.681528 kg per square inch. We therefore adjust the volumes reported by the EIA by factor of 0.9483 to convert to standard temperature and pressure of 0°C and 1 atm then convert cubic feet to cubic meters. The Bakken natural gas production for May 2014 was 10.2×10^8 m³ according to the EIA when converted to standard cubic meters. Similarly, the sum of individual well production data from the North Dakota Oil and Gas Division is 9.7×10^8 m³, within 5% of the EIA estimate. Each airborne transect analyzed here was downwind of wells that accounted for greater than 97% of total production. We therefore estimate natural gas production upwind of any given transect in the region to be $(9.9 \pm 0.3) \times 10^8$ m³ for the month of May 2014 (Table 2).

5.3. Other Sources of CH₄ in the Bakken Study Region

The Bakken region encompasses landfills and livestock; these sources emit CH₄ but are not known to emit C₂H₆. The total CH₄ emission from point sources in the 2013 EPA GHGRP inventory not related to oil and natural gas processing in the Bakken region, two natural gas-fired electrical generating stations, is 0.23×10^3 kg/yr, which averages to 0.03 kg CH₄/h, which we treat as negligible. Here we use bottom-up methods to estimate CH₄ emissions from livestock in the Bakken region and then compare these emissions to the total CH₄ emission from the region to confirm that livestock emissions are also not a major source of CH₄ to the atmosphere in the Bakken region. We estimate that the Twin Otter flight track encompassed 9% of the cattle and calves in North Dakota, based on a geographical apportionment of USDA NASS county data. This results in an estimated enteric fermentation emission from 161,000 cattle and calves of 1.3×10^3 kg CH₄/h. CH₄ emissions from manure are estimated at 0.003×10^3 kg CH₄/h. We estimate approximately 32,000 bison emit 0.3×10^3 kg CH₄/h, based on statewide bison populations scaled by estimated cattle and calf populations in the Bakken region. The combined point source and livestock emission, 1.5×10^3 kg CH₄/h (Table 2), accounts for approximately 5% of the 28×10^3 kg CH₄/h emission calculated for May.

Coal mines, another possible source of CH₄, are located southeast of the Bakken region. The state of North Dakota has four active coal mines (EIA, <http://www.eia.gov/coal/annual/pdf/table2.pdf>), all of which are surface mines. These mines are represented in the 2013 EPA GHGRP inventory and are located between 47.0° and 47.5°N latitude and 101° and 102°W longitude (Figure 2). CH₄ emissions from these mines were further downwind of the 13 and 14 May transects, and therefore do not affect the mass balance calculations

Table 2. Summary of CH₄ Emissions From the Bakken Study Region

Quantity Evaluated	Amount
CH ₄ flux	$(28 \pm 5) \times 10^3$ kg/h
CH ₄ estimated from nonoil and gas point sources and livestock	$\sim 1.5 \times 10^3$ kg/h
CH ₄ emissions attributed to the oil and gas industry	$(20\text{--}33) \times 10^3$ kg/h
Natural gas production in May 2014	$(9.9 \pm 0.3) \times 10^8$ m ³
CH ₄ abundance in natural gas ^a	47 ± 13%
CH ₄ extraction rate in May 2014	$(420 \pm 120) \times 10^3$ kg/h
Natural gas loss rate to the atmosphere	6.3 ± 2.1%

^aBrandt et al. [2015].




Figure 6. C_2H_6 and CH_4 are plotted for each of the five transects used to calculate a CH_4 flux downwind of the Bakken region. Also, plotted are lines with a slope equal to the C_2H_6 to CH_4 molar composition ratio (C_2/C_1) from 710 samples of natural gas from the Bakken region [Brandt et al., 2015], assuming a background of 1.5 ppb C_2H_6 and 1900.5 ppb CH_4 . The slope of the yellow line represents the geometric mean C_2H_6 to CH_4 composition ratio of the samples.

on these days, but may have contributed to the CH_4 enhancements measured on the 22 May flight. This is discussed in more detail in the next section.

The small contribution of livestock to the total CH_4 emission in the Bakken region suggested by the USDA inventory is confirmed using airborne data. Enhancements of C_2H_6 in the boundary layer along the mass balance transects correlate with enhancements of CH_4 similar to the composition ratios of C_2H_6 to CH_4 in natural gas extracted from the region (Figure 6). The lines representing the composition ratios in Figure 6 are offset to 1900.5 ppb CH_4 and 1.5 ppb C_2H_6 , which represents the approximate background mixing ratios for the mass balance transects on the 13 and 14 May flights. For the 22 May flight, we plot the CH_4 enhancement above the sloping background in Figure 4 then offset the difference to the background values as for the composition ratios. The C_2H_6 to CH_4 enhancement ratio was on the order of 0.4–0.5, an atypically high ratio compared to previously studied oil and gas-producing regions. As a result, this ratio provides a unique source fingerprint and distinguishes Bakken region oil and gas emissions from natural gas in the supply chain or coal mine emissions, which typically have C_2H_6 to CH_4 ratios of less than 0.05. Finally, an approximately 10 km wide CH_4 plume on 22 May sampled shortly before 17:00 CST not highly correlated with C_2H_6 represents 6% of the CH_4 flux calculation for that day. These data were sampled along the eastern edge of the 22 May transects, where emissions from coal mines to the south-southeast may have influenced the CH_4 measurements.

We attribute CH_4 emissions to the oil and gas industry in the Bakken region as follows. We subtract the estimated livestock emission and the plume not correlated with C_2H_6 on the 22 May flight from the total CH_4 emission estimated for the region. Since we cannot confirm quantitatively with our measurements the impact that livestock and coal mine emissions may have had on the CH_4 emissions, we assign a 100% uncertainty to these subtractions. Consequently, CH_4 emissions attributed to the oil and gas industry have larger error bars than the estimate for the total CH_4 emissions from the region, but both emissions estimates share the same upper limit (Table 2).

Our estimate for oil and gas emissions of $(20\text{--}33) \times 10^3$ kg CH_4 /h is a factor of 5–8 greater than the 2013 EPA GHGRP estimate for the entire Williston Basin of 0.0047 Tg CH_4 /yr, which averages to 4.1×10^3 kg CH_4 /h. However, under this program, emissions from many small facilities may not meet the threshold required to report (<http://www2.epa.gov/sites/production/files/2015-07/documents/subpartwinformation-sheet.pdf>), so we would expect the GHGRP value to be a lower limit to the total emissions. However, the EPA also estimates U.S. CH_4 emissions from petroleum systems of 1.009 Tg CH_4 /yr (2013 EPA GHG inventory, Tables 3–37). Assuming inventory CH_4 emissions scale with oil production, the Bakken region would account for 12.5% of the total U.S. emission, which averages to 14.4×10^3 kg CH_4 /h. Therefore, our estimate of CH_4 emissions from the Bakken region oil and gas industry is a factor of 1.4–2.3 greater than the 2013 EPA GHG inventory's emissions from petroleum systems scaled by oil production.

5.5. Loss Rates to the Atmosphere From Oil and Natural Gas Production in the Bakken Study Region

CH_4 may be emitted, or lost, to the atmosphere from oil and natural gas operations through a variety of ways: intentional venting, partially combusted flaring, fugitive losses, etc. An atmospheric loss rate from oil and

natural gas operations in the Bakken region for May 2014 is estimated by taking the CH₄ mass emission rate attributed to the oil and gas industry and dividing by the average CH₄ mass extraction rate, or production rate, of natural gas for the month. We use 22.414 L/mol (the EIA standard 23.636 L/mol times 0.9483 to convert to standard L) of natural gas in the conversion from volume to mass. The mean CH₄ abundance in the 710 Bakken natural gas samples is $47 \pm 13\%$ [Brandt *et al.*, 2015] (Table 2). The resulting estimated natural gas loss rate from the Bakken region is 4.2%–8.4% based on this analysis of airborne in situ data. This loss rate is not significantly different than that calculated for another oil-producing region, the Denver-Julesburg Basin in northeastern Colorado, where Pétron *et al.* [2014] found loss rates of 2.6%–5.6%.

We convert this loss rate to an “energy loss rate” in order to compare to that reported by Schneising *et al.* [2014]. We estimate an energy loss rate by converting the natural gas and oil extracted from the Bakken formation to energy extracted using the following conversions: 5.8×10^7 J/m³ for natural gas and 3.464×10^{10} J/m³ for oil [Brandt *et al.*, 2015]. We assume the change in density of oil into standard units is <1% and therefore negligible. In May 2014, natural gas accounted for 25.4% of the energy extracted in the Bakken region (EIA), which results in an energy loss rate of $1.6 \pm 0.5\%$, a factor of approximately 6 lower than that reported by Schneising *et al.* [2014] for 2006–2008 to 2009–2011. Alternatively, we convert the energy loss rate found by Schneising *et al.* [2014] to a natural gas loss rate. Natural gas in the North Dakota portion of the Bakken accounted for 23% of the energy extracted from 2009 to 2011. Assuming that this energy extraction ratio holds true for production in the Saskatchewan portion of the Bakken, the $10.1 \pm 7.3\%$ energy loss rate reported by Schneising *et al.* [2014] would imply an emission to the atmosphere of approximately $44 \pm 32\%$ of the natural gas extracted from the Bakken formation. Finally, where we derive a total CH₄ emission of $(28 \pm 5) \times 10^3$ kg/h from the Bakken region, Schneising *et al.* [2014] found an increase in emissions of $(110 \pm 70) \times 10^3$ kg/h from western North Dakota and southeastern Saskatchewan. We conclude that the exceptionally high atmospheric loss rate of CH₄ reported by Schneising *et al.* [2014] for 2009–2011 is inconsistent with our airborne data from May 2014.

6. Conclusions

We report the first in situ measurements of CH₄ to estimate atmospheric CH₄ emissions from the Bakken oil and natural gas production region of northwestern North Dakota. We calculate CH₄ fluxes from the Bakken region for May 2014 from three flights and five transects downwind of the region and find CH₄ emissions to the atmosphere of $(28 \pm 5) \times 10^3$ kg CH₄/h. Both aircraft measurements of C₂H₆ and a bottom-up assessment derived here indicate that livestock are not a dominant source of CH₄ to the atmosphere in this region, nor are landfills and other CH₄ sources not related to the oil and gas industry. Instead, atmospheric enhancement ratios of C₂H₆ to CH₄ are consistent with the composition ratio of natural gas extracted in the Bakken region. We therefore attribute CH₄ emissions of $(20\text{--}33) \times 10^3$ kg CH₄/h to the oil and gas production and distribution activities in this region. This is approximately a factor of 1.4–2.3 greater than the 2013 EPA GHG inventory emission rate from petroleum systems when scaled by oil production. Stated in terms of the natural gas extracted from the ground, we estimate atmospheric loss rates from natural gas production of 4.2–8.4%, similar to the rate found for the Denver-Julesburg oil-producing region [Pétron *et al.*, 2014].

Note Added in Proof

The EPA GHG inventory report for 2014 was released in April 2016 (EPA 430-R-16-002, <https://www3.epa.gov/climatechange/ghgemissions/usinventoryreport.html>), while this paper was under final review; inventoried emissions from petroleum systems increased by a factor of 2.5 compared to the GHG inventory for 2013 that was used for comparison above. The CH₄ emissions attributed to oil and gas operations in this paper are now a factor of 0.51–0.85 times the 2014 EPA GHG inventory for petroleum systems scaled by oil production in the Bakken.

References

- Brandt, A. R., *et al.* (2014), Methane leaks from North American natural gas systems, *Science*, 343, 733–735, doi:10.1126/science.1247045.
- Brandt, A. R., T. Yeskoo, S. McNally, K. Vafi, H. Cai, and M. Q. Wang (2015), Energy intensity and greenhouse gas emissions from crude oil production in the Bakken Formation: Input data and analysis methods, Department of Energy, Argonne National Laboratory. [Available at <https://greet.es.anl.gov/publication-bakken-oil>.]

Acknowledgments

J. Peischl, K. Aikin, M. Trainer, and T. Ryerson were supported in part by the NOAA Climate Program Office. A. Karion, C. Sweeney, S. Wolter, E. Kort, M. Smith, and A. Gvakharia were supported in part by the NOAA Atmospheric Chemistry, Carbon Cycle, and Climate Program under grant NA14OAR0110139. Data are publicly available at <http://esrl.noaa.gov/csd/groups/csd7/measurements/2014topdown/>.

- Conley, S. A., I. C. Faloona, D. H. Lenschow, A. Karion, and C. Sweeney (2014), A low-cost system for measuring horizontal winds from single-engine aircraft, *J. Atmos. Oceanic Technol.*, *31*, 1312–1320, doi:10.1175/JTECH-D-13-00143.1.
- Karion, A., et al. (2013), Methane emissions estimate from airborne measurements over a western United States natural gas field, *Geophys. Res. Lett.*, *40*, 4393–4397, doi:10.1002/grl.50811.
- Karion, A., et al. (2015), Aircraft-based estimate of total methane emissions from the Barnett Shale Region, *Environ. Sci. Technol.*, *49*, 8124–8131, doi:10.1021/acs.est.5b00217.
- Kelliher, F. M., and H. Clark (2010), Methane emissions from bison—An historic herd estimate for the North American Great Plains, *Agric. For. Meteorol.*, *150*(3), 473–477, doi:10.1016/j.agrformet.2009.11.019.
- Kort, E. A., M. L. Smith, L. T. Murray, A. Gvakharia, A. R. Brandt, J. Peischl, T. B. Ryerson, C. Sweeney, and K. Travis (2016), Fugitive emissions from the Bakken shale illustrate role of shale production in global ethane shift, *Geophys. Res. Lett.*, *43*, doi:10.1002/2016GL068703.
- Peischl, J., et al. (2013), Quantifying sources of methane using light alkanes in the Los Angeles basin, California, *J. Geophys. Res. Atmos.*, *118*, 4974–4990, doi:10.1002/jgrd.50413.
- Peischl, J., et al. (2015), Quantifying methane emissions from the Haynesville, Fayetteville, and Marcellus shale play regions, *J. Geophys. Res. Atmos.*, *120*, 2119–2139, doi:10.1002/2014JD022697.
- Pétron, G., et al. (2012), Hydrocarbon emissions characterization in the Colorado Front Range: A pilot study, *J. Geophys. Res.*, *117*, D04304, doi:10.1029/2011JD016360.
- Pétron, G., et al. (2014), A new look at methane and nonmethane hydrocarbon emissions from oil and natural gas operations in the Colorado Denver-Julesburg Basin, *J. Geophys. Res. Atmos.*, *119*, 6836–6852, doi:10.1002/2013JD021272.
- Schneising, O., J. P. Burrows, R. R. Dickerson, M. Buchwitz, M. Reuter, and H. Bovensmann (2014), Remote sensing of fugitive methane emissions from oil and gas production in North American tight geologic formations, *Earth's Future*, *2*, doi:10.1002/2014EF000265.
- Schwarz, J. P., J. S. Holloway, J. M. Katich, S. McKeen, E. A. Kort, M. L. Smith, T. B. Ryerson, C. Sweeney, and J. Peischl (2015), Black carbon emissions from the Bakken Oil and Gas Development region, *Environ. Sci. Technol. Lett.*, doi:10.1021/acs.estlett.5b0025.
- Smith, M. L., E. A. Kort, A. Karion, C. Sweeney, S. C. Herndon, and T. I. Yacovitch (2015), Airborne ethane observations in the Barnett Shale: Quantification of ethane flux and attribution of methane emissions, *Environ. Sci. Technol.*, *49*, 8128–8166, doi:10.1021/acs.est.5b00219.
- Stein, A. F., R. R. Draxler, G. D. Rolph, B. J. B. Stunder, M. D. Cohen, and F. Ngan (2015), NOAA's HYSPLIT atmospheric transport and dispersion modeling system, *Bull. Am. Meteorol. Soc.*, *96*, 2059–2077, doi:10.1175/BAMS-D-14-00110.1.
- White, W. H., J. A. Anderson, D. L. Blumenthal, R. B. Husar, N. V. Gillani, J. D. Husar, and W. E. Wilson Jr. (1976), Formation and transport of secondary air pollutants: Ozone and aerosols in the St. Louis urban plume, *Science*, *194*, 187–189, doi:10.1126/science.959846.